

Open Research Online

The Open University's repository of research publications and other research outputs

Volcanic constraints on the unzipping of Africa from South America: Insights from new geochronological controls along the Angola margin

Journal Item

How to cite:

Jerram, D.A.; Sharp, I.R.; Torsvik, T.H.; Poulsen, R.; Watton, T.; Freitag, U.; Halton, A.; Sherlock, S.C.; Malley, J.A.S.; Finley, A.; Roberge, J.; Swart, R.; Fabregas, P.; Ferreira, C.H. and Machado, V. (2019). Volcanic constraints on the unzipping of Africa from South America: Insights from new geochronological controls along the Angola margin. *Tectonophysics*, 760 pp. 252–266.

For guidance on citations see [FAQs](#).

© 2018 Elsevier B.V.

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1016/j.tecto.2018.07.027>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

Accepted Manuscript


Volcanic constraints on the unzipping of Africa from South America: Insights from new geochronological controls along the Angola margin

D.A. Jerram, I.R. Sharp, T.H. Torsvik, R. Poulsen, T. Watton, U. Freitag, A. Halton, S.C. Sherlock, J.A.S. Malley, A. Finley, J. Roberge, R. Swart, P. Fabregas, C.H. Ferreira, V. Machado

PII: S0040-1951(18)30270-1
DOI: doi:[10.1016/j.tecto.2018.07.027](https://doi.org/10.1016/j.tecto.2018.07.027)
Reference: TECTO 127903
To appear in: *Tectonophysics*
Received date: 23 December 2017
Revised date: 27 July 2018
Accepted date: 30 July 2018

Please cite this article as: D.A. Jerram, I.R. Sharp, T.H. Torsvik, R. Poulsen, T. Watton, U. Freitag, A. Halton, S.C. Sherlock, J.A.S. Malley, A. Finley, J. Roberge, R. Swart, P. Fabregas, C.H. Ferreira, V. Machado , Volcanic constraints on the unzipping of Africa from South America: Insights from new geochronological controls along the Angola margin. *Tecto* (2018), doi:[10.1016/j.tecto.2018.07.027](https://doi.org/10.1016/j.tecto.2018.07.027)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Volcanic constraints on the unzipping of Africa from South America: insights from new geochronological controls along the Angola margin.

Jerram, D.A.^{1,2,3}, Sharp, I. R.⁴, Torsvik, T. H.^{1,5,6}, Poulsen, R.⁴, Watton, T.⁴, Freitag, U.⁴, Halton, A.⁷, Sherlock, S. C.⁷, Malley, J. A. S.⁷, Finley, A.⁸, Roberge, J.⁹, Swart, R.¹⁰, Fabregas, P.¹¹, Ferreira, C., H.¹¹, Machado, V.¹¹

¹ Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Norway

² DougalEARTH Ltd.1, Solihull, UK

³ Visiting research fellow, Earth, Environmental and Biological Sciences, Queensland University of Technology, Brisbane, Queensland, Australia

⁴ Statoil ASA, Sandslivegen 90, Bergen, Norway

⁵ Geological Survey of Norway, 7040 Trondheim, Norway.

⁶ School of Geosciences, University of Witwatersrand, Johannesburg 2050, South Africa.

⁷ Department of Earth and Environmental Sciences, The Open University, CEPSAR, Walton Hall, Milton Keynes MK7 6AA, UK

⁸ Origin Analytical Ltd, 1Ravenscroft Court, Buttington Cross Enterprise Park, Welshpool Powys SY21 8SL UK

⁹ ESIA-Ticomán, Instituto Politécnico Nacional (IPN), Av. Ticoman #600, Mexico

¹⁰ BlackGold Geosciences cc, P.O. Box 24287, Windhoek, Namibia

¹¹ Sonangol A.S. Direcção de Exploração (DEX), Rua Rainha Ginga n. 29/31, C.P. 1316, Luanda, Angola

Abstract

The breakup of Africa from South America is associated with the emplacement of the Paraná-Etendeka flood basalt province from around 134 Ma and the Tristan da Cunha plume. Yet many additional volcanic events occur that are younger than the main pulse of the Paraná-Etendeka and straddle the rift to drift phases of the main breakup. This contribution reports on new geochronological constraints from the Angolan part of the African Margin. Three coastal and one inland section have been sampled stretching across some 400 Km, with ³⁹Ar/⁴⁰Ar, U-Pb and Palaeontology used to provide age constraints. Ages from the new data range from ~100 to 81 Ma, with three main events (cr. 100, 91 and 82-81 Ma). Volcanic events are occurring within the Early to Late Cretaceous, along this part of the margin with a general younging towards Namibia. With the constraints of additional age information both onshore and offshore Angola, a clear younging trend at the early stages of rift to drift is recorded in the volcanic events that unzip from North to South. Similar age volcanic events are reported from the Brazilian side of the conjugate margin, and highlight the need to fully incorporate these relatively low volume volcanic pulses into the plate tectonic breakup models of the South Atlantic Margin.

Introduction

The break-up between Africa and South America marks one of the classic pictures of continental separation. Due to the good geometrical fit between these two land masses, already recognized by Wegener (1915), much work has been undertaken to corroborate the pre-break up configuration, intraplate geometries, the early rifting phases, and the final separation of the continents (e.g., Untemehr et al. 1988; Mohriak *et al.* 1995; Jackson et al. 2000; Torsvik et al., 2009; Moulin et al., 2010; Heine et al. 2013; Gaina et al. 2013; Pérez-Díaz and Eagles, 2014; Kukla et al., 2018).

Magmatism associated with the break-up is also important as it relates to the emplacement of the Paraná-Etendeka flood basalts and the volcanism associated with the volcanic rifted margin (e.g. Jerram and Widdowson 2005), yet the exact duration of magmatic events seem somewhat more complicated than a single pulse of activity associated with the Tristan da Cunha plume, and maybe more pulsed like that seen at other rifted margins such as Ethiopia-Yemen (e.g. Rooney 2017). On a general scale this should be a relatively simple break up story, but much debate still continues about the exact nature and timing of the rifting and separation of the land masses leading to the eventual onset of sea floor spreading. This is somewhat complicated due to the lack of clear reflectors in the Early stages of rifting (Cande & Kent 1995). Work aimed at reconstructing the conjugate salt basins of the margin has helped to constrain some of the major factors controlling break-up basin configuration (e.g. Torsvik et al., 2009; Strozyk et al., 2017; Kukla et al., 2018), though geochronological constraints in the later stages are sparse. A factor that further fuels the need to understand the very specifics of the separation, is the discovery of, and exploration for, major hydrocarbon reserves that are found within the sedimentary basins formed during the rift to drift separation of the continents (e.g. Szatmari and Milani, 2016; Masse and Laurent, 2016).

Concomitant with the onset of rifting of Africa from South America is the occurrence of the Paraná-Etendeka large igneous province (Jerram et al., 1999; Marsh et al 2001; Jerram & Widdowson 2005). This major volcanic event occurring at around 134 Ma (e.g. Dodd et al.2015; Svensen et al., 2017; Marsh and Swart, 2018), is associated with a large mantle anomaly, the trail of which can be seen today running along the Walvis ridge towards Tristan de Cuna. Although the main pulse of magmatism associated with this event is relatively well constrained to a 3-4 Ma period during the early phases of rifting, later stage magmatic activity related to the waning phase of flood volcanism (e.g. Jerram & Widdowson 2005), rift climax, break-up and sag phases, or indeed any volcanism unrelated directly with the rift to drift process, are poorly constrained on both margins. This is partly due to limited access to key exposures (buried in the offshore), and the nature of the magmatic events being of a relatively low volume in the proximal onshore sections, which leads to restricted availability of good geochronological constraints. Igneous events younger than the Paraná-Etendeka LIP have long been known to be present on the Brazilian margin (e.g. Nascimento et al., 2003; Mizusaki et al., 2002; Enrich et al., 2009; Guedes et al., 2005). Additional age constraints based on commercial well and seismic data in offshore Angola and Brazil can also be found (e.g. Mohriak and Leroy, 2012), and appear to be volumetrically significant in the latter phases of rifting (e.g. Higgins et al Submitted), but at present publications that address these younger units, particularly on the Angolan side, are limited.

The Angolan margin, north of Namibia, provides a key area where, until recently, access to important outcrops had been limited. Renewed exploration efforts along the margin, as well as

improved access, particularly along coastal stretches, has allowed significant improvements to be made to our understanding of the preserved geology and tectono-stratigraphic timing relationships in Angola. This in turn has implications for understanding the rift to drift transition in Africa. The extensive salt deposits on the Angolan side as well as those found along the conjugate margin in Brazil, provide valuable information to determine the basin settings prior to and during the break up (Torsvik et al. 2009).

This short contribution reports on new geochronological constraints for Early to Late Cretaceous magmatism found along the Angolan margin as part of an extensive onshore basin analysis and stratigraphic study undertaken jointly by Statoil and Sonangol. Three main locations of magmatism have been identified and studied along ~400 km of the SW Angolan margin (Figure 1). The main goal of this study is to provide new age and geochemical data that can be utilized to constrain the rift to drift evolution of Africa from South America. The locations, stratigraphic context and general rock types of the three main study localities will first be introduced, as well as the methods used to constrain their ages. The results and a summary of the age significance of these magmatic events are then presented as well as palaeo-reconstructions of the rifting in order to place their spatial and temporal evolution in the context of continental break-up.

[Figure 1] a) Location maps of SW Angola showing the 3 main study locations. b) Casa Branca beach section, Sumbe, Kwanza Basin. c) Baía dos Elefantes and Serra de Neve, Benguela Basin. d) Bentiaba, Ponta Negra, Canico, Piambo and Uah sections, Namibe Basin. Additional features offshore indicated.

Geological constraints and sampling

Mesozoic volcanic rocks found within the onshore Angolan section of the West African margin are found almost exclusively within Cretaceous aged units. These have been identified as either pertaining to the main Paraná-Etendeka flood volcanics (e.g. Marzoli et al., 1999; Sharp et al., 2012; Gindre-Chanu et al., 2014; Marsh and Swart 2018), or as younger lower volume events with an alkali affinity (Marzoli et al., 1999). Eroded remnants of the Paraná-Etendeka Igneous Province occur as fault bounded blocks or isolated sill and dyke systems. Sections are exposed in the onshore Kwanza, Benguela and Namibe basins, and these sections are unconformably overlain by Aptian aged mixed carbonate-clastic evaporitic sections of the latest "Pre-Salt" (Cuvo Fm) and "Salt" succession (Loeme Fm.) in all 3 areas (Sharp et al., 2012; Gindre-Chanu et al., 2014). The Early to late Cretaceous succession, dominated by mixed carbonate-clastic and salt sequences, are grouped into; a pre-salt Aptian, a salt Aptian, and post salt succession ranging from Albian to Maastrichtian (See figure 2).

Volcanic sections with clear Parana-Etendeka affinities have recently been described at the base of the Cretaceous on-shore Namibe Basin in SW Angola, termed the Bero Volcanic Complex, where they unconformably overly Precambrian basement or local fluvio-lacustrine sediments of assumed Jurassic to Permian age (Sharp et al., 2012; Gindre-Chanu 2014; Marsh & Swart 2018). Additional weathered and to some extent isolated segments of volcanic and intrusive units are also found in patches further north in Angola, including volumetrically significant sill and dyke systems stretching from Namibe to Benguela, and a large area of highly faulted outcrops SE of the town of Sumbe in the Kwanza Basin, both of which are interpreted to represent erosional remnants of the originally extensive Paraná-Etendeka LIP (cf. Marzoli et al., 1999).

[Figure 2] Stratigraphic context of the Cretaceous lithologies present along the Angolan margin. Section are generalised for the Kwanza Basin (left) and onshore Namibe Basin (right) (see also Gindre-Chanu et al., 2014).

Outcrops of volcanics of a younger age have been reported from onshore locations within the Kwanza Basin which produce low volume flows that are interbedded within stratigraphic units assigned to the Albian-aged Pinda Group. These flows have alkaline compositions (Marzoli et al., 1999). Comparable low volume flows have also been documented from the Namibe Basin (Carvalho and Soares, 1961; Sharp et al., 2012; Stragnac et al., 2014), where they are interbedded with strata of reported Santonian age. As part of a study to characterise the tectono-stratigraphic development of these younger volcanics along the Angolan margin, significant occurrences of volcanic units were identified, and over a period of several focused field campaigns between 2010 and 2014 were systematically described, sampled and correlated. Focus was on detailed facies description of igneous rock types present and facies and faunal content of sediments that are intercalated within, under and over the volcanic units. Three main sections are introduced here (e.g. figure 1). These sections run along a significant section (400 km+) of the SW Angolan coast and also slightly inland, and as such provide an extremely good spatial control (strike section) on late stage magmatism in Angola. The three key sections will briefly be described below.

Site 1. Casa Branca Beach Section, Sumbe, Kwanza Basin

The first studied set of exposures are located south of the town of Sumbe at a location known locally as Casa Branca (figure 1b). This is the same coastal section as briefly described by Marzoli et al. (1999). Exceptional coastal outcrops along the Casa Branca section, in particular the previously undescribed sections that extend 2km north and 1km south of Casa Branca (figure 3a), highlight a sub-marine depositional environment associated with shallow shelf to predominantly slope mixed carbonate clastic system characterized by a range of fossiliferous wackestone, grainstones to rudstones. Deposition in water depths of no more than 200m is envisaged. The section underlying the main igneous unit is characterized by a relatively monotonous shelfal succession of interbedded marlstones and micritic carbonates characterized by infaunal bivalves, echinoderms, pervasive grazing to dwelling trace fossils and occasional ammonites. Into this environment a submarine volcanic sequence (up to 70m thick) was emplaced consisting of pillow lavas and hyaloclastite deposits. In some instances there is significant interaction between the volcanics and the sediments including; inter pillow sediments, a minor volcanic hiatus with sediment interbed (ammonite bearing micrites), and spectacular sediment filled fissures that cut the volcanic pile.

The section overlying the main volcanic interval is characterized by a mixed carbonate-clastic succession of turbidites and slope channels rich in redeposited shallow marine allochems (peloids, ooids) and fauna (oysters, rudist bivalves, Nerinid gastropods), but also well-preserved ammonites. This succession passively onlaps and oversteps the volcanic succession with a passive drape character, and ultimately shallows up to high energy tidal to shoreface carbonate grainstones. Water depths are again interpreted to range from circa 200m (base of section) to a few m's (top of section). The sediment filled fissures that cut the volcanic pile are likely related to halokinesis, with the carbonates and volcanics interpreted as having slid on rafts which detached on the underlying Aptian-aged salt. Lithostratigraphically, the sedimentary succession below the volcanics is assigned to the Catumbela Fm, whilst the unit above the volcanics is predominantly assigned to the Quissonde Fm (Fig 2).

The sediment/volcanic interaction within this section provide excellent evidence of the emplacement environment as well as good stratigraphic control which will be discussed further. The volcanic rocks themselves consist of olivine/pyroxene rich basalts/alkali basalts, which in some instances are exceptionally fresh (e.g. Figure 3b).

Site 2. 2a – Serra de Neve. 2b - Baía dos Elefantes, Benguela Basin

The studied Benguela Basin sections comprises 2 distinct locations that are here grouped for brevity of documentation. These encompasses a large inland central volcanic complex (Serra de Neve), as well as a spectacular coastal section that extends along strike for over 10km between Baía dos Elefantes and Binga (see figure 1). The central Serra de Neve complex rises as a number of NW-SE striking high hills (over 2300 m elevation) with a main dominant central area at Serra de Neve (figure 3c). The complex comprises a number of cross cutting intrusive units including granites, phonolites, and mafic intrusions. The region was targeted for study as it falls along a clear NW-SE trending lineament from the coastal exposures of lava flows and shallow intrusions at Baía dos Elefantes. Importantly, the continuation of the NW-SE trending volcanic province is also evident in offshore seismic and potential fields data (gravity and magnetics). In this context it was thought that the volcanic centre at Serra de Neve, acted as an eruption centre feeding the coastal lava sequences. Within the central part of the main mountain large feldspar rich intrusive rocks were found (figure 3d). A number of rock samples were taken in and around the complex for study. No sediments were found to be associated with the Serra de Neve complex, with the exception of very poorly exposed lacustrine sediments within the central crater. No fossil content was recovered from these sediments.

Along the coastal section between Elefantes and Binga, excellent exposures of lava sequences entering a palaeo-coastline can be seen (figure 3e). These lavas are predominantly made up of large feldspar rich flows similar to the facies found within the volcanic centre (figure 3f). The lava sequences themselves are mainly sub-aerial flows but can be seen entering a shore-face environment and in places fragmenting to hyaloclastites. The nature of the sediments directly beneath the flows can be established as submarine slope and shallow shelf carbonates and marls, and are well exposed. Immediately underlying the main volcanic section is a 50-100m thick carbonate succession including high energy grainstone shoals, indicative of deposition in a shoreface to tidal setting. Shore face deposits immediately underlying the subaerial to sub-aqueous transition in the volcanics. Then a 100 m thick slope succession comprising infaunal echinoderms and a locally abundant well preserved ammonite fauna is found. The succession has an overall progradational character, and locally includes very well exposed slope-channel systems, some of which include volcanic derived clasts, indicative of coeval volcanism. Below this sequence is a mixed carbonate clastic shallow marine shelfal succession (Dondo and Catumbela Fm, see figure 2) comprising high energy grainstone shoals and subordinate thrombolite and rhod-algal units. The platform is grain dominated, with ooids, peloids and rhodoliths dominating. The sandstones are quartz arenites. Growth faults are locally very well developed, interpreted as related to gravity gliding on the underlying salt. Shallow laccolith-like intrusives are evident intruding into the upper parts of the section associated with pepperite and soft sediment deformation – indicative of emplacement into relatively unconsolidated sediment. Seismic scale clinoform geometries within units exposed at Binga indicate depositional water depths at the time of deposition were circa 200 m, with accommodation space gradually being filled to sea level due to northwards progradation.

A 4km long North-South depositional dip section is exceptionally well exposed and accessible on the western side of Baía dos Elefantes, where the main volcanic units were sampled for geochronology in this study (figure 3e).

Site 3. Namibe volcanics section (Bentiaba, Ponta Negra, Piambo, Cangulo, Uah, Mariquita).

The third main section of volcanics sampled in the present study comprises a number of coastal and inland sections exposed between the towns of Lucira in the north and Namibe in the south (Figure 1). The igneous units occur as a number of isolated flows, volcanic centres and subordinate intrusions. These are often found as cliff forming units along the coast north of Namibe (see figure 1), forming key stratigraphic markers within the sediments (e.g figure 3g). The lava flows can again be shown to be transitional from sub-aerial flows into shallow marine settings. At a location called Ponta Negra, one of the small volcanic centres occurs as an emergent volcano that feeds first hyaloclastites and then lava flows as it evolves (figure 3G). Here glassy basaltic fragments are quenched in the hyaloclastites (e.g. figure 3h), or can be sampled as microcrystalline lavas. Sediments interbedded with the volcanics include fluvial to shallow marine mixed carbonate-clastic successions. In a similar fashion to the Benguela coastal outcrops, east to west dip sections record a transition from subaerial to subaqueous conditions for both the sediments and extrusives. At Uah, Canico, Piambo and Mariquita the immediately underlying sediments are assigned to the Ponta Salinas Fm (Carvalho, 1961; Cooper, 1972, 1978), which include ammonite and shelly fauna bearing shelfal to shoreface successions. Pepperite successions are well exposed at Uah, Canico and Maraquita valleys, indicative of lava – wet sediment interaction. At Canico, invasive flows are also evident associated with soft sediment deformation, highlighting unequivocal evidence that the flows are within sequence and emplaced into the active sedimentary environment. These volcanics represent the southern most samples collected in this study. Stgnac *et al* (2014) termed these volcanics the Ombe Fm, whilst Sharp *et al* (2013) informally termed them the Bentiaba Basanite Fm. Carvalho (1961) was the first to document their occurrence, using the sections around Bentiaba as his type location.

At all three studied locations the volcanic sections contain relatively fresh rocks and can be seen to be of Early to Late Cretaceous age (Late Albian to Santonian) in the case of the lava sequences that are found within stratigraphy. The relative age of the Serra de Neve complex is less obvious to constrain as it appears to cut through basement rocks. The range of compositions for these volcanics is presented in Figure 4 along with the distribution of regional rocks compiled recently by Comin-Chiaromonte *et al.* (2011). They cover a wide range of compositions with many of the units showing an alkali affinity with alkali basalts, basanites and phonolites (figure 3). The next stage is to better constrain their geochronology in order to understand how these magmatic events fit within the context of break-up.

[Figure 3] a) View looking North along the Sumbe volcanics section at Casa Branca, Kwanza Basin. In view are pillow basalts and hyaloclastites interbedded with and overlain by a mixed carbonate-clastic slope succession of Latest Albian age. b) Example of exceptionally fresh olivine rich basalt from the Sumbe volcanics, Casa Branca section, Kwanza Basin. c) View from the edge of the Serra de Neve complex, Southern Benguela Basin. d) Large feldspar rich phonolite from the central intrusions within the Serra de Neve complex, Southern Benguela Basin . e) Coastal exposure at Baía dos Elefantes, Benguela Basin. At this location lavas overly shelfal to slope ammonite bearing carbonates of Latest Aptian age. f) close up of feldspar rich phonolite lava flow, as similar facies to that found in d. Baía dos Elefantes sections, Benguela Basin. g) Coastal section at Ponte Negra, Namibe Basin, comprising reworked hyaloclastites, invasive flows,

pillows and emergent volcanics immediately overlying sediments of the Salinas Fm of Coniacian to Santonian age. h) Hyaloclastites exposed at the Ponta Negra section, Namibe Basin.

[Figure 4] Total alkalis vs. silica diagrams for all analysed TAS classification diagram of the samples from this contribution. Overlain on top of a selection of representative rocks from the margin (after Comin-Chiaramonti et al., 2011). Numbered fields (volcanic-intrusive): 1) picrobasalt-picrogabbro; 2) basaltgabbro; 3) basaltic andesite-gabbrodiorite; 4) andesite-diorite; 5) dacite-granodiorite; 6) rhyolite-granite; 7) trachybasalt- monzogabbro; 8) basaltic trachyandesite–monzodiorite; 9) trachyandesite-monzonite; 10) trachydacite–quartz monzonite; 11) trachyte-syenite; 12) basanite/ tephrite–foïd gabbro; 13) phonotephrite–foïd monzogabbro; 14) tephriphonolite–foïd monzosyenite; 15) phonolite–foïd syenite; 16) foïdite-foïdolite (see also Comin-Chiaramonti et al., 2011) .

Geochronological methods

In order to place the Angola volcanic sections identified into the context of their evolution with respect to the separation of Africa from South America, samples were taken for geochronological control. The general geochemical affinity of the sample areas is presented in Figure 4, which highlights that a range of compositions exist from basalts to alkaline as well as some more evolved samples. A large part of the sections are dominated by basic composition rocks, and as such the majority of geochronological dating attempted was using the Ar/Ar system. Some of the evolved samples from the SN-BE area were also targeted for U/Pb analysis. Additionally, the stratigraphic and palaeontological knowledge of the area was used to help confirm confidence in the analysis results. The techniques used will be briefly summarised in the following subsections.

⁴⁰Ar/³⁹Ar analysis

⁴⁰Ar/³⁹Ar dating was carried out at the Open University. The samples were crushed using a pestle and mortar and the crushate was sieved and washed repeatedly in de-ionised water to remove dust and clay particles from the surfaces of all the size fractions. Using a binocular microscope, whole rock basalt chips were picked, selecting pieces free, from alteration, from samples with large, fresh and inclusion free feldspars, a feldspar separate was also picked for analysis from samples where this was possible. The picked separates were cleaned ultrasonically in acetone and de-ionised water, dried using a hot plate, and packaged in aluminium foil packets of ca. 10mm x 10mm in size prior to irradiation. Samples were irradiated at the McMaster Nuclear Reactor (McMaster University, Canada) for 100 hours. Cadmium shielding was used and the samples were held in position 8B. Neutron flux was monitored using biotite mineral standard GA1550, which has an age of 99.44 ± 0.17 Ma (Renne et al. 1998; Schmitz 2012).

The samples were step-heated using a 1059nm CSI fibre laser with the extracted gases passing through a liquid nitrogen trap and two SAES AP-10 getters (at 450°C and room temperature). After this, the gases were let into a MAP215-50 mass spectrometer for measurement. The mass discrimination was measured at 283 for ⁴⁰Ar/³⁶Ar. System blanks were measured before and after every one or two analyses, and an average of the day's blanks subtracted from the data. Results were corrected for ³⁷Ar and ³⁹Ar decay since irradiation and neutron-induced interference reactions, using the following correction factors: (³⁹Ar/³⁷Ar)Ca = 0.00065 ± 0.00000325 , (³⁶Ar/³⁷Ar)Ca = 0.000264 ± 0.000001325 , and (⁴⁰Ar/³⁹Ar)K = 0.0085 ± 0.0000425 (based on analysis of K and Ca salts). The decay constants of Min et al. (2000) were used and all corrections carried out using in house

data processing software and ages calculated using Isoplot (Ludwig, 2003). All age are reported at 2 σ level and include the errors in the J value.

U/Pb analysis

U-Pb dating of zircons was carried out by Origin Analytical Ltd. using their standard methodology. Briefly, samples were disaggregated using a Retsch Jaw crusher and had heavy minerals separated from the 40-250 micron grain-size fraction using LST Heavy Liquid (2.95 g/mL). The dense fraction was then run through a Frantz magnetic separator and the non-magnetic fraction was mounted in araldite resin block and polished. The resin block was then placed into a Scanning Electron Microscope and mapped using electron dispersive spectroscopy to identify zircons in the sample. The identified zircons were then dated using quadrupole LA-ICPMS.

In order to correct for mass and instrumental bias, data were corrected against the Plesovice zircon standard (Sláma et al., 2008). During each run three Temora-2 (Black et al., 2004) and between three and ten 91500 (Woodhead et al., 2004) zircon standards were run as unknowns. Raw data were reduced using Iolite V3 software including correction for down-hole fractionation effects. To ensure that final data set contains U-Pb ages from zircons only, all data were screened for 178Hf counts as Hf is incorporated into zircon crystal lattice and is therefore abundant at the percent level in zircon (e.g. Belousova et al., 2006). All zircon ages were calculated using Isoplot (Ludwig, 2003) and ages with >10% discordance were rejected.

Stratigraphic and palaeontological control

The onshore sections of the central to southern coastal area of Angola have been subject to detailed field campaigns between 2010 and 2014, as part of a broader regional Statoil South Atlantic study addressing the onshore to offshore development of the South Atlantic margin in Angola and beyond. As part of this work, detailed stratigraphic correlations were undertaken on a basin by basin basis addressing the main syn, pre and post rift successions. A lithostratigraphic and chronostratigraphic template was established, and where possible, controlled using available biomarkers. A representative stratigraphic column for the onshore Namibe section is outlined in figure 2, and compared to the established Kwanza Basin stratigraphy. Key lithostratigraphic units are highlighted. In many of the Post Salt stratigraphic units ammonite biomarkers are present and help provide good general age constraints. Also by placing units within an Angolan margin scale sequence stratigraphic context, the relative timing of key basin wide events (e.g. salt basin development, base Pinda transgression, top Pinda flooding event etc.) are also better constrained. As each of the 3 main study locations within this contribution contain erupted volcanic sequences interbedded with fossiliferous sediments, then their relationships with-in the regional stratigraphy provides additional constraints and corroboration of any geochronological data.

Geochronological results

As introduced above the main geochronological data used within this contribution include the Ar/Ar and U/Pb systems as well as published and new stratigraphic/palaeontological control on within-stratigraphy volcanic flow units. Results from the three main study sections will be presented and then summarised before a consideration of these new results in the context of the unzipping of Africa from South America.

Casa Branca section, Kwanza Basin; $^{40}\text{Ar}/^{39}\text{Ar}$ -palaeontology

The Casa Branca Sumbe section provided some 3 km of outcrop with sediments intermingle with volcanic units emplaced in a submarine environment. Outcrops just inland from this section were also visited. Samples of the basalts were taken for $^{40}\text{Ar}/^{39}\text{Ar}$ analysis and additional stratigraphic biomarker information was also available to help build confidence in the results. Figure 5 a&b presents the $^{40}\text{Ar}/^{39}\text{Ar}$ result of a fresh whole rock analysis from the coastal section. This suggest that the age of this sequence is around 101.5 Ma, with the plateaux and invers isochron overlapping. An additional age from a sample from just inland gave an age of ~98.5 with more significant error in both the plateaux and inverse isochron age, and errors overlapping (see table 1).

One of the most striking observations made at this section was the occurrence of ammonites below, within and above the volcanic succession. Figure 5 c and d highlight an ammonite preserved within the inter-pillow section (unfortunately too poorly preserved to date). The ammonite zones within this area are relatively well known (e.g. Tavares et al 2007) and can help to provide additional information in consideration of the $^{40}\text{Ar}/^{39}\text{Ar}$ results. In short, well preserved ammonites collected during the course of this study in the section below the main volcanic section have typical Mid Albian ages, whilst new ammonite data (figure 5 e&f) from the sediments sitting immediately above the volcanics can be assigned with confidence to the last stage of the Late Albian (*Mortoniceras (Angolaites) gregory* (Spath, 1992). Upper Upper Albian, *M. (S.) rosratum* subzone, *vicina* horizon of Tavares et al., 2007 – giving an age of 99Ma). This age sits well with the volcanic age of 101 Ma (figure 5 a&b, table 1). Lithostratigraphically, the succession below the volcanics is assigned to the Catumbela Fm, whilst the succession that sits above the volcanics is assigned to the Quissonde Fm (Fig 2). It is interesting to note that the contact between the two formations is associated with a marked deepening and rotational faulting event in both onshore and offshore sections. The new data presented here indicates that the Sumbe volcanics are developed exactly at the boundary of the 2 formations, and could represent the driver for the basin wide change in depositional setting.

[Figure 5] Geochronology of the Casa Branca Beach Section, Sumbe. a & b) $^{40}\text{Ar}/^{39}\text{Ar}$ age with good overlapping plateaux and inverse isochron ~101.5 Ma. C & D) Ammonites within interpillow sediments. e, f & g) Ammonites from sediments directly above the volcanics - *Mortoniceras (Angolaites) gregoryi* (Spath, 1922), Upper Upper Albian, *M. (S.) rosratum* subzone (*vicina* horizon of Tavares et al., 2007 – giving an age of 99 Ma).

Serra de Neve – Baía dos Elefantes lineament: $^{40}\text{Ar}/^{39}\text{Ar}$ – U/Pb

The SN-BE lineament consists of two main sampling areas, the interior volcanic centre of Serra de Neve and the coastal sections around Baía dos Elefantes and Binga. Both areas are well exposed and a large range of good samples were available for analysis. Representative $^{40}\text{Ar}/^{39}\text{Ar}$ data are presented in figure 6 (a-d), and highlight a clustering of ages around 91 Ma (see also table 1 in summary section). The evolved samples within the Serra de Neve were also targeted to see if there were zircons present for U/Pb control. Figure 6e highlights some of the mineral separates and age analysis for these samples providing an additional U/Pb control age of 90.5Ma. Both these ages overlap providing good confidence in the geochronological data for this event. The stratigraphic control provided by the lava flows around the Baía dos Elefantes area suggest eruption from a subaerial into a submarine transition. New ammonite dating of samples taken within Unit 2, which sits some 50-110 m below the main lava flows, indicated a very tight age of the last stage of the Late Albian (*Mortoniceras (Angolaites) gregory* (Spath, 1992). Upper Upper Albian, *M. (S.) rosratum*

subzone, *vicina* horizon of Tavares et al., 2007 – giving an age of 99 Ma). This is the same fauna and ammonite stage that was identified to sit **above** the main lava flow succession at Casa Branca. When the geochronological and palaeontological data are combined it appears clear that the volcanic event within the studied Benguela Basin section is several Myrs younger than that documented in the Kwanza Basin.

[Figure 6] Geochronology of the Serra de Neve – Baía dos Elefantes lineament. A-D) representative $^{40}\text{Ar}/^{39}\text{Ar}$ ages from the Elefantes section. E & F) Zircon separates with Concordia age from Serra de Neve.

Namibe flows; $^{40}\text{Ar}/^{39}\text{Ar}$

In the Southern Namibe sections an area around an emergent sub-marine to subaerial volcano situated at Ponta Negra was chosen for sampling fresh material for analysis. $^{40}\text{Ar}/^{39}\text{Ar}$ data for two samples is presented in Figure 7, highlighting a younger age still at around 81-82 Ma. K-Ar dating of lava flows slightly further north in the Namibe Basin (inland from Bentiaba section) recorded slightly older dates of 88Ma \pm 1.3Ma (Statoil internal report). The majority of the Namibe section volcanics are lava flows, hyaloclastites and small eruptive centres, often nucleated along NW-SE trending segmented fault arrays, like that at Ponta Negra which feed lava/hyaloclastite sequences. Again the direct relationship with the stratigraphy is very evident. The flows are found sitting immediately above the regionally extensive transgressive Ponta Salinas Fm in most studied sections, and ammonites collected during the course of this study confidently assign this unit to the latest Cenomanian into Turonian, possibly extending into the Early Coniacian. Fauna present and dated include *Gaudryceras isovokyense* (Collignon, 1964), *Calycoceras (Calycoceras) naviculare* (Mantell, 1822) *Metoicoceras geslinianum* zone, *Metoicoceras geslinianum* (d'Orbigny, 1850) *Metoicoceras geslinianum* zone (Figure 8). Sediments and fauna found sitting transgressively above the volcanic flows at all Namibe locations have been confidently dated to the Latest Santonian to Early Campanian.

The fauna and age assignments recorded in our study are in keeping with existing stratigraphic studies in the Namibe Basin (e.g. Carvalho and Soares, 1961; Cooper, 1972, 1973, 1978, 2003 a&b; Strganac *et al* 2014). Cooper reports a Late Turonian to Early Coniacian ammonite fauna from the Salinas Fm at Bentiaba immediately below the volcanics, and a Middle Santonian to Early Campanian ammonite fauna sitting immediately above the volcanics (Cooper 2003 a&b). A recent study by Strganac *et al* (2014) have further constrained the age of the volcanics in Namibe based on a detailed section at Bentiaba using carbon isotope, magnetostratigraphy and $^{40}\text{Ar}/^{39}\text{Ar}$ methods. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of the main flow at this section gave a radiometric age of 84.6 \pm 1.5 Ma, and identification of the Cenomanian-Turonian Boundary Event in the sediments directly underneath giving an age of 93.9Ma (Ocean Anoxic Event 2). Sediments sitting above the volcanics are dated from the Santonian to Early Campanian. The dating of the volcanic centre at Ponta Negra in this study extends the duration of volcanism to 81-82 Ma, with the younging of the volcanics as we move south.

[Figure 7] $^{40}\text{Ar}/^{39}\text{Ar}$ age analysis of the Ponta Negra volcanic centre and associated flows.

[Figure 8] Ammonites found in Namibe section below volcanics: a) 11 *Calycoceras (Calycoceras) naviculare* (Mantell, 1822) Upper Cenomanian, *Metoicoceras geslinianum* Zone. b) 12 *Metoicoceras geslinianum* (d'Orbigny, 1850) Upper Cenomanian, *Metoicoceras geslinianum* Zone.

Summary of geochronological data

The three main sections sampled during this study have shown a progressive younging of ages from the northernmost Sumbe section in the Kwanza Basin through to the southernmost sections in the Namibe Basin. Table 1 provides a summary of the age data from this study. A plot of age vs distance from Namibe as well as the approximate locations of the volcanic events within the stratigraphy is given in figure 9. This provides a useful visual assessment of this data in the context of the margin. Volcanic events are occurring within the Early to Late Cretaceous, along this margin with a general younging towards Namibia and the site of the emerging Walvis Ridge relating to the continued expression of the Tristan da Cunha hotspot trail away from the Parana-Entendeka large igneous province. These volcanic events, although small, provide vital age information about the margin and are worthy of further consideration in respect of their distribution in time and space in the context of the separation of Africa with South America.

[Table 1] Summary of geochronological data presented in this contribution.

[Figure 9] Younging age progression of the Geochronology data presented in this study with stratigraphic location indicated. 1 – Kwanza Basin, 2 Benguela Basin, 3 Namibe Basin

Plate reconstruction, unzipping and volcanic distribution

Ocean basin geometries are best reconstructed by a combination of magnetic isochrons and fracture zones. Unfortunately, vast amounts of seafloor spreading in the South Atlantic occurred during the Kiaman Normal Superchron between 121 Ma (chron MO) and 83.5 Ma (anomaly 24) (Cande & Kent 1995). The opening history of the South Atlantic is therefore partly based on only fracture zone geometries and interpolation between 121 and 83.5 Ma. Figure 10 shows plate reconstructions for the time periods indicated by the volcanic events presented in this study. Also indicated are the main distribution of the Paraná-Etendeka preserved onshore exposures, and the distribution of the younger volcanic rocks from this study.

[Figure 10] Left hand diagrams: Plate reconstructions of the central South Atlantic (updated from Torsvik et al. 2009) in an absolute mantle reference frame (Dobrovine et al. 2012) Onshore volcanic sections highlighted with a star. Right hand images highlight the relative distribution of the volcanic events and locations of offshore volcanic anomalies.

The Paraná-Etendeka large igneous province (LIP) erupted above the plume generation zone (e.g. the PGZ, thick red stippled line on Figure 10, which is the 0.9% slow contour in the *s10mean* tomography model of Dobrovine et al (2016) at around 134-135 Ma (see Svensen et al. 2017). Seafloor spreading started south of the Florianopolis Fracture Zone at ~130 Ma whilst the area to the north underwent pre-break extension until ~112-110 Ma when a single subtropical syn-rift Aptian salt basin (preserved on the Brazilian, Angolan, Congo and Gabon continental shelves) was

dissected by seafloor spreading (Torsvik et al., 2009; Kukla et al., 2018). At ~100 Ma, the salt basins (Brazilian-Angolan margins) were separated by ~700 km of oceanic crust and South America is dominated by a strong component of westerly motion relative to the mantle (black thick arrows). From 100 to 81 Ma around 1200 km of westerly drift of South America (subduction rollback) corresponds 6.3 cm/yr of longitudinal motion whilst Africa is almost 'fixed' in longitude over the same time interval (see black absolute velocity arrows in figure 10). The white arrows at 92 and 81 Ma (figure 10) show the relative motion paths for three points on the African margin (relative South America) and denote the motion paths from 100 to 92 Ma (92 Ma reconstruction) and 90 to 81 Ma (81 Ma reconstruction). Additionally the NW-SE trending volcanic bodies are also developed on ocean crust picked out using seismic and potential field data. Seismic stratigraphy also clearly shows the offshore Sumbe volcanics young southwards (*in-house* Statoil data). These observations are consistent with the timings of separation as listed above, and the ages presented in this study.

The offshore structural domain and seismic stratigraphy work (e.g. Higgins et al submitted) clearly shows that the rift offshore Angola is "unzipping" and propagating from north to south. The age of this unzipping seems to be post salt (Albian) in age south of the city of Namibe, but is older than that in the north. Interestingly, the Inner Santos rift is clearly propagating from south to north around this time, with the Sao Paulo plateau rifted fragment caught in between (cf. Scotchman et al 2011). There are younger than Paraná-Etendeka volcanic events found on the Brazilian side. Ages around 100-102 Ma are reported for the Cabo Magmatic Province (Nascimento et al., 2003), which is somewhat to the north of this region when looking at the reconstructions. This seems to also be corroborated by palaeomagnetic data (e.g. Font et al., 2009). Younger ages of ~90Ma (mostly K-Ar) have been noted in the Potiguar Basin (equatorial region) and as some ages from south of Brazil (Mizusaki et al., 2002; Souza et al., 2003). The alkaline centre at Monte de Trigo Island in a geographically very close proximity section on the Brazilian margin has a reported age of 86.6 Ma ($^{40}\text{Ar}/^{39}\text{Ar}$, Enrich et al., 2009). Other alkaline intrusions (dykes and plugs) with ages of 82 Ma as well as younger examples are found intersecting the basement of the Santos Basin (Guedes et al., 2005), which would also be geographically at a similar section along the Brazilian side of the South Atlantic Margin. Additionally, kimberlites and phlogopite bearing picrites are reported with age ranges from 91-78 Ma within the Paranaíba Igneous Province (Guarino et al., 2013), which are occurring inland from the rifting margin, with some similar young kimberlites pulses (90 ± 10 Ma and 60 ± 10 Ma) preserved inland on the African margin (e.g. Tappe et al., 2018). These all give testament of a series of younger than Paraná-Etendeka magmatic events, albeit of relatively low volumes, that occur as the rifting continents parted. The new ages presented in this contribution clearly highlight that more geochronological information can be yielded from the Late Cretaceous and can provide valuable geographical constraints on magmatic events in the context of the continental separation of Africa from South America.

Conclusions and closing remarks

This contribution has introduced the geochronological and plate tectonic significance of Early to Late Cretaceous magmatic events found along the Angolan section of the West African margin. Three main igneous events are found and have been placed into their spatial and temporal occurrences along the margin with the following main conclusions and comments:

- The oldest of the events is found along a coastal section (Casa Branca) south of the city of Sumbe, termed the Sumbe Volcanics, which erupted in a shallow sub-marine environment at around 100 Ma. The main rock types consist of pillow basalts and hyaloclastites which interact with and are also interbedded between fossiliferous mixed carbonate-clastic shelfal sediments of Late Albian age. The volcanics at this location are “sealed” by sediments dated as latest Albian.
- The next event occurred at around 91 Ma within the Benguela Basin and is found spanning an area from a large volcanic centre, Sierra de Neve 65 km inland from the coast, out to a coastal section between Baia dos Elefantes and Binga. The section is termed the Sierra de Neve-Baia dos Elefantes lineament. The rocks contain both basic to evolved units, with a full suite from intrusives to lava flows that have flowed into a palaeo coastal environment of shore face and shallow marine sediments deposited in water depths of between 0-200 m. Sediments underlying the main volcanic events at the coast have been dated as Latest Albian.
- The final event is as young as 82-81 Ma considering the new $^{40}\text{Ar}/^{39}\text{Ar}$ ages presented here. With the additional associated age constraints this pulse spans between 88-81Ma, and is found as lava flows, hyaloclastites and localised plugs, in a coastal zone north of the town Namibe, termed the Namibe volcanics in this contribution. These commonly transition from lava flows from sub-aerial into shoreface and shallow marine environments as well as submarine eruptions that emerge to form sub-aerial volcanoes. These sections are underlain by well dated fossiliferous sediments of Turonian-Early Coniacian age, and overlain by sediments dated as Middle Santonian to Early Campanian age
- $^{40}\text{Ar}/^{39}\text{Ar}$, U/Pb and paleontological constraints have been used in combination with a regionally understood stratigraphic sequence to place these three magmatic events into order. With a trend from north to south along the coast in terms of relative emplacement age.
- Plate reconstructions, including the location of the main salt basins, have been presented to highlight the unzipping of the margin and the location of these magmatic events that migrate southwards to the proto Walvis ridge.
- Similar constraints on the Early to Late Cretaceous volcanics found along the equivalent sections along the Brazilian side of the South American Margin will be needed to fully assess the role of this late stage magmatism as the African and South American continents separate.

Acknowledgements

Field discussions, help and general camaraderie was greatly appreciated from; Eric Blanc, Laurent Gindre-Chanu, Marco Snidero, Mark Scott, Makutulu Dongala, Miguel B. Gamela, and Julio Pacheco. Rafael and the camp organising team are thanked for their great hospitality and food during our field seasons. Equinor (formally Statoil) are gratefully acknowledged for permission to publish our results. The manuscript benefited from two anonymous reviews and with editorial guidance from Philippe Agard and the Tectonophysics team. Dougal Jerram is partly funded through a Norwegian Research Council Centres of Excellence project (project number 223272, CEED).

Figure and table captions

Figure 1. a) Location maps of SW Angola showing the 3 main study locations. b) Casa Branca beach section, Sumbe, Kwanza Basin. c) Baía dos Elefantes and Serra de Neve, Benguela Basin. d) Bentiaba, Ponta Negra, Canico, Piambo and Uah sections, Namibe Basin. Additional features offshore indicated.

Figure 2 Stratigraphic context of the Cretaceous lithologies present along the Angolan margin. Section are generalised for the Kwanza Basin (left) and onshore Namibe Basin (right) (see also Gindre-Chanu et al., 2014).

Figure 3. a) View looking North along the Sumbe volcanics section at Casa Branca, Kwanza Basin. In view are pillow basalts and hyaloclastites interbedded with and overlain by a mixed carbonate-clastic slope succession of Latest Albian age. b) Example of exceptionally fresh olivine rich basalt from the Sumbe volcanics, Casa Branca section, Kwanza Basin. c) View from the edge of the Serra de Neve complex, Southern Benguela Basin. d) Large feldspar rich phonolite from the central intrusions within the Serra de Neve complex, Southern Benguela Basin. e) Coastal exposure at Baía dos Elefantes, Benguela Basin. At this location lavas overly shelfal to slope ammonite bearing carbonates of Latest Aptian age. f) close up of feldspar rich phonolite lava flow, as similar facies to that found in d. Baía dos Elefantes sections, Benguela Basin. g) Coastal section at Ponte Negra, Namibe Basin, comprising reworked hyaloclastites, invasive flows, pillows and emergent volcanics immediately overlying sediments of the Salinas Fm of Coniacian to Santonian age. h) Hyaloclastites exposed at the Ponta Negra section, Namibe Basin.

Figure 4 Total alkalis vs. silica diagrams for all analysed TAS classification diagram of the samples from this contribution. Overlain on top of a selection of representative rocks from the margin (after Comin-Chiaramonti et al., 2011). Numbered fields (volcanic-intrusive): 1) picrobasalt-picrogabbro; 2) basaltgabbro; 3) basaltic andesite-gabbrodiorite; 4) andesite-diorite; 5) dacite-granodiorite; 6) rhyolite-granite; 7) trachybasalt- monzogabbro; 8) basaltic trachyandesite–monzodiorite; 9) trachyandesite-monzonite; 10) trachydacite–quartz monzonite; 11) trachyte-syenite; 12) basanite/ tephrite–foiid gabbro; 13) phonotephrite–foiid monzogabbro; 14) tephriphonolite–foiid monzosyenite; 15) phonolite–foiid syenite; 16) foidite-foiidolite (see also Comin-Chiaramonti et al., 2011) .

Figure 5. Geochronology of the Casa Branca Beach Section, Sumbe. a & b) $^{40}\text{Ar}/^{39}\text{Ar}$ age with good overlapping plateaux and inverse isochron ~ 101.5 Ma. c & d) Ammonites within interpillow sediments. e, f & g) Ammonites from sediments directly above the volcanics - *Mortoniceras (Angolaites) gregoryi* (Spath, 1922), Upper Upper Albian, M. (S.) *rosratum* subzone (vicina horizon of Tavares et al., 2007 – giving an age of 99 Ma).

Figure 6. Geochronology of the Serra de Neve – Baía dos Elefantes lineament. A-D) representative $^{40}\text{Ar}/^{39}\text{Ar}$ ages from the Elefantes section. E & F) Zircon separates with Concordia age from Serra de Neve.

Figure 7 $^{40}\text{Ar}/^{39}\text{Ar}$ age analysis of the Ponta Negra volcanic centre and associated flows

Figure 8 Ammonites found in Namibe section below volcanics: a) 11 *Calycoceras (Calycoceras) naviculare* (Mantell, 1822) Upper Cenomanian, *Metoicoceras geslinianum* Zone. b) 12 *Metoicoceras geslinianum* (d'Orbigny, 1850) Upper Cenomanian, *Metoicoceras geslinianum* Zone.

Figure 9 Younging age progression of the Geochronology data presented in this study with stratigraphic location indicated. 1 – Kwanza Basin, 2 Benguela Basin, 3 Namibe Basin

Figure 10 – Left hand diagrams: Plate reconstructions of the central South Atlantic (updated from Torsvik et al. 2009) in an absolute mantle reference frame (Dobrovine et al. 2012) Onshore volcanic sections highlighted with a star. Right hand images highlight the relative distribution of the volcanic events and locations of offshore volcanic anomalies.

Table 1 Summary of geochronological data presented in this contribution.

ACCEPTED MANUSCRIPT

References:

- Belousova, E.A., Griffin, W.L., O'Reilly, S.Y., 2006. Zircon morphology, trace element signatures and Hf-isotope composition as a tool for petrogenetic modeling: examples from eastern Australian granitoids. *J. Petrol.* 47, 329–353. <https://doi.org/10.1093/petrology/egi077>
- Black, Lance P., Kamo, Sandra L., Allen, Charlotte M., Davis, Donald W., Aleinikoff, John N., Valley, John W., Mundil, Roland, Campbell, Ian H., Korsch, Russell J., Williams, Ian S., Foudoulis, Chris, 2004. Improved $^{206}\text{Pb}/^{238}\text{U}$ microprobe geochronology by the monitoring of a trace-element-related matrix effect; SHRIMP, ID-TIMS, ELA-ICP-MS and oxygen isotope documentation for a series of zircon standards. *Chem. Geol.*, 205(1-2), 115-140. <https://doi.org/10.1016/j.chemgeo.2004.01.003>
- Cande, S.C., Kent, D.V., 1995. Revised calibration of the geomagnetic polarity timescale for the Late Cretaceous and Cenozoic. *J. Geophys. Res.*, 100, 6093–6095. <https://doi.org/10.1029/94JB03098>
- Carvalho, G., Soares de, 1961. Geologia do deserto de Mocamedes (Angola); Uma contribuição para o conhecimento das problemas da orla sedimentar de Mocamedes. *Mem. Junta Invest. Ultramar*, Lisbon 26.
- Cooper, M.R., 1972. The Cretaceous stratigraphy of San Nicolau and Salinas, Angola. *Annals S. Afr. Mus.* 60 (8), 245–264.
- Cooper, M.R., 1973. Cenomanian ammonites from Novo Redondo, Angola. *Ann. S. Afr. Mus.* 62 (2), 41–67.
- Cooper, M.R., 1978. Uppermost Cenomanian-Basal Turonian ammonites from Salinas. Angola. *Ann. S. Afr. Afr. Mus.* 75 (5), 51–152.
- Cooper, M.R., 2003a. Upper Cretaceous Ammonites (Turonian-Coniacian) from São Nicolau, Angola. *Ann. S. Afr. Afr. Mus.* 110, 89–146.
- Cooper, M.R., 2003b. Stratigraphy and paleontology of the Upper Cretaceous (Santonian) Baba formation at São Nicolau, Angola. *Ann. S. Afr. Afr. Mus.* 110, 147–170.
- Comin-Chiaramonti, P., De Min, A., Girardi, V.A.V., Ruberti, E., 2011. Post-Paleozoic magmatism in Angola and Namibia: A review, in Beccaluva, L., Bianchini, G., and Wilson, M., eds., *Volcanism and Evolution of the African Lithosphere: Geol. Soc. Am. Special Paper 478*, pp. 223–247. [https://doi.org/10.1130/2011.2478\(12\)](https://doi.org/10.1130/2011.2478(12))
- Dodd, S.C., Mac Niocaill, C., Muxworthy, A.R., 2015. Long duration (>4 Ma) and steady-state volcanic activity in the early Cretaceous Paraná–Etendeka Large Igneous Province: New palaeomagnetic data from Namibia. *Earth Planet. Sci. Lett.*, 414 (15), 16–29. <https://doi.org/10.1016/j.epsl.2015.01.009>
- Dobrovine, P.V., Steinberger, B., Torsvik, T.H., 2012. Absolute plate motions in a reference frame defined by moving hotspots in the Pacific, Atlantic and Indian oceans. *J. Geophys. Res.*, 117: B09101. doi:10.1029/2011JB009072. <https://doi.org/10.1029/2011JB009072>

- Doubrovine, P.V., Steinberger, B., and Torsvik, T.H. 2016. A failure to reject: Testing the correlation between large igneous provinces and deep mantle structures with EDF statistics. *Geochem. Geophys. Geosyst.*, 17, 1130–1163, doi:10.1002/2015GC006044.
- Enrich, G.E.R., Ruberti E., Gomes C.B., 2009. Geology and geochronology of Monte de Trigo island alkaline suite, southeastern Brazil. *Revista Brasileira de Geociências*, 39(1), 67-80. DOI | 10.5327/rbg.v39i1.1441
- Font, E., Ernesto, M., Silva, P. F., Correia, P. B., Nascimento, M. A. L., 2009. Palaeomagnetism, rock magnetism and AMS of the Cabo Magmatic Province, NE Brazil, and the opening of South Atlantic. *Geophys. J. Int.*, 179, 905–922. <https://doi.org/10.1111/j.1365-246X.2009.04333.x>
- Gindre-Chanu, L., Warren, K.J., Puigdefabregas, C., Sharp, I., Peacock, C.P., Swart, R., Poulsen, R., Ferreira, H., Henrique, L., 2014. Diagenetic evolution of Aptian evaporates in the Namibe basin, South-West Angola. *Sedimentol.* 62 (1), 204–233. <https://doi.org/10.1111/sed.12146>
- Gindre-Chanu Laurent, Perri Edoardo, Sharp R. Ian, D.C.P. Peacock, Swart Roger, Poulsen Ragnar, Ferreira Hercinda, Machado Vladimir. 2016. Origin and diagenetic evolution of gypsum and microbialitic carbonates in the Late Sag of the Namibe Basin (SW Angola). *Sed. Geol.*, 342, 2016, 133-153. <https://doi.org/10.1016/j.sedgeo.2016.06.015>.
- Guarino, V., Fu-Yuan Wu, Michele Lustrino, Leone Melluso, Pietro Brotzu, Celso de Barros Gomes, Excelso Ruberti, Colombo Celso Gaeta Tassinari, Darcy Pedro Svisero, 2013. U–Pb ages, Sr–Nd- isotope geochemistry, and petrogenesis of kimberlites, kamafugites and phlogopite-picrites of the Alto Paranaíba Igneous Province, Brazil. *Chem. Geol.*, 353, 65-82. <https://doi.org/10.1016/j.chemgeo.2012.06.016>.
- Guedes, E., Monica Heilbron, Paulo M. Vasconcelos, Claudio de Morisson Valeriano, Julio Cesar Horta de Almeida, Wilson Teixeira, Antonio Thomaz Filho, 2005. K–Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ ages of dikes emplaced in the onshore basement of the Santos Basin, Resende area, SE Brazil: implications for the south Atlantic opening and Tertiary reactivation. *J. South Am. Earth Sci.*, 18, 371–382. <https://doi.org/10.1016/j.jsames.2004.11.008>
- Heine, C., Zoethout, J., Müller, R.D., 2013. Kinematics of the South Atlantic Rift. *Solid Earth Discussions* 5, 41–115. <https://doi.org/10.5194/se-4-215-2013>.
- Jackson, M.P.A., Cramez, C. & Fonck, J-M., 2000. Role of subaerial volcanic rocks and mantle plumes in creation of South Atlantic margins: implications for salt tectonics and source rocks. *Mar. Petrol. Geol.*, 17, 477–498. [https://doi.org/10.1016/S0264-8172\(00\)00006-4](https://doi.org/10.1016/S0264-8172(00)00006-4)
- Jerram, D.A., Svensen, H.H., Planke, S., Polozov, A.G., and Torsvik, T.H., 2016. The onset of flood volcanism in the north-western part of the Siberian traps: Explosive volcanism versus effusive lava flows. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 441, 38-50. <https://doi.org/10.1016/j.palaeo.2015.04.022>
- Jerram, D.A., Single, R.T., Hobbs, R.W., Nelson, C.E., 2009. Understanding the offshore flood basalt 1507 sequence using onshore volcanic facies analogues: an example from the Faroe-Shetland basin. *Geol. Mag.*, 146, 353-367. <https://doi.org/10.1017/S0016756809005974>

- Jerram, D. a. & Widdowson, M., 2005. The anatomy of Continental Flood Basalt Provinces: Geological constraints on the processes and products of flood volcanism. *Lithos*, 79, 385–405. <https://doi.org/10.1016/j.lithos.2004.09.009>
- Jerram, D.A., Mountney, N., Holzförster, F., Stollhofen, H., 1999. Internal stratigraphic relationships in the Etendeka Group in the Huab Basin, NW Namibia: Understanding the onset of flood volcanism. *J. Geodynamics*, 28, 393-418. [https://doi.org/10.1016/S0264-3707\(99\)00018-6](https://doi.org/10.1016/S0264-3707(99)00018-6)
- Kukla, P. A., Frank Strozyk, Webster U. Mohriak, 2018. South Atlantic salt basins – Witnesses of complex passive margin evolution. *Gondwana Research*, 53, 41-57. <https://doi.org/10.1016/j.gr.2017.03.012>
- Le Maitre, R.W., 1989, *A Classification of Igneous Rocks and Glossary of Terms*: Oxford, Blackwell Scientific Publications, 193 p.
- Ludwig, K. R., 2003. Isoplot/Ex 3. A geochronological toolkit for Microsoft Excel. Berkeley Geochronology Center. Special publication 4.
- Marsh, J.S., Swart, R., 2018. The Bero Volcanic Complex: Extension of the Paraná-Etendeka Igneous Province into SW Angola, *J. Volcanol. Geotherm. Res.*, 355, 21-31. <http://dx.doi.org/10.1016/j.jvolgeores.2016.10.011>
- Marsh, J. S., Ewart, A., Milner, S. C., Duncan, A. R., Miller, R. McG., 2001. The Etendeka Igneous Province: magma types and their stratigraphic distribution with implications for the evolution of the Paraná–Etendeka flood basalt province. *Bull. Volc.*, 62, 464–486. <https://doi.org/10.1007/s004450000115>
- Marzoli, A., Melluso, L., Morra, V., Renne, P.R., Sgrosso, I., D'Antonio, M., Duarte Morais, L., Morais, E.A.A., Ricci, G., 1999. Geochronology and petrology of Cretaceous basaltic magmatism in the Kwanza basin (western Angola), and relationships with the Paraná-Etendeka continental flood basalt province. *J. Geodyn.* 28, 341–356. [https://doi.org/10.1016/S0264-3707\(99\)00014-9](https://doi.org/10.1016/S0264-3707(99)00014-9)
- Menzies, M.A., Klemperer, S.L., Ebinger, C.J., Baker, J., 2002. Characteristics of volcanic rifted margins. In: *Volcanic Rifted Margins* (eds Menzies, M. A., Klemperer, S. L., Ebinger, C. J. & Baker, J.), Geological Society of America Special Paper 362, pp. 1-14. <https://doi.org/10.1130/0-8137-2362-0.1>
- Milner, S.C., Duncan, A.R., Whittingham, A.M., Ewart, A., 1995. Trans-Atlantic correlation of eruptive sequences and individual silicic units within the Paraná-Etendeka igneous province. *J. Volcanol. Geotherm. Res.*, 69, 137–157. [https://doi.org/10.1016/0377-0273\(95\)00040-2](https://doi.org/10.1016/0377-0273(95)00040-2)
- Min, K.W., Mundil, R., Renne, P.R., Ludwig, K.R., 2000. A test for systematic errors in $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology through comparison with U/Pb analysis of a 1.1-Ga rhyolite. *Geochim. Cosmochim. Acta*, 64, 73-98. [https://doi.org/10.1016/S0016-7037\(99\)00204-5](https://doi.org/10.1016/S0016-7037(99)00204-5)
- Mizusaki, A. M. P., Thomaz-Filho, A., Milani, E. J., de Césero, P., 2002. Mesozoic and Cenozoic igneous activity and its tectonic control in northeastern Brazil. *J. South Am. Earth Sci.*, 15(2), 183-198. [https://doi.org/10.1016/S0895-9811\(02\)00014-7](https://doi.org/10.1016/S0895-9811(02)00014-7)

- Mohriak, W.U., Macedo, J.M., Castellani, R.T., Rangel, H.D., Barros, A.Z.N., Latg, M.A.L., Mizusaki, A.M.P., Szatmari, P., Demercian, L.S., Rizzo, J.G., Aires, J.A., 1995. Salt tectonics and structural styles in the deep-water province of the Cabo Frio region, Rio de Janeiro, Brazil. In: Jackson, M.P.A., Roberts, D.G., Snelson, S. (Eds.), *Salt Tectonics: A Global Perspective*. AAPG Memoir 65, pp. 273–304. <https://doi.org/10.1306/M65604C13>
- Mohriak, W., Leroy, S., 2012. Architecture of rifted continental margins and break-up evolution: insights from the South Atlantic, North Atlantic and Red Sea-Gulf of Aden conjugate margins. In: Mohriak, W.U., Danforth, A., Post, P.J., Brown, D.E., Tari, G.C., Nemčok, M., Sinha, S.T. (Eds.), *Conjugate Divergent Margins*. Geol. Soc. Lond., Spec. Publ., 369, 497-535. <http://dx.doi.org/10.1144/SP369.17>.
- Moulin, M., Aslanian, D., Unternher, P., 2010. A new starting point for the South and Equatorial Atlantic Ocean. *Earth-Sci. Rev.*, 98, 1–37. doi: 10.1016/j.earscirev.2009.08.001
- Nascimento, M.A.L., Vasconcelos, P.M., Souza, Z.S., Jardim de Sá, E.F., Carmo, I.O., and Thied, D. 2003. 40Ar-39Ar geochronology of the Cabo Magmatic Province, Pernambuco Basin, NE Brazil. *Short Papers IV South American Symposium on Isotope Geology*, 624-627.
- Nelson, C.E., Jerram, D.A., Clayburn, J.A.P., Halton, A.M., Roberge, J., 2015. Eocene volcanism in offshore southern Baffin Bay. *Mar. Pet. Geol.*, 67, 678-691. <https://doi.org/10.1016/j.marpetgeo.2015.06.002>
- Nürnberg, D., Müller, R.D., 1991. The tectonic evolution of the South Atlantic from late Jurassic to present. *Tectonophysics*, 191, 27-53. [https://doi.org/10.1016/0040-1951\(91\)90231-G](https://doi.org/10.1016/0040-1951(91)90231-G)
- Peate, D.W., 1997. The Parana-Etendeka Province. In: Large Igneous Provinces. :Mahoney, J.J.:Coffin, M.F.: (Editors), American Geophysical Union, Washington D.C.. 217-245. <https://doi.org/10.1029/GM100p0217>
- Pérez-Díaz, L., Eagles, G., 2014. Constraining South Atlantic growth with seafloor spreading data. *Tectonics*, 33. doi: 10.1002/(ISSN)1944-9194.2014TC003644
- Masse, P., Laurent, O., 2016. Geological exploration of Angola from Sumbe to Namibe: A review at the frontier between geology, natural resources and the history of geology, In *Comptes Rendus Geoscience*, 348(1), 80-88. <https://doi.org/10.1016/j.crte.2015.09.007>
- Renne, P.R., Swisher, C.C., Deino, A.L., Karner, D.B., Owens, T.L., DePaolo, D.J., 1998. Intercalibration of standards, absolute ages and uncertainties in 40Ar/39Ar dating: *Chem. Geol.*, 145, 117-152. [https://doi.org/10.1016/S0009-2541\(97\)00159-9](https://doi.org/10.1016/S0009-2541(97)00159-9)
- Renne, P. R., Mundil, R., Balco, G., Min, K. & Ludwig, K. R., 2010. Joint determination of 40K decay constants and 40Ar*/40K for the Fish Canyon sanidine standard, and improved accuracy for 40Ar/39Ar geochronology. *Geochimica et Cosmochimica Acta*, 74(18), 5349–5367. <https://doi.org/10.1016/j.gca.2010.06.017>
- Rooney, T. O., 2017. The Cenozoic magmatism of East-Africa: Part I — Flood basalts and pulsed magmatism, In *Lithos*, Volumes 286–287, 264-301. <https://doi.org/10.1016/j.lithos.2017.05.014>.

- Schmitz, M.D., 2012. Radiogenic isotope geochronology. In: Gradstein, F.M., Ogg, J.G., Schmitz, M. & Ogg, G. (eds), *The geologic time scale*, pp. 115-126. Amsterdam: Elsevier. DOI: 10.1016/B978-0-444-59425-9.00006-8
- Scotchman, I.C., Gilchrist, G., Kusznir, N.J., Roberts, A.M., Fletcher, R., 2011. The breakup of the South Atlantic Ocean: formation of failed spreading axes and blocks of thinned continental crust in the Santos Basin, Brazil and its consequences for petroleum system development. In: Vining, B.A., Pickering, S.C. (Eds.), *Petroleum Geology: From Mature Basins to New Frontiers—Proceedings of the 7th Petroleum Geology Conference*, Geological Society, London, *Petroleum Geology Conference Series*, 7, 855–866. <https://doi.org/10.1144/0070855>
- Sharp, I., Verwer, K., Ferreira, H., Snidero, M., Machado, V., Holtar, E., Swart, R., Marsh, J., Gindre, L., Puigdefabregas, C., Fejerskov, M., 2012. In: *Pre- and Post-Salt Non-Marine Carbonates of the Namibe Basin Angola*, AAPG Search and Discovery Article #90142 (2012 AAPG Annual Conf).
- Slama, Jiri; Kosler, Jan; Condon, Daniel J.; Crowley, James L.; Gerdes, Axel; Hanchar, John M.; Horstwood, Matthew S. A.; Morris, George A.; Nasdala, Lutz; Norberg, Nicholas; Schaltegger, Urs; Schoene, Blair; Tubrett, Michael N.; Whitehouse, Martin J., 2008. Plesovice zircon - A new natural reference material for U-Pb and Hf isotopic microanalysis. *Chem. Geol.*, 249(1-2), 1-35. <https://doi.org/10.1016/j.chemgeo.2007.11.005>
- Souza, Z.S., Vasconcelos, P.M., Nascimento, M.A.L., Silveira, F.V., Paiva, H.S., Dias, L.G.D., Thiede, D., and Carmo, I.O., 2003. $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of Mesozoic and Cenozoic magmatism in NE Brazil. *short Papeers, IV South American Symposium on Isotope Geology*, 691-694.
- Spath, L. F., 1922. On Cretaceous Ammonoidea from Angola, collected by Professor J. W. Gregory, D.Sc., FR.S. *Trans. R. Soc. Edinburgh*, 53, 91-160.
- Strozyk, F., Stefan Back, Peter A. Kukla, 2017. Comparison of the rift and post-rift architecture of conjugated salt and salt-free basins offshore Brazil and Angola/Namibia, South Atlantic, In *Tectonophysics*, 716, 204-224. <https://doi.org/10.1016/j.tecto.2016.12.012>.
- Strganac, Ch., Salminen, J., Jacobs, L.L., Polcyn, M.J., Ferguson, K.M., Mateus, O., Schulp, A., Morais, M.L., Tavares de Silva, T., Gonçalves, A.O., 2014. Carbon isotope stratigraphy, magnetostratigraphy, $^{40}\text{Ar}/^{39}\text{Ar}$ age of the Cretaceous South Atlantic coast, Namibe basin, Angola. *J. Afr. Earth Sci.*, 99 (2), 452–462. <https://doi.org/10.1016/j.jafrearsci.2014.03.003>
- Svensen, H. H., T. H. Torsvik, S. Callegaro, L. Augland, T. H. Heimdal, D. A. Jerram, S. Planke and E. Pereira 2017. Gondwana Large Igneous Provinces: plate reconstructions, volcanic basins and sill volumes. *Geol. Soc. Lond., Spec. Publ.*, 463, 17-40. <https://doi.org/10.1144/SP463.7>
- Szatmari, P., Milani, E.J., 2016. Tectonic control of the oil-rich large igneous-carbonate-salt province of the South Atlantic rift. *Mar. Pet. Geol.*, 77, 567-596. <https://doi.org/10.1016/j.marpetgeo.2016.06.004>

- Tappe, S., Katie Smart, Trond Torsvik, Malcolm Massuyeau, Mike de Wit, 2018. Geodynamics of kimberlites on a cooling Earth: Clues to plate tectonic evolution and deep volatile cycles, *Earth Planet. Sci. Lett.*, 484, 1-14. <https://doi.org/10.1016/j.epsl.2017.12.013>.
- Tavares, T., Meister, Ch., Duarte-Morais, David, B., 2007. Albian ammonites of the Benguela Basin (Angola): a biostratigraphic framework. *S. Afr. J. Geol.*, 110, 137–156. <https://doi.org/10.2113/gssajg.110.1.137>
- Torsvik, T.H., Cocks, L.R.M., 2013. Gondwana from top to base in space and time. GR focus review, *Gondwana Research*, 24, 999-1030. <https://doi.org/10.1016/j.gr.2013.06.012>
- Torsvik, T.H., Sonia Rouse, S., Labails, C. and Smethurst M. A. 2009. A new scheme for the opening of the South Atlantic Ocean and the dissection of an Aptian salt basin. *Geophys. J. Int.*, 177, 1315–1333. doi: 10.1111/j.1365-246X.2009.04137.x
- Torsvik, T.H., Müller, R.D., Van der Voo, R., Steinberger, B., Gaina, C., 2008. Global plate motion frames: Toward a unified model. *Reviews of Geophysics*, 46, RG3004, pp. 44. <https://doi.org/10.1029/2007RG000227>
- Torsvik, T.H., Smethurst, M.A., 1999. Plate tectonic modelling: virtual reality with GMAP (GMAP Standard is available as a freeware on <http://www.ngu.no/geophysics1>). *Comp. Geosci.*, 25, 395-402. [https://doi.org/10.1016/S0098-3004\(98\)00143-5](https://doi.org/10.1016/S0098-3004(98)00143-5)
- Torsvik, T.H., Steinberger, B., Gurnis, M., Gaina, C., 2010. Plate tectonics and net lithosphere rotation over the past 150 My. *Earth Planet. Sci. Lett.*, 291, 106-112. <https://doi.org/10.1016/j.epsl.2009.12.055>
- Torsvik, T.H., Steinberger, B., Ashwal, L.D, Doubrovine, P.V. & Trønnes, R.C. 2016. Earth Evolution and Dynamics – A tribute to Kevin Burke. *Canadian Journal of Earth Sciences Special Issue: Uniformitarianism and Plate Tectonics: A tribute to Kevin C. Burke and John F. Dewey*. *Canadian Journal of Earth Sciences*, 53(11): 1073-1087. <https://doi.org/10.1139/cjes-2015-0228>
- Untemehr, P., Curie, D., Olivet, J.L., Goslin, J., Beuzart, P., 1988. South Atlantic fits and intraplate boundaries in Africa and South America. *Tectonophysics*, 155, 169-179. [https://doi.org/10.1016/0040-1951\(88\)90264-8](https://doi.org/10.1016/0040-1951(88)90264-8)
- Watton, T.J., Jerram, D.A., Thordarson, T., Davies, R.J., 2013. Three-dimensional lithofacies variations in hyaloclastite deposits. *J. Volcanol. Geotherm. Res.*, 250, 19-33. <https://doi.org/10.1016/j.jvolgeores.2012.10.011>
- Woodhead, J., Hergt, J., Shelley, M., Eggins, S., Kemp, R., 2004. Zircon Hf-isotope analysis with an excimer laser, depth profiling, ablation of complex geometries and concomitant age estimation. *Chemical Geology*, 209, 121–135. <https://doi.org/10.1016/j.chemgeo.2004.04.026>
- Wright, K.A., Davies, R.J., Jerram, D.A., Morris, J., Fletcher, R., 2012. Application of seismic and 1809 sequence stratigraphic concepts to a lava-fed delta system in the Faroe-Shetland Basin, UK

and 1810 Faroes. Basin Research, 24, 91-106. <https://doi.org/10.1111/j.1365-2117.2011.00513.x>

ACCEPTED MANUSCRIPT





Figure 1



Key to main units

- Iabe Group. Late Cretaceous mixed carbonate-clastic shelfal succession.
- Albian Pinda Group. Mixed carbonate-clastic-evaporitic marine to marginal/non marine succession.
- Loeme Fm. evaporites. Halite, gypsum & anhydrite.
- Mixed carbonate-clastic, non marine to marginal marine succession. Cuvo-Chela Fm. equivalent. "pre-Salt" Aptian.

- ⑨ Maastrichtian and Campanian mixed carbonate-clastic, non marine to marine. Fluvial-shoreface-shelf.
- ⑧ Mixed igneous, clastic, carbonate Sumbe volcanics. Non marine to marginal marine with local travertines.
- ⑦ Transgressive Cenomanian-Turonian Salinas Fm. Fluvial-shoreface-shelf.
- ⑥ Mid Albian Douga Fm. equivalent. Fluvial to marginal marine clastics. Braided river to fan delta. Passing into Pinda carbonates to the west.
- ⑤ Early Albian Binga Fm. equivalent. Marginal marine mixed carbonate-clastic succession, rhod-algal.
- ④ Loeme/Bambata Fm. equivalent. Evaporites - gypsum + anhydrite. Halite in subsurface.
- ③ "Sag" lacustrine to marginal marine mixed carbonate-clastic succession. Including sabkha, fluvial-tidal valley fills, lacustrine-marginal marine mouth bar, fan deltas, braided river + alluvial fans. Thin interbedded lacustrine marine + non marine carbonates. Overall transgressive/fining upwards motif. Cuvo/Chela Fm. equivalent.
- ② "Sag" carbonates (lacustrine + travertines) + clastics (lacustrine + fluvial)
- ① Etendeka volcanics with interbedded aeolian/fluvial sediments. Some Karoo sediments beneath (Perm.-Jur.), Pan African basement rocks.

Figure 2



Figure 3




Figure 4




Figure 5









Figure 6







Figure 7



Figure 8





Figure 9




Figure 10