

---

RAFFLES BULLETIN OF ZOOLOGY Supplement No. 30: 97–108

Date of publication: 25 December 2014

<http://zoobank.org/urn:lsid:zoobank.org:pub:95535B0A-560E-415D-B0BB-4C92D7003F63>

## Salt and water balance in the land crabs of Christmas Island: a review

Lucy M. Turner

**Abstract.** Work conducted on the land crabs of Christmas Island has advanced our knowledge of the ecophysiology of ion and water balance not only in terrestrial crabs, but across the Crustacea. For instance, studies undertaken on endemic Christmas Island species, such as *Discoplax celeste*, which is unique in its evolutionary position and ecology as a freshwater/land crab, have allowed us to increase our understanding of the mechanisms that have underpinned the evolution of these animals from the sea to the terrestrial environment. Recent studies have focused on understanding the neuroendocrine control of osmoregulation in terrestrial crabs on Christmas Island, with crustacean hyperglycaemic hormone being confirmed to have significant roles in the control of seasonal osmoregulation in gecarcinids.

**Key words.** Christmas Island, osmoregulation, physiology, crab, crustacean

---

### INTRODUCTION

The ability to control internal osmolarity is fundamental to the success of all crabs, and enables these crustaceans to colonise not only the full spectrum of aquatic habitats (i.e., marine, brackish and freshwater environments) (Mantel & Farmer, 1983; Pequeux, 1995; Freire et al., 2008; McNamara & Faria, 2012), but also to successfully invade the terrestrial realm (Greenaway, 1988). There is perhaps nowhere that this is more obvious than Christmas Island (CI), in the Indian Ocean. This relatively small (135 km<sup>2</sup>), remote island of volcanic origin is renowned for the diversity of the terrestrial crabs that inhabit it (Hicks et al., 1990; Gray, 1995) (separate comprehensive discussions of the landscape and natural history, fauna and flora of Christmas Island are included in this Special Supplement of the Raffles Bulletin of Zoology and the reader is directed to these for further detail). Recent surveys on CI have put the number of terrestrial crab species at 20 (including four endemics), and that of marine species at over 100 (Ng et al., 2010; Orchard, 2012), with this number likely to be revised upwards with further taxonomic study. Since the first survey of the natural history of the island, which was conducted in collaboration with the British Museum (Andrews, 1900), the island's land crabs have attracted the interest of ecophysiologicals and evolutionary biologists, with the resultant body of work contributing to many of the major advances in our understanding of land crab physiology, including in the fields of respiratory gas exchange (i.e., Greenaway et al., 1988; Adamczewska & Morris, 1998, 2000), acid base balance (i.e., Adamczewska

& Morris, 1996), nitrogenous excretion (i.e., Greenaway & Morris, 1989; Greenaway & Nakamura, 1991), exercise physiology (i.e., Adamczewska & Morris, 1994a, 1994b; Morris et al., 2010), digestion (i.e., Greenaway & Raghaven, 1998; Linton & Greenaway, 2004), olfaction (i.e., Krieger et al., 2010) and osmoregulation (i.e., Morris & Ahern, 2003; Turner et al., 2013).

Maintaining salt and water balance is problematic, and thus is a significant limiting factor for crabs when living in the terrestrial environment. The reasons for this are twofold: firstly, in aquatic crustaceans it is the gills that are used for salt transport. In terrestrial species these are no longer in contact with water, removing the ability for adjustments in ion content to be made via this method (McNamara & Faria, 2012). Secondly, the fact that these crabs are no longer submerged in an aquatic medium, but instead surrounded by air, means that maintaining water balance is also difficult due to evaporation (Greenaway, 1988). Osmoregulatory mechanisms in terrestrial species have thus been modified to cope with these changes. Land crabs have evolved through several routes (exemplified by several extant lineages) from the fully marine to terrestrial habitat, via the intertidal or through a variety of brackish and freshwater life histories (Little, 1983, 1990). Land crabs, therefore, are truly useful and interesting subjects for the study of the evolution of the physiological mechanisms, such as osmoregulation, that have allowed the terrestrial invasion to take place. CI with its abundance of species that occupy various positions along this continuum of forms, together with the phenomenal numbers of some species present on the island (for example, the population of the endemic CI red crab, *Gecarcoidea natalis*, has been estimated at 43.7 million (Adamczewska & Morris, 2001)) has facilitated the study of the ecophysiology of these unique organisms.

---

Marine Biology and Ecology Research Centre, School of Marine Science and Engineering, Davy Building, Plymouth University, Drake Circus, Plymouth, Devon PL4 8AA, United Kingdom; Email: [lucy.m.turner@plymouth.ac.uk](mailto:lucy.m.turner@plymouth.ac.uk)

© National University of Singapore  
ISSN 2345-7600 (electronic) | ISSN 0217-2445 (print)

Work conducted on the land crabs of Christmas Island has significantly contributed to our understanding of osmoregulation in terrestrial crabs (for reviews see Morris (2002) and Greenaway (2003)). The aims of this review are to summarise the current knowledge in the field of CI land crab osmoregulation and suggest directions for future research.

### ION AND WATER BALANCE IN CRUSTACEANS

There are obvious osmoregulatory challenges encountered by crabs when living in the terrestrial environment, specifically the need to avoid desiccation as well as the loss of salt from the body (Greenaway, 1988; Morris, 2001, 2002). In aquatic crabs, the gills are the primary site of salt transport, although the mechanisms for this are different when marine species are compared to those that reside in freshwater environments (Pequeux, 1995; Freire et al., 2008). Freshwater crabs are able to supplement branchial pumping with the utilisation of the antennal glands to reclaim salts from the urine. However, due to the large difference in salt concentration between the crab's tissues and the external freshwater medium, the gills of freshwater crabs must work 'harder' than those of marine species to maintain body salt concentration and prevent salt loss (Morris, 2001). At the gills, euryhaline marine crabs utilise apical membrane branchial  $\text{Na}^+/\text{H}^+$  and  $\text{Cl}^-/\text{HCO}_3^-$  exchange (Lucu, 1990) powered by a basal membrane  $\text{Na}^+/\text{K}^+$ -ATPase to maintain salt balance (Lucu & Towle, 2003; Tsai & Lin, 2007). However, this is not a stand-alone mechanism and there are fundamentally important linkages between salt balance,  $\text{CO}_2$  excretion, acid-base balance and nitrogen excretion in the gills of aquatic crustaceans (Henry, 1988; Morris, 2001; Weihrauch et al., 2004; Freire et al., 2008). In the case of  $\text{CO}_2$  excretion, this occurs as  $\text{HCO}_3^-$  through the apical  $\text{Cl}^-/\text{HCO}_3^-$  exchanger (Truchot, 1983; Henry & Wheatly, 1992; Wheatly & Henry, 1992). However, this process is facilitated by carbonic anhydrase (CA) which at the basal membrane assists the dehydration of  $\text{HCO}_3^-$  to  $\text{CO}_2$  which then diffuses into the cytoplasm where an additional CA facilitates the rehydration to  $\text{HCO}_3^-$  and  $\text{H}^+$  (Henry, 1984a, 1988). This provides the counterions for the  $\text{Cl}^-$  and  $\text{Na}^+$  exchangers involved in salt balance as well as managing acid-base balance (Fig. 1).

In decapod freshwater crustaceans however, rather different physiologies are used. Firstly, the antennal gland plays a significant role in osmoregulation. In freshwater crabs (e.g., *Potamon* spp., *Metopaulias depressus*, *Pseudothelphusa jouyi*, *Holthuisana transversa* and *Potamonautes warreni*), it has been shown that salts and water are reabsorbed at the antennal gland (Shaw, 1959; Thompson, 1970; Harris, 1975; Greenaway, 1980, 1981; Morris & Van Aardt, 1998). In addition, it has been demonstrated in the crayfish, *Pacifastacus leniusculus*, that the antennal gland utilises  $\text{Na}^+/\text{K}^+$ -ATPase and CA to reclaim urinary salts, resulting in urine which is hypo-ionic *before* it is passed into the gill chambers (Wheatly & Henry, 1987; Wheatly & Toop, 1989). In freshwater,  $\text{NaCl}$  concentration is lower than that of the gill epithelium, meaning that straightforward apical exchange of  $\text{H}^+$  and  $\text{Na}^+$  is no longer possible at the gills. Using the Chinese mitten crab, *Eriocheir sinensis*, as a model




Fig. 1. Model of gill epithelial ion exchange mechanisms typical for a marine euryhaline crab, but also the basic model for terrestrial crustaceans of marine ancestry (after Morris, 2001).




Fig. 2. Generalised model for branchial transport mechanisms in freshwater decapods (after Morris, 2001).

(Onken & Graszynski, 1989; Onken et al., 1991; Zeiske et al., 1992; Riestenpatt et al., 1994; Onken & Putzenlechner, 1995; Riestenpatt et al., 1995; Onken & Riestenpatt, 1998; Onken, 1999), it is now known that an electrogenic force (now known to be a vacuolar (V) type  $\text{H}^+$ -ATPase, or V-ATPase) is necessary to drive  $\text{Cl}^-$  transport, pumping  $\text{H}^+$  into the water independently of  $\text{Na}^+$  uptake. This V-ATPase provides for electrogenic uptake of  $\text{Cl}^-$  in exchange for  $\text{HCO}_3^-$  (Onken et al., 1991; Onken & Putzenlechner, 1995). The  $\text{HCO}_3^-$  is provided by CA facilitating  $\text{CO}_2$  excretion while  $\text{NH}_4^+$  can substitute for  $\text{K}^+$  in the basal ATPase and for  $\text{H}^+$  in the apical exchange (Henry, 1984a, 1984b) (Fig. 2). Perhaps, even more so in freshwater crabs, it is clear that the maintenance of salt balance,  $\text{CO}_2$  excretion, acid-base balance and nitrogen excretion in crustaceans relies on all of these processes.

In land crabs the gills are retained, but are reduced in number, as well as having a reduced planar gill surface area compared to those of water-breathing crabs (Farrelly & Greenaway, 1992). In these crabs the gills were originally thought to play a role in gas exchange, but it is now known that they are the organs used primarily for ion regulation and urine reprocessing rather than respiration (Greenaway,

1989; Adamczewska & Morris, 1996; DelaCruz & Morris, 1997a, 1997b; Morris & Dela-Cruz, 1998; Morris, 2005) and are thus used as a filtration-reabsorption system, 'similar to that of the vertebrate kidney' to control urinary salt loss (Fig. 3) (Morris, 2001).

It is now understood that essentially the same ion-exchange mechanisms of their aquatic ancestors are utilised in land crabs, but are employed instead to facilitate exchanges within their own urine (Morris, 2001). The gills as a site of ion uptake are retained in terrestrial crabs, which results in a very dilute final excretory product 'P' (Wolcott & Wolcott, 1985). In general terms, land crabs with a marine origin conserve water and salts by extrarenal urinary reprocessing and by decreasing urinary output (Greenaway, 1988), i.e., *Cardisoma guanhumii* (Harris, 1977; Wolcott & Wolcott, 1984; Wolcott, 1991), *Ocypode quadrata* (Wolcott & Wolcott, 1985), *Birgus latro* (Harris & Kormanik, 1981; Greenaway & Morris, 1989; Greenaway et al., 1990; Morris et al., 1991; Taylor et al., 1993; Morris et al., 2000; Greenaway, 2001), *Gecarcinus lateralis* (Harris, 1977; Wolcott, 1991; Wolcott & Wolcott, 1991), *Geograpsus grayi* (Varley & Greenaway, 1994), *Gecarcoidea lalandii* (Harris & Kormanik, 1981), *Discoplax celeste* (DelaCruz & Morris, 1997b; Dela-Cruz, 1998), *Cardisoma carnifex* (Harris & Kormanik, 1981; Dela-Cruz, 1998), and *Gecarcoidea natalis* (Greenaway, 1994; Morris, 2001; Taylor & Greenaway, 2002; Greenaway, 2003; Morris & Ahern, 2003). These species can redirect their primary urine from the antennal gland to the gill chambers where salt reabsorption takes place at the gills producing a final hypo-ionic excretory product 'P' (Wolcott & Wolcott, 1985). This is very dilute, containing less than 30% of the ions measured in the primary urine and less than 5% of the ions measured in the haemolymph (Morris, 2001). In addition, some terrestrial crabs also utilise the ion exchange mechanisms of the antennal gland to reclaim urinary salts (DeVries & Wolcott, 1993; DeVries et al., 1994), a process which is similar to that employed by freshwater crustaceans.

#### **DISCOPLAX CELESTE: THE PERFECT CRUSTACEAN MODEL TO STUDY OSMOREGULATION?**

Previously misidentified on CI as *Discoplax hirtipes* (Dana, 1851) (and before that as *Cardisoma hirtipes* – for a full discussion see Ng & Davie, 2012), this crab has been recognised as a distinct new species endemic to CI, the Christmas Island blue crab, *Discoplax celeste* (Ng & Davie, 2012). Unlike the other two most obvious and abundant land crab species on CI, *G. natalis* and *B. latro*, *D. celeste* is restricted in its distribution (especially during the dry season (April–October)) to areas of freshwater seepages and springs, usually well inland (>1 km from and 150 m above the ocean) (DelaCruz & Morris, 1997a, 1997b; Morris, 2005). Due to the underlying geology (Barrett, 2001), these areas of freshwater and thus established populations of *D. celeste* are found at only a few places on the island. When the wet season begins (November–March) *D. celeste* disperse over a wider area (DelaCruz & Morris, 1997a, 1997b), and only when there




Fig. 3. Two part model: A, the antennal gland and B, the branchial epithelial layer (after Morris, 2001).

is an adequate supply of standing water (February–March) do they undertake their breeding migration to the ocean.

We now know that *D. celeste* is an ammoniotelic species and it is the need for long term nitrogen, ion and water balance that drives this crab into association with freshwater, therefore explaining the local and seasonal variations in the distribution of this species (Greenaway, 1989; Adamczewska & Morris, 1996; DelaCruz & Morris, 1997b). When this crab submerges in freshwater the gills are exposed to an enormous dilution gradient (~400 times lower than that of the crab's tissues) yet *D. celeste* is still able to maintain salt balance in solutions containing concentrations as low as 0.12  $\text{mmol.L}^{-1}$   $\text{Na}^+$  (Greenaway, 1989). This is due to the fact that although *D. celeste* has a high  $\text{Na}^+$  permeability (1.609  $\mu\text{mol.g}^{-1}.\text{h}^{-1}$ ), this is compensated by a high  $\text{Na}^+$  affinity uptake mechanism ( $K_m = 0.264 \text{ mmol.L}^{-1}$ ) (Greenaway, 1989), and also that the stored nitrogen is excreted in rapid, intense bursts during brief immersion episodes (DelaCruz & Morris, 1997b). However, it was noted by the same authors that although *D. celeste* manages  $\text{Na}^+$ ,  $\text{Cl}^-$  and  $\text{K}^+$  in the same way as other species of gecarcinid,  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  are conserved using similar methods to those of freshwater crabs, i.e., that they are reabsorbed in the antennal gland/bladder structure. The importance of behaviour in the osmoregulatory ability of this crab was also highlighted—the salts lost by *D. celeste* in their urine are passed into the burrow water which remains as a large reservoir of ions that potentially the crab can later draw on (Greenaway, 1989).

Urine flow rates in *D. celeste* have been found to be three times higher than those of other gecarcinid species (but still less than marine crabs), as well as other species of freshwater crustaceans (Greenaway, 1989). These high urine flow rates mean that the amount of salt lost in the urine is much greater compared to these other species. A high production rate of concentrated urine is very energetically expensive to produce. Therefore, although *D. celeste* is able to successfully osmoregulate in a freshwater environment, it appears to be doing so at the very limits of its sodium regulatory system (Greenaway, 1989). In addition, this crab is also able to maintain salt balance when supplied with only freshwater or seawater to drink (Greenaway, 1989). These factors indicate that this species occupies a unique position in evolutionary terms and can be considered an intermediary between marine and freshwater species (Greenaway, 1989). *Discoplax celeste* therefore makes an excellent model species for studies of the evolution of osmoregulation and ultimately the evolution of life on land (Little, 1983, 1990; Morris, 2001). See section on 'Hormonal control of osmoregulation' for further details of the control of osmoregulation in this species.

### ***BIRGUS LATRO***

The Robber Crab or Coconut Crab, *B. latro*, is the largest land arthropod, weighing up to 4 kg (Drew et al., 2010). Currently listed as data deficient on the International Union for Conservation of Nature Red List, this once widely distributed crab was previously common throughout the islands of the Indo-Pacific as well as on large land masses including Madagascar and the tropical north-east coast of Australia. *Birgus latro* is now only present on remote islands and rapidly declining throughout its range primarily due to overharvesting for human consumption and as a result of habitat loss (Drew et al., 2010). However, a significant population is still present on CI and this species is found all over the island (Hicks et al., 1990; Orchard, 2012). This crab is classed as a true terrestrial crab and does not require periodic immersion in fresh or saltwater for the purposes of osmoregulation (Hartnoll, 1988; Greenaway, 2003).

When compared to gecarcinids, in *B. latro* it is the gills in the branchial chambers that play the greater role in ion and water balance rather than the antennal gland, with water reabsorption also occurring in the gut (Greenaway et al., 1990; Morris et al., 1991). Ion uptake at the gills is driven by ATPases (Morris et al., 1991). It is apparent that the ion-transport system in the gills of *B. latro* is reminiscent of those of marine species and thus demonstrates the evolutionary route that this species has taken. *Birgus latro* is also able to select drinking water of a suitable salinity to regulate body osmolarity although it is the physiological up or down regulation of ion uptake from the urine, instead of any kind of behavioural mechanism, by which this crab ultimately regulates its ion uptake (Taylor et al., 1993). Field studies of free-ranging *B. latro* on CI have also demonstrated this species' ability to be a competent osmoregulator under natural conditions and have confirmed that the drinking of saline water is not necessary for maintaining salt and water balance (Greenaway, 2001).

In *B. latro*, dopamine and cAMP both decrease branchial  $\text{Na}^+/\text{K}^+$ -ATPase activity as well as ion ( $\text{Na}^+$  and  $\text{Cl}^-$ ) uptake (Morris et al., 2000). However, this inhibition of ion uptake in *B. latro* is in contrast to what is seen in similar pharmacology-based experiments in aquatic brachyurans (for a recent review see McNamara & Faria (2012)). However, this mechanistic control makes sense in the context of the ecology of these crabs, which despite their ability to cope in very dry conditions will at no time of the year be completely restricted from the permanent freshwater seepages on CI (except by geographical distance) from which they can take up ions. This coupled with the fact that these crabs are able to reabsorb ions from their own urine via the gills explains the evolution of a mechanism to down-regulate ion uptake in this way (Morris et al., 2000).

### ***GECARCOIDEA NATALIS***

The Christmas Island red crab, *G. natalis*, is endemic to CI and is widely distributed throughout the island (Hicks et al., 1990; Adamczewska & Morris, 2001; Orchard, 2012). *Gecarcoidea natalis* is also classified as a true terrestrial crab as this species is independent of water (fresh or saline) for osmoregulatory purposes (Morris, 2001). This species participates in a spectacular annual breeding migration to the ocean, and the osmoregulatory system has been shown to be able to cope successfully with the energetic demands of exercise that this entails (Greenaway, 1994). Upon arrival at the shoreline these crabs are observed to 'dip' into the ocean before the commencement of breeding activities, thus replenishing lost salts (Hicks, 1985; Greenaway, 1994).

In terms of control of osmoregulation via biogenic amines, in this species, in contrast to what has been observed in *B. latro*, dopamine appears to stimulate the up-regulation of branchial chloride transport and increases the amount of urine produced (Taylor & Greenaway, 2002). Serotonin also stimulates  $\text{Na}^+$  uptake at the gills, but the exact mechanism for this varies depending on the salinity of the available drinking water. When crabs are given freshwater to drink, serotonin stimulates  $\text{Na}^+$  uptake via increased gill  $\text{Na}^+/\text{K}^+$ -ATPase activity, whereas dopamine has no effect. Serotonin stimulation also appears to be independent of cAMP and therefore is in contrast to what is observed in marine species where the control of branchial ion uptake is normally via a dopamine/cAMP pathway. In comparison, when *G. natalis* are given 50% seawater to drink, serotonin and dopamine appear to control ion uptake via the control of branchial leak permeability (the rate of ion pumping, as well as gill  $\text{Na}^+/\text{K}^+$ -ATPase activity, remains unchanged) (Morris & Ahern, 2003). Furthermore, serotonin appears to increase urine production in the dry season in *G. natalis* (when urine production rates were lowered), whereas dopamine had no effect (Morris & Ahern, 2003). Thus it appears that the specific effects of monoamines on osmoregulation in *G. natalis*, as well as the overall control of osmoregulation in land crabs, is far more complex than previously thought and has led directly to the work carried out in recent years to more fully understand the neuroendocrine control of osmoregulation in terrestrial crabs (see next section 'Hormonal control of osmoregulation').

## HORMONAL CONTROL OF OSMOREGULATION

Crustacean hyperglycaemic hormone (CHH) has recently been shown to have osmoregulatory effects in CI land crabs, including on a seasonal basis (Turner, 2010; Turner et al., 2013). Hormones in the CHH family are known to have pivotal roles in various physiological processes, including the control of energy metabolism, moulting and reproduction (see Webster et al. (2012) for a recent review) and there is a growing body of evidence implicating involvement of CHH in ionic homeostasis in decapod crustaceans (Charmantier-Daures et al., 1994; Charmantier et al., 1999; Chung et al., 1999; Townsend et al., 2001; Serrano et al., 2003; Chung & Webster, 2006). Therefore CHH seemed a prime candidate for investigating the hormonal control of osmoregulatory process in these crabs.

Crustacean hyperglycaemic hormone is able to adjust gill  $\text{Na}^+$  uptake in a seasonally dependent manner in gecarcinids (Turner et al., 2013). In *G. natalis* there is a clear downregulation of  $\text{Na}^+$  uptake in the dry season and for both *D. celeste* and *G. natalis* an upregulation of  $\text{Na}^+$  uptake in the wet season following injection with CHH. Therefore, it seems that CHH is important in driving the response to high water throughput in the wet season. It is imperative that land crabs do not become depleted of salts, something that can easily happen if urine and 'P' production rates are high at this time of year. The similar observations recorded for these closely related land crab species suggest a universal ion uptake mechanistic effect of CHH amongst terrestrial crabs. CHH was also shown to increase urine production at the antennal gland on a seasonally specific basis in *D. celeste*, but had no effect on urine production in *G. natalis*, perhaps as a result of the contrasting life histories of these two species: *D. celeste* spends much of its life in association with freshwater, and therefore faces rapid water influx and salt loss compared to *G. natalis* which is much more independent of water and terrestrial throughout its adult life history (Turner et al., 2013).

Phylogenetic analyses of CHH suggests that this hormone may have had a fundamental role in the invasion of land by decapod crustaceans (Table 1, Fig. 4). When compared with the phylogeny produced from the analysis of moult inhibiting hormone (MIH) (Fig. 5.), which is also in the CHH-superfamily (Webster et al., 2012), it is clear that while CHH has diverged several times during the evolution of the Decapoda, MIH has remained highly conserved (Fig. 5). Thus it appears that different selection pressures may have shaped the evolution of CHH and MIH. It is well recognised that CHH sensu stricto exhibits many pleiotropic functions (Webster et al., 2012), whereas the physiological function of MIH is constrained to that of involvement with the control of the timing of ecdysis (Nakatsuji et al., 2009). Recent molecular phylogenetic evidence has confirmed the position of ITP at the base of the CHH family in terms of ecdysozoan evolution (Montagne et al., 2010) and this finding appears to support the hypothesis that ionic and water regulation could be the ancestral function of CHH/ITP family peptides (Toullec et al., 2006). Indeed, the present extant land crab

fauna have evolved through a number of routes; moving to a terrestrial habitat directly from the ocean, through the intertidal, or through a variety of brackish and freshwater habitats (Greenaway, 1988). Consequently, in line with their evolutionary route into a terrestrial existence, land crabs have evolved a number of mechanisms to cope with the physiological demands of life on land, including those to deal with salt transport and ion regulation. Therefore it is possible that the divergent evolution of CHH as well as the conserved evolution of MIH reflect this.

Despite work demonstrating the involvement of CHH in osmoregulation in land crabs, as yet the exact osmoregulatory mode of action of CHH in these crabs has not been identified: intriguingly this hormone appeared to have no effect on gill  $\text{Na}^+/\text{K}^+$ -ATPase or V-ATPase activity, suggesting unknown mechanisms for this hormone's action on  $\text{Na}^+$  transport across gill epithelia in terrestrial crabs (Turner et al., 2013). Nevertheless, these results clearly suggest that much more work is required to examine the hormonal controls on osmoregulation in land crabs, with one obvious avenue for future research being to compare and contrast the situation in the anomurans with that in the brachyurans.

## CONCLUSIONS

Work conducted on the land crabs of CI has undoubtedly advanced our knowledge of the ecophysiology of ion and water balance not only in terrestrial crustaceans, but in crustaceans as a whole. For instance, studies undertaken on CI species, such as *D. celeste* with its unique evolutionary position and ecology have allowed us to increase our understanding of the mechanisms that have underpinned the evolution of these animals, including the routes they have taken to enable them to successfully exploit the terrestrial landscape (Little, 1990).

It is imperative that we continue to learn all we can about these unique animals, especially as the number of described and/or endemic CI species continues to rise (Orchard, 2012). It appears likely that when these kinds of studies are extended to investigate the patterns and processes of osmoregulation in additional (endemic) species of CI land crab, our knowledge and understanding of crustacean osmoregulation will be further enhanced. With the application of new molecular and analytical techniques we have the tools available that will enable us to learn much more about the molecular mechanisms that underpin the ecophysiology and ultimately drives the ecology of these crabs. However, work must continue with some urgency due to the unfortunate fact that many of these species are at risk from the challenges imposed on the CI ecosystem by anthropogenic influences, for example, habitat destruction (Morris-Pocock et al., 2012), increased numbers of invasive species (Thomas et al., 2010; Green et al., 2011), as well as global climate change (Marshall & McCulloch, 2001). It is a worrying thought that it appears possible that further species of land crab could potentially be lost forever before their biology has been studied by a new generation of carcinologists. The prevention of this scenario must continue to be taken extremely seriously and




Fig. 4. Phylogeny of crustacean hyperglycaemic hormones (CHH) based on maximum-likelihood analysis of the CHH amino acid data set (39 taxa, 71 sequences, 78 residues (Table 1)) using a JTT+I+G model of protein evolution (tree shown is 50% majority rule consensus tree). Maximum-likelihood tree reconstruction was performed using PhyML (Abascal et al., 2005). The best-fit model for protein evolution was determined using ProtTest 1.3 (Jones et al., 1992; Guindon & Gascuel, 2003). The locust *S. gregaria* ITP was assigned as the outgroup. Sequences were acquired from GenBank or directly sequenced (Turner, 2010). See Turner (2010) for methodologies used. Numbers at nodes are bootstrap values (100 replicates). Only bootstrap values  $\geq 70$  are shown.




Fig. 5. Phylogeny of crustacean moult inhibiting hormones (MIH) based on maximum-likelihood analysis of the MIH amino acid data set (26 taxa, 34 sequences, 114 residues) using a JTT+I+G model of protein evolution (tree shown is 50% majority rule consensus tree). Maximum-likelihood tree reconstruction was performed using PhyML (Abascal et al., 2005). The best-fit model for protein evolution was determined using ProtTest 1.3 (Jones et al., 1992; Guindon & Gascuel, 2003). The locust *S. gregaria* ITP was assigned as the outgroup. Sequences were acquired from GenBank or directly sequenced (Turner, 2010). See Turner (2010) for methodologies used. Numbers at nodes are bootstrap values (100 replicates). Only bootstrap values  $\geq 70$  are shown.

efforts must continue to conserve and protect not only the crabs, but the whole CI ecosystem for future generations.

#### ACKNOWLEDGEMENTS

This paper is dedicated to the memory of Steve Morris (1956–2009) in recognition and appreciation of his 25 years of research on the physiology of the land crabs of Christmas Island. The Government conservators and staff of Parks Australia, Christmas Island are thanked for their logistical assistance, hospitality and enthusiasm over the years. Thanks also go to the many volunteer field assistants who have helped out, as well as other research groups, as without

their hard work and dedication much of this work would not have been possible. Thanks go to the two reviewers who provided constructive and helpful suggestions that improved this manuscript.

#### LITERATURE CITED

- Abascal F, Zardoya R & Posada D (2005) ProtTest: selection of best-fit models of protein evolution. *Bioinformatics*, 21: 2104–2105.
- Adamczewska AM & Morris S (1994a) Exercise in the terrestrial Christmas Island red crab *Gecarcoidea natalis* .1. Blood-gas transport. *Journal of Experimental Biology*, 188: 235–256.
- Adamczewska AM & Morris S (1994b) Exercise in the terrestrial Christmas Island red crab *Gecarcoidea natalis* .2. Energetics of locomotion. *Journal of Experimental Biology*, 188: 257–274.

Table 1. Amino acid sequences used in this study.

Species	Sequence Identifier	GenBank Accession Number
<b>Arthropod: Crustacean</b>		
<i>Astacus astacus</i>	Asa-CHH	P83800
<i>Bythograea thermydron</i>	Byt-CHH	AF241264
<i>Callinectes sapidus</i>	Cas-CHH-I	Q6QJL8
	Cas-CHH-II	ACH85179
	Cas-MIH	P55321
<i>Chionoecetes bairdi</i>	Chb-CHH	O61389
<i>Cancer magister</i>	Crn-MIH	ACG50068
<i>Cancer pagurus</i>	Cap-CHH	P81032
	Cap-MIH	CAC39425
<i>Cancer productus</i>	Crp-CHH-I	ABQ41269
	Crp-CHH-II-A	ABQ41270
	Crp-CHH-II-B	ABQ41271
	Crp-CHH-III	ABQ41272
<i>Carcinus maenas</i>	Cam-CHH	P14944
	Cam-MIH	Q27225
<i>Charybdis feriatius</i>	Chf-MIH	O96605
<i>Charybdis japonica</i>	Chj-MIH	ACD11361
<i>Cherax destructor</i>	Chd-CHH-A	P83485
	Chd-CHH-B	P83486
<i>Cherax quadricarinatus</i>	Chq-CHH	AAZ03612
<i>Discoplax celeste</i>	Dic-CHH	AEM45614
	Dic-MIH	AEM45616
<i>Eriocheir sinensis</i>	Ers-CHH	ABA42603
	Ers-MIH	ABC68517
<i>Fenneropenaeus chinensis</i>	Fec-MIH	AAL55258
<i>Galathea strigosa</i>	Gas-CHH	ABS01332
<i>Gecarcinus lateralis</i>	Gel-CHH-A	ABF48652
	Gel-CHH-B	ABF58090
	Gel-MIH	DQ473354
<i>Gecarcoidea natalis</i>	Gen-CHH	ABL09570
	Gen-MIH	(Webster et al., unpublished)
<i>Grapsus tenuicrustatus</i>	Grt-CHH	AER27832
<i>Homarus americanus</i>	Hoa-CHH-A	CAA38611
	Hoa-CHH-B	CAA38612
	Hoa-CHH-I	2105187A
	Hoa-CHH-II	2105187B
	Hoa-CHH	Q25154
	Hog-CHH-A	Q3HXZ6
<i>Homarus gammarus</i>	Hog-CHH-B	Q3HXZ5
	Jal-CHH	P56687
<i>Jasus lalandii</i>	Jal-MIH	P83220
	Lss-CHH	P59685
<i>Litopenaeus schmitti</i>	Lis-MIH	AF312976
<i>Litopenaeus stylirostris</i>	Liv-CHH	CAA68067
	Liv-MIH-I	AAR04348
	Liv-MIH-II	AAR04349
<i>Macrobrachium lanchesteri</i>	Mal-CHH	AAC36310
<i>Macrobrachium rosenbergii</i>	Mar-CHH	AAL40915
	Mar-SGP-A	AAL37948
	Mar-SGP-B	AAL37949
<i>Marsupenaeus japonicus</i>	Maj-CHH-I	O15980
	Maj-CHH-II	Q9U5D2
	Maj-CHH-III	BAA13481
	Maj-CHH-IV	P55847



Species	Sequence Identifier	GenBank Accession Number
	Maj-CHH-V	O15981
	Maj-CHH-VI	P81700
	Maj-CHH-VII	O15982
	Maj-MIH-A	P55847
	Maj-MIH-B	BAD36757
	Maj-MIH-C	BAE78494
<i>Metapenaeus ensis</i>	Mee-CHH-A	AAD45233
	Mee-CHH-B	AAF63028
	Mee-MIH-A	AAC27452
	Mee-MIH-B	AAL33882
<i>Nephrops norvegicus</i>	Nen-CHH-A	Q6WGR4
	Nen-CHH-B	AAQ22392
<i>Orconectes limosus</i>	Orl-CHH	CAA56674
	Orl-CHH-A	1802398A
	Orl-MIH	P83636
<i>Pachygrapsus marmoratus</i>	Pam-CHH-A	Q6Y5A6
	Pam-CHH-B	AAO27805
	Pam-CHH	AAM21927
<i>Pagurus bernhardus</i>	Pab-CHH	ABE02191
<i>Penaeus monodon</i>	Pem-CHH	AAC84143
	Pem-CHH-I	O97383
	Pem-CHH-II	O97384
	Pem-CHH-III	O97385
	Pem-CHH-IV	O97386
	Pem-CHH-V	O97387
	Pem-CHH-A	AAD03606
	Pem-CHH-B	AAD03607
	Pem-MIH-I	AAR89516
	Pem-MIH-II	AAR89517
	Pem-SGP-I	BAB69829
	Pem-SGP-II	BAB69830
<i>Pontastacus leptodactylus</i>	Pol-CHH	Q1RN83
	Pol-CHH-I	AAS45406
<i>Portunus pelagicus</i>	Pop-CHH	ABM74398
	Pop-MIH	ABM74397
<i>Portunus trituberculatus</i>	Pot-CHH	ACB46189
	Pot-MIH	ABZ04547
<i>Potamon ibericum</i>	Poi-CHH	Q2VF26
<i>Procambarus bouvieri</i>	Prb-CHH	P55845
	Prb-CHH-I	AAB2553
	Prb-MIH	Q10987
<i>Procambarus clarkii</i>	Prc-CHH	BAA89003
	Prc-MIH	P55848
<i>Ptychognathus pusillus</i>	Ptp-CHH	AER27833
<i>Rimicaris kairei</i>	Rik-CHH-I	ACS35346
	Rik-CHH-II	ACS35347
<i>Scylla olivacea</i>	Sco-CHH	AAQ75760
<i>Scylla serrata</i>	Scs-MIH	AY083797
<i>Trachypenaeus curvirostris</i>	Trc-MIH	AF312978
<b>Outgroup</b>		
<b>Arthropod: Insect</b>		
<i>Schistocerca gregaria</i>	Scg-ITP	Q26491

- Adamczewska AM & Morris S (1996) The respiratory gas transport, acid-base state, ion and metabolite status of the Christmas Island blue crab, *Cardisoma hirtipes* (Dana) assessed in situ with respect to immersion. *Physiological Zoology*, 69: 67–92.
- Adamczewska AM & Morris S (1998) The functioning of the haemocyanin of the terrestrial Christmas Island red crab *Gecarcoidea natalis* and roles for organic modulators. *Journal of Experimental Biology*, 201: 3233–3244.
- Adamczewska AM & Morris S (2000) Respiratory gas transport, metabolic status, and locomotor capacity of the Christmas Island red crab *Gecarcoidea natalis* assessed in the field with respect to dichotomous seasonal activity levels. *Journal of Experimental Zoology*, 286: 552–562.
- Adamczewska AM & Morris S (2001) Ecology and behavior of *Gecarcoidea natalis*, the Christmas Island red crab, during the annual breeding migration. *Biological Bulletin*, 200: 305–320.
- Andrews C (1900) A Monograph of Christmas Island (Indian Ocean). British Museum (Natural History), Department of Geology, London.
- Barrett PJ (2001) Searching for water on Christmas Island. *Helictite*, 37: 37–39.
- Charmantier-Daures M, Charmantier G, Janssen KPC, Aiken DE & Vanherp F (1994) Involvement of eyestalk factors in the neuroendocrine control of osmoregulation in adult American lobster *Homarus americanus*. *General and Comparative Endocrinology*, 94: 281–293.
- Charmantier G, Spanings-Pierrot C, Soyex D & Van Herp F (1999) Involvement of CHH in the control of osmoregulation in a grapsid crab. *American Zoologist*, 39: 68A–68A.
- Chung JS, Dirksen H & Webster SG (1999) A remarkable, precisely timed release of hyperglycemic hormone from endocrine cells in the gut is associated with ecdysis in the crab *Carcinus maenas*. *Proceedings of the National Academy of Sciences of the United States of America*, 96: 13103–13107.
- Chung JS & Webster SG (2006) Binding sites of Crustacean Hyperglycemic Hormone and its second messengers on gills and hindgut of the green shore crab, *Carcinus maenas*: a possible osmoregulatory role. *General and Comparative Endocrinology*, 147: 206–213.
- Dela-Cruz J (1998) Ion and Water Regulation in *Cardisoma* (Brachyura, Gecarcinidae). MSc Thesis, University of Sydney.
- DelaCruz J & Morris S (1997a) Respiratory, acid-base, and metabolic responses of the Christmas Island blue crab, *Cardisoma hirtipes* (Dana), during simulated environmental conditions. *Physiological Zoology*, 70: 100–115.
- DelaCruz J & Morris S (1997b) Water and ion balance and nitrogenous excretion as limitations to terrestrial excursion in the Christmas Island Blue Crab, *Cardisoma hirtipes* (Dana). *Journal of Experimental Zoology*, 279: 537–548.
- DeVries MC & Wolcott DL (1993) Gaseous ammonia evolution is coupled to reprocessing of urine at the gills of ghost crabs. *Journal of Experimental Zoology*, 267: 97–103.
- DeVries MC, Wolcott DL & Holliday CW (1994) High ammonia and low pH in the urine of the ghost crab, *Ocypode quadrata*. *Biological Bulletin*, 186: 342–348.
- Drew MM, Harzsch S, Stensmyr M, Erland S & Hansson BS (2010) A review of the biology and ecology of the Robber Crab, *Birgus latro* (Linnaeus, 1767) (Anomura: Coenobitidae). *Zoologischer Anzeiger – A Journal of Comparative Zoology*, 249: 45–67.
- Farrelly CA & Greenaway P (1992) Morphology and ultrastructure of the gills of terrestrial crabs (Crustacea, Gecarcinidae and Grapsidae): adaptations for air-breathing. *Zoomorphology*, 112: 39–49.
- Freire CA, Onken H & McNamara JC (2008) A structure-function analysis of ion transport in crustacean gills and excretory organs. *Comparative Biochemistry and Physiology A-Molecular and Integrative Physiology*, 151: 272–304.
- Gray HS (1995) Christmas Island Naturally: The Natural History of an Isolated Oceanic Island the Australian Territory of Christmas Island Indian Ocean. Christmas Island Natural History Association, Christmas Island, Indian Ocean.
- Green PT, O'Dowd DJ, Abbott KL, Jeffery M, Retallick K & MacNally R (2011) Invasional meltdown: invader-invader mutualism facilitates a secondary invasion. *Ecology*, 92: 1758–1768.
- Greenaway P (1980) Water-balance and urine production in the Australian arid-zone crab *Holthuisana transversa*. *Journal of Experimental Biology*, 87: 237–246.
- Greenaway P (1981) The fate of glomerular-filtration markers injected into the hemolymph of the amphibious crab *Holthuisana transversa*. *Journal of Experimental Biology*, 91: 339–347.
- Greenaway P (1988) Ion and water balance. In: Burggren WW & McMahon BR (eds.) *Biology of the Land Crabs*. Cambridge University Press, Cambridge.
- Greenaway P (1989) Sodium-balance and adaptation to fresh-water in the amphibious crab *Cardisoma hirtipes*. *Physiological Zoology*, 62: 639–653.
- Greenaway P (1994) Salt-balance and water-balance in-field populations of the terrestrial crab *Gecarcoidea natalis*. *Journal of Crustacean Biology*, 14: 438–453.
- Greenaway P (2001) Sodium and water balance in free-ranging robber crabs, *Birgus latro* (Anomura: Coenobitidae). *Journal of Crustacean Biology*, 21: 317–327.
- Greenaway P (2003) Terrestrial adaptations in the Anomura (Crustacea: Decapoda). *Memoirs of Museum Victoria*, 60: 13–26.
- Greenaway P & Morris S (1989) Adaptations to a terrestrial existence by the robber crab, *Birgus latro* L. 3. Nitrogenous excretion. *Journal of Experimental Biology*, 143: 333–346.
- Greenaway P, Morris S & McMahon BR (1988) Adaptations to a terrestrial existence by the robber crab *Birgus latro*. 2. *In vivo* respiratory gas-exchange and transport. *Journal of Experimental Biology*, 140: 493–509.
- Greenaway P & Nakamura T (1991) Nitrogenous excretion in 2 terrestrial crabs (*Gecarcoidea natalis* and *Geograpsus grayi*). *Physiological Zoology*, 64: 767–786.
- Greenaway P & Raghaven S (1998) Digestive strategies in two species of leaf-eating land crabs (Brachyura: Gecarcinidae) in a rain forest. *Physiological Zoology*, 71: 36–44.
- Greenaway P, Taylor HH & Morris S (1990) Adaptations to a terrestrial existence by the robber crab *Birgus latro*. 6. The role of the excretory system in fluid balance. *Journal of Experimental Biology*, 152: 505–519.
- Guindon S & Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, 52: 696–704.
- Harris RR (1975) Urine production-rate and urinary sodium loss in freshwater crab *Potamon edulis*. *Journal of Comparative Physiology*, 96: 143–153.
- Harris RR (1977) Urine production-rate and water-balance in terrestrial crabs *Gecarcinus lateralis* and *Cardisoma guanhumi*. *Journal of Experimental Biology*, 68: 57–67.
- Harris RR & Kormanik GA (1981) Salt and water-balance and antennal gland-function in three Pacific species of terrestrial crab (*Gecarcoidea lalandii*, *Cardisoma carnifex*, *Birgus latro*). 2. The effects of desiccation. *Journal of Experimental Zoology*, 218: 107–116.
- Hartnoll RG (1988) Evolution, systematics and geographical distribution. In: Burggren WW & McMahon BR (eds.) *Biology of the Land Crabs*. Cambridge University Press, Cambridge, pp. 6–54.
- Henry RP (1984a) The function of invertebrate carbonic-anhydrase in ion-transport. *Annals of the New York Academy of Sciences*, 429: 544–546.

- Henry RP (1984b) The role of carbonic-anhydrase in blood ion and acid-base regulation. *American Zoologist*, 24: 241–251.
- Henry RP (1988) Multiple functions of carbonic-anhydrase in the crustacean gill. *Journal of Experimental Zoology*, 248: 19–24.
- Henry RP & Wheatly MG (1992) Interaction of respiration, ion regulation, and acid-base-balance in the everyday life of aquatic crustaceans. *American Zoologist*, 32: 407–416.
- Hicks J, Rumpff H & Yorkston H (1990) Christmas Crabs. Christmas Island Natural History Association, Christmas Island, Indian Ocean.
- Hicks JW (1985) The breeding behaviour and migrations of the terrestrial crab *Gecarcoidea natalis* (Decapoda: Brachyura). *Australian Journal of Zoology*, 33: 127–142.
- Jones DT, Taylor WR & Thornton JM (1992) The rapid generation of mutation data matrices from protein sequences. *Computer Applications in the Biosciences*, 8: 275–282.
- Krieger J, Sandeman RE, Sandeman DC, Hansson BS & Harzsch S (2010) Brain architecture of the largest living land arthropod, the Giant Robber Crab *Birgus latro* (Crustacea, Anomura, Coenobitidae): evidence for a prominent central olfactory pathway? *Frontiers in Zoology*, 7: 25.
- Linton SM & Greenaway P (2004) Presence and properties of cellulase and hemicellulase enzymes of the gecarcinid land crabs *Gecarcoidea natalis* and *Discoplax hirtipes*. *Journal of Experimental Biology*, 207: 4095–4104.
- Little C (1983) *The Colonisation of Land*. Cambridge University Press, Cambridge.
- Little C (1990) *The Terrestrial Invasion: An Ecophysiological Approach to the Origins of Land Animals*. Cambridge University Press, Cambridge.
- Lucu C (1990) Ionic regulatory mechanisms in crustacean gill epithelia. *Comparative Biochemistry and Physiology A-Physiology*, 97: 297–306.
- Lucu C & Towle DW (2003) Na<sup>+</sup>/K<sup>+</sup>-ATPase in gills of aquatic crustacea. *Comparative Biochemistry and Physiology A-Molecular and Integrative Physiology*, 135: 195–214.
- Mantel LH & Farmer LL (1983) Osmotic and ionic regulation. In: Mantel LH (ed.) *The Biology of Crustacea*. Academic Press, New York, pp. 53–161.
- Marshall JF & McCulloch MT (2001) Evidence of El Nino and the Indian Ocean Dipole from Sr/Ca derived SSTs for modern corals at Christmas Island, Eastern Indian Ocean. *Geophysical Research Letters*, 28: 3453–3456.
- McNamara JC & Faria SC (2012) Evolution of osmoregulatory patterns and gill ion transport mechanisms in the decapod Crustacea: a review. *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology*, 182: 997–1014.
- Montagne N, Desdevises Y, Soyeux D & Toullec JY (2010) Molecular evolution of the Crustacean Hyperglycemic Hormone family in ecdysozoans. *BMC Evolutionary Biology*, 10: 62.
- Morris-Pocock JA, Hennicke JC & Friesen VL (2012) Effects of long-term isolation on genetic variation and within-island population genetic structure in Christmas Island (Indian Ocean) seabirds. *Conservation Genetics*, 13: 1469–1481.
- Morris S (2001) Neuroendocrine regulation of osmoregulation and the evolution of air-breathing in decapod crustaceans. *Journal of Experimental Biology*, 204: 979–989.
- Morris S (2002) The ecophysiology of air-breathing in crabs with special reference to *Gecarcoidea natalis*. *Comparative Biochemistry and Physiology B-Biochemistry and Molecular Biology*, 131: 559–570.
- Morris S (2005) Respiratory and acid-base responses during migration and to exercise by the terrestrial crab *Discoplax (Cardisoma) hirtipes*, with regard to season, humidity and behaviour. *Journal of Experimental Biology*, 208: 4333–4343.
- Morris S & Ahern MD (2003) Regulation of urine reprocessing in the maintenance of sodium and water balance in the terrestrial Christmas Island red crab *Gecarcoidea natalis* investigated under field conditions. *Journal of Experimental Biology*, 206: 2869–2881.
- Morris S & Dela-Cruz J (1998) The ecophysiological significance of lung-air retention during submersion by the air-breathing crabs *Cardisoma carnifex* and *Cardisoma hirtipes*. *Experimental Biology Online*, 3: 1–19.
- Morris S, Greenaway P, Adamczewska AM & Ahern MD (2000) Adaptations to a terrestrial existence in the robber crab *Birgus latro* L. IX. Hormonal control of post-renal urine reprocessing and salt balance in the branchial chamber. *Journal of Experimental Biology*, 203: 389–396.
- Morris S, Postel U, Mrinalini, Turner LM, Palmer J & Webster SG (2010) The adaptive significance of crustacean hyperglycaemic hormone (CHH) in daily and seasonal migratory activities of the Christmas Island red crab *Gecarcoidea natalis*. *Journal of Experimental Biology*, 213: 3062–3073.
- Morris S, Taylor HH & Greenaway P (1991) Adaptations to a terrestrial existence by the robber crab *Birgus latro*. 7. The branchial chamber and its role in urine reprocessing. *Journal of Experimental Biology*, 161: 315–331.
- Morris S & WJ Van Aardt (1998) Salt and water relations, and nitrogen excretion, in the amphibious African freshwater crab *Potamonautes warreni* in water and in air. *Journal of Experimental Biology*, 201: 883–893.
- Nakatsuji T, Lee CY & Watson RD (2009) Crustacean Molt-Inhibiting Hormone: structure, function, and cellular mode of action. *Comparative Biochemistry and Physiology, Part A*, 152: 139–148.
- Ng PKL & Davie PJF (2012) The blue crab of Christmas Island, *Discoplax celeste*, new species (Crustacea: Decapoda: Brachyura: Gecarcinidae). *Raffles Bulletin of Zoology*, 60: 89–100.
- Ng PKL, Davie P, Tan SH, Tan K-x & Tan HH (2010) Report on Christmas Island Research Permit AU-COM 2009059: Decapod crustaceans and inshore fishes of Christmas Island, Indian Ocean Territory, Australia. Raffles Museum of Biodiversity Research, National University of Singapore, Singapore.
- Onken H (1999) Active NaCl absorption across split lamellae of posterior gills of Chinese crabs (*Eriocheir sinensis*) adapted to different salinities. *Comparative Biochemistry and Physiology A-Molecular and Integrative Physiology*, 123: 377–384.
- Onken H & Graszynski K (1989) Active Cl<sup>-</sup> absorption by the Chinese crab (*Eriocheir sinensis*) gill epithelium measured by trans-epithelial potential difference. *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology*, 159: 21–28.
- Onken H, Graszynski K & Zeiske W (1991) Na<sup>+</sup>-independent, electrogenic Cl<sup>-</sup> uptake across the posterior gills of the Chinese crab (*Eriocheir sinensis*) – voltage-clamp and microelectrode studies. *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology*, 161: 293–301.
- Onken H & Putzenlechner M (1995) A V-ATPase drives active, electrogenic and Na<sup>+</sup>-independent Cl<sup>-</sup> absorption across the gills of *Eriocheir sinensis*. *Journal of Experimental Biology*, 198: 767–774.
- Onken H & Riestenpatt S (1998) NaCl absorption across split gill lamellae of hyperregulating crabs: transport mechanisms and their regulation. *Comparative Biochemistry and Physiology A-Molecular and Integrative Physiology*, 119: 883–893.
- Orchard M (2012) Crabs of Christmas Island. Christmas Island Natural History Association, 287 pp.
- Pequeux A (1995) Osmotic regulation in crustaceans. *Journal of Crustacean Biology*, 15: 1–60.

- Riessenpatt S, Petrasch G & Siebers D (1995) Cl<sup>-</sup> Influx across posterior gills of the Chinese crab (*Eriocheir sinensis*) – Potential energization by a V-Type H<sup>+</sup>-ATPase. *Comparative Biochemistry and Physiology A-Physiology*, 110: 235–241.
- Riessenpatt S, Zeiske W & Onken H (1994) Cyclic-AMP stimulation of electrogenic uptake of Na<sup>+</sup> and Cl<sup>-</sup> across the gill epithelium of the Chinese crab *Eriocheir sinensis*. *Journal of Experimental Biology*, 188: 159–174.
- Serrano L, Blanvillain G, Soyez D, Charmantier G, Grousset E, Aujoulat F & Spanings-Pierrot C (2003) Putative involvement of Crustacean Hyperglycemic Hormone isoforms in the neuroendocrine mediation of osmoregulation in the crayfish *Astacus leptodactylus*. *Journal of Experimental Biology*, 206: 979–988.
- Shaw J (1959) Salt and water balance in the east African fresh-water crab, *Potamon niloticus* (M Edw). *Journal of Experimental Biology*, 36: 157–176.
- Taylor HH & Greenaway P (2002) Osmoregulation in the terrestrial Christmas Island red crab *Gecarcoidea natalis* (Brachyura: Gecarcinidae): modulation of branchial chloride uptake from the urine. *Journal of Experimental Biology*, 205: 3251–3260.
- Taylor HH, Greenaway P & Morris S (1993) Adaptations to a terrestrial existence by the robber crab *Birgus latro*. 8. Osmotic and ionic regulation on fresh-water and saline drinking regimens. *Journal of Experimental Biology*, 179: 93–113.
- Thomas ML, Becker K, Abbott K & Feldhaar H (2010) Supercolony mosaics: two different invasions by the yellow crazy ant, *Anoplolepis gracilipes*, on Christmas Island, Indian Ocean. *Biological Invasions*, 12: 677–687.
- Thompson LC (1970) Osmoregulation of the freshwater crabs *Metopaulius depressus* (Grapsidae) and *Pseudothelphusa jouyi* (Pseudothelphusidae). University of California, Berkeley, CA.
- Toullec JY, Serrano L, Lopez P, Soyez D & Spanings-Pierrot C (2006) The Crustacean Hyperglycemic Hormones from an euryhaline crab *Pachygrapsus marmoratus* and a fresh water crab *Potamon ibericum*: eyestalk and pericardial isoforms. *Peptides*, 27: 1269–1280.
- Townsend K, Spanings-Pierrot C, Hartline DK, King S, Henry RP & Towle DW (2001) Salinity-related changes in Crustacean Hyperglycemic Hormone (CHH) mRNA in pericardial organs of the shore crab *Carcinus maenas*. *American Zoologist*, 41: 1608–1609.
- Truchot JP (1983) Regulation of acid-base balance. In: Mantel LH (eds.) *The Biology of the Crustacea*. Academic Press, New York, pp. 431–457.
- Tsai JR & Lin HC (2007) V-type H<sup>+</sup>-ATPase and Na<sup>+</sup>/K<sup>+</sup>-ATPase in the gills of 13 euryhaline crabs during salinity acclimation. *Journal of Experimental Biology*, 210: 620–627.
- Turner LM (2010) A role for crustacean hyperglycaemic hormone (CHH) in the regulation of kidney-like function in freshwater land crabs: a study of the Christmas Island blue Crab, *Discoplax hirtipes*. PhD Thesis, University of Bristol.
- Turner LM, Webster SG & Morris S (2013) Roles of crustacean hyperglycaemic hormone in ionic and metabolic homeostasis in the Christmas Island Blue crab *Discoplax celeste*. *Journal of Experimental Biology*, 216: 1191–1201.
- Varley DG & Greenaway P (1994) Nitrogenous excretion in the terrestrial carnivorous crab *Geograpsus grayi* – site and mechanism of excretion. *Journal of Experimental Biology*, 190: 179–193.
- Webster SG, Keller R & Dirksen H (2012) The CHH-superfamily of multifunctional peptide hormones controlling crustacean metabolism, osmoregulation, moulting, and reproduction. *General and Comparative Endocrinology*, 175: 217–233.
- Weihrauch D, Morris S & Towle DW (2004) Ammonia excretion in aquatic and terrestrial crabs. *Journal of Experimental Biology*, 207: 4491–4504.
- Wheatly MG & Henry RP (1987) Branchial and antennal gland Na<sup>+</sup>/K<sup>+</sup>-dependent ATPase and carbonic-anhydrase activity during salinity acclimation of the euryhaline crayfish *Pacifastacus leniusculus*. *Journal of Experimental Biology*, 133: 73–86.
- Wheatly MG & Henry RP (1992) Extracellular and intracellular acid-base regulation in crustaceans. *Journal of Experimental Zoology*, 263: 127–142.
- Wheatly MG & Toop T (1989) Physiological-responses of the crayfish *Pacifastacus leniusculus* to environmental hyperoxia. 2. Role of the antennal gland in acid-base and ion regulation. *Journal of Experimental Biology*, 143: 53–70.
- Wolcott DL (1991) Nitrogen-excretion is enhanced during urine recycling in two species of terrestrial crab. *Journal of Experimental Zoology*, 259: 181–187.
- Wolcott TG & Wolcott DL (1984) Extra-renal urine modification for salt conservation in the land crab *Cardisoma guanhumi*. *American Zoologist*, 24: A78–A78.
- Wolcott TG & Wolcott DL (1985) Extrarenal modification of urine for ion conservation in ghost crabs, *Ocypode quadrata* (Fabricius). *Journal of Experimental Marine Biology and Ecology*, 91: 93–107.
- Wolcott TG & Wolcott DL (1991) Ion conservation by reprocessing of urine in the land crab *Gecarcinus lateralis* (Fremenville). *Physiological Zoology*, 64: 344–361.
- Zeiske W, Onken H, Schwarz HJ & Graszynski K (1992) Invertebrate epithelial Na<sup>+</sup> channels – Amiloride-induced current-noise in crab gill. *Biochimica Et Biophysica Acta*, 1105: 245–252.