



MINIREVIEW

Genetic advances in dermatophytes

Maria Grumbt¹, Michel Monod² & Peter Staib¹

¹Leibniz Institute for Natural Product Research and Infection Biology – Hans Knoell Institute, Junior Research Group Fundamental Molecular Biology of Pathogenic Fungi, Jena, Germany; and ²Department of Dermatology, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland

Correspondence: Peter Staib, Leibniz Institute for Natural Product Research and Infection Biology – Hans Knoell Institute, Junior Research Group Fundamental Molecular Biology of Pathogenic Fungi, Beutenbergstr. 11a, D-07745 Jena, Germany. Tel.: +49 3641 532 1600; fax: +49 3641 532 0809; e-mail: peter.staib@hki-jena.de

Received 8 March 2011; accepted 30 March 2011.

Final version published online 9 May 2011.

DOI:10.1111/j.1574-6968.2011.02276.x

Editor: Derek Sullivan

Keywords

transformation; selection marker; gene targeting; *Arthroderma*; *Trichophyton*; filamentous fungi.

Abstract

Millions of superficial fungal infections are annually observed in humans and animals. The majority of these mycoses are caused by dermatophytes, a specialized group of filamentous fungi that exclusively infect keratinized host structures. Despite the high prevalence of the disease, dermatophytosis, little is known about the pathogenicity mechanisms of these microorganisms. This drawback may be related to the fact that dermatophytes have been investigated poorly at the molecular level. In contrast to many other pathogenic fungi, they grow comparatively slowly under *in vitro* conditions, and in the last decades, only a limited number of molecular tools have been established for their manipulation. In recent years, however, major promising approaches were undertaken to improve genetic analyses in dermatophytes. These strategies include efficient systems for targeted gene inactivation and gene silencing, and broad transcriptional profiling techniques, which have even been applied in sophisticated infection models. As a fundamental prerequisite for future genetic analyses, full genome sequences of seven different dermatophyte species have become available recently. Therefore, it appeared timely to review the available molecular tools and methodologies in dermatophyte research, which may provide future insights into the virulence of these clinically important pathogens.

Introduction

Genetic approaches have allowed fundamental insights into almost all areas of microbial pathogenesis research. Yet, today, such methodologies have only rarely been established in dermatophytes, in contrast to other clinically important fungal pathogens, for example *Candida albicans*, *Aspergillus fumigatus* or *Cryptococcus neoformans*. Consequently, little is known about the pathogenicity of dermatophytes at the molecular level. Dermatophytes constitute a group of highly specialized filamentous fungi that share the peculiar ability to digest and grow on keratinized host structures such as skin stratum corneum, hair and nails (Fig. 1) (Ajello, 1974). Keratin utilization by these microorganisms as the sole carbon and nitrogen source has been linked to extracellular proteolysis, and a large number of secreted proteases were identified in different dermatophyte species (reviewed in Monod, 2008). Despite these major efforts, however, the role of individual proteases during infection remains almost elusive. Moreover, dermatophyte pathogenicity likely tends to be more complex and involves fungal mechanisms that still have to be identified. At the same time, it appears to be of particular note that the adaptation of dermatophytes to

specific host niches is associated with variable clinical signs, i.e. chronic vs. inflammatory disease, suggesting distinct, almost unknown pathophysiological reactions. Therefore, studies on dermatophyte pathogenicity focus not only on fungal attributes but also on host immune response mechanisms (Vermout *et al.*, 2008; Brasch, 2009).

Comprehensive up-to-date review articles covering dermatophyte epidemiology and clinical importance as well as genetic approaches in taxonomy and diagnosis are already available (Binstock, 2007; Abdel-Rahman, 2008; Gräser *et al.*, 2008; Kanbe, 2008; Seebacher *et al.*, 2008; Ameen, 2010). These topics will not be a part of the present overview. Nevertheless, some basic information on species diversity and medical impact will be provided in order to better convey the recent achievements in molecular genetic research in this fascinating group of microorganisms.

Dermatophytes: clinical importance and taxonomy

Dermatophytoses belong to the most common infectious diseases in humans, affecting 10–20% of the population




Fig. 1. Scanning electron micrograph of the dermatophyte *Arthroderma benhamiae* colonizing human hair. Sterilized hair was infected with *A. benhamiae* microconidia and incubated for 14 days at 30 °C. Scale bar = 10 µm.

worldwide. These infections are usually not life threatening, but occur even in immunocompetent hosts, and in many cases, are long lasting, recurrent and difficult to cure (Borgers *et al.*, 2005). Depending on their predominant natural reservoir, dermatophyte species are classified into three groups: anthropophilic, zoophilic and geophilic (Weitzman & Summerbell, 1995). The natural hosts of anthropophilic and zoophilic species are humans and animals, respectively, whereas geophilic dermatophytes are soil saprophytes. Symptoms of dermatophytosis can vary from chronic to highly inflammatory, depending on the causative agent and the body location affected. The given disease is described with the word 'tinea,' followed by a term referring to the infected body site, for example tinea pedis (feet), tinea capitis (scalp or head), tinea corporis (body or trunk) and tinea unguium (nails, also called onychomycosis) (Degreef, 2008). Major prominent anthropophilic species, for example, *Trichophyton rubrum*, *Trichophyton interdigitale* and *Trichophyton tonsurans*, are mostly associated with more chronic, less inflammatory infections. In contrast, zoophilic species, for example, *Microsporum canis*, *Arthroderma benhamiae*, *Arthroderma vanbreuseghemii*, *Trichophyton erinacei* and *Trichophyton verrucosum* as well as geophilic dermatophytes such as *Microsporum gypseum* often induce highly inflamed lesions in humans.

Dermatophytes are ascomycete fungi. The anamorphs (asexual forms) are classified into three genera: *Trichophyton*, *Microsporum* and *Epidermophyton*. Teleomorphs (sexual forms) belong to the *Arthroderma* genus in the Ascomycotina subphylum. Dermatophytes are heterothallic (mating types are designated as either '+' or '-'); however, in many zoophilic and anthropophilic species, sexual reproduction has not been observed. Recent progress in molecular taxonomy and insights into mating revealed that *Trichophyton mentagrophytes* was a complex of anthropophilic and zoophilic species that produce different teleomorphs, leading to

a current confusion in species denomination. For example, *A. benhamiae* is the teleomorph obtained by mating isolates from rodents (Ajello & Cheng, 1967), whereas *A. vanbreuseghemii* is the teleomorph from strains isolated from humans and certain rodents (Takashio, 1979). Both zoophilic species *A. benhamiae* and *A. vanbreuseghemii* cause highly inflammatory tinea capitis, tinea corporis and tinea faciei. They are designated *T. mentagrophytes* and *T. mentagrophytes* var. *asteroides* in many textbooks and publications. The anthropophilic strains of the *T. mentagrophytes* species complex produce noninflammatory tinea pedis and tinea unguium. Sexual reproduction has not been observed and the fungus is still called by the anamorph name *T. interdigitale* (or *T. mentagrophytes* var. *interdigitale*) (Symoens *et al.*, 2011). Therefore, the formerly widely used species description, *T. mentagrophytes*, should nowadays only be used for isolates referring to the reference strain designated as a neotype (Gräser *et al.*, 1999). This hint appears to be noteworthy, because many of the genetic studies in dermatophytes were performed using species of the *T. mentagrophytes* complex, i.e. *A. benhamiae* and *A. vanbreuseghemii*. However, in the case of the latter species, the name *T. mentagrophytes* was used (e.g. Yamada *et al.*, 2005, 2008, 2009a, b; Alshahni *et al.*, 2011).

Broad-scale gene discovery in dermatophytes

Transcriptional profiling

Broad-scale gene discovery by differential cDNA analysis, expressed sequence tag (EST) sequencing and cDNA-based microarrays allows global insights into cellular adaptation at the level of gene expression. In dermatophytes, such techniques were recently established and revealed the transcriptional response of these fungi under different biologically interesting and also pathogenicity-related conditions. A comprehensive *T. rubrum* Expression Database was launched by Wang *et al.* (2004, 2006), offering a platform for ESTs and cDNA microarray-based transcriptional profiles (<http://www.mgc.ac.cn/TrED/>). Documented in a number of publications, this approach resulted in the identification of *T. rubrum* genes, whose expression is linked to distinct developmental growth phases or the presence of selected drugs (Liu *et al.*, 2007; Yang *et al.*, 2007; Yu *et al.*, 2007; Zhang *et al.*, 2007, 2009). Broad transcriptional analyses were also performed in our work on *T. rubrum* and *A. benhamiae*, with a focus on genes putatively implicated in extracellular proteolysis. Herein, ESTs from *T. rubrum* grown on protein as the sole carbon and nitrogen source were analysed and used for the construction of a cDNA microarray containing at least 23 protease genes (Zaugg *et al.*, 2009). Major dermatophyte-secreted

keratinases have been known before and were correlated with the degradation of hard compact keratin (for a review, see Monod, 2008). Notably, dermatophytes were shown to secrete multiple serine proteases of the subtilisin family (Sub) as well as metalloproteases of the fungalyisin family (Mep) [S8 and M36 family, respectively, in the MEROPS proteolytic enzyme database (<http://merops.sanger.ac.uk>)]. Microarray analysis during the growth of *T. rubrum* or *A. benhamiae* on soy and keratin protein confirmed the activation of particular *SUB* and *MEP* genes as well as genes encoding secreted exoproteases such as leucine aminopeptidases and dipeptidylpeptidases. In addition, other specifically induced factors playing a potential role in protein utilization were identified, including heat shock proteins, various transporters, metabolic enzymes, transcription factors and hypothetical proteins with unknown functions (Zaugg *et al.*, 2009; Staib *et al.*, 2010). Similar approaches were also supported by the analysis of suppression subtractive hybridization libraries, applied for the identification of novel dermatophyte genes specifically expressed by *T. rubrum* cells upon contact with keratin, in response to varying pH or to other environmental stimuli (Kaufman *et al.*, 2005; Baeza *et al.*, 2007; Maranhao *et al.*, 2007, 2009; Peres *et al.*, 2010; Silveira *et al.*, 2010). A comparative transcriptional analysis in the two closely related species *T. tonsurans* and *Trichophyton equinum* detected differential, species-specific expression levels of selected genes encoding secreted proteases upon growth on keratin (Preuett *et al.*, 2010).

Gene expression profiling in *A. benhamiae* during infection

In order to unravel pathogenicity-related adaptation mechanisms of dermatophytes during infection, we explored the transcriptional response of the fungal cells in an animal model. For this approach, the zoophilic dermatophyte *A. benhamiae* was selected as an appropriate species for several reasons (Fig. 2). *Arthroderma benhamiae* is zoophilic and causes inflammatory cutaneous infections not only in humans but also in guinea-pigs, allowing the establishment of an animal model (Staib *et al.*, 2010). Under laboratory conditions, *A. benhamiae* grows relatively fast and produces abundant microconidia, single-nucleated round-oval cells that are useful for transformation. Cleistothecia formation further facilitates genetic analyses and allows to shed light on the basis of sexual development in dermatophytes. As a major additional prerequisite, the genome of our *A. benhamiae* strain, which had been isolated from a patient with highly inflammatory tinea faciei (Fumeaux *et al.*, 2004), has recently been decoded and annotated (Burmester *et al.*, 2011) (Fig. 2). Transcriptional analysis in *A. benhamiae* cells isolated during experimental cutaneous infection of guinea-pigs uncovered a distinct protease gene expression profile, which is essentially different from the pattern displayed during *in vitro* growth on keratin. Most notably, a differential expression of genes coding for members of the Sub and Mep protease families was detected. Instead of the




Fig. 2. Summary of the basic characteristics that make *Arthroderma benhamiae* a useful model for molecular research in dermatophytes. The centre shows a typical colony of *A. benhamiae* on Sabouraud glucose agar after 5 days of growth at 30 °C. Detailed explanations are given in the text.

major keratinase genes expressed *in vitro*, others were activated specifically during infection, suggesting functions that are not necessarily associated with the degradation of keratin. Future studies will address the strong *in vivo* activation of the gene encoding the serine protease Sub6, a known major allergen in the related dermatophyte *T. rubrum*. The broad *A. benhamiae* *in vivo* gene expression profile further revealed other putatively pathogenicity-related factors, whose role has to be studied by straightforward functional analysis. Other interesting, putatively pathogenicity-related dermatophyte genes have been identified recently in a broad transcriptome approach in *A. benhamiae* during the interaction with human keratinocytes (Burmester et al., 2011).

Transformation and gene targeting in dermatophytes

Transformation and selection markers

In comparison with many other fungi, dermatophytes have been shown to be less amenable to genetic manipulation. As a result, site-directed mutagenesis in dermatophyte species has been evidenced only in a very small number of cases. This drawback is assumed to be a result of both low transformation frequency and inefficient homologous integration, processes that are indispensable for targeted genetic manipulations. The first successful transformation of a dermatophyte has been described in 1989 by Gonzalez et al. (1989) in *T. mentagrophytes* (Table 1). The transformation protocol applied was based on a standard protoplast/polyethylene glycol (PEG)-mediated procedure that has been established widely in filamentous fungi, for example *Asper-*

gillus nidulans, *Neurospora crassa* and others (for a review, see Fincham, 1989; Weld et al., 2006). As a marker for the selection of *T. mentagrophytes* transformants, the system used the bacterial hygromycin B phosphotransferase gene *hph*. Plasmid DNA was stably integrated into the fungal genome with varying integration sites and numbers of insertions in the resulting transformants. Thereafter, no further attempts on dermatophyte transformation have been reported until 2004, when Kaufman et al. (2004) described PEG-mediated protoplast transformation and restriction-enzyme-mediated integration in *T. mentagrophytes*, using the *hph* gene as a selectable marker and the gene encoding the enhanced green fluorescent protein (eGFP) as a reporter. PEG-mediated transformation and transformant selection via hygromycin resistance was further demonstrated in *M. canis* (Yamada et al., 2005, 2006; Vermout et al., 2007) and *T. rubrum* (Fachin et al., 2006; Ferreira-Nozawa et al., 2006). Different other drugs/dominant markers have meanwhile been proven successful for the selection of transformants in *T. mentagrophytes*, i.e. two other aminoglycoside antibiotics/resistance genes, nourseothricin/*Streptomyces noursei* nourseothricin acetyltransferase gene *nat1* (Alshahni et al., 2010) and geneticin (G-418)/*Escherichia coli* neomycin phosphotransferase gene *neo* (Yamada et al., 2008). The latter marker as well as *hph* were also used successfully in *A. benhamiae* (Grumbt et al., 2011). Besides PEG-mediated protoplast transformation, other techniques facilitating gene transfer were also meanwhile adopted in dermatophytes. A promising *Agrobacterium tumefaciens*-mediated transformation (ATMT) system was established recently for *T. mentagrophytes* (Yamada et al., 2009b). ATMT has already strongly advanced functional genomics in various filamentous fungi before (for a

Table 1. Chronological overview of successful genetic transformation experiments in dermatophytes

Species	Method	Resistance gene	Specific integration	References
<i>T. mentagrophytes</i>	PEG	<i>hph</i>	No	Gonzalez et al. (1989)
<i>T. mentagrophytes</i>	PEG/REMI	<i>hph</i>	No	Kaufman et al. (2004)
<i>T. mentagrophytes</i> / <i>M. canis</i>	PEG	<i>hph</i>	No	Yamada et al. (2005)
<i>M. canis</i>	PEG	<i>hph</i>	Yes	Yamada et al. (2006)
<i>T. rubrum</i>	PEG	<i>hph</i>	Yes	Fachin et al. (2006)
<i>T. rubrum</i>	PEG	<i>hph</i>	Yes	Ferreira-Nozawa et al. (2006)
<i>M. canis</i>	PEG	<i>hph</i>	No	Vermout et al. (2007)
<i>T. mentagrophytes</i>	PEG	<i>neo</i>	No	Yamada et al. (2008)
<i>T. rubrum</i>	Electroporation	<i>hph</i>	No	Dobrowolska & Staczek (2009)
<i>T. mentagrophytes</i>	ATMT	<i>hph</i>	Yes/no	Yamada et al. (2009b)
<i>T. mentagrophytes</i>	ATMT	<i>hph/neo</i>	Yes	Yamada et al. (2009a)
<i>T. mentagrophytes</i>	ATMT	<i>nat1</i>	Yes	Alshahni et al. (2010)
<i>T. mentagrophytes</i>	ATMT	<i>hph/neo</i>	Yes	Alshahni et al. (2011)
<i>A. benhamiae</i>	PEG	<i>hph/neo</i>	Yes	Grumbt et al. (2011)

REMI, restriction-enzyme-mediated integration; *hph*, hygromycin phosphotransferase gene; *neo*, neomycin phosphotransferase gene; *nat1*, nourseothricin acetyltransferase gene.

review, see Michielse *et al.*, 2005). Notably, Yamada and colleagues used the system for both random integration of T- (transferred) DNA and targeted insertion, for example disruption of the *areA/nit-2* gene. As another alternative transformation technique, electroporation of germinated conidia was applied in *T. rubrum*, allowing the random integration of *hph* and *eGFP* (Dobrowolska & Staczek, 2009). Although not many comparative data on transformation efficiency are available – some species have not even been addressed at all – different dermatophyte species appear to be more or less amenable to DNA uptake and/or stable integration. Therefore, transformation protocols established for a selected species are not necessarily transferable to another, but require precise modifications. From our own work, we know for example that our standard PEG-protocol for the efficient transformation of *A. benhamiae* was not directly applicable for *T. rubrum* or *M. canis*. The reasons for this observation are likely multifactorial, including differential protoplast stability, cell wall composition, microconidia production, etc.

Targeted gene inactivation

Filamentous fungi are known to only poorly support site-directed insertion of linear DNA cassettes in the genome by homologous recombination, in contrast to yeasts such as *Saccharomyces cerevisiae* or the opportunistic pathogen *C. albicans*. Therefore, in filamentous fungi, identification of transformants with a desired genetic alteration has proven laborious in many cases. In order to circumvent this obstacle, parental strains were generated in diverse species that lack the nonhomologous end joining (NHEJ) recombination pathway, for example in *N. crassa* (Ninomiya *et al.*, 2004), *Aspergillus* spp. (da Silva Ferreira *et al.*, 2006; Krappmann *et al.*, 2006; Nayak *et al.*, 2006), and since recently, also in *T. mentagrophytes* (Yamada *et al.*, 2009a) and *A. benhamiae* (Grumbt *et al.*, 2011) (Table 1). Mutants deficient in NHEJ processes allow a strongly increased frequency of targeted insertions; however, an altered risk of unforeseen genetic variations cannot be excluded. In dermatophyte species, only a small number of genes have so far been analysed by targeted inactivation, for example *pacC* and *MDR2* in *T. rubrum* (Fachin *et al.*, 2006; Ferreira-Nozawa *et al.*, 2006), *Ku80*, *areA* and *Trim4* in *T. mentagrophytes* (Yamada *et al.*, 2009a, b), *areA* in *M. canis* (Yamada *et al.*, 2006) and *Ku70* and *AcuE* in *A. benhamiae* (Grumbt *et al.*, 2011). Interestingly, *A. benhamiae* has been shown in our work to allow efficient targeted gene deletion not only in a *ku70* mutant background but also in the wild-type strain. This has been demonstrated by the construction of mutants in malate synthase *AcuE*, *KU70* and other candidates (Grumbt *et al.*, 2011; M. Grumbt and P. Staib, unpublished data). The use of two different dominant

selection markers, *hph* and *neo*, even allowed for the first time the site-directed complementation of knockout mutant strains. Because the deletion of *KU70* had no adverse effect on the virulence of *A. benhamiae* in a guinea-pig infection model, both the wild type and the *ku70* mutant appear to be suitable parental strains for future pathogenicity research. In general, isogenic strain construction is assumably facilitated in species such as *A. benhamiae* and *T. mentagrophytes* by the fact that they easily allow the production of abundant single nucleated cells in the form of microconidia as a starting material.

RNA silencing

RNA interference, originally described in the nematode *Caenorhabditis elegans*, is based on a cellular process by which an introduced double-stranded RNA induces the degradation of specific mRNAs of interest (Fire *et al.*, 1998). RNA silencing was widely applied as an efficient tool to address gene function in multiple research areas, especially when conventional site-directed gene inactivation is difficult or, due to knockout lethality, impossible. As another advantage, the technique offers the possibility to inhibit several genes at the same time, a characteristic that might be useful for the functional analysis of homologous genes within large families, for example those encoding secreted endoproteases in dermatophytes. Here, the system was first established by Vermout *et al.* (2007) by the construction of *M. canis* transformants in which the expression of genes encoding secreted proteases Sub3 and dipeptidyl peptidase IV, respectively, was suppressed. Using the *SUB3* RNA-silenced strain, the authors revealed a contribution of this protease in the adherence of *M. canis* to feline epidermis, whereas a function in epidermal invasion and virulence of the fungus during cutaneous guinea-pig infection was not assigned (Baldo *et al.*, 2010).

Genome sequencing projects

Given the fact that powerful tools have meanwhile become available for the genetic manipulation of dermatophytes, the advent of dermatophyte genome sequencing projects now offers a fundamental basis for future research. Annotated genome sequences of seven different dermatophyte species have become available recently (http://www.broadinstitute.org/annotation/genome/dermatophyte_comparative/MultiHome.html), provided by projects headed by the Broad Institute (Cambridge) and the Hans Knoell Institute (Jena, Germany), respectively. The latter institution has recently published the first report on dermatophyte genomes, presenting a comparative study on the two closely related zoophilic, human pathogenic species *A. benhamiae* (major reservoir are guinea-pigs) and *T. verrucosum* (major

reservoir are cattle) (Burmester *et al.*, 2011). The genome sequences identified were compared not only with each other but also with those of other species of the *Onygenales*, i.e., *Coccidioides posadasii* and *Coccidioides immitis*, and with the mould *A. fumigatus*. The 22–23 Mb genomes of *A. benhamiae* and *T. verrucosum*, containing 7980 and 8024 predicted protein-encoding genes, respectively, were found to be smaller than those of *Aspergillus* (e.g. 28 and 37.3 Mb for *Aspergillus clavatus* and *Aspergillus niger*, respectively), *Coccidioides* spp. (27–29 Mb) or *Histoplasma capsulatum* (30–39 Mb). Special attention was paid not only to the analysis of genes that are putatively associated with host adaptation, for example genes encoding secreted proteases. Genes involved in the biosynthesis of secondary metabolites and mating were also found to be of future interest (Burmester *et al.*, 2011). Additional insights are expected from the envisaged genome comparison including the other five sequenced human pathogenic dermatophyte species. The species selection was based on different biological parameters and pathogenicity-related hypotheses (White *et al.*, 2008), and the basic traits of the selected strains such as growth rate and resistance to diverse antibiotics were already monitored (Achterman *et al.*, 2011). Because these species encompass anthropophilic (*T. rubrum*, the most common inducer of dermatophytosis in humans worldwide; *T. tonsurans*, often associated with tinea capitis in America), zoophilic (*T. equinum*, associated with horses; *M. canis*, associated with cats and dogs) and geophilic (*M. gypseum*) dermatophytes, a comparative genome analysis will, among other topics, address factors that are potentially involved in host preference, adaptation during chronic vs. inflammatory infection and saprophytic growth.

Conclusion

An increasing, lively interest in the molecular biology of dermatophytes combined with the establishment of fundamental genetic approaches has strongly advanced the research in these filamentous fungi. Basic prerequisites have been launched, such as genome sequencing projects, expression profile data sets and efficient targeted gene inactivation techniques. Nevertheless, molecular research is still preliminary in these genetically less amenable microorganisms. Therefore, further efforts have to be undertaken for the improvement of existing and the establishment of additional genetic tools and methodologies. Such efforts will be worthwhile, given the fact that dermatophytoses are widespread and of particular clinical interest. Using the available techniques, now fundamental questions can be addressed in dermatophytes, related to the pathogenicity as well as general host and environmental adaptation mechanisms, sexual development, basic biology and evolution.

Acknowledgements

We are sorry that space limitations did not allow us to cite all important papers. We thank Axel A. Brakhage, Christoph Heddergott and the electron microscopy centre at the University Hospital Jena for providing the scanning electron micrograph in Fig. 1, and Bernard Mignon for the photograph visualizing the guinea-pig animal model in Fig. 2. Work in our laboratory is supported by the Deutsche Forschungsgemeinschaft and the Hans Knoell Institute.

References

- Abdel-Rahman SM (2008) Strain differentiation of dermatophytes. *Mycopathologia* **166**: 319–333.
- Achterman RR, Smith AR, Oliver BG & White TC (2011) Sequenced dermatophyte strains: growth rate, conidiation, drug susceptibilities, and virulence in an invertebrate model. *Fungal Genet Biol* **48**: 335–341.
- Ajello L (1974) Natural history of the dermatophytes and related fungi. *Mycopath Mycol Appl* **53**: 93–110.
- Ajello L & Cheng SL (1967) The perfect state of *Trichophyton mentagrophytes*. *Sabouraud* **5**: 230–234.
- Alshahni MM, Makimura K, Yamada T, Takatori K & Sawada T (2010) Nourseothricin acetyltransferase: a new dominant selectable marker for the dermatophyte *Trichophyton mentagrophytes*. *Med Mycol* **48**: 665–668.
- Alshahni MM, Yamada T, Takatori K, Sawada T & Makimura K (2011) Insights into a nonhomologous integration pathway in the dermatophyte *Trichophyton mentagrophytes*: efficient targeted gene disruption by use of mutants lacking ligase IV. *Microbiol Immunol* **55**: 34–43.
- Ameen M (2010) Epidemiology of superficial fungal infections. *Clin Dermatol* **28**: 197–201.
- Baeza LC, Bailao AM, Borges CL, Pereira M, Soares CM & Mendes Giannini MJ (2007) cDNA representational difference analysis used in the identification of genes expressed by *Trichophyton rubrum* during contact with keratin. *Microbes Infect* **9**: 1415–1421.
- Baldo A, Mathy A, Tabart J *et al.* (2010) Secreted subtilisin Sub3 from *Microsporum canis* is required for adherence to but not for invasion of the epidermis. *Brit J Dermatol* **162**: 990–997.
- Binstock JM (2007) Molecular biology techniques for identifying dermatophytes and their possible use in diagnosing onychomycosis in human toenail: a review. *J Am Podiat Med Assn* **97**: 134–144.
- Borgers M, Degreef H & Cauwenbergh G (2005) Fungal infections of the skin: infection process and antimycotic therapy. *Curr Drug Targets* **6**: 849–862.
- Brasch J (2009) Current knowledge of host response in human tinea. *Mycoses* **52**: 304–312.
- Burmester A, Shelest E, Glockner G *et al.* (2011) Comparative and functional genomics provide insights into the pathogenicity of dermatophytic fungi. *Genome Biol* **12**: R7.

- da Silva Ferreira ME, Kress MR, Savoldi M *et al.* (2006) The *akuB*(KU80) mutant deficient for nonhomologous end joining is a powerful tool for analyzing pathogenicity in *Aspergillus fumigatus*. *Eukaryot Cell* **5**: 207–211.
- Degreef H (2008) Clinical forms of dermatophytosis (ringworm infection). *Mycopathologia* **166**: 257–265.
- Dobrowolska A & Staczek P (2009) Development of transformation system for *Trichophyton rubrum* by electroporation of germinated conidia. *Curr Genet* **55**: 537–542.
- Fachin AL, Ferreira-Nozawa MS, Maccheroni W Jr & Martinez-Rossi NM (2006) Role of the ABC transporter TruMDR2 in terbinafine, 4-nitroquinoline *N*-oxide and ethidium bromide susceptibility in *Trichophyton rubrum*. *J Med Microbiol* **55**: 1093–1099.
- Ferreira-Nozawa MS, Silveira HC, Ono CJ, Fachin AL, Rossi A & Martinez-Rossi NM (2006) The pH signaling transcription factor PacC mediates the growth of *Trichophyton rubrum* on human nail *in vitro*. *Med Mycol* **44**: 641–645.
- Fincham JR (1989) Transformation in fungi. *Microbiol Rev* **53**: 148–170.
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE & Mello CC (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**: 806–811.
- Fumeaux J, Mock M, Ninet B *et al.* (2004) First report of *Arthroderma benhamiae* in Switzerland. *Dermatology* **208**: 244–250.
- Gonzalez R, Ferrer S, Buesa J & Ramon D (1989) Transformation of the dermatophyte *Trichophyton mentagrophytes* to hygromycin B resistance. *Infect Immun* **57**: 2923–2925.
- Gräser Y, El Fari M, Vilgalys R, Kuijpers AF, De Hoog GS, Presber W & Tietz H (1999) Phylogeny and taxonomy of the family *Arthrodermataceae* (dermatophytes) using sequence analysis of the ribosomal ITS region. *Med Mycol* **37**: 105–114.
- Gräser Y, Scott J & Summerbell R (2008) The new species concept in dermatophytes – a polyphasic approach. *Mycopathologia* **166**: 239–256.
- Grumbt M, Defaweux V, Mignon B, Monod M, Burmester A, Wöstemeyer J & Staib P (2011) Targeted gene deletion and *in vivo* analysis of putative virulence gene function in the pathogenic dermatophyte *Arthroderma benhamiae*. *Eukaryot Cell*, DOI: 10.1128/EC.00273-10.
- Kanbe T (2008) Molecular approaches in the diagnosis of dermatophytosis. *Mycopathologia* **166**: 307–317.
- Kaufman G, Horwitz BA, Hadar R, Ullmann Y & Berdicevsky I (2004) Green fluorescent protein (GFP) as a vital marker for pathogenic development of the dermatophyte *Trichophyton mentagrophytes*. *Microbiology* **150**: 2785–2790.
- Kaufman G, Berdicevsky I, Woodfolk JA & Horwitz BA (2005) Markers for host-induced gene expression in *Trichophyton* dermatophytosis. *Infect Immun* **73**: 6584–6590.
- Krappmann S, Sasse C & Braus GH (2006) Gene targeting in *Aspergillus fumigatus* by homologous recombination is facilitated in a nonhomologous end-joining-deficient genetic background. *Eukaryot Cell* **5**: 212–215.
- Liu T, Zhang Q, Wang L *et al.* (2007) The use of global transcriptional analysis to reveal the biological and cellular events involved in distinct development phases of *Trichophyton rubrum* conidial germination. *BMC Genomics* **8**: 100.
- Maranhao FC, Paiao FG & Martinez-Rossi NM (2007) Isolation of transcripts over-expressed in human pathogen *Trichophyton rubrum* during growth in keratin. *Microb Pathogenesis* **43**: 166–172.
- Maranhao FC, Paiao FG, Fachin AL & Martinez-Rossi NM (2009) Membrane transporter proteins are involved in *Trichophyton rubrum* pathogenesis. *J Med Microbiol* **58**: 163–168.
- Michiels CB, Hooykaas PJ, van den Hondel CA & Ram AF (2005) *Agrobacterium*-mediated transformation as a tool for functional genomics in fungi. *Curr Genet* **48**: 1–17.
- Monod M (2008) Secreted proteases from dermatophytes. *Mycopathologia* **166**: 285–294.
- Nayak T, Szewczyk E, Oakley CE *et al.* (2006) A versatile and efficient gene-targeting system for *Aspergillus nidulans*. *Genetics* **172**: 1557–1566.
- Ninomiya Y, Suzuki K, Ishii C & Inoue H (2004) Highly efficient gene replacements in *Neurospora* strains deficient for nonhomologous end-joining. *P Natl Acad Sci USA* **101**: 12248–12253.
- Peres NT, Sanches PR, Falcao JP *et al.* (2010) Transcriptional profiling reveals the expression of novel genes in response to various stimuli in the human dermatophyte *Trichophyton rubrum*. *BMC Microbiol* **10**: 39.
- Preuett BL, Schuenemann E, Brown JT, Kovac ME, Krishnan SK & Abdel-Rahman SM (2010) Comparative analysis of secreted enzymes between the anthropophilic–zoophilic sister species *Trichophyton tonsurans* and *Trichophyton equinum*. *Fungal Biol* **114**: 429–437.
- Seebacher C, Bouchara JP & Mignon B (2008) Updates on the epidemiology of dermatophyte infections. *Mycopathologia* **166**: 335–352.
- Silveira HC, Gras DE, Cazzaniga RA, Sanches PR, Rossi A & Martinez-Rossi NM (2010) Transcriptional profiling reveals genes in the human pathogen *Trichophyton rubrum* that are expressed in response to pH signaling. *Microb Pathogenesis* **48**: 91–96.
- Staib P, Zaugg C, Mignon B *et al.* (2010) Differential gene expression in the pathogenic dermatophyte *Arthroderma benhamiae* *in vitro* versus during infection. *Microbiology* **156**: 884–895.
- Symoens F, Jousson O, Planard C, Fratti M, Staib P, Mignon B & Monod M (2011) Molecular analysis and mating behaviour of the *Trichophyton mentagrophytes* species complex. *Int J Med Microbiol* **301**: 260–266.
- Takashio M (1979) Taxonomy of dermatophytes based on their sexual states. *Mycologia* **71**: 968–976.
- Vermout S, Tabart J, Baldo A, Monod M, Losson B & Mignon B (2007) RNA silencing in the dermatophyte *Microsporium canis*. *FEMS Microbiol Lett* **275**: 38–45.

- Vermout S, Tabart J, Baldo A, Mathy A, Losson B & Mignon B (2008) Pathogenesis of dermatophytosis. *Mycopathologia* **166**: 267–275.
- Wang L, Ma L, Leng W *et al.* (2004) Analysis of part of the *Trichophyton rubrum* ESTs. *Sci China Ser C* **47**: 389–395.
- Wang L, Ma L, Leng W *et al.* (2006) Analysis of the dermatophyte *Trichophyton rubrum* expressed sequence tags. *BMC Genomics* **7**: 255.
- Weitzman I & Summerbell RC (1995) The dermatophytes. *Clin Microbiol Rev* **8**: 240–259.
- Weld RJ, Plummer KM, Carpenter MA & Ridgway HJ (2006) Approaches to functional genomics in filamentous fungi. *Cell Res* **16**: 31–44.
- White TC, Oliver BG, Gräser Y & Henn MR (2008) Generating and testing molecular hypotheses in the dermatophytes. *Eukaryot Cell* **7**: 1238–1245.
- Yamada T, Makimura K, Uchida K & Yamaguchi H (2005) Reproducible genetic transformation system for two dermatophytes, *Microsporum canis* and *Trichophyton mentagrophytes*. *Med Mycol* **43**: 533–544.
- Yamada T, Makimura K & Abe S (2006) Isolation, characterization, and disruption of *dnr1*, the *areA/nit-2*-like nitrogen regulatory gene of the zoophilic dermatophyte, *Microsporum canis*. *Med Mycol* **44**: 243–252.
- Yamada T, Makimura K, Hisajima T, Ito M, Umeda Y & Abe S (2008) Genetic transformation of the dermatophyte, *Trichophyton mentagrophytes*, based on the use of G418 resistance as a dominant selectable marker. *J Dermatol Sci* **49**: 53–61.
- Yamada T, Makimura K, Hisajima T, Ishihara Y, Umeda Y & Abe S (2009a) Enhanced gene replacements in Ku80 disruption mutants of the dermatophyte, *Trichophyton mentagrophytes*. *FEMS Microbiol Lett* **298**: 208–217.
- Yamada T, Makimura K, Satoh K, Umeda Y, Ishihara Y & Abe S (2009b) *Agrobacterium tumefaciens*-mediated transformation of the dermatophyte, *Trichophyton mentagrophytes*: an efficient tool for gene transfer. *Med Mycol* **47**: 485–494.
- Yang L, Wang L, Peng J *et al.* (2007) Comparison between gene expression of conidia and germinating phase in *Trichophyton rubrum*. *Sci China Ser C* **50**: 377–384.
- Yu L, Zhang W, Wang L *et al.* (2007) Transcriptional profiles of the response to ketoconazole and amphotericin B in *Trichophyton rubrum*. *Antimicrob Agents Ch* **51**: 144–153.
- Zaugg C, Monod M, Weber J *et al.* (2009) Gene expression profiling in the human pathogenic dermatophyte *Trichophyton rubrum* during growth on proteins. *Eukaryot Cell* **8**: 241–250.
- Zhang W, Yu L, Leng W *et al.* (2007) cDNA microarray analysis of the expression profiles of *Trichophyton rubrum* in response to novel synthetic fatty acid synthase inhibitor PHS11A. *Fungal Genet Biol* **44**: 1252–1261.
- Zhang W, Yu L, Yang J, Wang L, Peng J & Jin Q (2009) Transcriptional profiles of response to terbinafine in *Trichophyton rubrum*. *Appl Microbiol Biot* **82**: 1123–1130.