

**HHS PUBLIC ACCESS**

Author manuscript

*Endocrinol Metab Clin North Am.* Author manuscript; available in PMC 2018 March 01.

Published in final edited form as:

*Endocrinol Metab Clin North Am.* 2017 March ; 46(1): 1–18. doi:10.1016/j.ecl.2016.09.003.

## The Role of the Osteocyte in Bone and Non-bone Disease

**Lynda F. Bonewald, PhD [Executive Director]**

Indiana Center for Musculoskeletal Health, Professor, Anatomy and Cell Biology and Orthopaedics, VanNuys Medical Science Bldg, MS 5055S, 635 Barnhill Drive, Indianapolis, IN 46202, Office: 317 278-5155, Fax: 317 278-2040, lbonewal@iu.edu

### Synopsis

When normal physiological functions go awry, disorders and disease occurs. This is universal, even for the osteocyte, a cell embedded within the mineralized matrix of bone. It was once thought that this cell was simply a place-holder in bone. However, within the last decade, the number of studies of osteocytes has dramatically increased leading to the discovery of novel functions of these cells. But with the discovery of novel physiological functions came the discoveries of how these cells can also be responsible for not only bone diseases and disorders, but also those of kidney, heart, and potentially muscle.

### Keywords

osteocyte; bone disease; sclerostin; FGF23; therapeutics

### Introduction

Before osteocytes were recognized as active essential bone cells necessary for bone health, it was assumed that all the action took place on the bone surface and not within the bone. Osteoblasts and osteoclasts were the major players, osteoblasts making bone and osteoclasts resorbing bone to maintain bone homeostasis. It was assumed that osteoblasts and osteoclasts were regulated by external factors such as parathyroid hormone, PTH or 1,25 dihydroxyvitamin D3, and other external regulatory factors. It has also been proposed that osteoblasts make factors that regulate osteoclast activity and conversely that osteoclasts make factors that could regulate osteoblast activity. Therapeutics were generated that would target either osteoclasts or osteoblasts. Osteocytes were left out of the picture.

With new technology and new tools, it became possible to study osteocytes. The normal functions of osteocytes expanded rapidly to include: regulation of osteoblast and osteoclast activity to control bone remodeling, as regulators of both phosphate and calcium homeostasis, as mechanosensory cells that coordinate the skeleton's response to loading or

Correspondence to: Lynda F. Bonewald.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The Author has nothing to disclose.

unloading, and as endocrine cells targeting other tissues such as kidney. These cells are also one of the longest lived cell types in the body, some living for decades, therefore survival and normal function is paramount. (See box 1). A number of pathologic or disease conditions can now be ascribed to abnormal or missing osteocyte functions including sclerosteosis, hypophosphatemic rickets, osteoporosis, necrotic bone, aging and others. (See Figure 1). Now therapeutics are being generated that target osteocyte factors. (For reviews see<sup>1,2</sup>)

## Normal osteocyte functions

### Mechanosensation

When early histomorphomists peered through their microscopes and began to visualize osteocytes in bone, the morphology and connectedness suggested a network perhaps similar to the neural system. One of the earliest functions ascribed to osteocytes was mechanosensation based on Julius Wolff's descriptions of the capacity of bone to adapt to mechanical loading or lack of loading by adding or removing bone<sup>3</sup>. Over a century later, experiments have been performed supporting the hypothesis that osteocytes are responsible for bone adaption in response to loading. By performing targeted deletion of osteocytes in mice expressing the diphtheria toxin receptor specifically in osteocytes, it was shown that these mice were resistant to unloading-induced bone loss<sup>4</sup>.

The osteocyte and its dendritic processes are constantly exposed to canalicular fluid that flows through the lacunocanalicular system. A baseline flow of canalicular fluid flow is driven by the extravascular pressure and intermittent mechanical loading superimposes rapid alterations in canalicular fluid flow<sup>5</sup>. This results in the cells being exposed to different types and magnitudes of fluid flow shear stress. Almost every cell responds to mechanical loading, however osteocytes appear to be most sensitive when compared to osteoblasts and fibroblasts<sup>6,7</sup>.

In both primary osteocytes and MLO-Y4 osteocyte-like cells, fluid flow shear stress has been shown to have numerous sequential effects<sup>2</sup>. The first event is the release of intracellular calcium followed by the release of nitric oxide, ATP and prostaglandins, and the opening of connexin 43 hemichannels enhancing gap junction functions. Soon after the rapid change in calcium signaling (within seconds) nitric oxide, ATP, and prostaglandin (within seconds to minutes) are released. Deleting any one of these three early small molecules will inhibit bone's anabolic response to loading. Shear stress has also been shown to induce the bending of osteocyte cilia, and to initiate signaling pathways such as the wnt/ $\beta$ -catenin and PKA pathways. Shear stress also activates gene transcription and translation, and promotes dendrite elongation. One very important effect of fluid flow shear stress is the protection of osteocytes against apoptosis and cell death. Ideally, it would be important to identify early regulators of anabolic signaling in osteocytes in addition to calcium, nitric oxide, ATP, and PGE<sub>2</sub> in order to develop new therapeutics. (For review see<sup>8</sup>).

A major source of prostaglandin in the body appears to be from bone. Osteocytes are prodigious producers of prostaglandin in response to loading<sup>9</sup>, PGE<sub>2</sub> has paracrine effects on osteocytes to enhance gap junction function<sup>10</sup>, it protects and maintains osteocyte viability<sup>11</sup>

and PGE<sub>2</sub> appears to be one of the key initiators of anabolic bone formation. Administration of prostaglandin increases bone mass and inhibitors of prostaglandin production, such as indomethacin, block the effects of anabolic loading<sup>12,13</sup>. One of the most important effects of prostaglandin released in response to loading may be to activate a very important signaling pathway in the osteocyte, the Wnt/ $\beta$ -catenin signaling pathway.

Fluid flow shear stress activates the Wnt/ $\beta$ -catenin signaling pathway through the rapid release of prostaglandin which acts through EP receptors to bypass Low Density Lipoprotein receptor, LRP receptor, activation<sup>14,15</sup>. Components of the  $\beta$ -catenin pathway are essential for osteocyte viability, mechanosensation and transduction, and release of important factors essential for bone homeostasis. The central molecule through which all molecules must go is  $\beta$ -catenin.  $\beta$ -catenin regulates expression of both the positive activators of this pathway, the wnts, and the negative regulators of this pathway, sclerostin and Dkk1 (For review see<sup>16</sup>). Global deletion of  $\beta$ -catenin is embryonically lethal, but deletion in osteocytes using the Dmp1-Cre results in dramatic bone loss characterized by perforated cortices<sup>17</sup>. Interestingly, deletion of only one allele in osteocytes results in mice with a normal skeleton but a completely abrogated response to anabolic loading<sup>18</sup>.  $\beta$ -catenin plays an important role in bone integrity, osteocyte communication, osteocyte viability, but also in bone response to loading. This extends to other components of this signaling pathway.

Two of the most famous and well-studied components of this pathway are the Lrp5 receptor and the negative regulator of the  $\beta$ -catenin pathway, sclerostin encoded by the gene *Sost*. Lrp5, a major co-receptor for Wnt signalling is expressed by many cells in the body, but sclerostin is relatively osteocyte specific. Deletion of Lrp5 results in mice with impaired response to anabolic loading<sup>19</sup>. As Sclerostin is expressed in mature osteocytes and mechanical loading reduces sclerostin levels downregulation of sclerostin most likely creates a permissive environment in which Wnt proteins already present can activate the Wnt/ $\beta$ -catenin pathway. The role of the  $\beta$ -catenin pathway in disease is discussed below.

It has been shown that osteocyte specific or selective genes are regulated by loading or unloading. These genes include the markers for early osteocytes, E11/gp38, Phosphate Regulating Neutral Endopeptidase on Chromosome X, (PHEX), Dentin Matrix Protein 1, (DMP1), and the markers for late osteocytes, sclerostin, Matrix Extracellular phosphoglycoprotein, (MEPE), and Fibroblast Growth Factor, (FGF23). (The function of these and their relationship to disease will be discussed below.) Regulators of mineralization and phosphate homeostasis such as Phex, MEPE, and DMP1 are upregulated in response to mechanical loading<sup>20–22</sup> as is E11/gp38, a marker for the early osteocyte<sup>23</sup>. It would be expected that genes involved in bone formation would be upregulated in response to anabolic load and genes responsible for resorption would be downregulated. It has been shown that unloading increases RANKL, an essential promoter of osteoclast formation, in osteocytes<sup>24</sup>, that may be responsible for the bone loss associated with unloading. *Sost*/sclerostin, a marker for the late osteocyte and an inhibitor of osteoblast function, is down-regulated by anabolic mechanical loading and is increased in response to hindlimb unloading<sup>25,26</sup>.

In summary, the osteocyte's response to mechanical loading may be one of the major cellular mechanisms responsible for the positive effects of exercise not only on bone but on the function of other tissues and organs in the body.

### Calcium homeostasis

Probably the earliest proposed function of osteocytes was a capacity to remove their perilacunar matrix, a process referred to by Belanger as 'osteocytic osteolysis'<sup>27</sup>. In 1910, over 100 years ago, Von Recklinghausen described enlarged lacunae in patients with rickets or osteomalacia suggesting 'pericellular digestion'<sup>28</sup>. Belanger created the term "osteocytic osteolysis" for the enlarged lacunae induced by parathyroid hormone or a by low-calcium diet. 'Osteocytic osteolysis' was viewed as being a feature of pathological conditions, especially due to high or continuous PTH.<sup>27,29,30</sup> The stimulating effects of parathyroid hormone on lysosomal vesicles in osteocytes was described in the 1970s<sup>31,32</sup> and in 1977, "perilacunar osteolysis" was described in rats sent into space<sup>33</sup> and alveolar bone of hibernating ground squirrels<sup>34</sup>. However, the number of publications began to decrease for various reasons<sup>35</sup> until technology had advanced sufficiently to address critics of this concept. Baylink and Wergedal had described Tartrate Resistant Acid Phosphatase, TRAP, activity in osteocytes in 1969<sup>36</sup>, which was criticized as being a diffusion artifact from osteoclasts but later validated by *in situ* hybridization, a technology no available in the 60s and 70s<sup>37</sup>.

Baylink also showed tetracycline binding to the perilacunar matrix, which led to the hypothesis that osteocytes can replace their perilacunar matrix<sup>38</sup> which was later reproduced in egg-laying hens<sup>39</sup>. This suggested that under non-pathological conditions osteocytes could remove and replace their perilacunar matrix. Qing and colleagues proposed that the term 'perilacunar modeling' be used in place of 'osteocytic osteolysis' for non-pathological conditions such as lactation<sup>40</sup>. These investigators showed an increase in lacunar area with lactation, that the PTH type 1 receptor was responsible and described a return to normal lacunar area with weaning. They showed that genes thought to be osteoclast specific such as TRAP and Cathepsin K were elevated in osteocytes during lactation and returned to normal with weaning. This study shows that *healthy* osteocytes can both remove and replace their perilacunar matrix thereby playing a role in mineral homeostasis during calcium demanding conditions. Recently it has been shown that the Calcitonin Receptor may also play a role by inhibiting perilacunar remodeling with lactation<sup>41</sup>.

As the PTH type 1 receptor is most highly expressed in osteocytes, the osteocyte may be the target of PTH in hyperparathyroidism and conversely, the positive effects of intermittent PTH on bone formation may also be due to effects on the osteocyte. The target of the therapeutic *Forteo* may be osteocytes in the mature skeleton.

### Bone Repair

Repair of microdamage and bone fatigue in bone is a normal, physiological process. Bone is constantly sustaining damage in the form of microcracks that is repaired by osteoclasts targeting the damaged bone. Osteoclasts are responsible for initiating a cutting cone, but how does the osteoclast know the location of the microdamage? It appears that the osteocyte

sends signals to the osteoclast providing information on where to resorb and where not to resorb bone (For review see<sup>42</sup>). Microdamage and bone fatigue is associated with of osteocyte apoptosis where an anti-apoptotic factor, BAX, is found in osteocytes around the cutting cone, while the pro-apoptotic factor, Bcl2, is found in osteocytes in the path of the cutting cone<sup>43</sup>. This suggests that the osteocytes in the path are undergoing programmed cell death, while those in the periphery are preserving viability<sup>44</sup>. A number of *in vitro* studies using MLO-Y4 osteocyte-like cells have investigated potential mechanisms. Apoptotic bodies are released by MLO-Y4 cells and primary osteocytes, but not osteoblasts<sup>45</sup>, serum starved MLO-Y4 cells will secrete soluble RANKL which is necessary for osteoclast formation<sup>46</sup>, and damaged MLO-Y4 cell networks in 3 dimensional gels express elevated RANKL and lower osteoprotegerin, OPG, an inhibitor of the RANK receptor<sup>47</sup>.

Pathological osteocyte cell death is associated with thiazolidinediones<sup>48</sup>, high dose alcohol<sup>49</sup>, and methotrexate used for cancer treatment<sup>50</sup>. Osteocytes express markers of apoptosis in response to withdrawal of estrogen<sup>51</sup>, to oxygen deprivation as occurs during immobilization<sup>52</sup>, and in response to glucocorticoid treatment<sup>53</sup>. TNF $\alpha$  and Interleukin-1 (IL-1) are potent inducers of osteocyte apoptosis<sup>54</sup>. Osteonecrosis, or dead bone, is due to osteocyte cell death but the mechanisms responsible are still debated. Aging is associated with increased numbers of empty osteocyte lacunae (See below). Therefore, a major research focus has been on osteocyte viability and approaches to prevent osteocyte cell death.

### **Osteocytes are endocrine cells**

Potentially osteoblasts have the capacity to release factors into the circulation, but they compose approximately 3–5% of bone cells compared to 1% osteoclasts, whereas 90–95% of bone cells in the adult human skeleton are osteocytes. It has not been appreciated that the total mass of osteocytes and their dendritic processes in bone that are equivalent to or greater than the mass of the brain<sup>55</sup>, therefore these cells are most likely a major source of circulating bone factors. Bone is highly vascularized and secretes factors such as FGF23 into the bloodstream to affect distant targets<sup>56</sup>, it must be defined as an endocrine organ<sup>2</sup>. Interestingly, FGF23 is also able to act on the parathyroid gland to decrease PTH secretion, identifying the parathyroid gland as another endocrine target of osteocyte signaling<sup>57,58</sup>. The vascular system has a close, connecting association with the osteocyte lacuna-canalicular system with its bone fluid. Osteocytes also produce other circulating factors such as sclerostin. Osteocytes may also target muscle (See below). It has recently been shown that two factors, prostaglandin E<sub>2</sub> and Wnt3a, both produced by osteocytes in response to shear stress support myogenesis and muscle function<sup>59–62</sup>. Therefore, mechanical loading of the skeleton especially in the form of exercise is important to ensure that osteocyte factors are released into the circulation.

In addition to cross talk with muscle, osteocytes may also send signals to hematopoietic cells. Studies showed that osteocytes and GPCR signaling were important in controlling myeloid cells proliferation<sup>63</sup> and mice lacking osteocytes were shown to have defective hematopoietic stem cell mobilization and lymphopenia<sup>64,65</sup>. Osteocyte may also have a role in regulating fat. Using a mouse model in which osteocytes can be ablated by use of

diphtheria toxin, it was shown that osteocytes may also regulate adipose tissue<sup>65</sup>. Studies have shown that sclerostin may play a role in inducing adipocyte differentiation<sup>66</sup>. Osteocytes may be an important reservoir of factors that target other unknown organs and tissues.

## Role of osteocytes in bone disease

### Osteoporosis

Estrogen deficiency, glucocorticoid treatment, oxidative stress caused by disuse and oxidative stress with aging may be responsible for osteocyte cell death and therefore bone fragility and osteoporosis<sup>67</sup>. As described above, osteocyte cell death is important for repair of damaged bone<sup>43,44,68</sup> so any condition that compromises osteocyte health and function most likely compromises the skeleton. A number of factors and cytokines have been shown to induce osteocyte cell death including glucocorticoids, IL-1 and TNFalpha and conversely, a number of molecules have been shown to protect osteocytes from cell death such as estrogen, Parathyroid Hormone, bisphosphonates (For review see<sup>69-71</sup>), and secreted muscle factors<sup>59</sup>. Osteocyte viability is crucial not only for the normal functioning of the skeleton but for other organs such as kidney as discussed above and muscle as discussed below. As osteocytes appear to have multiple, very important functions, it is important to maintain normal viability and function of these cells<sup>2</sup>.

### Aging

The osteocyte is a long lived, non-dividing, aging cell headed for senescence. While some osteocytes are removed from bone with remodeling many osteocytes can reside in human bone for decades in contrast to osteoblasts and osteoclasts that live for only days or weeks. Recently it has been shown that primary osteocytes from old 22 mo mice have 6 fold more telomere dysfunction-induced foci than osteocytes from young 6 mo old mice<sup>72</sup>. Senescent osteocytes predominately develop the Senescence Associated Secretory Phenotype, SASP, compared to other bone cells types which may contribute to age related bone loss. Once the cell dies, micropetrosis can result where mineral fills the lacuna resulting in a cell that becomes a 'living fossil'<sup>73</sup>. Fewer numbers of osteocyte lacunae were found in patients suffering from fractures compared to controls<sup>74</sup> and an age-dependent decrease occurs in osteocyte lacunar density with an increased amount of hypermineralized calcium phosphate occlusions caused by micropetrosis<sup>75</sup>. The aging and dying osteocyte in a compromised lacuno-canalicular system is less likely to produce secretory factors, less likely to repair bone, and less likely to respond to anabolic load. Therefore, it is important to maintain osteocyte viability with age.

### Hypophosphatemic Rickets

Several phosphate regulating hormones and enzymes, Phosphate Regulating Neutral Endopeptidase on Chromosome X, PHEX, Dentin Matrix Protein 1, DMP1, Matrix Extracellular phosphoglycoprotein MEPE and Fibroblast Growth Factor 23, FGF23, are mainly produced by late osteoblasts and by early and late osteocytes. PHEX is the earliest regulator of phosphate homeostasis to be expressed in the late osteoblast/early osteocyte. PHEX is the mutated gene in the Hyp mouse, which is also widely used as a model of X-



linked hypophosphatemic rickets<sup>76</sup> which leads to increased levels of FGF23 and hypophosphatemia<sup>77</sup>. Hypophosphatemic Rickets in humans is caused by inactivating mutations of *Pex* (*HYP Consortium Nat Gen 1995*). PHEX also interacts with DMP1 to regulate phosphate homeostasis by mechanisms that still have not been clearly elucidated. DMP1 is produced by early osteocytes and mice lacking DMP-1 are hypophosphatemic and have increased FGF23 levels<sup>56</sup>. Two teams of investigators independently showed that Autosomal Recessive Hypophosphatemic Rickets (ARHR) was caused by a mutation in DMP1 that affected FGF23 circulating levels<sup>56,78</sup>. Both PHEX and DMP1 are negative regulators of FGF23 but again the mechanisms have not been clearly determined. MEPE is predominantly expressed by osteocytes<sup>79</sup> and is not believed to act on FGF23 directly but through Phex<sup>80</sup>. The ASARM peptide of MEPE can bind to PHEX inhibiting its activity which results in an increase in FGF23<sup>80,81</sup>.

The molecule at the center of phosphate regulation is FGF23. FGF-23 was identified in 2000<sup>82</sup> as the phosphate-regulating hormone responsible for Autosomal Dominant Hypophosphatemic Rickets (ADHR), and for phosphate-wasting in Tumor-Induced-Osteomalacia (TIO) and X-linked hypophosphatemia (XLH). Osteocytes are also a main source of FGF23 and recent work showed that targeted ablation of FGF23 in bone cells, recapitulates the hypophosphatemia observed in FGF23 null mice<sup>83</sup>. Clinkenbeard<sup>84,85</sup> targeted FGF23 deletion using Col2.3-cre in osteoblasts and using DMP1-cre in early osteocytes and showed that most likely both osteoblasts and osteocytes are the physiological source of FGF23. FGF23 is not normally expressed at high levels in osteocytes in the healthy state but is dramatically upregulated in both DMP1 and PHEX associated hypophosphatemic rickets<sup>77</sup> and osteocytes appear to be the main source of the elevated circulating levels of FGF23.

### Genetic High and Low Bone Mass Diseases

Many of the mutations resulting in high or low bone mass are due to mutations in components of the wnt/b-catenin signaling pathway. One of the most well known are mutations in *SOST* which is highly expressed in mature osteocytes, but also expressed in articular chondrocytes (Hinton, 2009, Chan et al,2011). As stated above, the protein, sclerostin, is an inhibitor of bone formation and the gene, *Sost* is increased in response to unloading, decreased in response to loading, but also decreased in response to PTH<sup>86</sup>. Patients with sclerosteosis carry a point mutation in the *SOST* gene whereas patients with van Buchem disease are characterized by a 52kb deletion downstream of the gene<sup>87</sup>. Recently, craniodiaphyseal dysplasia, a rare and severe bone dysplasia characterized by sclerosis of the skull and facial bones has also been linked to a “de novo” mutation in the *SOST* gene<sup>88</sup>. The pathological role of mutations in sclerostin has been reproduced in knock-out and transgenic animal models that also show the high-bone mass phenotype of sclerosteosis and van Buchem patients<sup>26,89</sup>. Sclerostin binds to low-density lipoprotein (LDL)-related protein 5, 6 and 4 to inhibit Wnt/ $\beta$ -catenin signaling<sup>90</sup>.

Genetic mutations in the receptors for sclerostin have also been shown to result in bone disease. Loss-of-function of LRP5 results in the condition Osteoporosis Pseudoglioma, OPPG<sup>91</sup>. The condition is homozygous recessive and affected individuals had a Z-score of

–4.7. Conversely, gain-of-function of LRP5 results in highly increased bone mass<sup>92</sup>. The condition is autosomal dominant and individuals had BMDs 5–8, yet had normally shaped bones. These individuals never broke bones but could not float in water. Mutations in Lrp4 have also resulted in bone overgrowth<sup>93</sup>. Genetically modified mice have replicated these human phenotypes.

## Role of osteocytes in non-bone disease

### Chronic Kidney Disease

FGF23 secreted from osteocytes plays a pathological role in Chronic Kidney Disease, CKD. FGF23 is elevated in osteocytes in CKD<sup>94</sup> and serum levels of FGF23 are increased, particularly in the later stages of the disease<sup>95,96</sup>. FGF23 levels predict cardiovascular events before but not after dialysis and FGF23 is a risk factor for adverse outcomes in patients with CKD and End Stage Renal Disease (for review see<sup>97</sup>). Therefore, considerable effort is being put towards blocking or reducing FGF23 levels in these patients.

### Cardiac Function

High circulating levels of FGF23 have negative effects on cardiac muscle (For review see<sup>98</sup>). Elevated levels of circulating FGF23 have been linked to increased risk of heart disease and independently associated with left ventricular hypertrophy in human population studies<sup>99,100</sup>. Increased serum FGF23 has also been linked with impaired vascular function<sup>99</sup>, vascular calcification<sup>101</sup> and increased fat mass<sup>102</sup>.

### Sarcopenia?

It is not clear if osteoporosis and sarcopenia are concurrent or if one precedes the other. Dogma has been that the main interaction between muscle and bone is the mechanical loading of bone through muscle contraction. Osteocytes secrete factors that regulate muscle mass and function. MLO-Y4 osteocyte-like cells and primary osteocyte factors induce muscle myogenesis and activate the Wnt/ $\beta$ catenin pathway<sup>60,61</sup>. Two factors produced by osteocytes in response to shear stress, PGE<sub>2</sub> and Wnt3a, were found to enhance myogenesis and ex vivo primary muscle function<sup>62</sup>. The hypothesis that osteocytes can support myogenesis and muscle function is now supported by several lines of evidence.

Very recently, *in vivo* data has been published to support the concept that bone regulates muscle mass and function. In 2015 it was shown that osteocalcin partially restored muscle mass in a model of deletion of Cx43 in osteocytes<sup>103</sup>. In 2016 it was shown that osteocalcin can have positive effects on muscle mass<sup>104</sup>. Also in 2015, it was shown that cancer induced release of Transforming Growth Factor beta, TGF $\beta$  from bone was responsible for muscle cachexia<sup>105</sup>. That same year it was shown that deletion of a protease, MBTPS1, in osteocytes had little effect on bone but significantly increased muscle mass and function with aging<sup>106</sup>. A reduction in members of the TGF $\beta$  superfamily was observed. This suggests that bone produces osteocalcin which has positive effects on muscle, but also that bone and in particular osteocytes can produce negative regulators of muscle. It is not known if osteocyte dysfunction will play a role in sarcopenia.



Conversely, muscle also appears to secrete factors that affect osteocyte viability and function. In 2007 Pedersen coined the term 'myokines,' for muscle secreted factors opening the door to new concepts regarding muscle interaction with bone. Conversely, secreted factors from electrically stimulated skeletal muscle and from myotubes but not from myoblasts was shown to protect MLO-Y4 osteocytelike cells from dexamethasone induced cell death<sup>107</sup>. Other muscle factors are being identified that have positive effects on bone.

## Therapeutics targeting osteocyte factors

### Anti-sclerostin antibody

In animal studies, anti-sclerostin antibody has consistently been shown to increase bone mass. Inhibition of sclerostin by monoclonal antibody increases bone formation, bone mass, and bone strength in aged male rats<sup>108</sup>. Neutralizing antibody to Sclerostin is being developed as a therapeutic to treat osteoporosis (For review see<sup>109</sup>). The antibody blocks or reduces bone loss and supports bone formation-promotes fracture healing<sup>110</sup> and is a potential therapeutic for a number of conditions of low bone mass such as OI<sup>111</sup>. Romosozumab (AMG 785) has been tested in Phase 1 and II and Blososumab has been tested in Phase 1 clinical trial with good results. These antibodies increase bone mass to a greater extent than any previously developed therapeutic for osteoporosis including bisphosphonates, Forteo, and Denosumab. Phase III clinical trials in progress.

### Anti-RANKL antibody

MLO-Y4 osteocyte like cells support osteoclast formation<sup>112</sup> and apoptotic bodies released from MLO-Y4 cells express RANKL<sup>113</sup>. Primary osteocytes express RANKL<sup>114</sup>. Osteocytes express greater amounts of RANKL than osteoblasts and are better supporters of osteoclast formation<sup>115,116</sup>. Deletion of RANKL using the 10kb Dmp1-Cre results in mice with increased bone mass. This suggests that anti-RANKL antibody is mainly targeting osteocytes. Human anti-RANKL monoclonal antibody, Denosumab, is now available for treatment of osteoporosis. The antibody decreases osteoclast differentiation, function and survival. It reduces risk of spine, hip and nonvertebral fractures and does not require dose adjustment for decreased kidney function. For treatment of osteoporosis, SQ dosing every 6 months is applied and the effect is reversible within 6–12 months of stopping<sup>117</sup>.

### Anti-FGF 23 antibody

There are several conditions that could potentially benefit from treatment with anti-FGF23 antibody. In bone, Autosomal dominant hypophosphatemic rickets, caused by gain of function mutations in FGF23 that prevent proteolytic cleavage<sup>118</sup> and homozygous mutation in FAM20, a regulatory molecule of FGF23, resulting in hypophosphatemic osteomalacia<sup>119</sup> may benefit from anti-FGF23 antibody. In kidney, as FGF23 is elevated in osteocytes<sup>94</sup> and in serum in Chronic Kidney Disease, CKD<sup>95,96</sup>, these would also benefit from reducing FGF23 circulating levels. In heart disease, studies have linked raised levels of circulating FGF23 to an increased risk of heart disease, left ventricular hypertrophy in human population studies<sup>100,120</sup>, impaired vascular function, vascular calcification, and increased fat mass<sup>101</sup>. All of these conditions could potentially benefit from FGF receptor inhibitors and anti-FGF23 antibody. Treatment with FGF23 antibody restored serum phosphate levels

and corrects bone defects in the Hyp mouse and Dmp1 null model<sup>121</sup> and treatment with anti-FGF23 antibody KR23 increases serum phosphate in X-linked hypophosphatemic rickets<sup>122</sup>. (For review see<sup>123</sup>).

### **Other therapeutics to treat bone disease**

Could other therapeutics targeted to either osteoclasts or osteoblasts be also having effects on osteocytes? There is a distinct possibility. Calcitonin, bisphosphonates (Fosamax, Boniva, etc.), Anti-RANKL (Denosumab) and Cathepsin K inhibitors (odanacatinb) have been developed to target osteoclasts. However, bisphosphonates have also been shown to reduce osteocyte apoptosis<sup>69</sup>, osteocytes can also express Cathepsin K under certain conditions<sup>40</sup>, and as shown above, the anti-RANKL antibody may be targeting osteocytes. Hormone replacement therapy, Selective Estrogen Receptor Modulators (Evista), and Parathyroid Hormone peptides (Forteo) have been thought to mainly target osteoblasts, but these could also be having significant effects on osteocytes.

### **Are there other osteocyte factors?**

Much attention has focused on osteocalcin produced by osteoblasts. This bone specific factor has been shown to have effects on glucose metabolism, fertility, calcification and others<sup>124</sup>. The osteoblast has been ascribed the production of osteocalcin and the osteoclast as the releaser of uncarboxylated, the 'active' form of osteocalcin from the bone matrix to target other tissues. However, in the adult skeleton, osteoclasts make up less than 1% of bone cells, and osteoblasts less than 5%. Osteocytes have been shown to also produce osteocalcin so it will be interesting to see if osteocalcin production by osteocytes has a role in normal physiology and pathophysiology. Gene arrays have been performed on primary osteocytes and osteocyte cell lines. As with any gene arrays analysis, it is difficult to identify specific genes. It is likely that there are osteocyte factors responsible for normal function that may also play a role in disease that have not yet been identified.

### **Summary**

When normal physiological functions go awry, disorders and disease occurs. This is universal, and as discussed here, even for the osteocyte, a cell thought to not be in contact with the rest of the body. It was once thought that this cell was simply a place-holder in bone. The early functions proposed for this type of bone cell were mechanosensation and the capacity to remove their perilacunar matrix called "osteocytic osteolysis". Considerable skepticism existed even for these ascribed functions for decades. Within the last decade, the number of studies of osteocytes has dramatically increased leading to the discovery of novel functions of these cells. Along with these discoveries came the discoveries of how these cells can also be responsible for not only bone diseases and disorders, but also those of kidney, heart, and potentially muscle. Osteocytes have entered the realm of therapeutic targets for bone disease and now potentially kidney disease. It will be important not to overlook these cells with regards to the health of other systems and organs.

## Acknowledgments

The author is funded by NIH NIA grants PO1AG039355.

## References

1. Bonewald LF. The Amazing Osteocyte. *J Bone Miner. Res.* 2011; 26(2):229–238. [PubMed: 21254230]
2. Dallas SL, Prideaux M, Bonewald LF. The osteocyte: an endocrine cell ... and more. *Endocr. Rev.* 2013; 34(5):658–690. [PubMed: 23612223]
3. Wolff, J. *Das Gesetz der Transformation der Knochen.* Berlin: A Hirschwald; 1982.
4. Tatsumi S, Ishii K, Amizuka N, et al. Targeted ablation of osteocytes induces osteoporosis with defective mechanotransduction. *Cell metabolism.* 2007; 5(6):464–475. [PubMed: 17550781]
5. Weinbaum S, Cowin SC, Zeng Y. A model for the excitation of osteocytes by mechanical loading-induced bone fluid shear stresses. *J Biomech.* 1994; 27(3):339–360. [PubMed: 8051194]
6. Klein-Nulend J, van der Plas A, Semeins CM, et al. Sensitivity of osteocytes to biomechanical stress in vitro. *Faseb J.* 1995; 9(5):441–445. [PubMed: 7896017]
7. Klein-Nulend J, Semeins CM, Ajubi NE, Nijweide PJ, Burger EH. Pulsating fluid flow increases nitric oxide (NO) synthesis by osteocytes but not periosteal fibroblasts--correlation with prostaglandin upregulation. *Biochem Biophys Res Commun.* 1995; 217(2):640–648. [PubMed: 7503746]
8. Klein-Nulend J, Bakker AD, Bacabac RG, Vatsa A, Weinbaum S. Mechanosensation and transduction in osteocytes. *Bone.* 2013; 54(2):182–190. [PubMed: 23085083]
9. Kamel MA, Picconi JL, Lara-Castillo N, Johnson ML. Activation of beta-catenin signaling in MLOY4 osteocytic cells versus 2T3 osteoblastic cells by fluid flow shear stress and PGE (2): Implications for the study of mechanosensation in bone. *Bone.* 2010; 47(5):872–881. [PubMed: 20713195]
10. Cherian PP, Siller-Jackson AJ, Gu S, et al. Mechanical strain opens connexin 43 hemichannels in osteocytes: a novel mechanism for the release of prostaglandin. *Mol. Biol. Cell.* 2005; 16:3100–3106. [PubMed: 15843434]
11. Kitase Y, Jiang JX, Johnson ML, Bonewald LF. The Anti-Apoptotic Effects of Mechanical Strain on Osteocytes Are Mediated by PGE2 and Monocyte Chemoattractant Protein-3 (MCP-3); Selective protection by MCP-3 against Glucocorticoid (GC), but Not TNF- $\alpha$  Induced Apoptosis. *J. Bone Miner. Res.* 2006; 21(Suppl 1):S48.
12. Forwood MR. Inducible cyclo-oxygenase (COX-2) mediates the induction of bone formation by mechanical loading in vivo. *J Bone Miner. Res.* 1996; 11(11):1688–1693. [PubMed: 8915776]
13. Pead MJ, Lanyon LE. Indomethacin modulation of load-related stimulation of new bone formation in vivo. *Calcif Tissue. Int.* 1989; 45(1):34–40. [PubMed: 2504461]
14. Bonewald LF, Johnson ML. Osteocytes, mechanosensing and Wnt signaling. *Bone.* 2008; 42(4):606–615. [PubMed: 18280232]
15. Johnson ML, Picconi JL, Recker RR. The Gene for High Bone Mass. *The Endocrinologist.* 2002; 12:445–453.
16. Baron R, Kneissel M. WNT signaling in bone homeostasis and disease: from human mutations to treatments. *Nat. Med.* 2013; 19(2):179–192. [PubMed: 23389618]
17. Kramer I, Halleux C, Keller H, et al. Osteocyte Wnt/beta-catenin signaling is required for normal bone homeostasis. *Mol Cell Biol.* 2010; 30(12):3071–3085. [PubMed: 20404086]
18. Javaheri B, Dallas M, Zhao H, Bonewald L, Johnson M.  $\beta$ -catenin Haploinsufficiency in Osteocytes Abolishes the Osteogenic Effect of Mechanical Loading in Vivo. *J. Bone. Min Res.* 2011:S24.
19. Sawakami K, Robling AG, Pitner ND, et al. Site-specific osteopenia and decreased mechanoreactivity in Lrp5 mutant mice. *J. Bone. Min. Res.* 2004; 19:S1–S38.

20. Yang W, Lu Y, Kalajzic I, et al. Dentin matrix protein 1 gene cis-regulation: use in osteocytes to characterize local responses to mechanical loading in vitro and in vivo. *J Biol Chem.* 2005; 280(21):20680–20690. [PubMed: 15728181]
21. Gluhak-Heinrich J, Ye L, Bonewald LF, et al. Mechanical loading stimulates dentin matrix protein 1 (DMP1) expression in osteocytes in vivo. *J Bone Miner. Res.* 2003; 18(5):807–817. [PubMed: 12733719]
22. Gluhak-Heinrich J, Pavlin D, Yang W, Macdougall M, Harris SE. MEPE expression in osteocytes during orthodontic tooth movement. *Arch Oral Biol.* 2007; 52(7):684–690. [PubMed: 17270144]
23. Zhang K, Barragan-Adjemian C, Ye L, et al. E11/gp38 selective expression in osteocytes: regulation by mechanical strain and role in dendrite elongation. *Mol Cell Biol.* 2006; 26(12):4539–4552. [PubMed: 16738320]
24. Xiong J, Onal M, Jilka RL, Weinstein RS, Manolagas SC, O'Brien CA. Matrix-embedded cells control osteoclast formation. *Nat. Med.* 2011; 17(10):1235–1241. [PubMed: 21909103]
25. Robling AG, Niziolek PJ, Baldrige LA, et al. Mechanical stimulation of bone in vivo reduces osteocyte expression of Sost/sclerostin. *The Journal of biological chemistry.* 2008; 283(9):5866–5875. [PubMed: 18089564]
26. Lin C, Jiang X, Dai Z, et al. Sclerostin mediates bone response to mechanical unloading through antagonizing Wnt/beta-catenin signaling. *J Bone Miner. Res.* 2009; 24(10):1651–1661. [PubMed: 19419300]
27. Belanger LF, Belanger C, Semba T. Technical approaches leading to the concept of osteocytic osteolysis. *Clin Orthop Relat. Res.* 1967; 54:187–196. [PubMed: 4173071]
28. Recklinghausen FV. Untersuchungen uber Rachitis und Osteomalacia. Jena: Gustav Fischer. 1910
29. Belanger LF. Osteocytic osteolysis. *Calcif Tissue. Res.* 1969; 4(1):1–12. [PubMed: 4310125]
30. Belanger LF, Robichon J. Parathormone-induced osteolysis in dogs. A microradiographic and alphanradiographic survey. *J. Bone Joint Surg. Am.* 1964; 46:1008–1012. [PubMed: 14192496]
31. Bonucci E, Gherardi G. Osteocyte ultrastructure in renal osteodystrophy. *Virchows Arch A Pathol. Anat. Histol.* 1977; 373(3):213–231. [PubMed: 140505]
32. Bonucci E, Gherardi G, Mioni G, et al. Clinico-morphological correlations in uremic osteodystrophy of patients with conservative and hemodialytic treatment with special regard to the ultrastructure. *Minerva nefrologica.* 1975; 22(2–3):99–108. [PubMed: 1226257]
33. Iagodovskii VS, Trifanidi LA, Gorokhova GP. Effect of space flight on rat skeletal bones (an optical light and electron microscopic study). *Kosmicheskaja biologija i aviakosmicheskaja meditsina.* 1977; 11(1):14–20.
34. Haller AC, Zimny ML. Effects of hibernation on interradicular alveolar bone. *J Dent. Res.* 1977; 56(12):1552–1557. [PubMed: 277478]
35. Qing H, Bonewald L. Osteocyte Remodeling of the Perilacunar and Pericanalicular Matrix. *International Journal of Oral Science.* 2009; 1(2):59–65. [PubMed: 20687297]
36. Wergedal JE, Baylink DJ. Distribution of acid and alkaline phosphatase activity in undemineralized sections of the rat tibial diaphysis. *J Histochem Cytochem.* 1969; 17(12):799–806. [PubMed: 4189437]
37. Nakano Y, Toyosawa S, Takano Y. Eccentric localization of osteocytes expressing enzymatic activities, protein, and mRNA signals for type 5 tartrate-resistant acid phosphatase (TRAP). *J Histochem Cytochem.* 2004; 52(11):1475–1482. [PubMed: 15505342]
38. Baylink DJ, Wergedal JE. Bone formation by osteocytes. *Am J Physiol.* 1971; 221(3):669–678. [PubMed: 5570322]
39. Zambonin Zallone A, Teti A, Primavera MV, Pace G. Mature osteocytes behaviour in a repletion period: the occurrence of osteoplastic activity. *Basic and applied histochemistry.* 1983; 27(3):191–204. [PubMed: 6196018]
40. Qing H, Ardeshirpour L, Pajevic PD, et al. Demonstration of osteocytic perilacunar/canalicular remodeling in mice during lactation. *J. Bone Miner. Res.* 2012; 27(5):1018–1029. [PubMed: 22308018]
41. Clarke MV, Russell PK, Findlay DM, et al. A role for the calcitonin receptor to limit bone loss during lactation in female mice by inhibiting osteocytic osteolysis. *Endocrinology.* 2015 <http://dx.doi.org/10.1210/en.2015-1345>.

42. Schaffler MB, Cheung WY, Majeska R, Kennedy O. Osteocytes: Master Orchestrators of Bone. *Calcif. Tissue. Int.* 2013; 94(1):5–24. [PubMed: 24042263]
43. Verborgt O, Gibson GJ, Schaffler MB. Loss of osteocyte integrity in association with microdamage and bone remodeling after fatigue in vivo. *J Bone Miner. Res.* 2000; 15(1):60–67. [PubMed: 10646115]
44. Verborgt O, Tatton NA, Majeska RJ, Schaffler MB. Spatial distribution of Bax and Bcl-2 in osteocytes after bone fatigue: complementary roles in bone remodeling regulation? *J Bone Miner. Res.* 2002; 17(5):907–914. [PubMed: 12009022]
45. Kogianni G, Mann V, Noble BS. Apoptotic bodies convey activity capable of initiating osteoclastogenesis and localized bone destruction. *J Bone Miner. Res.* 2008; 23(6):915–927. [PubMed: 18435576]
46. Al-Dujaili SA, Lau E, Al-Dujaili H, Tsang K, Guenther A, You L. Apoptotic osteocytes regulate osteoclast precursor recruitment and differentiation in vitro. *J Cell Biochem.* 2011; 112(9):2412–2423. [PubMed: 21538477]
47. Mulcahy LE, Taylor D, Lee TC, Duffy GP. RANKL and OPG activity is regulated by injury size in networks of osteocyte-like cells. *Bone.* 2011; 48(2):182–188. [PubMed: 20854946]
48. Mabileau G, Mieczkowska A, Edmonds ME. Thiazolidinediones induce osteocyte apoptosis and increase sclerostin expression. *Diabet. Med.* 2010; 27(8):925–932. [PubMed: 20653751]
49. Maurel DB, Jaffre C, Rochefort GY, et al. Low bone accrual is associated with osteocyte apoptosis in alcohol-induced osteopenia. *Bone.* 2011; 49(3):543–552. [PubMed: 21689804]
50. Shandala T, Ng YS, Hopwood B, Yip YC, Foster BK, Xian CJ. The role of osteocyte apoptosis in cancer chemotherapy-induced bone loss. *J Cell Physiol.* 2011
51. Tomkinson A, Reeve J, Shaw RW, Noble BS. The death of osteocytes via apoptosis accompanies estrogen withdrawal in human bone. *Journal of Clinical Endocrinology & Metabolism.* 1997; 82(9):3128–3135. [PubMed: 9284757]
52. Dodd JS, Raleigh JA, Gross TS. Osteocyte hypoxia: a novel mechanotransduction pathway. *Am J Physiol.* 1999; 277(3 Pt 1):C598–C602. [PubMed: 10484347]
53. Weinstein RS, Nicholas RW, Manolagas SC. Apoptosis of osteocytes in glucocorticoid-induced osteonecrosis of the hip. *J Clin Endocrinol Metab.* 2000; 85(8):2907–2912. [PubMed: 10946902]
54. Pacifici R, Brown C, Puscheck E, et al. Effect of surgical menopause and estrogen replacement on cytokine release from human blood mononuclear cells. *Proc Natl Acad Sci U S A.* 1991; 88(12):5134–5138. [PubMed: 2052592]
55. Buenzli PR, Sims NA. Quantifying the osteocyte network in the human skeleton. *Bone.* 2015
56. Feng JQ, Ward LM, Liu S, et al. Loss of DMP1 causes rickets and osteomalacia and identifies a role for osteocytes in mineral metabolism. *Nat. Genet.* 2006; 38(11):1310–1315. [PubMed: 17033621]
57. Krajisnik T, Bjorklund P, Marsell R, et al. Fibroblast growth factor-23 regulates parathyroid hormone and 1 $\alpha$ -hydroxylase expression in cultured bovine parathyroid cells. *J Endocrinol.* 2007; 195(1):125–131. [PubMed: 17911404]
58. Ben-Dov IZ, Galitzer H, Lavi-Moshayoff V, et al. The parathyroid is a target organ for FGF23 in rats. *J Clin Invest.* 2007; 117(12):4003–4008. [PubMed: 17992255]
59. Jahn K, Lara-Castillo N, Brotto L, et al. Skeletal muscle secreted factors prevent glucocorticoid-induced osteocyte apoptosis through activation of beta-catenin. *Eur Cell Mater.* 2012; 24:197–210. [PubMed: 22972510]
60. Mo C, Romero-Suarez S, Bonewald L, Johnson M, Brotto M. Prostaglandin E2: from clinical applications to its potential role in bone-muscle crosstalk and myogenic differentiation. *Recent patents on biotechnology.* 2012; 6(3):223–229. [PubMed: 23092433]
61. Mo CL, Romero-Suarez S, Bonewald LF, Johnson ML, Brotto M. Evidence for Biochemical and Functional Communication from the MLO-Y4 Osteocyte-like Cell to the C2C12 Muscle Cells. *Cell Cycle.* 2012 In Submission.
62. Huang J, Mo C, Bonewald L, Brotto M. Wnt3a potentiates myogenesis in C2C12 myoblasts through changes of signaling pathways including Wnt and NF $\kappa$ B. *ASBMR 2014 Annual Meeting.* 2014; SU0190:s266.

63. Fulzele K, Krause DS, Panaroni C, et al. Myelopoiesis is regulated by osteocytes through Gsalphadependent signaling. *Blood*. 2013; 121(6):930–939. [PubMed: 23160461]
64. Asada N, Katayama Y, Sato M, et al. Matrix-embedded osteocytes regulate mobilization of hematopoietic stem/progenitor cells. *Cell Stem Cell*. 2013; 12(6):737–747. [PubMed: 23746979]
65. Sato M, Asada N, Kawano Y, et al. Osteocytes regulate primary lymphoid organs and fat metabolism. *Cell Metab*. 2013; 18(5):749–758. [PubMed: 24140021]
66. Ukita M, Yamaguchi T, Ohata N, Tamura M. Sclerostin Enhances Adipocyte Differentiation in 3T3-L1 Cells. *J Cell Biochem*. 2016; 117(6):1419–1428. [PubMed: 26553151]
67. Almeida M. Aging mechanisms in bone. *Bonekey. Rep*. 2012; 1
68. Kennedy OD, Herman BC, Laudier DM, Majeska RJ, Sun HB, Schaffler MB. Activation of resorption in fatigue-loaded bone involves both apoptosis and active pro-osteoclastogenic signaling by distinct osteocyte populations. *Bone*. 2012; 50(5):1115–1122. [PubMed: 22342796]
69. Bellido T, Plotkin LI. Novel actions of bisphosphonates in bone: Preservation of osteoblast and osteocyte viability. *Bone*. 2011; 49:50–55. [PubMed: 20727997]
70. Jilka RL, Noble B, Weinstein RS. Osteocyte apoptosis. *Bone*. 2013; 54(2):264–271. [PubMed: 23238124]
71. Jilka RL, O'Brien CA. The Role of Osteocytes in Age-Related Bone Loss. *Curr. Osteoporos. Rep*. 2016
72. Farr JN, Fraser DG, Wang H, et al. Identification of Senescent Cells in the Bone Microenvironment. *J Bone Miner. Res*. 2016
73. Bell LS, Kayser M, Jones C. The mineralized osteocyte: a living fossil. *American journal of physical anthropology*. 2008; 137(4):449–456. [PubMed: 18615594]
74. Qiu S, Rao DS, Palnitkar S, Parfitt AM. Reduced iliac cancellous osteocyte density in patients with osteoporotic vertebral fracture. *J. Bone Miner. Res*. 2003; 18(9):1657–1663. [PubMed: 12968675]
75. Qiu S, Rao DS, Palnitkar S, Parfitt AM. Age and distance from the surface but not menopause reduce osteocyte density in human cancellous bone. *Bone*. 2002; 31(2):313–318. [PubMed: 12151084]
76. Strom TM, Francis F, Lorenz B, et al. Pex gene deletions in Gy and Hyp mice provide mouse models for X-linked hypophosphatemia. *Hum Mol Genet*. 1997; 6(2):165–171. [PubMed: 9063736]
77. Liu S, Tang W, Zhou J, Vierthaler L, Quarles LD. Distinct Roles For Intrinsic Osteocyte Abnormalities and Systemic Factors in Regulation Of FGF23 and Bone Mineralization In Hyp Mice. *Am J Physiol Endocrinol Metab*. 2007
78. Lorenz-Depiereux B, Bastepe M, Benet-Pages A, et al. DMP1 mutations in autosomal recessive hypophosphatemia implicate a bone matrix protein in the regulation of phosphate homeostasis. *Nat Genet*. 2006; 38(11):1248–1250. [PubMed: 17033625]
79. Igarashi M, Kamiya N, Ito K, Takagi M. In situ localization and in vitro expression of osteoblast/osteocyte factor 45 mRNA during bone cell differentiation. *Histochem J*. 2002; 34(5):255–263. [PubMed: 12588003]
80. Liu S, Rowe PS, Vierthaler L, Zhou J, Quarles LD. Phosphorylated acidic serine-aspartate-rich MEPE-associated motif peptide from matrix extracellular phosphoglycoprotein inhibits phosphate regulating gene with homologies to endopeptidases on the X-chromosome enzyme activity. *J Endocrinol*. 2007; 192(1):261–267. [PubMed: 17210763]
81. Rowe PS, Garrett IR, Schwarz PM, et al. Surface plasmon resonance (SPR) confirms that MEPE binds to PHEX via the MEPE-ASARM motif: a model for impaired mineralization in X-linked rickets (HYP). *Bone*. 2005; 36(1):33–46. [PubMed: 15664000]
82. Consortium A. Autosomal dominant hypophosphataemic rickets is associated with mutations in FGF23. *Nat Genet*. 2000; 26(3):345–348. [PubMed: 11062477]
83. Kim SJ, Bieganski T, Sohn YB, et al. Identification of signal peptide domain SOST mutations in autosomal dominant craniodiaphyseal dysplasia. *Hum. Genet*. 2011; 129(5):497–502. [PubMed: 21221996]
84. Clinkenbeard EL, Cass TA, Ni P, et al. Conditional Deletion of Murine Fgf23: Interruption of the Normal Skeletal Responses to Phosphate Challenge and Rescue of Genetic Hypophosphatemia. *J Bone Miner. Res*. 2016; 31(6):1247–1257. [PubMed: 26792657]



85. Clinkenbeard EL, White KE. Systemic Control of Bone Homeostasis by FGF23 Signaling. *Curr Mol Biol. Rep.* 2016; 2(1):62–71. [PubMed: 27134818]
86. Bellido T, Ali AA, Gubrij I, et al. Chronic elevation of PTH in mice reduces expression of sclerostin by osteocytes: a novel mechanism for hormonal control of osteoblastogenesis. *Endocrinology.* 2005; 146(11):4577–4583. [PubMed: 16081646]
87. Balemans W, Patel N, Ebeling M, et al. Identification of a 52 kb deletion downstream of the SOST gene in patients with van Buchem disease. *Journal of medical genetics.* 2002; 39(2):91–97. [PubMed: 11836356]
88. Kim SJ, Bieganski T, Sohn YB, et al. Identification of signal peptide domain SOST mutations in autosomal dominant craniiodiaphyseal dysplasia. *Hum Genet.* 2011; 129(5):497–502. [PubMed: 21221996]
89. Li X, Ominsky MS, Niu QT, et al. Targeted deletion of the sclerostin gene in mice results in increased bone formation and bone strength. *J. Bone Miner. Res.* 2008; 23(6):860–869. [PubMed: 18269310]
90. Li X, Ominsky MS, Warmington KS, et al. Sclerostin antibody treatment increases bone formation, bone mass, and bone strength in a rat model of postmenopausal osteoporosis. *J Bone Miner. Res.* 2009; 24(4):578–588. [PubMed: 19049336]
91. Gong Y, Slee RB, Fukai N, et al. LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. *Cell.* 2001; 107(4):513–523. [PubMed: 11719191]
92. Little RD, Carulli JP, Del Mastro RG, et al. A Mutation in the LDL Receptor-Related Protein 5 Gene Results in the Autosomal Dominant High-Bone-Mass Trait. *Am. J. Hum. Genet.* 2002; 70(1):11–19. [PubMed: 11741193]
93. Leupin O, Piters E, Halleux C, et al. Bone overgrowth-associated mutations in the LRP4 gene impair sclerostin facilitator function. *J. Biol. Chem.* 2011; 286(22):19489–19500. [PubMed: 21471202]
94. Pereira RC, Juppner H, Azucena-Serrano CE, Yadin O, Salusky IB, Wesseling-Perry K. Patterns of FGF-23, DMP1, and MEPE expression in patients with chronic kidney disease. *Bone.* 2009; 45(6):1161–1168. [PubMed: 19679205]
95. Larsson T, Nisbeth U, Ljunggren O, Juppner H, Jonsson KB. Circulating concentration of FGF-23 increases as renal function declines in patients with chronic kidney disease, but does not change in response to variation in phosphate intake in healthy volunteers. *Kidney. Int.* 2003; 64(6):2272–2279. [PubMed: 14633152]
96. Imanishi Y, Inaba M, Nakatsuka K, et al. FGF-23 in patients with end-stage renal disease on hemodialysis. *Kidney. Int.* 2004; 65(5):1943–1946. [PubMed: 15086938]
97. Kovesdy CP, Alrifai A, Gosmanova EO, et al. Age and Outcomes Associated with BP in Patients with Incident CKD. *Clin J Am Soc Nephrol.* 2016; 11(5):821–831. [PubMed: 27103623]
98. Bonewald LF, Kiel DP, Clemens TL, et al. Forum on bone and skeletal muscle interactions: summary of the proceedings of an ASBMR workshop. *J Bone Miner. Res.* 2013; 28(9):1857–1865. [PubMed: 23671010]
99. Mirza MA, Larsson A, Lind L, Larsson TE. Circulating fibroblast growth factor-23 is associated with vascular dysfunction in the community. *Atherosclerosis.* 2009; 205(2):385–390. [PubMed: 19181315]
100. Faul C, Amaral AP, Oskouei B, et al. FGF23 induces left ventricular hypertrophy. *J Clin Invest.* 2011; 121(11):4393–4408. [PubMed: 21985788]
101. Desjardins L, Liabeuf S, Renard C, et al. FGF23 is independently associated with vascular calcification but not bone mineral density in patients at various CKD stages. *Osteoporos. Int.* 2011
102. Mirza MA, Alsio J, Hammarstedt A, et al. Circulating fibroblast growth factor-23 is associated with fat mass and dyslipidemia in two independent cohorts of elderly individuals. *Arterioscler Thromb Vasc Biol.* 2011; 31(1):219–227. [PubMed: 20966399]
103. Shen H, Grimston S, Civitelli R, Thomopoulos S. Deletion of connexin43 in osteoblasts/osteocytes leads to impaired muscle formation in mice. *J Bone Miner. Res.* 2015; 30(4):596–605. [PubMed: 25348938]

104. Mera P, Laue K, Ferron M, et al. Osteocalcin Signaling in Myofibers Is Necessary and Sufficient for Optimum Adaptation to Exercise. *Cell Metab.* 2016; 23(6):1078–1092. [PubMed: 27304508]
105. Waning DL, Mohammad KS, Reiken S, et al. Excess TGF-beta mediates muscle weakness associated with bone metastases in mice. *Nat. Med.* 2015; 21(11):1262–1271. [PubMed: 26457758]
106. Gorski JP, Huffman NT, Vallejo J, et al. Deletion of Mbtps1 (Pcsk8, S1p, Ski-1) Gene in Osteocytes Stimulates Soleus Muscle Regeneration and Increased Size and Contractile Force with Age. *J Biol Chem.* 2016; 291(9):4308–4322. [PubMed: 26719336]
107. Brakenhoff JPJ, Hart M, De Groot ER, Di Padova F, Aarden LA. Structure-function analysis of human IL-6: Epitope mapping of neutralizing monoclonal antibodies with amino- and carboxylterminal deletion mutants. *J. Immunol.* 1990; 145:561–568. [PubMed: 1694882]
108. Li X, Warmington KS, Niu QT, et al. Inhibition of sclerostin by monoclonal antibody increases bone formation, bone mass, and bone strength in aged male rats. *J Bone Miner. Res.* 2010; 25(12):2647–2656. [PubMed: 20641040]
109. Clarke BL. Anti-sclerostin antibodies: utility in treatment of osteoporosis. *Maturitas.* 2014; 78(3): 199–204. [PubMed: 24842796]
110. Morse A, Yu NY, Peacock L, et al. Endochondral fracture healing with external fixation in the Sost knockout mouse results in earlier fibrocartilage callus removal and increased bone volume fraction and strength. *Bone.* 2015; 71:155–163. [PubMed: 25445453]
111. Roschger A, Roschger P, Keplingter P, et al. Effect of sclerostin antibody treatment in a mouse model of severe osteogenesis imperfecta. *Bone.* 2014; 66:182–188. [PubMed: 24953712]
112. Zhao S, Zhang YK, Harris S, Ahuja SS, Bonewald LF. MLO-Y4 osteocyte-like cells support osteoclast formation and activation. *J Bone Miner. Res.* 2002; 17(11):2068–2079. [PubMed: 12412815]
113. Kogianni G, Mann V, Noble BS. Apoptotic bodies convey activity capable of initiating osteoclastogenesis and localised bone destruction. *J. Bone Miner. Res.* 2008; 23(6):915–927. [PubMed: 18435576]
114. Kramer I, Halleux C, Keller H, et al. Osteocyte Wnt/beta-catenin signaling is required for normal bone homeostasis. *Mol. Cell Biol.* 2010; 30(12):3071–3085. [PubMed: 20404086]
115. Xiong J, Onal M, Jilka RL, Weinstein RS, Manolagas SC, O'Brien CA. Matrix-embedded cells control osteoclast formation. *Nat. Med.* 2011; 17(10):1235–1241. [PubMed: 21909103]
116. Nakashima T, Hayashi M, Fukunaga T, et al. Evidence for osteocyte regulation of bone homeostasis through RANKL expression. *Nat. Med.* 2011; 17(10):1231–1234. [PubMed: 21909105]
117. Miller PD. A review of the efficacy and safety of denosumab in postmenopausal women with osteoporosis. *Therapeutic advances in musculoskeletal disease.* 2011; 3(6):271–282. [PubMed: 22870485]
118. Autosomal dominant hypophosphataemic rickets is associated with mutations in FGF23. *Nat Genet.* 2000; 26(3):345–348. [PubMed: 11062477]
119. Takeyari S, Yamamoto T, Kinoshita Y, et al. Hypophosphatemic osteomalacia and bone sclerosis caused by a novel homozygous mutation of the FAM20C gene in an elderly man with a mild variant of Raine syndrome. *Bone.* 2014; 67:56–62. [PubMed: 24982027]
120. Mirza MA, Hansen T, Johansson L, et al. Relationship between circulating FGF23 and total body atherosclerosis in the community. *Nephrol Dial Transplant.* 2009; 24(10):3125–3131. [PubMed: 19429932]
121. Aono Y, Yamazaki Y, Yasutake J, et al. Therapeutic effects of anti-FGF23 antibodies in hypophosphatemic rickets/osteomalacia. *J. Bone Miner. Res.* 2009; 24(11):1879–1888. [PubMed: 19419316]
122. Carpenter TO, Imel EA, Ruppe MD, et al. Randomized trial of the anti-FGF23 antibody KRN23 in X-linked hypophosphatemia. *J. Clin. Invest.* 2014; 124(4):1587–1597. [PubMed: 24569459]
123. Bonewald LF, Wacker MJ. FGF23 production by osteocytes. *Pediatr Nephrol.* 2013; 28(4):563–568. [PubMed: 22983423]
124. Wei J, Karsenty G. An overview of the metabolic functions of osteocalcin. *Reviews in endocrine & metabolic disorders.* 2015

125. Plotkin LI, Bellido T. Osteocytic signalling pathways as therapeutic targets for bone fragility. *Nature Reviews Endocrinology*. 2016
126. Yuan B, Takaiwa M, Clemens TL, et al. Aberrant Phex function in osteoblasts and osteocytes alone underlies murine X-linked hypophosphatemia. *J. Clin. Invest.* 2008; 118(2):722–734. [PubMed: 18172553]
127. Gowen LC, Petersen DN, Mansolf AL, et al. Targeted disruption of the osteoblast/osteocyte factor 45 gene (OF45) results in increased bone formation and bone mass. *J Biol Chem*. 2003; 278(3): 1998–2007. [PubMed: 12421822]
128. Addison WN, Nakano Y, Loisel T, Crine P, McKee MD. MEPE-ASARM peptides control extracellular matrix mineralization by binding to hydroxyapatite: an inhibition regulated by PHEX cleavage of ASARM. *J Bone Miner. Res.* 2008; 23(10):1638–1649. [PubMed: 18597632]
129. Clinkenbeard EL, Cass TA, Ni P, et al. Conditional Deletion of Murine Fgf23: Interruption of the Normal Skeletal Responses to Phosphate Challenge and Rescue of Genetic Hypophosphatemia. *J. Bone Miner. Res.* 2016
130. Divieti PP. PTH and osteocytes. *J. Musculoskelet. Neuronal. Interact.* 2005; 5(4):328–330. [PubMed: 16340124]
131. Winkler DG, Sutherland MK, Geoghegan JC, et al. Osteocyte control of bone formation via sclerostin, a novel BMP antagonist. *EMBO J.* 2003; 22(23):6267–6276. [PubMed: 14633986]
132. Van Bezooijen RL, Roelen BA, Visser A, et al. Sclerostin is an osteocyte-expressed negative regulator of bone formation, but not a classical BMP antagonist. *J. Exp. Med.* 2004; 199(6):805–814. [PubMed: 15024046]
133. Bonewald LF, Johnson ML. Osteocytes, mechanosensing and Wnt signaling. *Bone*. 2008; 42(4): 606–615. [PubMed: 18280232]
134. Duan P, Bonewald LF. The role of the wnt/beta-catenin signaling pathway in formation and maintenance of bone and teeth. *Int J Biochem Cell Biol.* 2016; 77(Pt A):23–29. [PubMed: 27210503]
135. Klein-Nulend J, Van Der Plas A, Semeins CM, et al. Sensitivity of osteocytes to biomechanical stress in vitro. *FASEB J.* 1995; 9(5):441–445. [PubMed: 7896017]
136. Lara-Castillo N, Kim-Weroha NA, Kamel MA, et al. In vivo mechanical loading rapidly activates beta-catenin signaling in osteocytes through a prostaglandin mediated mechanism. *Bone*. 2015; 76:58–66. [PubMed: 25836764]
137. Cheng B, Kato Y, Zhao S, et al. PGE(2) is essential for gap junction-mediated intercellular communication between osteocyte-like MLO-Y4 cells in response to mechanical strain. *Endocrinology*. 2001; 142(8):3464–3473. [PubMed: 11459792]
138. Lieben L, Masuyama R, Torrekens S, et al. Normocalcemia is maintained in mice under conditions of calcium malabsorption by vitamin D-induced inhibition of bone mineralization. *J. Clin. Invest.* 2012; 122(5):1803–1815. [PubMed: 22523068]
139. Torreggiani E, Matthews BG, Pejda S, et al. Preosteocytes/osteocytes have the potential to dedifferentiate becoming a source of osteoblasts. *PLoS. One*. 2013; 8(9):e75204. [PubMed: 24040401]
140. Brotto M, Bonewald LF. Bone and Muscle: Interactions beyond Mechanical. *Bone*. 2015 in press.
141. Gorski JP, Huffman NT, Vallejo J, et al. Deletion of Mbtps1 (PCSK8, S1P, SKI-1) in Osteocytes Stimulates Soleus Muscle Regeneration and Increased Size and Contractile Force with Age. *J. Biol. Chem.* 2015
142. Touchberry CD, Green TM, Tchikrizov V, et al. FGF23 is a novel regulator of intracellular calcium and cardiac contractility in addition to cardiac hypertrophy. *American Journal of Physiology - Endocrinology and Metabolism*. 2013; 304(8):E863–E873. [PubMed: 23443925]
143. Singh S, Grabner A, Yanucil C, et al. Fibroblast growth factor 23 directly targets hepatocytes to promote inflammation in chronic kidney disease. *Kidney. Int.* 2016
144. Lau KW, Rundle CH, Zhou XD, Baylink DJ, Sheng MH. Conditional deletion of IGF-I in osteocytes unexpectedly accelerates bony union of the fracture gap in mice. *Bone*. 2016; 92:18–28. [PubMed: 27519969]

145. Sheng MH, Zhou XD, Bonewald LF, Baylink DJ, Lau KH. Disruption of the insulin-like growth factor-1 gene in osteocytes impairs developmental bone growth in mice. *Bone*. 2013; 52(1):133–144. [PubMed: 23032105]
146. Lim J, Shi Y, Karner CM, et al. Dual function of *Bmpr1a* signaling in restricting preosteoblast proliferation and stimulating osteoblast activity in mouse. *Development*. 2016; 143(2):339–347. [PubMed: 26657771]
147. Canalis E, Bridgewater D, Schilling L, Zanotti S. Canonical Notch activation in osteocytes causes osteopetrosis. *Am J Physiol Endocrinol Metab*. 2016; 310(2):E171–E182. [PubMed: 26578715]
148. Kondoh S, Inoue K, Igarashi K, et al. Estrogen receptor alpha in osteocytes regulates trabecular bone formation in female mice. *Bone*. 2014; 60:68–77. [PubMed: 24333171]
149. Windahl SH, Borjesson AE, Farman HH, et al. Estrogen receptor-alpha in osteocytes is important for trabecular bone formation in male mice. *Proc. Natl. Acad. Sci. U. S. A.* 2013; 110(6):2294–2299. [PubMed: 23345419]
150. Zhou JZ, Riquelme MA, Gu S, et al. Osteocytic connexin hemichannels suppress breast cancer growth and bone metastasis. *Oncogene*. 2016

**Key Points**


- Within the last decade, the number of studies of osteocytes has dramatically increased leading to the discovery of novel functions of these cells.
- Along with these discoveries came the discoveries of how these cells can also be responsible for not only bone diseases and disorders, but also those of kidney, heart, and potentially muscle.
- Osteocytes have entered the realm of therapeutic targets for bone disease and now potentially kidney disease.
- It will be important not to overlook these cells with regards to the health of other systems and organs.

**Box 1****Normal Functions of Osteocytes**

- Control mineralization through **Phex**<sup>126</sup>, **Dmp1**<sup>56</sup>, and **MEPE**<sup>127,128</sup>
- Regulate phosphate homeostasis through **FGF23**<sup>56,129</sup>
- Play a role in calcium homeostasis in response to PTH/PTHrP<sup>40,130</sup>
- Can recruit osteoclasts through expression of **RANKL** with or without cell death<sup>112,115,116</sup>
- Can regulate osteoblast activity through **Sclerostin**<sup>131,132</sup>
- Are mechanosensory cells through  $\beta$ -catenin signaling<sup>133,134</sup>
- Have autocrine/paracrine effects through **prostaglandin** production<sup>135–137</sup>.
- Under calcium restriction, osteocytes remove calcium from bone through the Vitamin D Receptor<sup>138</sup>
- Osteocytes regulate myelopoiesis/hematopoiesis through **G-CSF**<sup>63</sup>
- **G-CSF** targets osteocytes that mediate mobilization of Hematopoietic Stem/Progenitor Cells and is prevented by surgical sympathectomy<sup>64</sup>
- Osteocytes regulate primary lymphoid organs and fat metabolism<sup>65</sup>
- Osteocytes can dedifferentiate to become a source of matrix-producing osteoblasts<sup>139</sup>
- Can increase muscle myogenesis and muscle function<sup>60,62,140</sup> and can inhibit muscle mass with aging<sup>141</sup>
- Can have effects on heart<sup>123,142</sup> and liver<sup>143</sup> through FGF23.
- Play a role in fracture healing through IGF-1<sup>144,145</sup>
- Regulate bone formation through Bmpr1a signaling<sup>146</sup>, Notch activation<sup>147</sup>, and ER $\alpha$  signaling<sup>148,149</sup>
- Suppress breast cancer growth and bone metastasis<sup>150</sup>

Data from references 1, 2, 125.





**Figure 1.**  
Defective Osteocyte Function and Disease.