


# Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: C. Virumbrales, S. Suárez-Pantiga, M. Solas, M. Á. Fernández-Rodríguez and R. Sanz, *Org. Biomol. Chem.*, 2018, DOI: 10.1039/C8OB00406D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## Gold(I)-catalyzed diastereoselective synthesis of 1- $\alpha$ -oxybenzyl-1*H*-indenes†

Received 00th January 20xx,  
Accepted 00th January 20xx

Cintia Virumbrales, Samuel Suárez-Pantiga, Marta Solas, Manuel A. Fernández-Rodríguez and Roberto Sanz\*


DOI: 10.1039/x0xx00000x

www.rsc.org/

The gold(I)-catalyzed oxycyclization of  $\beta$ -aryl monosubstituted *o*-(alkynyl)styrenes gives rise to 1- $\alpha$ -methoxy or 1- $\alpha$ -hydroxybenzyl-1*H*-indenes in a diastereospecific way. In contrast with  $\beta$ , $\beta$ -disubstituted *o*-(alkynyl)styrenes, the stereochemical outcome of the process, diastereospecific reactions, supports the higher contribution of a gold intermediate with cyclopropylcarbene-like character.

Gold-catalyzed cycloisomerization reactions of 1,*n*-enynes have received a good deal of attention in the last years providing versatile routes to interesting carbocyclic structures not easily prepared by classical methods.<sup>1</sup> In this context, 1,5-enyne cycloisomerizations involving almost always 5-*endo-dig* cyclizations<sup>2</sup> have been less developed, despite their synthetic usefulness for the preparation of interesting cyclopentene derivatives.<sup>3</sup> In this field, our group and others have been specially interested in the study of the reactivity of *o*-(alkynyl)styrenes,<sup>4</sup> as a particular type of 1,3-dien-5-ynes.<sup>5</sup> Although most of the previously reported cycloisomerizations of *o*-(alkynyl)styrenes led to naphthalene derivatives through formal 6-*endo* cyclizations,<sup>6</sup> considering Gagosz and co-workers' report about the synthesis of cyclopentenes through 5-*endo* alkoxy cyclization of 3-acetoxy-1,5-enynes,<sup>7</sup> we first demonstrated that the corresponding 5-*endo* cyclization pathway could be induced by a proper selection of the olefin substitution pattern. We found that  $\beta$ , $\beta$ -disubstituted *o*-(alkynyl)styrenes undergo selectively 5-*endo* cyclizations, using cationic gold complexes bearing triphenyl phosphine as ligand both in the presence or absence of silver salts.<sup>8</sup> In this way, indene derivatives, which are privileged scaffolds mainly due to their applications in materials science and as ligands for metallic catalysts,<sup>9</sup> are easily accessed from simple starting materials.<sup>10</sup> After our seminal work, different authors have reported related 5-*endo* cyclizations of  $\beta$ , $\beta$ -disubstituted *o*-(alkynyl)styrenes under gold- and other transition metal-

catalysis.<sup>11</sup> Using these  $\beta$ , $\beta$ -disubstituted *o*-(alkynyl)styrenes, and external oxygen nucleophiles, we have been able to prepare 1-alkoxy- or 1-hydroxy-dialkylmethyl-1*H*-indenes. However, when these starting materials possess two different groups at the  $\beta$ -position the stereoselectivity of the cyclization had not been properly established in our previous work, and we assumed that the oxycyclization was not stereospecific likely due to a high contribution of a gold-stabilized homoallylic carbocation species as intermediate (Scheme 1a).<sup>8c</sup> At this point, we decided to explore the gold-catalyzed nucleophilic additions to  $\beta$ -aryl-monosubstituted *o*-(alkynyl)styrenes.<sup>12</sup> In this case two major issues, regio- and stereoselectivity, should be addressed. Firstly, for the initial



**Scheme 1** Gold(I)-catalyzed alkoxy cyclization of  $\beta$ , $\beta$ -disubstituted *o*-(alkynyl)styrenes (previous work) and  $\beta$ -monosubstituted *o*-(alkynyl)styrenes (this work).

Área de Química Orgánica, Departamento de Química, Facultad de Ciencias, Universidad de Burgos, Pza. Misael Bañuelos s/n, 09001 Burgos, Spain.  
E-mail: rsd@ubu.es

† Electronic Supplementary Information (ESI) available: Experimental details, characterization data, NMR spectra of all products. CCDC 1585617 for **3i**. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/x0xx00000x

olefin attack, triggered by the gold-activation of the alkyne, a regioisomeric 6-*endo* cyclization could be competitive with the previously reported 5-*endo* as the positive charge would be located at secondary carbons in the carbocationic-like intermediate for both types of ring closures (Scheme 1b). Secondly, the stereochemical outcome of these reactions, i.e. stereospecific or non-stereospecific nature of the process, could contribute to shed light about the controversial character of the involved gold intermediates, which could be also influenced by the nature of the ligands on the gold catalyst.<sup>13</sup> Herein, we want to report our results about the nucleophilic additions to  $\beta$ -aryl-monosubstituted *o*-(alkynyl)styrenes under gold-catalysis, which take place selectively through 5-*endo* cyclizations and in a diastereospecific manner leading to a new family of functionalized indene derivatives (Scheme 1c).


To perform our goal, we selected (*E*)- $\beta$ -phenyl *o*-(alkynyl)styrene **1a** as model substrate and treated it with an excess of MeOH in CH<sub>2</sub>Cl<sub>2</sub> under the catalysis of a selection of gold complexes (Table 1). In a first attempt, **1a** was treated with Ph<sub>3</sub>PAuNTf<sub>2</sub> and after 1 h we observed the partial but stereospecific formation of 1- $\alpha$ -methoxybenzyl-1*H*-indene **2a** (entry 1). A similar result was obtained when using the catalytic system Ph<sub>3</sub>PAuCl/AgSbF<sub>6</sub> that we had previously used for the methoxycyclization of  $\beta,\beta$ -disubstituted *o*-(alkynyl)styrenes (entry 2). A slightly better result was displayed by gold catalyst bearing a phosphite ligand (entry 3), whereas complete conversions were obtained when gold complexes with bulky biarylphosphines as ligands were employed (entries 4–6). However, the best yield was obtained using the NHC-gold(I) complex IPrAuNTf<sub>2</sub>, possessing the weakly coordinating counteranion Tf<sub>2</sub>N<sup>-</sup> and a highly electron-donating ligand (entry 7). Moreover, the catalyst load could be

halved without a serious impact on the reaction outcome (entry 8), supporting its remarkably high activity for a wide variety of Au-catalyzed transformations that had been elegantly shown by Gagosz and co-workers.<sup>14</sup> A brief silver salt screening showed no beneficial silver effect or improvement for this process (entries 9–11). Remarkably, in all the tests only one diastereoisomer of indene **2a** was observed in the <sup>1</sup>H NMR of the crude reaction mixture, whose stereochemistry was initially assigned based on previous literature reports.<sup>7a</sup>

Echavarren and co-workers had established that gold complexes with highly electron-donating ligands, such as NHCs, enhance the carbene-like character of the gold intermediates in cyclizations of 1,5-enynes.<sup>13</sup> Considering this fact in the context of the ligand effects in gold catalysis,<sup>15</sup> these preliminary results seem to support that the reactions of these  $\beta$ -aryl monosubstituted *o*-(alkynyl)styrenes **1** would take place through an intermediate that requires a higher retrodonation of gold to stabilize the initially generated carbocation and so, they proceed more efficiently with complexes bearing donor ligands.


With the optimized reaction conditions in hand, we proceeded to evaluate the behaviour of a series of  $\beta$ -aryl *o*-(alkynyl)styrenes **1**. As shown in Table 2, the transformation was suitable for a variety of starting substrates **1a–e** bearing different substituents at the triple bond, including (hetero)aromatic, alkyl, alkenyl, and arylthio groups (entries 1–7). Interestingly, starting from geometrically pure *E*-isomers of the *o*-(alkynyl)styrenes **1a–e**, the corresponding methoxybenzyl indenenes **2a–e** were obtained in high yields, and as single diastereoisomers in all the cases, showing the potential of this reaction as a wide variety of groups are allowed to be present in the starting alkyne. In addition, we have also checked that the stereoinformation of the alkene is finely transferred to the final cyclized product, as using **1a** as a ca. 1/1 mixture of geometrical isomers led to the corresponding indene **2a** as a ca. 1/1 mixture of diastereoisomers (entry 2). Moreover, using (*Z*)-**1b** as starting material the other diastereoisomer of **2b** (**diast-2b**) was selectively formed (entries 3 vs. 4), thus proving the stereospecificity of the reaction. Regarding the aryl substituent at the  $\beta$ -position of the olefin, *p*-tolyl- as well as 1-naphthyl-substituted *o*-(alkynyl)styrenes **1f–i** behaved the same way as parent **1a** providing high yields of the corresponding indene derivatives **2f–i** and with complete control of the stereochemical outcome of the reactions determined by the stereochemistry of the starting enyne (entries 8–11). Moreover, substrates bearing functionalized aryl groups, such as **1j** and **1k**, also methoxycyclized leading to the corresponding indenenes **2j** and **2k** in good yields and in a diastereospecific manner (entries 12–13). Interestingly, with *o*-(alkynyl)styrene **1k** possessing a butyl group as alkyne substituent the methoxycyclization of the *E*-isomer was found to be slower than the *Z*-isomer leading, in the first case, to competitive addition of methanol to the triple bond (entry 13). Finally, *o*-(alkynyl)styrenes **1l** and **1m**, functionalized with an electron-rich 4-methoxyphenyl group at the  $\beta$ -position, were

**Table 1** Effect of the gold catalyst on the methoxycyclization of (*E*)- $\beta$ -phenyl *o*-(alkynyl)styrene **1a**<sup>a</sup>



Entry	Catalyst	Conversion (%) <sup>b</sup>	Yield (%) <sup>c</sup>
1	Ph <sub>3</sub> PAuNTf <sub>2</sub>	76	51
2	Ph <sub>3</sub> PAuCl/AgSbF <sub>6</sub>	66	54
3	[(3,5- <i>t</i> -Bu) <sub>2</sub> C <sub>6</sub> H <sub>3</sub> O <sub>3</sub> ] <sub>3</sub> AuCl/AgNTf <sub>2</sub>	97	65
4	XPhosAuNTf <sub>2</sub>	100	76
5	BrettPhosAuNTf <sub>2</sub>	100	62
6	JohnPhosAuNTf <sub>2</sub>	100	84
7	IPrAuNTf <sub>2</sub>	100	89
8 <sup>d</sup>	IPrAuNTf <sub>2</sub>	100	84
9	IPrAuCl/AgNTf <sub>2</sub>	100	63
10	IPrAuCl/AgSbF <sub>6</sub>	100	79
11	IPrAuCl/AgBF <sub>4</sub>	100	86

<sup>a</sup> Reactions conducted using 0.1 mmol of **1a** in CH<sub>2</sub>Cl<sub>2</sub> (0.8 mL). <sup>b</sup> Determined by <sup>1</sup>H NMR analysis of the crude reaction mixture. <sup>c</sup> Determined by <sup>1</sup>H NMR using CH<sub>2</sub>Br<sub>2</sub> as internal standard. When the NMR yield is significantly lower than the conversion several unidentified products were observed. <sup>d</sup> Carried out with 2.5 mol% of catalyst.

**Table 2** Gold-catalyzed methoxycyclization of  $\beta$ -aryl *o*-(alkynyl)styrenes **1**. Diastereospecific synthesis of 1-methoxybenzyl-1*H*-indenes **2**<sup>a</sup>


Entry	<b>1</b>	Ar	R	<i>E/Z</i>	<b>2</b>	d.r. <sup>b</sup>	Yield (%) <sup>c</sup>
1	<b>1a</b>	Ph	Ph	>20/1	<b>2a</b>	>20/1	93
2	<b>1a</b>	Ph	Ph	1/1	<b>2a</b>	1/1	86
3	<b>1b</b>	Ph	<i>n</i> -Bu	>20/1	<b>2b</b>	>20/1	91
4	<b>1b</b>	Ph	<i>n</i> -Bu	<1/20	<b>diast-2b</b>	>20/1	87
5 <sup>d</sup>	<b>1c</b>	Ph	<i>c</i> -C <sub>6</sub> H <sub>9</sub> <sup>e</sup>	>20/1	<b>2c</b>	>20/1	82
6	<b>1d</b>	Ph	3-Th <sup>f</sup>	>20/1	<b>2d</b>	>20/1	87
7	<b>1e</b>	Ph	SPh	>20/1	<b>2e</b>	>20/1	90
8	<b>1f</b>	4-MeC <sub>6</sub> H <sub>4</sub>	Ph	>20/1	<b>2f</b>	>20/1	86
9	<b>1g</b>	4-MeC <sub>6</sub> H <sub>4</sub>	<i>n</i> -Bu	>20/1	<b>2g</b>	>20/1	75
10	<b>1h</b>	1-naphthyl	Ph	>20/1	<b>2h</b>	>20/1	88
11	<b>1i</b>	1-naphthyl	<i>n</i> -Bu	>20/1	<b>2i</b>	>20/1	91
12	<b>1j</b>	4-ClC <sub>6</sub> H <sub>4</sub>	Ph	>20/1	<b>2j</b>	>20/1	85
13 <sup>g</sup>	<b>1k</b>	4-ClC <sub>6</sub> H <sub>4</sub>	<i>n</i> -Bu	<1/20	<b>diast-2k</b>	>20/1	88
14	<b>1l</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Ph	>20/1	<b>2l</b>	16/1	86
15	<b>1l</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Ph	1/8	<b>2l</b>	1/5	83
16	<b>1m</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	<i>n</i> -Bu	9/1	<b>2m</b>	9/1	70


<sup>a</sup> Reactions conducted using 0.3 mmol of *o*-(alkynyl)styrene derivative **1** in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) at RT. <sup>b</sup> Determined by <sup>1</sup>H NMR analysis of the crude reaction mixture. <sup>c</sup> Isolated yield of product based on starting material **1**. <sup>d</sup> Reaction time = 48 h. <sup>e</sup> Cyclohexen-1-yl. <sup>f</sup> 3-Thienyl. <sup>g</sup> Reaction time = 4 h. When the reaction was carried out with (*E*)-**1k** sub-products derived from competitive attack of MeOH to the triple bond were also formed, although **2k** was also formed with d.r. >20/1.

subjected to the standard conditions giving rise to the methoxy-functionalized indene derivatives **2l** and **2m** in high yields (entries 14–16). The reactions of **1l** take place with a minor loss of the stereospecificity (entries 14–15), likely due to a competitive contribution of the carbocationic like-character of the gold intermediate considering the ability of the 4-methoxyphenyl group to stabilize carbocations at the benzylic position. A similar observation had been reported by Echavarren et al. in the alkoxylation of 1,6-enynes.<sup>16</sup>

As we had previously demonstrated that water could also be a useful nucleophile for the hydroxycyclization reactions of  $\beta,\beta$ -disubstituted *o*-(alkynyl)styrenes,<sup>8c</sup> we decided to check if this process also works with  $\beta$ -aryl *o*-(alkynyl)styrenes **1**. After some experimentation with model **1a** we found that the use of a mixture of CH<sub>2</sub>Cl<sub>2</sub> and dioxane as solvent allows external water to participate in the cyclization leading to the formation of 1-( $\alpha$ -hydroxybenzyl)-1*H*-indene **3a** that was obtained in good yield and as a single diastereoisomer (Table 3, entry 1). Remarkably, in the optimization study for this process we observed no hydroxycyclization for **1a** in the presence of Ph<sub>3</sub>PAuNTf<sub>2</sub> or (PhO)<sub>3</sub>PAuCl/AgSbF<sub>6</sub> as catalysts,<sup>17</sup> showing again the superior activity of the gold complex bearing the NHC ligand for the cyclizations of  $\beta$ -monosubstituted *o*-(alkynyl)styrenes. Taking into account the potential synthetic interest of 1-( $\alpha$ -hydroxybenzyl)-1*H*-indenes for further transformations, we decided to prepare a variety of hydroxycyclized products **3**. A selection of starting (*E*)- $\beta$ -aryl *o*-(alkynyl)styrenes **1** efficiently underwent the

hydroxycyclization under the standard conditions, resulting in formation of the expected alcohols **3** in high yields and in a stereospecific manner. As established above, the nucleophilic addition of water to (*E*)-**1a** afforded one diastereoisomer of **3a**, whereas starting from a mixture of geometrical isomers of **1a** led to hydroxyindene derivative **3a** as a mixture of diastereoisomers (entries 1 vs. 2). A similar result was observed for *o*-(alkynyl)styrene **1b** bearing a butyl group as substituent of the triple bond (entries 3 vs. 4). Starting enynes **1c–e**, possessing different groups at the triple bond, also efficiently underwent the hydroxycyclization reaction giving rise to the corresponding alcohols as single diastereoisomers although with **1e** no complete conversion was achieved under these conditions (entries 5–7). In addition, selected starting *o*-(alkynyl)styrenes **1**, bearing different aryl groups at the  $\beta$ -position of the olefin, **1h,i,l,m** also provided access to the corresponding 1-( $\alpha$ -hydroxybenzyl)indenes **3** in good yields (entries 8–11), although with butyl-substituted starting *o*-(alkynyl)styrene **1i** complete conversion was again not achieved (entry 9). Analogous to what was observed in its methoxycyclization, when *o*-(alkynyl)styrene **1l**, bearing an electron-rich 4-methoxyphenyl group as aryl substituent of the alkene, was used as starting material the diastereospecificity was practically lost and the corresponding hydroxyindene derivative **3l** was obtained as a mixture of diastereoisomers (entry 10). Moreover, in order to unambiguously establish the relative stereochemistry of the diastereoisomer generated in these gold-catalyzed methoxy- and hydroxycyclizations of *o*-(alkynyl)styrenes **1**, the structure of **3i** was further assigned by X-ray analysis.<sup>18</sup>

Having determined that the methoxycyclization reaction of  $\beta$ -aryl *o*-(alkynyl)styrenes **1** is stereospecific, i.e. the stereoinformation is completely transferred in most of the

**Table 3** Gold-catalyzed hydroxycyclization of  $\beta$ -aryl *o*-(alkynyl)styrenes **1**. Synthesis of 1-(1-hydroxybenzyl)-1*H*-indenes **3**<sup>a</sup>


Entry	<b>1</b>	Ar	R	<i>E/Z</i>	<b>3</b>	d.r. <sup>b</sup>	Yield (%) <sup>c</sup>
1 <sup>d</sup>	<b>1a</b>	Ph	Ph	>20/1	<b>3a</b>	>20/1	76
2	<b>1a</b>	Ph	Ph	1/1	<b>3a</b>	1/1	76
3	<b>1b</b>	Ph	<i>n</i> -Bu	>20/1	<b>3b</b>	>20/1	77
4	<b>1b</b>	Ph	<i>n</i> -Bu	1/1	<b>3b</b>	1/1	72
5	<b>1c</b>	Ph	<i>c</i> -C <sub>6</sub> H <sub>9</sub> <sup>e</sup>	>20/1	<b>3c</b>	>20/1	70
6	<b>1d</b>	Ph	3-Th <sup>f</sup>	>20/1	<b>3d</b>	>20/1	75
7	<b>1e</b>	Ph	SPh	>20/1	<b>3e</b>	>20/1	60 <sup>g</sup>
8	<b>1h</b>	1-naphthyl	Ph	>20/1	<b>3h</b>	>20/1	79
9	<b>1i</b>	1-naphthyl	<i>n</i> -Bu	>20/1	<b>3i</b>	>20/1	53 <sup>h</sup>
10	<b>1l</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Ph	>20/1	<b>3l</b>	3.5/1	65
11	<b>1m</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	<i>n</i> -Bu	>20/1	<b>3m</b>	>20/1	63

<sup>a</sup> Reactions conducted using 0.3 mmol of *o*-(alkynyl)styrene derivative **1** in a 1:1 mixture of CH<sub>2</sub>Cl<sub>2</sub> and 1,4-dioxane (1.6 mL) at rt. <sup>b</sup> Determined by <sup>1</sup>H NMR analysis of the crude reaction mixture. <sup>c</sup> Isolated yield of product based on starting material **1**. <sup>d</sup> Reaction time = 4 h. <sup>e</sup> Cyclohexen-1-yl. <sup>f</sup> 3-Thienyl. <sup>g</sup> ~80% conversion. <sup>h</sup> ~65% conversion despite an additional load of 2.5 mol % of the catalyst was added.




## COMMUNICATION

cases from the alkene to the final indene, we wondered if this fact is also true for  $\beta,\beta$ -disubstituted *o*-(alkynyl)styrenes **4** as it had not been properly established in our previous reports (Table 4).<sup>8c</sup> When (*Z*)- $\beta$ -methyl- $\beta$ -phenyl *o*-(alkynyl)styrene **4a** was used as starting enyne the corresponding methoxycyclized indene **5a** was obtained as a single diastereoisomer, under the same conditions described for **1** (entry 1). Moreover, almost the same result was obtained using gold complexes bearing Ph<sub>3</sub>P as ligand in the presence or absence of silver salts (entries 2–3). In this case, the similar reactivity of substrate **4a** in the presence of the gold catalyst, bearing both the NHC and the phosphine ligands, is not surprising due to the higher nucleophilic nature of the trisubstituted alkene that would contribute to stabilize the carbocationic-like character of the intermediate. However, submission of *o*-(alkynyl)styrene **4b**, with an ethyl substituent instead of a methyl one at the  $\beta$ -position, to the same conditions resulted in an almost complete loss of stereospecificity (entry 4). Intrigued by this result, other *o*-(alkynyl)styrenes **4** with different substituents at the  $\beta$ -positions were also tested. With **4c**, bearing a propyl group as R substituent, the corresponding methoxycyclized product **5c** was obtained as a mixture of diastereoisomers clearly different from the mixture of geometrical isomers of the starting material (entry 5). A similar behaviour was observed for substrates **4d** (Ar = 4-FC<sub>6</sub>H<sub>4</sub>, R = Et), **4e** (Ar = 4-MeOC<sub>6</sub>H<sub>4</sub>, R = Me), and **4f** (Ar = 4-MeOC<sub>6</sub>H<sub>4</sub>, R = Et) (entries 6–8). These results seem to indicate that when starting from  $\beta,\beta$ -disubstituted *o*-(alkynyl)styrenes **4**, possessing a more nucleophilic trisubstituted alkene, their methoxycyclization reactions are not completely stereospecific in all cases, with variable degrees of stereoinformation loss that could also be influenced by steric hindrance issues.

By analogy with our previously reported mechanistic proposal for the methoxycyclization of  $\beta,\beta$ -disubstituted *o*-(alkynyl)styrenes<sup>8c</sup> and on the basis of other known gold-


**Table 4** Gold-catalyzed methoxycyclization of  $\beta,\beta$ -disubstituted *o*-(alkynyl)styrenes **4**<sup>a</sup>



Entry	<b>4</b>	Ar	R	<i>E/Z</i>	<b>5</b>	d.r. <sup>b</sup>	Yield (%) <sup>c</sup>
1	<b>4a</b>	Ph	Me	<1/20	<b>5a</b>	>20/1	87
2 <sup>d</sup>	<b>4a</b>	Ph	Me	<1/20	<b>5a</b>	16/1	89
3 <sup>e</sup>	<b>4a</b>	Ph	Me	<1/20	<b>5a</b>	16/1	86
4	<b>4b</b>	Ph	Et	17/1	<b>5b</b>	3/1	78
5 <sup>f</sup>	<b>4c</b>	Ph	<i>n</i> -Pr	8/1	<b>5c</b>	3/1	61
6	<b>4d</b>	4-FC <sub>6</sub> H <sub>4</sub>	Et	1/15	<b>5d</b>	5/1	60
7	<b>4e</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Me	<1/20	<b>5e</b>	13/1	88
8	<b>4f</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Et	1/18	<b>5f</b>	8/1	90

<sup>a</sup> Reactions conducted using 0.3 mmol of *o*-(alkynyl)styrene derivative **4** in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) at rt. <sup>b</sup> Determined by <sup>1</sup>H NMR analysis of the crude reaction mixture. <sup>c</sup> Isolated yield of product based on starting material **4**. <sup>d</sup> Carried out with Ph<sub>3</sub>PAuNTf<sub>2</sub> as catalyst. <sup>e</sup> Carried out with Ph<sub>3</sub>PAuCl/AgSbF<sub>6</sub> as catalytic system. <sup>f</sup> Carried out with an additional load of 2.5 mol % of catalyst and one hour more.

## Organic &amp; Biomolecular Chemistry



**Scheme 2** Mechanistic proposal

catalyzed enyne cyclizations,<sup>13</sup> a likely mechanism to account for the observed stereoselectivity issues is outlined in Scheme 2. An initial activation of the alkyne through coordination of cationic gold(I) triggers the attack of the olefin moiety through a 5-*endo* cyclization to give intermediate cyclopropyl carbene **A**, which has an alkenylgold carbocation mesomeric form **A'**. Whereas attack of the *O*-nucleophile to **A** would ensure a stereospecific process leading to the vinyl gold species **B** that would be subsequently protodemetalated to afford the functionalized indenenes **2** and **3**, the competitive contribution of **A'** to the intermediate gold complex structure would lead to a partial loss of the stereoinformation from the configurationally defined alkene. When this stereoinformation is totally transferred to the final indene it seems to indicate a greater, or almost exclusive, contribution of the cyclopropyl gold carbene **A**, as well as its configurational stability. However, when the final functionalized indene derivative is obtained as a mixture of diastereoisomers in spite of starting from only one geometrical isomer of the alkene, as it is the case with most of the  $\beta,\beta$ -disubstituted *o*-(alkynyl)styrenes **4**, it seems to point out that a competitive contribution of the carbocationic intermediate **A'** is occurring, likely due to the higher stabilization of the positive charge at the  $\beta$ -carbon of the substrate caused by its tertiary nature. In agreement with Echavarren's and Gagosz's proposals,<sup>7,13</sup> our results with  $\beta$ -monosubstituted *o*-(alkynyl)styrenes support that the intermediates in these Au-catalyzed cyclizations are gold-stabilized homoallylic carbocations and not open carbocations, and are probably better represented as cyclopropyl gold carbenes instead of simple carbocations.<sup>19</sup> In addition, the rapid and stereospecific attack of the nucleophile to configurationally stable cyclopropyl gold carbene **A** accounts for the relative stereochemistry of the obtained indene derivatives. For substrates **4b-f**, as well as with **1**, in which the reactions are not completely stereospecific, the bond rotation of the carbocationic intermediate should be faster than the attack of the methanol, likely due to the higher weakness of the cleaved bond in these cases. The stereospecific reaction of **4a** shows that the homoallylic stabilization of the carbocation is significant even for a substrate in which the positive charge is highly stabilized by methyl and phenyl groups.

## Conclusions

In conclusion, we have reported the gold(I)-catalyzed methoxy- and hydroxy-cyclizations of  $\beta$ -aryl-monosubstituted *o*-(alkynyl)styrenes that take place in a regio- and stereospecific way leading to interesting functionalized indene derivatives in high yields and as single diastereoisomers. The obtained results support that the intermediates in these gold-catalyzed cyclizations are better described as cyclopropyl gold carbenes in which the positive charge is significantly stabilized homoallylically. Further attack of the external nucleophile to these intermediates is faster than the competitive rotation around the weakest bond of the cyclopropyl ring. Only with highly activated and/or sterically-encumbered substrates, which place the positive charge in a tertiary carbon, the cyclizations are not completely stereospecific.

### Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

We gratefully acknowledge Ministerio de Economía y Competitividad (MINECO) and FEDER (CTQ2016-75023-C2-1-P), and Junta de Castilla y León and FEDER (BU076U16) for financial support. C.V. thanks Universidad de Burgos for a predoctoral contract. S.S.-P. thanks Junta de Castilla y León and FEDER for a postdoctoral contract.

### Notes and references

- For selected reviews, see: (a) A. S. K. Hashmi, *Pure Appl. Chem.*, 2010, **82**, 1517; (b) C. Obradors and A. M. Echavarren, *Acc. Chem. Res.*, 2014, **47**, 902; (c) R. Dorel and A. M. Echavarren, *Chem. Rev.*, 2015, **115**, 9028.
- We have classified all the cyclization reactions along the manuscript only considering the evolution of the triple bond of the enyne, although the more accurate way to name these electrophilic promoted cyclizations attending to the Baldwin's classification should reflect the ring closure to both  $\pi$ -bonds (5-*exo-trig*/5-*endo-dig* in these cases). See, for example: (a) R. K. Mohamed, S. Mondal, K. Jorner, T. Faria Delgado, H. Ottoson and I. V. Alabugin, *J. Am. Chem. Soc.*, 2015, **137**, 15441. For a recent review on the Baldwin rules, see: (b) K. Gilmore, R. K. Mohamen and I. V. Alabugin, *WIREs Comput. Mol. Sci.*, 2016, **6**, 487.
- (a) M. R. Luzung, J. P. Markham and F. D. Toste, *J. Am. Chem. Soc.*, 2004, **126**, 10858; (b) L. Zhang and S. A. Kozmin, *J. Am. Chem. Soc.*, 2004, **126**, 11806; (c) F. Gagosz, *Org. Lett.*, 2005, **7**, 4129.
- For other reactions of *o*-(alkynyl)styrenes leading to indene derivatives see, for instance: (a) Ru-catalyzed: R. J. Madhushaw, C.-Y. Lo, C.-W. Hwang, M.-D. Su, H.-C. Shen, S. Pal, I. R. Shaikh and R.-S. Liu, *J. Am. Chem. Soc.*, 2004, **126**, 15560; (b) Radical-mediated: S. Mondal, R. K. Mohamed, M. Manoharan, H. Phan and I. V. Alabugin, *Org. Lett.*, 2013, **15**, 5650; (c) Al-promoted: H. Kinoshita, N. Hirai and K. Miura, *J. Org. Chem.*, 2014, **79**, 8171; (d) FLP-catalyzed: S. Tamke, Z.-W. Qu, N. A. Sitte, U. Flörke, S. Grimme and J. Paradies, *Angew. Chem., Int. Ed.*, 2016, **55**, 4336.
- E. Aguilar, R. Sanz, M. A. Fernández-Rodríguez and P. García-García, *Chem. Rev.*, 2016, **116**, 8256.
- (a) T. Shibata, Y. Ueno and K. Kanda, *Synlett*, 2006, **411**. For related processes catalyzed by other transition-metal complexes, see: (b) J. W. Dankwardt, *Tetrahedron Lett.*, 2001, **42**, 5809; (c) V. Mamane, P. Hannen and A. Fürstner, *Chem. Eur. J.*, 2004, **10**, 4556.
- (a) A. K. Buzas, F. M. Istrate and F. Gagosz, *Angew. Chem., Int. Ed.*, 2007, **46**, 1141; (b) S. Böhringer and F. Gagosz, *Adv. Synth. Catal.*, 2008, **350**, 2617.
- (a) A. Martínez, P. García-García, M. A. Fernández-Rodríguez, F. Rodríguez and R. Sanz, *Angew. Chem., Int. Ed.*, 2010, **49**, 4633; (b) P. García-García, M. A. Rashid, A. M. Sanjuán, M. A. Fernández-Rodríguez and R. Sanz, *Org. Lett.*, 2012, **14**, 4778; (c) A. M. Sanjuán, M. A. Rashid, P. García-García, A. Martínez-Cuevza, M. A. Fernández-Rodríguez, F. Rodríguez and R. Sanz, *Chem. Eur. J.*, 2015, **21**, 3042; (d) A. M. Sanjuán, C. Virumbrales, P. García-García, M. A. Fernández-Rodríguez and R. Sanz, *Org. Lett.*, 2016, **18**, 1072. For related halocyclization reactions of these substrates, see: (e) R. Sanz, A. Martínez, P. García-García, M. A. Fernández-Rodríguez, M. A. Rashid and F. Rodríguez, *Chem. Commun.*, 2010, **46**, 7427; (f) P. García-García, A. M. Sanjuán, M. A. Rashid, A. Martínez-Cuevza, M. A. Fernández-Rodríguez, F. Rodríguez and R. Sanz, *J. Org. Chem.*, 2017, **82**, 1155.
- See, for instance: (a) M. Enders and R. W. Baker, *Curr. Org. Chem.*, 2006, **10**, 937; (b) J.-i. Nishida, S. Tsukaguchi and Y. Yamashita, *Chem. Eur. J.*, 2012, **18**, 8964.
- For reviews for the synthesis of indenenes, see: (a) G. Qiu and J. Wu, *Synlett*, 2014, **25**, 2703; (b) B. Gabriele, R. Mancuso and L. Veltri, *Chem. Eur. J.*, 2016, **22**, 5056. For selected recent reports, see: (c) A. Preinfalk, A. Misale and N. Maulide, *Chem. Eur. J.*, 2016, **22**, 14471; (d) B. Ma, Z. Wu, B. Huang, L. Liu and J. Zhang, *Chem. Commun.*, 2016, **52**, 9351. (e) R. Barroso, M. Paraja, M.-P. Cabal and C. Valdés, *Org. Lett.*, 2017, **19**, 4086.
- See, for instance: (a) D. Vasu, H.-H. Hung, S. Bhunia, S. A. Gawade, A. Das and R.-S. Liu, *Angew. Chem., Int. Ed.*, 2011, **50**, 6911; (b) C.-H. Chen, Y.-C. Tsai and R.-S. Liu, *Angew. Chem., Int. Ed.*, 2013, **52**, 4599; (c) M.-C. P. Yeh, C.-J. Liang, H.-F. Chen and Y.-T. Weng, *Adv. Synth. Catal.*, 2015, **357**, 3242; (d) W. Fang, Y. Wei and M. Shi, *Chem. Commun.*, 2017, **53**, 11666.
- Only particular examples for the alkoxy cyclization of  $\beta$ -monosubstituted *o*-(alkynyl)styrenes, such as enol ethers, had been previously reported: (a) Y. Liu, J. Guo, Y. Liu, X. Wang, Y. Wang, X. Jia, G. Wei, L. Chen, J. Xiao and M. Cheng, *Chem. Commun.*, 2014, **50**, 6243; (b) J. Wang, K. Huang, L. Liu, W. Chang and J. Li, *Tetrahedron Lett.*, 2015, **56**, 2659.
- (a) V. López-Carrillo, N. Huguet, A. Mosquera and A. M. Echavarren, *Chem. Eur. J.*, 2011, **17**, 10972. For a similar discussion with 1,6-enynes, see: (b) P. Pérez-Galán, N. J. A. Martin, A. G. Campaña, D. J. Cárdenas and A. M. Echavarren, *Chem. Asian J.*, 2011, **6**, 482; (c) R. Dorel and A. M. Echavarren, *J. Org. Chem.*, 2015, **80**, 7321.
- L. Ricard and F. Gagosz, *Organometallics*, 2007, **26**, 4704.
- For selected recent examples, see: (a) R. E. Ebu, D. Malhotra, G. B. Hammond and B. Xu, *Adv. Synth. Catal.*, 2016, **358**, 1478; (b) D. Ding, T. Mou, M. Feng and X. Jiang, *J. Am. Chem. Soc.*, 2016, **138**, 5218.
- E. Jiménez-Núñez, C. K. Claverie, C. Bour, D. J. Cárdenas and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2008, **47**, 7892.
- See ESI for further details.
- CCDC 1585617 (**3i**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- For recent reviews about the nature of cationic intermediates relevant to Au-catalyzed enyne cyclizations, see: (a) Y. Wang, M. E. Muratore and A. M. Echavarren,

## COMMUNICATION


## Organic &amp; Biomolecular Chemistry

*Chem. Eur. J.*, 2015, **21**, 7332; (b) R. J. Harris and R. A. Widenhoefer, *Chem. Soc. Rev.*, 2016, **45**, 4533.

View Article Online  
DOI: 10.1039/C8OB00406D

Organic & Biomolecular Chemistry Accepted Manuscript

Published on 19 March 2018. Downloaded by Universidad de Burgos on 19/03/2018 17:01:39.



40x21mm (600 x 600 DPI)