

Am J Transl Res 2017;9(8):3573-3585
www.ajtr.org /ISSN:1943-8141/AJTR0055033

Original Article

Posttranslational modifications of calcium/calmodulin-dependent protein kinase II δ and its downstream signaling in human failing hearts

Tomas Rajtik¹, Eva Goncalvesova², Zoltan V Varga³, Przemyslaw Leszek⁴, Mariusz Kusmierczyk⁴, Michal Hulman⁵, Jan Kyselovic¹, Peter Ferdinandy³, Adriana Adameova¹

¹Department of Pharmacology & Toxicology, Faculty of Pharmacy, Comenius University, Bratislava, Slovak Republic; ²Department of Heart Failure & Transplantation, ⁵Clinic of Heart Surgery, The National Institute of Cardiovascular Diseases, Bratislava, Slovak Republic; ³Department of Pharmacology & Pharmacotherapy, Semmelweis University, Budapest, Hungary; ⁴Institute of Cardiology, Warszawa, Poland

Received April 7, 2017; Accepted July 16, 2017; Epub August 15, 2017; Published August 30, 2017

Abstract: Background: In human failing hearts (HF) of different origin (coronary artery disease-CAD, dilated-DCM, restrictive and hypertrophic cardiomyopathy-OTHER), we investigated the active forms of Ca²⁺/calmodulin-dependent protein kinase II δ (p-Thr²⁸⁷-CaMKII δ , oxMet^{281/282}-CaMKII δ) and their role in phenotypes of the disease. Methods and results: Although basic diagnostic and clinical markers indicating the attenuated cardiac contractility and remodeling were comparable in HF groups, CaMKII δ -mediated axis was different. P-Thr²⁸⁷-CaMKII δ was unaltered in CAD group, whereas it was upregulated in non-ischemic cardiomyopathic groups. No correlation between the upregulated p-Thr²⁸⁷-CaMKII δ and QT interval prolongation was detected. Unlike in DCM, oxMet^{281/282}-CaMKII δ did not differ among HF groups. Independently of CaMKII δ phosphorylation/oxidation, activation of its downstreams-phospholamban and cardiac myosin binding protein-C was significantly downregulated supporting both diminished cardiac lusitropy and inotropy in all hearts. Content of sarcoplasmic reticulum Ca²⁺-ATPase 2a in all HF was unchanged. Protein phosphatase1 β was upregulated in CAD and DCM only, while 2A did not differ among groups. Conclusion: This is the first demonstration that the posttranslational activation of CaMKII δ differs in HF depending on etiology. Lower levels of downstream molecular targets of CaMKII δ do not correlate with either activation of CaMKII δ or the expression of major protein phosphatases in the HF. Thus, it is unlikely that these mechanisms exclusively underlie failing of the heart.

Keywords: Human heart failure, Ca²⁺/calmodulin-dependent protein kinase II, sarcoplasmic reticulum calcium handling, cardiac myosin binding protein-C

Introduction

Heart failure (HF) is a progressive cardiac disease ranging from stages which impair the patient's quality of life and finally progress to a stage that is characterized by symptoms resistant to treatment. HF can occur as a result of coronary artery disease (CAD), dilated cardiomyopathy (DCM) and other hereditary and idiopathic causes [1]. In contrast to the most prevalent HF due to CAD, the age-adjusted prevalence of other forms of HF is lower (1:500 to 1:2500) [1-3]. Regardless of etiology, all types of systolic HF are characterized by seriously impaired cardiac contractility leading into inadequate perfusion of organs. Great effort has

been undertaken to understand the pathomechanisms of systolic myocardial dysfunction. The abnormal Ca²⁺ cycling and defects in the sarcomeric proteins are proposed to be one of the crucial players in the adverse remodeling and dysfunction of the heart [4].

Ca²⁺/calmodulin-dependent protein kinase II δ (CaMKII δ) has emerged as an important Ca²⁺ handling protein regulating excitation-contraction coupling. Alterations in its activity have been proposed to exaggerate mishandling in Ca²⁺ homeostasis, and thereby underlie the depressed cardiac contractile function and arrhythmogenesis [5, 6]. Overactivation of CaMKII δ can arise from its capability to undergo the vari-

CaMKII active forms in failing hearts

ous posttranslational modifications, including autophosphorylation (pThr²⁸⁷-CaMKII δ) and oxidation (oxMet^{281/282}-CaMKII δ) due to oxidative stress [7]. As CaMKII δ phosphorylates a wide variety of proteins involved in the regulation of both Ca²⁺ handling (phospholamban-PLN), and contractile proteins (cardiac myosin binding protein-C-cMyBP-C), this protein kinase might be an interesting target in the management of HF treatment [8]. In fact, it has been previously reported that in failing hearts, sarcoplasmic reticulum (SR) Ca²⁺ uptake is significantly reduced due to the inhibitory effects of PLN on SERCA2a [9, 10]. This may occur as a result of a decreased activity/expression of SERCA2a [11] or a higher expression of PLN [12, 13]. In addition to Ca²⁺ dysregulation, attenuation of cMyBP-C function, which serves as a regulatory and structural protein of the sarcomere, due to its dephosphorylation and subsequent degradation of cMyBP-C is associated with the diminished cardiac contractile function and poor prognosis of patients with HF [14]. However, it is not known if the above-mentioned link involving SR Ca²⁺ proteins, cMyBP-C and CaMKII δ , as a central signal-transducing element, is altered depending on the certain type of systolic ventricular dysfunction. Therefore, the objective of this study was to provide a comprehensive analysis of the posttranslational activation of CaMKII δ and if it is associated with the alterations of downstream target proteins in the SR and sarcomere in human failing heart samples. We also investigated a potential role of protein phosphatases PP1 and PP2A counterbalancing the effects of CaMKII δ and its downstream proteins [15]. As CaMKII δ has been suggested as a pro-arrhythmogenic marker [16, 17], and arrhythmias are a common complication of HF [18], therefore we also analyzed a correlation between the levels of the active forms of CaMKII δ and QT interval duration.

Methods

Study design

All procedures were in accordance with ethical standards for human experiments based on Helsinki declaration. All experiments were approved by national and institutional ethical commissions (statement no. IK-NP-0021-24/1426/14 and NUSCH EK 126/180509). Samples of left ventricles were obtained from explanted hearts of patients with diagnosed ter-

минаl stadium of HF (NYHA III-IV) and reduced left ventricle ejection fraction (LVEF<25%). We employed the failing hearts due to coronary artery disease (CAD, n=6), dilated cardiomyopathy (DCM, n=10), and other cardiomyopathies being either restrictive or hypertrophied cardiomyopathy (OTHER, n=6). All patients underwent echocardiographical, hemodynamical and biochemical examination before transplantation. Additionally, the HF pharmacotherapy was analyzed from the last medical records. Samples of left ventricles from healthy donor patients (C, n=4), whose hearts cannot have been used for transplantation from various medical reasons (CMV infection, size/donor recipient mismatch, major damage during procedure), served as a control group. Tissues from the walls of left ventricles have been harvested in the time of explanation avoiding the scared, fibrotic and adipose tissues. They were afterwards rinsed, dried and snap-frozen in liquid nitrogen until further processing.

Immunoblotting

The samples of left ventricles were processed for the immunoblotting analysis based on our standard laboratory protocol for SDS-PAGE and Western Blotting [19]. Proteins were transferred on PVDF membranes (Immobilon-P, EMD Millipore, USA) and incubated with primary antibodies against p-Thr²⁸⁷-CaMKII (Cell Signaling, USA), oxMet^{281/282}-CaMKII (EMD Millipore, USA), total CaMKII δ (Santa Cruz, USA), p-Thr¹⁷-PLN (Badrilla, UK), p-Ser¹⁶-PLN (Badrilla, UK), total PLN (Badrilla, UK), p-Ser²⁸²⁽²⁸⁴⁾-cMyBP-C (Enzo, USA), total cMyBP-C (Santa Cruz, USA), total SERCA2a (Badrilla, UK), PP1 β (Abcam, USA) and PP2A (Sigma-Aldrich, USA). Afterwards, the incubation with HRP-conjugated secondary antibodies, donkey anti-rabbit IgG (GE Healthcare Life Sciences, UK) and rabbit anti-goat IgG (Sigma-Aldrich, USA) was performed. Signals were detected by enhanced chemiluminescence (Crescendo Luminata, EMD Millipore, USA) and chemiluminescence imaging system (myECL imager, Thermo Scientific, USA). Total protein staining with Coomassie Brilliant Blue R-250 or Reactive Brown 10B evaluated by scanning densitometry was used as the loading control as a substitution to the housekeeping protein immunodetection [20-22]. Intensity of a particular protein band was compared to a whole lane.

CaMKII active forms in failing hearts

Table 1. Echocardiographic, hemodynamic and biochemical parameters of probands

| | Controls | CAD | DCM | OTHER | |
|---|-------------|-------------|-------------|-------------|--------------|
| | | | | RCM | HCM |
| Age (years) | 30 ± 5 | 52 ± 4 | 47 ± 3 | 47 ± 6 | 60 ± 4 |
| LVEF (%) | n.a. | 21 ± 1 | 18 ± 1 | 34 ± 8 | 30 ± 5 |
| LVEDD (mm) | n.a. | 75 ± 4 | 72 ± 3 | 60 ± 3 | 59 ± 1 |
| RVEDD (mm) | n.a. | 34 ± 3 | 35 ± 1 | 33 ± 1 | n.a. |
| QT (s) | n.a. | 0.42 ± 0.01 | 0.41 ± 0.02 | 0.44 ± 0.02 | n.a. |
| sPAP (mm Hg) | n.a. | 44 ± 7 | 53 ± 5 | 52 ± 5 | 45 ± 6 |
| dPAP (mm Hg) | n.a. | 21 ± 4 | 27 ± 3 | 28 ± 3 | 21 ± 2 |
| CI (dm ³ ·min ⁻¹ ·m ⁻²) | n.a. | 2.2 ± 0.1 | 1.8 ± 0.1 | 1.8 ± 0.2 | n.a. |
| NT-proBNP (ng·dm ⁻³) | n.a. | 3541 ± 1035 | 6571 ± 1682 | 3394 ± 1007 | 10124 ± 4152 |
| Creatine kinase (μkat·dm ⁻³) | 3.09 ± 0.92 | 1.18 ± 0.26 | 1.17 ± 0.21 | 1.08 ± 0.47 | 0.97 ± 0.21 |

End-stage HF groups of ischemic and non-ischemic origin did not significantly differ in neither echocardiographic nor hemodynamic parameters. CAD-coronary artery disease; DCM-dilated cardiomyopathy; RCM-restrictive cardiomyopathy; HCM-hypertrophied cardiomyopathy. LVEF-left ventricular ejection fraction; LVEDD-left ventricular end-diastolic diameter; RVEDD-right ventricular end-diastolic diameter; QT-QT interval; sPAP-systolic pulmonary artery pressure; dPAP-diastolic pulmonary artery pressure; CI-cardiac index; NT-proBNP-N-terminal pro-B type natriuretic peptide. Data are expressed as the means ± SEM.

Statistical analysis

The results are expressed as means ± standard error of means (S.E.M.), unless stated otherwise. One-way ANOVA analysis with Newman-Keuls and Tukey's post-hoc tests and two-tailed unpaired Student's *t*-test were used for evaluation of group differences in variables with normal distribution. In case of the non-normal distribution, a Man-Whitney's analysis was used. Correlation between biochemical parameters, QT duration or N-terminal pro-B type natriuretic peptide (NT-proBNP) and the levels of certain proteins was analyzed by Pearson's test. All analyses were performed with GraphPad Prism 6.00 for Windows (GraphPad Software, USA). Differences between groups were considered significant when *P* < 0.05.

Results

Characteristics of study subjects

A summary of the main pre-transplant data of the study subjects are presented in **Table 1**. All patients irrespective of the etiology of HF were in either NYHA III or IV class. Among the HF groups, there were no significant differences in neither of the values of left ventricular ejection fraction (LVEF), parameters of cardiac remodeling (the size of left and right end-diastolic ventricular diameters-LVEDD, RVEDD), markers of pulmonary hypertension (systolic and diastolic pulmonary pressures-sPAP, dPAP) nor NT-proBNP, a common biochemical marker of HF. HF

patients were treated at least with ACE inhibitors or AT1R-blockers, beta-blockers and diuretics. The control subjects were treated with the infusion of noradrenaline and dopamine and fluid balance was maintained with desmopressin and hydroxyethyl starch, an intravenous colloid volume expander. They presented preserved systolic/diastolic function. No metabolic disorders were diagnosed in the study subjects.

Posttranslationally modified forms of CaMKIIδ in end-stage human HF show no association with the QT interval duration

Western blot analysis of the posttranslationally modified forms of CaMKIIδ by phosphorylation at Thr287 and oxidation at Met281/282 is shown in **Figures 1, 2**.

The content of p-Thr²⁸⁷-CaMKIIδ in the group of CAD did not differ from the values of the control subjects. On the other hand, in all cardiomyopathic groups of HF, the expression of p-Thr²⁸⁷-CaMKIIδ was significantly increased as compared with the control hearts and CAD (**Figures 1, 2A**). Of note, the levels of oxMet^{281/282}-CaMKIIδ did not differ among HF group. However, by using Student's *t*-test, significantly lower expression of oxMet^{281/282}-CaMKIIδ was observed in the DCM as compared to control group (**Figures 1, 2B**). This is the first analysis of the oxidized form of this protein kinase in human end-stage HF.

CaMKII active forms in failing hearts




Figure 1. Representative blots of evaluated proteins in control and human failing hearts. Total CaMKII δ -total Ca²⁺/calmodulin-dependent protein kinase 2 δ ; p-Thr²⁸⁷-CaMKII δ -phospho-Thr²⁸⁷-Ca²⁺/calmodulin-dependent protein kinase 2 δ ; ox-Met^{281/282}-CaMKII δ -oxidized-Met^{281/282}-Ca²⁺/calmodulin-dependent protein kinase 2 δ ; total PLN-total phospholamban; p-Thr¹⁷-PLN-phospho-Thr¹⁷-phospholamban; p-Ser¹⁶-PLN-phospho-Ser¹⁶-phospholamban; SERCA2a-sarcoplasmic/endoplasmic reticulum Ca²⁺-ATPase; total cMyBP-C-total cardiac myosin-binding protein C; p-Ser²⁸⁴-cMyBP-C-phospho-Ser²⁸⁴-cardiac myosin-binding protein C; PP1 β -protein phosphatase 1 β ; PP2A-protein phosphatase 2A.

The total CaMKII δ was comparable in all HF groups and did not differ from the values of controls (**Figures 1, 2C**). Thus, the ratio of both p-Thr²⁸⁷/total CaMKII δ and oxMet^{281/282}-CaMKII δ /total CaMKII δ mimicked the pattern of

changes of the particular posttranslational modifications (**Figures 1, 2D, 2E**).

QT interval prolongation is known as a predisposing factor for increased ventricular arrhythmia risk and a higher activity and/or expression of CaMKII δ has been associated with arrhythmia triggering [23, 24]. Therefore in the groups of HF with the altered expression of post-translationally modified form of CaMKII δ (DCM, OTHER), a potential correlation between the content of phosphorylated form and prolongation of QT interval was analyzed. However, we did not identify any link between p-Thr²⁸⁷-CaMKII δ and the prolongation of QT interval (**Figures 1, 2F**).

Regulation of SR Ca²⁺ uptake in end-stage human HF

As SR Ca²⁺ cycling is known to be diminished in HF [25] and CaMKII activates certain SR proteins [26] we further analyzed p-Thr¹⁷-PLN, a downstream of this protein kinase, and SERCA2a. In spite of the different state of the phosphorylation of CaMKII δ in the particular HF types (ischemic vs. non-ischemic) (**Figures 1, 2A**), the levels of p-Thr¹⁷-PLN were greatly downregulated in all HF groups in comparison to non-failing hearts (**Figures 1, 3A**). Likewise, PKA-dependent phosphorylation of PLN at Ser16 residue was decreased and there was no difference among the

diseased groups indicating that PKA does not substitute a role of CaMKII δ to maintain the function of PLN (**Figures 1, 3B**). The expression of the total PLN (**Figures 1, 3C**) was unchanged in the HF groups, thereby the ratio of both

CaMKII active forms in failing hearts





Figure 2. Phosphorylation (p-Thr²⁸⁷-CaMKIIδ) and oxidation (oxMet^{281/282}-CaMKIIδ) of CaMKIIδ in different types of terminal human HF. (A) Levels of p-Thr²⁸⁷-CaMKIIδ; (B) Content of oxMet^{281/282}-CaMKIIδ; (C) Expression of total CaMKIIδ; (D) Ratio of phosphorylated to total CaMKIIδ; (E) Ratio of oxidized to total CaMKIIδ; (F) Comparison between QT interval duration and phosphorylation of CaMKIIδ (n=12). Data for (A-E) are expressed as the means ± S.E.M. n=4-10 hearts per group. *P<0.05 vs. control hearts.



p-Thr¹⁷-PLN/total PLN and p-Ser¹⁶-PLN/total PLN showed the same pattern as the respective active forms of this protein (Figures 1, 3D, 3E). These data, as general characteristics of diastolic dysfunction, indicates no important role of the expression of either of active form of the upstream kinase in terminal HF.

Non-phosphorylated form of PLN produces inhibitory effects on SERCA2a activity and restricts Ca²⁺ filling back into the SR [27, 28]. Therefore, we estimated a ratio between the total PLN and SERCA2a. Similarly to the total PLN, the levels of total SERCA2a were unchanged in all HF groups when compared to control

CaMKII active forms in failing hearts




Figure 4. Content of SERCA2a in human failing hearts and ratio of total PLN to SERCA2a. (A) Expression of SERCA2a. (B) Ratio of total PLN to SERCA2a. Values for (A) and (B) are expressed as the means \pm S.E.M. n=4-10 hearts per group.




Figure 5. Expression and phosphorylation of cMyBP-C in human failing hearts. (A) Expression of p-Ser²⁸⁴-cMyBP-C; (B) Content of total cMyBP-C; (C) Ratio of p-Ser²⁸⁴-cMyBP-C to total protein. Values for (A-C) are expressed as the means \pm S.E.M. n=4-10 hearts per group. *P<0.05 vs. control hearts.




Figure 6. Correlation between the expression of p-Ser²⁸⁴-cMyBP-C and marker of volume overload and myocyte stretch NT-proBNP in failing hearts. n=13 hearts. Correlation was considered as positively significant (P=0.0181).

group (Figures 1, 4A). Accordingly, no significant changes were found in the PLN/SERCA2a ratio in HF groups as compared with non-failing hearts (Figures 1, 4B).

Cardiac MyBP-C in human HF

In addition to the altered Ca²⁺ cycling, the response of the contractile proteins is diminished in HF [29]. Therefore, we evaluated expression of total cMyBP-C and its phosphorylated form at Ser284. Unlike the expression of the total protein (Figure 5B), the levels of p-Ser²⁸⁴-cMyBP-C were significantly downregulated in all HF groups regardless the expression of particular posttranslationally modified forms of CaMKII δ (Figures 1, 5A, 5C). These results indicate serious systolic dysfunction in HF and similarly to the status of p-Thr¹⁷-PLN propose no role of the levels of the active forms of CaMKII δ in this context.

To further expand the knowledge about this active form of cMyBP-C in end-stage human HF, a correlation between the serum NT-proBNP and p-Ser²⁸⁴-cMyBP-C was evaluated. Interestingly, the levels of NT-proBNP positively corre-

CaMKII active forms in failing hearts




Figure 7. Expression of protein phosphatases PP1 β and PP2A in left ventricles of various types of human heart failure. (A) Expression of PP1 β ; (B) Expression of PP2A. Values for (A) and (B) are expressed as the means \pm S.E.M. n=4-10 hearts per group. * P <0.05 vs. control hearts.

Table 2. Expression of evaluated proteins in left ventricles of human failing hearts of different origin

| Proteins | CAD | DCM | OTHER |
|--|-----|-----|-------|
| Total CaMKII δ | ≈ | ≈ | ≈ |
| p-Thr ²⁸⁷ -CaMKII δ | ≈ | ↑ | ↑ |
| ox-Met ^{281/282} -CaMKII δ | ≈ | ↓ | ≈ |
| Total PLN | ≈ | ≈ | ≈ |
| p-Thr ¹⁷ -PLN | ↓ | ↓ | ↓ |
| p-Ser ¹⁶ -PLN | ↓ | ↓ | ↓ |
| SERCA2a | ≈ | ≈ | ≈ |
| Total cMyBP-C | ≈ | ≈ | ≈ |
| p-Ser ²⁸⁴ -cMyBP-C | ↓ | ↓ | ↓ |
| PP1 β | ↑ | ↑ | ≈ |
| PP2A | ≈ | ≈ | ≈ |

Table summarizes protein expression detected by immunoblot analysis between particular HF groups: CAD-coronary artery disease; DCM-dilated cardiomyopathy; RCM-restrictive cardiomyopathy; HCM-hypertrophied cardiomyopathy and control group. Total CaMKII δ , p-Thr²⁸⁶-CaMKII δ , ox-Met^{281/282}-CaMKII δ -total, phospho-Thr²⁸⁶ and oxidized-Met^{281/282}-Ca²⁺/calmodulin-dependent protein kinase II δ ; total PLN, p-Thr¹⁷-PLN, p-Ser¹⁶-PLN-total, phospho-Thr¹⁷ and phospho-Ser¹⁶-phospholamban; SERCA2a-sarcoplasmic/endoplasmic reticulum Ca²⁺-ATPase; total cMyBP-C, p-Ser²⁸⁴-cMyBP-C-total and phospho-Ser²⁸⁴-cardiac myosin-binding protein C; PP1 β -protein phosphatase 1 β ; PP2A-protein phosphatase 2A. (≈)-no difference in expression; (↑)-significantly increased expression in comparison to control group; (↓)-significantly decreased expression in comparison to controls. Differences between the groups were evaluated by ANOVA test and considered significant when P <0.05.

lated with the levels of p-Ser²⁸⁴-cMyBP-C in the HF groups (Figures 1, 6). This may indicate a compensatory increase in cMyBP-C phosphory-

lation in response to increased myocyte stretch documented by the higher levels of NT-proBNP.

Modulation of protein phosphorylation status by protein phosphatases in failing hearts

We also analyzed certain protein phosphatases which counterbalance the phosphorylation of PLN, cMyBP-C and CaMKII δ itself. Indeed, two major isoforms of protein phosphatases in human myocardium, PP1 subunit β (PP1 β) and PP2A were investigated. PP1 β , proposed to dephosphorylate the most of PLN as well as cMyBP-C [30], was increased in the CAD and DCM group but not in other cardiomyopathic groups (Figures 1, 7A). PP2A sharing similar catalytic sites with PP1 [31, 32] proposed also to dephosphorylate cMyBP-C [33, 34] and most of CaMKII in cytosolic and membrane fraction [35] did not differ among the diseased groups nor in comparison with control healthy hearts (Figures 1, 7B).

Discussion

In the present comprehensive study we have shown that the levels of both active, posttranslationally modified forms of CaMKII δ , p-Thr²⁸⁷-CaMKII δ and oxMet^{281/282}-CaMKII δ in failing hearts due to CAD were comparable to those of healthy hearts. On the other hand, in HF of non-ischemic origin (DCM, RCM and HCM), p-Thr²⁸⁷-CaMKII δ was upregulated whereas expression of oxMet^{281/282}-CaMKII δ was unchanged. Regardless the levels of these active forms of the kinase, the downstream targets of CaMKII δ , such as p-Thr¹⁷-PLN or p-Ser²⁸⁴-cMyBP-C were

CaMKII active forms in failing hearts

significantly downregulated in all HF types. Evaluation of protein phosphatases revealed the upregulated content of PP1 β in CAD and DCM but not in other types of HF. Expression of PP2A in failing hearts did not differ of the levels of controls. Thus, this study employing the samples of various types of HF (both of a high and low incidence) provides the first demonstration that i) the oxidative activation of CaMKII δ does not go hand in hand with its activation through phosphorylation, ii) CaMKII δ activation through phosphorylation and oxidation differs depending on the etiology of HF, iii) the lower phosphorylation of the downstream proteins indicating diminished inotropy and lusitropy in these diseased hearts is not paralleled with either active forms of CaMKII δ or with the expression of the main cardiac protein phosphatases (**Table 2**).

Under physiological conditions, the activation of CaMKII in heart is mainly regulated by auto-phosphorylation in the presence of Ca²⁺/CaM [36]. This mechanism allows the kinase to activate various Ca²⁺-sensitive proteins to ensure proper Ca²⁺ handling, including proteins regulating the function of the sarcoplasmic reticulum and sarcomere [37-39]. Therefore, the observed higher activation state of CaMKII δ and Ca²⁺ mishandling may contribute to the disturbances in contraction and relaxation in HF [40]. Considering these deleterious effects of the overactivated protein kinase, increased p-Thr²⁸⁷-CaMKII δ could be expected in the failing hearts. However, in spite of this logical hypothesis, data shown in this study as well as in others [41, 42] are unequivocal. In fact, all HF hearts characterized by depressed cardiac contractility and remodeling have shown the different pattern of oxidation and phosphorylation of CaMKII δ . Moreover, the pattern of changes in the oxidative activation of the kinase did not mimic the changes in phosphorylation and vice versa. Interestingly, in spite of this observation the markers of HF such as NT-proBNP, pulmonary arterial pressures, EF or cardiac indices did not differ among the types of HF.

So far published papers have mainly been dealing with CaMKII in human HF of a higher incidence. In study of Miyamoto et al. [43], in DCM, the increased phosphorylation of CaMKII has been reported what is in line with our data. On the other hand, similarly to Fisher et al [42] the

ratio of the phosphorylated form/total protein kinase has not indicated the changes among the groups. In another study, the activity of CaMKII measured by enzymatic assay was found to be increased in DCM while unchanged in CAD [41]. By predicting a linear link between phosphorylation and catalytic activity of the kinase this observation is in line with results reported in this study. In the present study, we have extended the current knowledge on the activation of CaMKII δ and reported for the first time the content of p-Thr²⁸⁷-CaMKII δ being also upregulated in HF types of a lower incidence. It has been suggested that the overactivated/hyperphosphorylated CaMKII is associated with electrical instability along with contractile dysfunction [19, 42, 44]. However, we have been unable to find any correlation between p-Thr²⁸⁷-CaMKII δ expression and the duration of QT interval in the HF groups of non-ischemic origin which exerted the higher phosphorylation of this protein kinase.

In addition to phosphorylation, CaMKII δ can undergo other posttranslational modifications [7]. In the presence of oxidative stress, which is commonly observed in HF [45], the activation of CaMKII δ through oxidation might be of a great relevance thereby suggesting the levels of oxMet^{281/282}-CaMKII δ being increased in our samples of human HF. Of note, the increased levels of oxMet^{281/282}-CaMKII δ have been shown to underlie electrical remodeling and hypertrophy in cardiac myocytes and isoprenaline-treated rats [46, 47]. Surprisingly, no changes in the expression of oxMet^{281/282}-CaMKII δ were found among HF groups. However, while comparing DCM and control group, oxMet^{281/282}-CaMKII δ was found to be decreased in the diseased hearts. It has been proposed that methionine residues at 281/282 can undergo oxidation by superoxide produced by NADPH oxidases what leads into the kinase activation [48, 49]. Another enzyme, methionine sulfoxide reductase A is able to reverse this effect and thereby decrease pro-oxidant potential of the kinase [48]. It would be interesting to measure the activity of this enzyme; however, at the time of performing the study it was impossible due to the unavailability of assay and antibody. Thus, the aforementioned data indicate that a role of CaMKII δ including its posttranslational forms should be carefully investigated and the particular etiology of HF should be taken into account while assessing

CaMKII active forms in failing hearts

its involvement in the pathologic mechanisms of the disease. Of note, in our recent study, we have shown that both p-Thr²⁸⁷-CaMKII δ and oxMet^{281/282}-CaMKII δ are not elevated, rather markedly decreased in the late phase of reperfusion of previously ischemic hearts which experienced contractile dysfunction and higher oxidative stress [19].

CaMKII δ is one of critical proteins regulating proper Ca²⁺ cycling. For instance, by phosphorylation of PLN at Thr¹⁷ it relieves the inhibitory effects on SERCA2a and thereby mediates lusitropic effects [50]. Since HF is characterized by significantly abolished cardiac contractility and relaxation we have hypothesized CaMKII δ -dependent phosphorylation of PLN being decreased. In agreement with this hypothesis, the expression of CaMKII δ -phosphorylated form of p-Thr¹⁷-PLN was decreased to almost non-detectable levels in all HF groups. Similarly to our study, Dash et al. [12] and Miyamoto et al. [43] have also reported CaMKII-mediated phosphorylation of PLN being decreased in DCM. The content of p-Thr¹⁷-PLN was parallel with the levels of p-Ser¹⁶-PLN regulated by PKA. Additionally, we have shown that SERCA2a levels did not differ among the groups what is in agreement with some previous studies [11, 51]. On the other hand, downregulation of SERCA2a in heart failure resulting from DCM has also been reported [12, 52]. The expression of the ratio of total PLN to SERCA2a was unaltered in all diseased groups. The levels of non-phosphorylated PLN have been suggested to act as a marker of decreased SERCA2a pumping activity [53]. However, our data and of others [39, 43, 54] do support the role of phosphorylated PLN rather than the non-phosphorylated protein in diminished Ca²⁺ SR handling in HF.

CaMKII δ is also known to directly phosphorylate some proteins of contractile apparatus and thereby regulates cardiac contractility independently of Ca²⁺ cycling modulation. In fact, it phosphorylates cMyBP-C, regulates actin-myosin cross-bridging, and controls force generation within the sarcomere [29]. Recently, cMyBP-C has been proposed to serve as a novel biomarker of cardiac injury [14, 55]. Dephosphorylation and subsequent degradation of this protein may indicate cardiac dysfunction and HF [56, 57]. Likewise, several gene mutations of this protein have been reported in CAD,

DCM and HCM [58, 59]. 17 phosphorylation sites of cMyBP-C being phosphorylated by four various kinases have been identified so far; however, Ser282 residue (Ser284 in humans), which can be phosphorylated by CaMKII [60], is the most frequent target of phosphorylation *in vivo* [61]. Here, we have shown that phosphorylation of Ser284 residue was significantly downregulated while the total protein was unaltered in all HF groups. In support, other studies dealing with genetic mutations promoting cardiac dysfunction have reported total cMyBP-C being downregulated [59]. In context with p-Thr²⁸⁷-CaMKII δ , these observations open several questions. Similarly to p-Thr¹⁷-PLN, p-Ser²⁸⁴-cMyBP-C was decreased independently of the phosphorylation/oxidation of the upstream protein kinase CaMKII δ . However, as indicated above, it should be mentioned that the Ser284 residue of cMyBP-C can also be phosphorylated by some other protein kinases such as PKA, PKC or ribosomal S6 kinase [29]. Nevertheless, the phosphorylation of Ser284 mediated by CaMKII seems to play an important role because its inhibition resulted in the decreased phosphorylation of cMyBP-C [62, 63].

As the ability of the kinase to phosphorylate the proteins is counterbalanced by phosphatases we investigated the expression levels of main cardiac protein phosphatases PP1 and PP2A. The PP1 has been proposed to serve as a negative regulator of cardiac function. It is able to dephosphorylate substrates of CaMKII δ and thus may play an important role in gradual blunting of heart function during the development of HF [15, 64]. In this and in previously published studies [30, 43, 65], the expression of β subunit of PP1 was increased only in CAD and DCM failing hearts. Interestingly, the most abundant PP, PP2A did not differ among the HF groups and was comparable to the value of controls. The data about PP2A are controversial. The increased mRNA levels have been reported in DCM [43], while the protein content of this phosphatase was unchanged [66] supporting the findings of our current study.

Although this study has reported several novel findings and raised some new questions, there are few limitations. Firstly, the conclusions are based on proteomic data only. Nevertheless, we aimed to investigate an association between certain posttranslational modifications of Ca-

MKII δ and its downstream proteins, with cardiac function of all known types of HF. Secondly, phosphorylation status of proteins can be affected by certain drugs, however, by assuming that all HF patients were given the standard, almost the same therapy, this is unlikely to influence our present data.

Conclusion

In failing hearts of ischemic and non-ischemic origin, not differing in any main diagnostic characteristics, we have detected a different phosphorylation and oxidation status of CaMKII δ . The changes in posttranslational status of the kinase have not reflected the phosphorylation of its downstream targets neither in p-Ser²⁸⁴-cMyBP-C nor p-Thr¹⁷-PLN which, however were significantly downregulated and thereby could underlie the diminished cardiac contractile and relaxation function in HF. By investigating a link between the duration of the QT interval and the posttranslationally modified levels of CaMKII δ we have been unable to confirm that the altered activation of CaMKII δ promotes pro-arrhythmogenic environment. Thus, the assessment of a role of CaMKII δ , its active forms and potential consequences on cardiac function in particular forms of HF needs more detailed investigations.

Acknowledgements

This study was supported by Slovak Scientific Grant Agency (VEGA 1-0271-16), Slovak Research and Development Agency (APVV 15-0607), Slovak Society of Cardiology and Comenius University in Bratislava (UK/402/2016) and by the National Research, Development, and Innovation Office of Hungary (OTKA K 109737, National Heart Program NVKP 16-1-2016-0017). Authors would like to thank Dr. Gabriel Doka for organizing clinical patient data and Mrs. Veronika Hassova for her technical assistance.

Disclosure of conflict of interest

None.

Address correspondence to: Dr. Adriana Adameova, Department of Pharmacology and Toxicology, Faculty of Pharmacy, Comenius University, Bratislava, Odbojarov 10, 832 32, Slovak Republic. Tel: +421 2 50117 366; Fax: +421 2 50 117 100; E-mail: adameova@fpharm.uniba.sk

References

- [1] Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE, McMurray JJ, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WH, Tsai EJ, Wilkoff BL; American College of Cardiology Foundation; American Heart Association Task Force on Practice Guidelines. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American college of cardiology foundation/American heart association task force on practice guidelines. *Circulation* 2013; 128: e240-327.
- [2] Gersh BJ, Maron BJ, Bonow RO, Dearani JA, Fifer MA, Link MS, Naidu SS, Nishimura RA, Ommen SR, Rakowski H, Seidman CE, Towbin JA, Udelson JE, Yancy CW; American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines; American Association for Thoracic Surgery; American Society of Echocardiography; American Society of Nuclear Cardiology; Heart Failure Society of America; Heart Rhythm Society; Society for Cardiovascular Angiography and Interventions; Society of Thoracic Surgeons. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American college of cardiology foundation/American heart association task force on practice guidelines. *Circulation* 2011; 124: e783-831.
- [3] Kushwaha SS, Fallon JT, Fuster V. Restrictive cardiomyopathy. *N Engl J Med* 1997; 336: 267-276.
- [4] Stienen GJ. Pathomechanisms in heart failure: the contractile connection. *J Muscle Res Cell Motil* 2015; 36: 47-60.
- [5] Hund TJ, Mohler PJ. Role of CaMKII in cardiac arrhythmias. *Trends Cardiovasc Med* 2015; 25: 392-397.
- [6] Swaminathan PD, Purohit A, Soni S, Voigt N, Singh MV, Glukhov AV, Gao Z, He BJ, Luczak ED, Joiner ML, Kutschke W, Yang J, Donahue JK, Weiss RM, Grumbach IM, Ogawa M, Chen PS, Efimov I, Dobrev D, Mohler PJ, Hund TJ, Anderson ME. Oxidized CaMKII causes cardiac sinus node dysfunction in mice. *J Clin Invest* 2011; 121: 3277-3288.
- [7] Erickson JR. Mechanisms of CaMKII activation in the heart. *Front Pharmacol* 2014; 5: 59.
- [8] Marks AR. Calcium cycling proteins and heart failure: mechanisms and therapeutics. *J Clin Invest* 2013; 123: 46-52.
- [9] Haghghi K, Bidwell P, Kranias EG. Phospholamban interactome in cardiac contractility and survival: a new vision of an old friend. *J Mol Cell Cardiol* 2014; 77: 160-167.

CaMKII active forms in failing hearts

- [10] Schmitt JP, Kamisago M, Asahi M, Li GH, Ahmad F, Mende U, Kranias EG, MacLennan DH, Seidman JG, Seidman CE. Dilated cardiomyopathy and heart failure caused by a mutation in phospholamban. *Science* 2003; 299: 1410-1413.
- [11] Frank K, Böck B, Bavendiek U, Schwinger R. Frequency dependent force generation correlates with sarcoplasmic calcium ATPase activity in human myocardium. *Basic Res Cardiol* 1998; 93: 405-411.
- [12] Dash R, Frank KF, Carr AN, Moravec CS, Kranias EG. Gender influences on sarcoplasmic reticulum Ca²⁺-handling in failing human myocardium. *J Mol Cell Cardiol* 2001; 33: 1345-1353.
- [13] Kadambi VJ, Ponniah S, Harrer JM, Hoit BD, Dorn GW 2nd, Walsh RA. Cardiac-specific overexpression of phospholamban alters calcium kinetics and resultant cardiomyocyte mechanics in transgenic mice. *J Clin Invest* 1996; 97: 533-539.
- [14] Baker JO, Tyther R, Liebetrau C, Clark J, Howarth R, Patterson T, Möllmann H, Nef H, Sicard P, Kailey B, Devaraj R, Redwood SR, Kunst G, Weber E, Marber MS. Cardiac myosin-binding protein C: a potential early biomarker of myocardial injury. *Basic Res Cardiol* 2015; 110: 23.
- [15] Weber S, Meyer-Roxlau S, Wagner M, Dobrev D, El-Armouche A. Counteracting protein kinase activity in the heart: the multiple roles of protein phosphatases. *Front Pharmacol* 2015; 6: 270.
- [16] Koval OM, Guan X, Wu Y, Joiner ML, Gao Z, Chen B, Grumbach IM, Luczak ED, Colbran RJ, Song LS, Hund TJ, Mohler PJ, Anderson ME. CaV1.2-subunit coordinates CaMKII-triggered cardiomyocyte death and afterdepolarizations. *Proc Natl Acad Sci U S A* 2010; 107: 4996-5000.
- [17] Sag CM, Wadsack DP, Khabbazzadeh S, Abesser M, Grefe C, Neumann K, Opiela MK, Backs J, Olson EN, Brown JH, Neef S, Maier SK, Maier LS. Calcium/calmodulin-dependent protein kinase II contributes to cardiac arrhythmogenesis in heart failure. *Circ Heart Fail* 2009; 2: 664-675.
- [18] Lip GY, Heinzel FR, Gaita F, Juanatey JR, Le Heuzey JY, Potpara T, Svendsen JH, Vos MA, Anker SD, Coats AJ, Haverkamp W, Manolis AS, Chung MK, Sanders P, Pieske B, Gorenek B, Lane D, Boriani G, Linde C, Hindricks G, Tsutsui H, Homma S, Brownstein S, Nielsen JC, Lainscak M, Crespo-Leiro M, Piepoli M, Seferovic P, Savelieva I; EP-Europace. European Heart Rhythm Association/Heart Failure Association joint consensus document on arrhythmias in heart failure, endorsed by the Heart Rhythm Society and the Asia Pacific Heart Rhythm Society. *Eurpace* 2016; 18: 12-36.
- [19] Rajtik T, Carnicka S, Szobi A, Giricz Z, O-Uchi J, Hassova V, Svec P, Ferdinandy P, Ravingerova T, Adameova A. Oxidative activation of CaMKII δ in acute myocardial ischemia/reperfusion injury: a role of angiotensin AT1 receptor-NOX2 signaling axis. *Eur J Pharmacol* 2016; 771: 114-122.
- [20] Romero-Calvo I, Ocón B, Martínez-Moya P, Suárez MD, Zarzuelo A, Martínez-Augustín O, de Medina FS. Reversible ponceau staining as a loading control alternative to actin in Western blots. *Anal Biochem* 2010; 401: 318-320.
- [21] Gilda JE, Gomes AV. Stain-Free total protein staining is a superior loading control to β -actin for Western blots. *Anal Biochem* 2013; 440: 186-188.
- [22] Welinder C, Ekblad L. Coomassie staining as loading control in Western blot analysis. *J Proteome Res* 2011; 10: 1416-1419.
- [23] Anderson ME, Braun AP, Wu Y, Lu T, Wu Y, Schulman H, Sung RJ. KN-93, an inhibitor of multifunctional Ca⁺⁺/calmodulin-dependent protein kinase, decreases early afterdepolarizations in rabbit heart. *J Pharmacol Exp Ther* 1998; 287: 996-1006.
- [24] Kirchhof P, Fabritz L, Kilić A, Begrow F, Breithardt G, Kuhn M. Ventricular arrhythmias, increased cardiac calmodulin kinase II expression, and altered repolarization kinetics in ANP receptor deficient mice. *J Mol Cell Cardiol* 2004; 36: 691-700.
- [25] Luo M, Anderson ME. Mechanisms of altered Ca²⁺ handling in heart failure. *Circ Res* 2013; 113: 690-708.
- [26] Mattiazzi A, Mundiña-Weilenmann C, Guoxiang C, Vittone L, Kranias E. Role of phospholamban phosphorylation on Thr17 in cardiac physiological and pathological conditions. *Cardiovasc Res* 2005; 68: 366-375.
- [27] Kimura Y, Kurzydowski K, Tada M, MacLennan DH. Phospholamban inhibitory function is activated by depolymerization. *J Biol Chem* 1997; 272: 15061-15064.
- [28] Smeazzetto S, Saponaro A, Young HS, Moncelli MR, Thiel G. Structure-function relation of phospholamban: modulation of channel activity as a potential regulator of SERCA activity. *PLoS One* 2013; 8: e52744.
- [29] Moss RL, Fitzsimons DP, Ralphe JC. Cardiac MyBP-C regulates the rate and force of contraction in mammalian myocardium. *Circ Res* 2015; 116: 183-192.
- [30] Aoyama H, Ikeda Y, Miyazaki Y, Yoshimura K, Nishino S, Yamamoto T, Yano M, Inui M, Aoki H, Matsuzaki M. Isoform-specific roles of protein phosphatase 1 catalytic subunits in sarcoplasmic reticulum-mediated Ca²⁺ cycling. *Cardiovasc Res* 2011; 89: 79-88.

CaMKII active forms in failing hearts

- [31] Terrak M, Kerff F, Langsetmo K, Tao T, Dominguez R. Structural basis of protein phosphatase 1 regulation. *Nature* 2004; 429: 780-784.
- [32] Cho US, Xu W. Crystal structure of a protein phosphatase 2A heterotrimeric holoenzyme. *Nature* 2007; 445: 53-57.
- [33] Schlender KK, Hegazy MG, Thysseril TJ. Dephosphorylation of cardiac myofibril C-protein by protein phosphatase 1 and protein phosphatase 2A. *Biochim Biophys Acta* 1987; 928: 312-319.
- [34] Noguchi T, Hünlich M, Camp PC, Begin KJ, El-Zaru M, Patten R, Leavitt BJ, Littleman FP, Alpert NR, LeWinter MM, VanBuren P. Thin-filament-based modulation of contractile performance in human heart failure. *Circulation* 2004; 110: 982-987.
- [35] Strack S, Barban MA, Wadzinski BE, Colbran RJ. Differential inactivation of postsynaptic density-associated and soluble Ca²⁺/calmodulin-dependent protein kinase II by protein phosphatases 1 and 2A. *J Neurochem* 1997; 68: 2119-2128.
- [36] Meyer T, Hanson PI, Stryer L, Schulman H. Calmodulin trapping by calcium-calmodulin-dependent protein kinase. *Science* 1992; 256: 1199-1202.
- [37] Guo T, Zhang T, Mestrlil R, Bers DM. Ca²⁺/calmodulin-dependent protein kinase II phosphorylation of ryanodine receptor does affect calcium sparks in mouse ventricular myocytes. *Circ Res* 2006; 99: 398-406.
- [38] Simmerman HK, Collins JH, Theibert JL, Wegener AD, Jones LR. Sequence analysis of phospholamban. Identification of phosphorylation sites and two major structural domains. *J Biol Chem* 1986; 261: 13333-13341.
- [39] Schlender KK, Bean LJ. Phosphorylation of chicken cardiac C-protein by calcium/calmodulin-dependent protein kinase II. *J Biol Chem* 1991; 266: 2811-2817.
- [40] Anderson ME, Brown JH, Bers DM. CaMKII in myocardial hypertrophy and heart failure. *J Mol Cell Cardiol* 2011; 51: 468-473.
- [41] Kirchhefer U, Schmitz W, Scholz H, Neumann J. Activity of cAMP-dependent protein kinase and Ca²⁺/calmodulin-dependent protein kinase in failing and nonfailing human hearts. *Cardiovasc Res* 1999; 42: 254-261.
- [42] Fischer TH, Eiringhaus J, Dybkova N, Förster A, Herting J, Kleinwächter A, Ljubojevic S, Schmitto JD, Streckfuß-Bömeke K, Renner A, Gummert J, Hasenfuss G, Maier LS, Sossalla S. Ca²⁺/calmodulin-dependent protein kinase II equally induces sarcoplasmic reticulum Ca²⁺ leak in human ischaemic and dilated cardiomyopathy. *Eur J Heart Fail* 2014; 16: 1292-1300.
- [43] Miyamoto SD, Stauffer BL, Nakano S, Sobus R, Nunley K, Nelson P, Sucharov CC. Beta-adrenergic adaptation in paediatric idiopathic dilated cardiomyopathy. *Eur Heart J* 2014; 35: 33-41.
- [44] Zhang T, Maier LS, Dalton ND, Miyamoto S, Ross J Jr, Bers DM, Brown JH. The deltaC isoform of CaMKII is activated in cardiac hypertrophy and induces dilated cardiomyopathy and heart failure. *Circ Res* 2003; 92: 912-919.
- [45] Dhalla NS, Temsah RM, Netticadan T. Role of oxidative stress in cardiovascular diseases. *J Hypertens* 2000; 18: 655-673.
- [46] Wagner S, Ruff HM, Weber SL, Bellmann S, Sowa T, Schulte T, Anderson ME, Grandi E, Bers DM, Backs J, Belardinelli L, Maier LS. Reactive oxygen species-activated Ca/calmodulin kinase II δ is required for late I(Na) augmentation leading to cellular Na and Ca overload. *Circ Res* 2011; 108: 555-565.
- [47] Velez Rueda JO, Palomeque J, Mattiazzi A. Early apoptosis in different models of cardiac hypertrophy induced by high renin-angiotensin system activity involves CaMKII. *J Appl Physiol* (1985) 2012; 112: 2110-2120.
- [48] Erickson JR, Joiner ML, Guan X, Kutschke W, Yang J, Oddis CV, Bartlett RK, Lowe JS, O'Donnell SE, Aykin-Burns N, Zimmerman MC, Zimmerman K, Ham AJ, Weiss RM, Spitz DR, Shea MA, Colbran RJ, Mohler PJ, Anderson ME. A dynamic pathway for calcium-independent activation of CaMKII by methionine oxidation. *Cell* 2008; 133: 462-474.
- [49] Palomeque J, Rueda OV, Sapia L, Valverde CA, Salas M, Petroff MV, Mattiazzi A. Angiotensin II-induced oxidative stress resets the Ca²⁺ dependence of Ca²⁺-calmodulin protein Kinase II and promotes a death pathway conserved across different species. *Circ Res* 2009; 105: 1204-1212.
- [50] Mundiña-Weilenmann C, Vittone L, Ortale M, de Cingolani GC, Mattiazzi A. Immunodetection of phosphorylation sites gives new insights into the mechanisms underlying phospholamban phosphorylation in the intact heart. *J Biol Chem* 1996; 271: 33561-3367.
- [51] Schwinger RH, Böhm M, Schmidt U, Karczewski P, Bavendiek U, Flesch M, Krause EG, Erdmann E. Unchanged protein levels of SERCA II and phospholamban but reduced Ca²⁺ uptake and Ca(2+)-ATPase activity of cardiac sarcoplasmic reticulum from dilated cardiomyopathy patients compared with patients with nonfailing hearts. *Circulation* 1995; 92: 3220-3228.
- [52] Meyer M, Schillinger W, Pieske B, Holubarsch C, Heilmann C, Posival H, Kuwajima G, Miko-shiba K, Just H, Hasenfuss G. Alterations of sarcoplasmic reticulum proteins in failing hu-

CaMKII active forms in failing hearts

- man dilated cardiomyopathy. *Circulation* 1995; 92: 778-784.
- [53] Luo W, Wolska BM, Grupp IL, Harrer JM, Haghghi K, Ferguson DG, Slack JP, Grupp G, Doetschman T, Solaro RJ, Kranias EG. Phospholamban gene dosage effects in the mammalian heart. *Circ Res* 1996; 78: 839-847.
- [54] Clarke JD, Caldwell JL, Horn MA, Bode EF, Richards MA, Hall MC, Graham HK, Briston SJ, Greensmith DJ, Eisner DA, Dibb KM, Trafford AW. Perturbed atrial calcium handling in an ovine model of heart failure: potential roles for reductions in the I-type calcium current. *J Mol Cell Cardiol* 2015; 79: 169-179.
- [55] Lynch TL, Sadayappan S. Surviving the infarct: a profile of cardiac myosin binding protein-C pathogenicity, diagnostic utility, and proteomics in the ischemic myocardium. *Proteomics Clin Appl* 2014; 8: 569-577.
- [56] Stathopoulou K, Wittig I, Heidler J, Piasecki A, Richter F, Diering S, van der Velden J, Buck F, Donzelli S, Schröder E, Wijnker PJ, Voigt N, Dobrev D, Sadayappan S, Eschenhagen T, Carrier L, Eaton P, Cuello F. S-glutathiolation impairs phosphoregulation and function of cardiac myosin-binding protein C in human heart failure. *FASEB J* 2016; 30: 1849-1864.
- [57] van Dijk SJ, Holewijn RA, Tebeest A, Dos Remedios C, Stienen GJ, van der Velden J. A piece of the human heart: variance of protein phosphorylation in left ventricular samples from end-stage primary cardiomyopathy patients. *J Muscle Res Cell Motil* 2009; 30: 299-302.
- [58] Gajendrarao P, Krishnamoorthy N, Selvaraj S, Girolami F, Cecchi F, Olivotto I, Yacoub M. An investigation of the molecular mechanism of double cMyBP-C mutation in a patient with end-stage hypertrophic cardiomyopathy. *J Cardiovasc Transl Res* 2015; 8: 232-243.
- [59] Kuster DW, Govindan S, Springer TI, Martin JL, Finley NL, Sadayappan S. A Hypertrophic Cardiomyopathy-associated MYBPC3 mutation common in populations of south Asian descent causes contractile dysfunction. *J Biol Chem* 2015; 290: 5855-5867.
- [60] Gautel M, Zuffardi O, Freiburg A, Labeit S. Phosphorylation switches specific for the cardiac isoform of myosin binding protein-C: a modulator of cardiac contraction? *EMBO J* 1995; 14: 1952-1960.
- [61] Kooij V, Holewinski RJ, Murphy AM, Van Eyk JE. Characterization of the cardiac myosin binding protein-C phosphoproteome in healthy and failing human hearts. *J Mol Cell Cardiol* 2013; 60: 116-120.
- [62] Sadayappan S, Gulick J, Osinska H, Barefield D, Cuello F, Avkiran M, Lasko VM, Lorenz JN, Maillet M, Martin JL, Brown JH, Bers DM, Molkentin JD, James J, Robbins J. A critical function for Ser-282 in cardiac myosin binding protein-C phosphorylation and cardiac function. *Circ Res* 2011; 109: 141-150.
- [63] Tong CW, Gaffin RD, Zawieja DC, Muthuchamy M. Roles of phosphorylation of myosin binding protein-C and troponin I in mouse cardiac muscle twitch dynamics. *J Physiol* 2004; 558: 927-941.
- [64] Carr AN, Schmidt AG, Suzuki Y, del Monte F, Sato Y, Lanner C, Breeden K, Jing SL, Allen PB, Greengard P, Yatani A, Hoit BD, Grupp IL, Hajjar RJ, DePaoli-Roach AA, Kranias EG. Type 1 phosphatase, a negative regulator of cardiac function. *Mol Cell Biol* 2002; 22: 4124-4135.
- [65] Hamdani N, Borbély A, Veenstra SP, Kooij V, Vrydag W, Zaremba R, Dos Remedios C, Niesen HW, Michel MC, Paulus WJ, Stienen GJ, van der Velden J. More severe cellular phenotype in human idiopathic dilated cardiomyopathy compared to ischemic heart disease. *J Muscle Res Cell Motil* 2010; 31: 289-301.
- [66] Wijnker PJ, Boknik P, Gergs U, Müller FU, Neumann J, dos Remedios C, Schmitz W, Sindermann JR, Stienen GJ, van der Velden J, Kirchhefer U. Protein phosphatase 2A affects myofilament contractility in non-failing but not in failing human myocardium. *J Muscle Res Cell Motil* 2011; 32: 221-233.