

# Ten-year inhospital mortality trends for patients with trauma in Japan: a multicentre observational study

著者	Nagata Isao, Abe Toshikazu, Uchida Masatoshi, Saitoh Daizoh, Tamiya Nanako
journal or publication title	BMJ Open
volume	8
number	2
page range	e018635
year	2018-02
権利	This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <a href="http://creativecommons.org/licenses/by-nc/4.0/">http://creativecommons.org/licenses/by-nc/4.0/</a> (C) Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.
URL	<a href="http://hdl.handle.net/2241/00153089">http://hdl.handle.net/2241/00153089</a>

doi: 10.1136/bmjopen-2017-018635

# BMJ Open Ten-year in-hospital mortality trends for patients with trauma in Japan: a multicentre observational study

Isao Nagata,<sup>1,2,3</sup> Toshikazu Abe,<sup>1,4</sup> Masatoshi Uchida,<sup>2</sup> Daizoh Saitoh,<sup>5</sup> Nanako Tamiya<sup>1</sup>

**To cite:** Nagata I, Abe T, Uchida M, *et al*. Ten-year in-hospital mortality trends for patients with trauma in Japan: a multicentre observational study. *BMJ Open* 2018;**8**:e018635. doi:10.1136/bmjopen-2017-018635

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-018635>).

Received 11 July 2017  
Revised 26 September 2017  
Accepted 8 November 2017



<sup>1</sup>Department of Health Services Research, Faculty of Medicine, University of Tsukuba, Tsukuba, Japan

<sup>2</sup>Graduate School of Comprehensive Human Sciences, Majors of Medical Sciences, University of Tsukuba, Tsukuba, Japan

<sup>3</sup>Department of Critical Care Medicine, Yokohama City Minato Red Cross Hospital, Yokohama, Japan

<sup>4</sup>Department of Emergency and Critical Care Medicine, Juntendo University Urayasu Hospital, Urayasu, Japan

<sup>5</sup>Department of Traumatology and Emergency Medicine, National Defense Medical College, Tokorozawa, Japan

## Correspondence to

Toshikazu Abe;  
abetoshi11@gmail.com

## ABSTRACT

**Objectives** Trauma is one of the main causes of death in Japan, and treatments and prognoses of these injuries are constantly changing. We therefore aimed to investigate a 10-year trend (2004–2013) in in-hospital mortality among patients with trauma in Japan.

**Design** Multicentre observational study.

**Setting** Japanese nationwide trauma registry (the Japan Trauma Data Bank) data.

**Participants** All patients with trauma whose Injury Severity Score (ISS) were 3 and above, who were aged 15 years or older, and whose mechanisms of injury (MOI) were blunt and penetrating between 2004 and 2013 (n=90 833).

**Outcome measures** A 10-year trend in in-hospital mortality.

**Results** In-hospital mortality for all patients with trauma significantly decreased over the study decade in our Cochran-Armitage test ( $P<0.001$ ). Similarly, in-hospital mortality for patients with ISS 16 or more and patients who scored 50% or better on the Trauma and Injury Severity Score (TRISS) probability of survival scale significantly decreased ( $P<0.001$ ). In addition, the OR for in-hospital mortality of these three patient groups decreased yearly after adjusting for age, gender, MOI, ISS, Glasgow Coma Scale, systolic blood pressure and respiratory rate on hospital arrival in multivariable logistic regression analyses. Furthermore, in-hospital mortality for patient with blunt trauma significantly decreased in injury mechanism-stratified Mantel-extension testing ( $P<0.001$ ). Finally, multivariable logistic regression analyses showed that the OR for in-hospital mortality of patients with ISS 16 and over decreased each year after adding and adjusting for means of transportation and usage of whole-body CT.

**Conclusion** In-hospital mortality for patients with trauma in Japan significantly decreased during the study decade after adjusting for patient characteristics, injury severity and the response environment after injury.

## INTRODUCTION

Unintentional injury has been in fifth or sixth place in cause of death for over a decade in Japan. Additionally, unintentional injury is a leading cause of death for Japanese youth (ages 1–29 years). Mostly, unintentional injury includes trauma.<sup>1</sup> Therefore, it is important to analyse patterns of injury and treatment

## Strengths and limitations of this study

- We showed detailed trauma trends for these 10 years (2004–2013) using Japanese nationwide trauma registry data.
- We analysed in-hospital mortality trends adjusting for patient characteristics, injury severity and the response environment after injury.
- We could not investigate regional and facility differences in our study.

to establish guidelines that can prevent trauma-related deaths.

Over the past 10 years, the causes of trauma have fluctuated with traffic accidents decreasing, but falls increasing.<sup>2</sup> Proactive trauma education has improved; the Japan Advanced Trauma Evaluation and Care association was started in 2002 to create trauma treatment guidelines, develop the off-the-job training based on these guidelines and enact treatment that meets such standards.<sup>3</sup> Also, the Japan Trauma Data Bank (JTDB) was established in 2003 by the Japanese Association for the Surgery of Trauma and the Japanese Association for Acute Medicine to improve and assure the quality of trauma care in Japan.<sup>4</sup> Furthermore, there are important yearly advancements in trauma care that may improve outcomes.

On the other hand, in Japan, ageing has accelerated rapidly and the elderly comprise the majority of patients with trauma who die due to traffic accident-related trauma.<sup>5</sup> This is supported by the literature; a systematic review and meta-analysis about older patients with trauma reported that older patients with trauma suffered higher mortality.<sup>6</sup> With such a complex and fluctuating subject, it is important to continuously update and improve trauma guidelines to reflect changes in patient injury types and demographics.

However, long-term detailed analyses concerning trauma prognoses for Japanese

patients are scarce in the literature. Annual fluctuations and critical turning points for patient with trauma prognoses are also unknown. Therefore, we aimed to investigate the trend in inhospital mortality for Japanese patients with trauma over a recent 10-year time period to indicate where improvements in future guidelines should be focused.

## METHODS

### Study design and data source

This was a multicentre retrospective observational study using JTDB data. The participation to the JTDB data collection started from 55 hospitals. Participant hospitals increased year over year, and a total of 221 hospitals, including 85% of tertiary emergency medical centres in Japan in March, 2013 (n=259), participated in the JTDB data collection in March, 2013. The inclusion criteria for the JTDB was patients with trauma who had Injury Severity Score (ISS) of 3 or above. The JTDB collected prehospital and hospital information such as patient demographics, comorbidities, injury types, mechanisms of injury (MOI), means of transportation, vital signs, Abbreviated Injury Scale score, prehospital treatment, inhospital procedures and inhospital mortality.<sup>7</sup>

Because of the anonymous and retrospective nature of this study, the need for informed consent was waived.

### Patients selection

We screened every patient who was registered in the JTDB from 1 January 2004 to 31 March 2013, and enrolled patients with trauma whose ISS were 3 or higher, who were 15 or more years old and whose MOI were blunt and penetrating traumas. Patients were excluded from analysis if they had out-of-hospital cardiopulmonary arrest (OHCA) (chest compression in prehospital setting, systolic blood pressure of 0mm Hg and pulse rate of

0/min at the scene). Also, we excluded any patients who had missing data for age, gender, MOI, Revised Trauma Score (RTS), Trauma and Injury Severity Score (TRISS), Glasgow Coma Scale (GCS), systolic blood pressure and respiratory rate on hospital arrival or mortality. Figure 1 shows participant selection data from this study.


### Data collection and end point

The JTDB data included the following information: age, gender, means of transportation, mechanism of injury, injury mechanism of blunt trauma, ISS, RTS, TRISS, GCS, systolic blood pressure and respiratory rate on hospital arrival, usage of whole-body CT and interventional radiology (IVR) for abdomen and pelvis, hospital location, length-of-hospital stay, inhospital mortality and disposition at discharge. Injury mechanism of blunt trauma was classified as motor vehicles, motorcycles, bicycles, pedestrian accidents, falls from high place, falls down and falls at same level. Falls from high place were defined as free falls from high place like a building. Falls down were defined as falls from stair or slope.

The primary outcome of this study was the trend in inhospital mortality and the data collection endpoint was determined by the end of the Japanese fiscal year in March 2013 for a total of 10 years of data (2004–2013).

### Statistical analysis

To reveal characteristics of patient demographics, backgrounds, examinations and treatments in relation to inhospital mortality and hospitalisation, non-normally distributed continuous data were presented as medians and IQR and categorical data were summarised as percentages with counts. As a primary analysis to find trends of inhospital mortality within the specified decade of data, the Cochran-Armitage test and multivariable logistic regression analyses were conducted. The Cochran-Armitage test is used to test for the trend among binominal



**Figure 1** Patients flow chart. GCS, Glasgow Coma Scale; ISS, Injury Severity Scores; JTDB, Japan Trauma Data Bank; OHCA, out of hospital cardiopulmonary arrest; RTS, Revised Trauma Score; TRISS, Trauma and Injury Severity Score.

proportions against ordinal explanatory variables. The following variables were applied to multivariable logistic regression analyses: age, gender, MOI, ISS, GCS, systolic blood pressure and respiratory rate on hospital arrival. In addition, the Cochran-Armitage test and multivariable logistic regression analyses were repeated for patients with ISS scores of 16 or higher and patients who scored 50% or better on the TRISS probability of survival scale. A three-step approach was used to analyse sensitivity. First, the Cochran-Armitage test was conducted among patients with trauma included any patients who had missing data to investigate the influence of it. Second, the Mantel-extension test was conducted to investigate the influence of injury mechanism blunt trauma on the 10-year trend in in-hospital mortality. Last, a multivariable logistic regression analysis was repeated for patients with ISS scores of 16 or higher after adjusting for means of transportation and usage of whole-body CT in addition to the same variables as the primary analysis. Data are presented as ORs with 95% CIs. P values for both the Cochran-Armitage and Mantel-extension tests were one sided, and P values for the multivariable logistic regression analyses were two sided.  $P < 0.05$  was considered statistically significant. All analyses were performed using SAS V.9.4.

## RESULTS

Out of a total of 159157 patients who were registered in the JTDB during the study period, 139463 patients with trauma who had ISS scores of 3 or higher from 2004 to 2013 were included. Cases were excluded if they were younger than 15 years old ( $n=7523$ ), if they had OHCA ( $n=5146$ ) or if they had any missing data for age, gender, MOI, RTS, TRISS, GCS, systolic blood pressure and respiratory rate on hospital arrival and mortality ( $n=31650$ ). Exclusions were also made for burn ( $n=3202$ ) or if the cause of injury mechanisms was unknown ( $n=1109$ ). After these adjustments, 90883 patients were eligible for inclusion in this study (figure 1).

Table 1 shows demographics and other data of patients with trauma by years. Mean age increased from 47 in 2004 to 64 in 2013, and rates in adult men showed a decreasing trend year by year. Helicopter transportation increased from 5.8% in 2004 to 8.0% in 2013. The ratio of blunt trauma shifted into the higher 90% range over the entire decade of observation. With regard to injury mechanisms of blunt trauma, traffic accidents saw a decrease, and falls down/falls at same level showed a yearly increase in trend. Trauma severity scores like ISS and RTS, as well as TRISS probability of survival scores of less than 50%, had a tendency to decrease.

Yearly examination, treatment and hospitalisation data are shown in table 2. In the examination data, the performance rate of whole-body CT increase in all patients as well as patients with ISS scores of 16 or higher. With regard to treatment, the performance rate of abdominal and pelvic IVR fluctuated between 3% and 5% over the entire decade. For hospitalisation data, the rate of patients

who were hospitalised in emergency and critical care centres decreased and the rate of general hospitalisation increased in a yearly fashion. The average length-of-hospital stay varied between 14 days and 18 days during the study period. Disposition at discharge classifications saw a steady trend of half the patients designated as 'home' and half designated as 'hospital transfer'.

The decade-long in-hospital mortality trend is shown in figure 2 and table 3. The results of Cochran-Armitage testing (figure 2) were that in-hospital mortality for all patients with trauma significantly decreased over the study decade (from 15.8% in 2004 to 6.6% in 2013,  $P < 0.001$ ). Similarly, in-hospital mortality for patients with ISS scores 16 or higher and patients with a 50% or better score on the TRISS probability of survival scale significantly decreased over the same period (from 28.5% in 2004 to 15.7% in 2013 among patients with ISS scores 16 or higher and from 8.4% in 2004 to 3.1% in 2013 among patients with a 50% or better score on the TRISS probability of survival scale,  $P < 0.001$ ). In multivariable logistic regression analyses with 2009 as the comparative control, all patients with trauma between 2004 and 2006 had higher ORs for in-hospital mortality and those between 2011 and 2013 had lower ORs. In a similar comparison in patients with ISS scores of 16 or higher and patients with a 50% or better TRISS score, we saw a higher OR for in-hospital mortality from 2004 to 2006 versus those from 2011 to 2013 when 2009 was used as the baseline.

With respect to sensitivity, the result of Cochran-Armitage testing conducted among patients with trauma included any patients who had missing data ( $n=112303$ ) was that in-hospital mortality significantly decreased over the study decade (from 15.2% in 2004 to 6.6% in 2013,  $P < 0.001$ ). In addition, the result of injury mechanism-stratified Mantel-extension testing was that in-hospital mortality for patients with blunt trauma significantly decreased over the study decade ( $P < 0.001$ ). Although means of transportation and usage of whole-body CT variables were included in our multivariable logistic regression analyses, yearly ORs for in-hospital mortality among patients in the ISS 16 or higher classification did not change significantly compared with primary analyses (table 4).

## DISCUSSION

### Brief summary

In this study, we saw a significant yearly decrease in in-hospital mortality for all patients with trauma in Japan over the study decade. Similarly, in-hospital mortality for patients with ISS 16 or more and patients with a 50% or better score on the TRISS probability of survival scale significantly decreased each year. In addition, multivariable logistic regression analyses saw decreasing ORs for in-hospital mortality of these three patient groups each year after adjusting for age, gender, MOI, ISS and vital signs.



**Table 1** Demographics and other data of patients with trauma by years

Year	2004, n=2209	2005, n=2708	2006, n=2766	2007, n=6674	2008, n=8020	2009, n=9788	2010, n=12544	2011, n=14907	2012, n=17124	2013, n=14093
Age, years (range)	47 (28–65)	50 (29–66)	53 (32–67)	56 (33–71)	58 (36–72)	60 (38–75)	60 (38–76)	62 (40–77)	63 (42–78)	64 (43–79)
Gender, n (%)										
Male	1610/2209 (72.9)	1881/2708 (69.5)	1892/2766 (68.4)	4491/6674 (67.3)	5451/8020 (68.0)	6303/9788 (64.4)	8002/12544 (63.8)	9526/14907 (63.9)	10 835/17124 (63.3)	8674/14093 (61.6)
Transportation, n (%)										
Ambulance	1975/2188 (90.3)	2402/2676 (89.8)	2399/2730 (87.9)	5584/6520 (85.6)	6838/7881 (86.8)	8004/9629 (83.1)	10403/12344 (84.3)	12134/14645 (82.9)	13692/16883 (81.1)	11270/13902 (81.1)
Ambulance with doctors	44/2188 (2.0)	70/2676 (2.6)	50/2730 (1.8)	201/6520 (3.1)	181/7881 (2.3)	239/9629 (2.5)	341/12344 (2.8)	425/14645 (2.9)	611/16883 (3.6)	397/13902 (2.8)
Helicopter	127/2188 (5.8)	133/2676 (4.9)	168/2730 (6.2)	465/6520 (7.1)	579/7881 (7.3)	748/9629 (7.8)	802/12344 (6.5)	1090/14645 (7.4)	1392/16883 (8.3)	1107/13902 (8.0)
Others*	42/2188 (1.9)	71/2676 (2.7)	113/2730 (4.1)	270/6520 (4.2)	283/7881 (3.6)	638/9629 (6.6)	798/12344 (6.4)	996/14645 (6.8)	1188/16883 (7.0)	1128/13902 (8.1)
Mechanism of injury, n (%)										
Blunt	2115/2209 (95.7)	2604/2708 (96.2)	2650/2766 (95.8)	6458/6674 (96.8)	7738/8820 (96.5)	9455/9788 (96.6)	12159/12544 (96.9)	14485/14907 (97.2)	16653/17124 (97.2)	13724/14093 (97.4)
Penetrating	94/2209 (4.3)	104/2708 (3.8)	116/2766 (4.2)	216/6674 (3.2)	282/8020 (3.5)	333/9788 (3.4)	385/12544 (3.1)	422/14907 (2.8)	471/17124 (2.8)	369/14093 (2.6)
Injury mechanism of blunt trauma, n (%)										
Traffic accident										
Motor vehicle	337/2030 (16.6)	455/2688 (16.9)	464/2746 (16.9)	950/6562 (14.5)	974/7738 (12.6)	1191/9415 (12.6)	1471/11974 (12.3)	1674/14520 (11.5)	2057/16839 (12.2)	1613/13918 (11.6)
Bike	434/2030 (21.4)	511/2688 (19.0)	473/2746 (17.2)	1192/6562 (18.2)	1297/7738 (16.7)	1447/9415 (15.4)	1737/11974 (14.5)	1931/14520 (13.3)	2128/16839 (12.7)	1723/13918 (12.4)
Bicycle	213/2030 (10.5)	251/2688 (9.4)	244/2746 (8.9)	503/6562 (7.7)	679/7738 (8.8)	821/9415 (8.7)	1028/11974 (8.6)	1197/14520 (8.2)	1251/16839 (7.4)	1025/13918 (7.3)
Pedestrian	207/2030 (10.2)	258/2688 (9.6)	262/2746 (9.5)	538/6562 (8.2)	734/7738 (9.5)	783/9415 (8.3)	949/11974 (7.9)	1085/14520 (7.5)	1242/16839 (7.4)	948/13918 (6.8)
Fall from high place†	300/2030 (14.8)	362/2688 (13.5)	379/2746 (13.8)	922/6562 (14.0)	997/7738 (12.9)	1065/9415 (11.3)	1335/11974 (11.1)	1608/14520 (11.1)	1678/16839 (10.0)	1250/13918 (9.0)
Fall down‡	145/2030 (7.1)	199/2688 (7.4)	254/2746 (9.3)	663/6562 (10.1)	858/7738 (11.1)	970/9415 (10.3)	1407/11974 (11.8)	1851/14520 (12.7)	2262/16839 (13.4)	1820/13918 (13.1)
Fall at same level	140/2030 (6.9)	291/2688 (10.8)	315/2746 (11.5)	985/6562 (15.0)	1306/7738 (16.9)	2022/9415 (21.5)	2749/11974 (23.0)	3610/14520 (24.9)	4450/16839 (26.4)	407/13918 (29.1)
Others§	254/2030 (12.5)	361/2688 (13.4)	355/2746 (12.9)	809/6562 (12.3)	893/7738 (11.5)	1116/9415 (11.9)	1298/11974 (10.8)	1564/14520 (10.8)	1771/16839 (10.5)	1492/13918 (10.7)
ISS	16 (9–25)	16 (9–25)	16 (9–25)	16 (9–24)	16 (9–25)	14 (9–22)	13 (9–22)	13 (9–22)	13 (9–21)	10 (9–19)
RTS	7.84 (6.61–7.84)	7.84 (6.90–7.84)	7.84 (6.90–7.84)	7.84 (7.10–7.84)	7.84 (6.90–7.84)	7.84 (7.11–7.84)	7.84 (7.55–7.84)	7.84 (7.55–7.84)	7.84 (7.55–7.84)	7.84 (7.55–7.84)

Continued

**Table 1** Continued

Year	2004, n=2209	2005, n=2708	2006, n=2766	2007, n=6674	2008, n=8020	2009, n=9788	2010, n=12544	2011, n=14907	2012, n=17124	2013, n=14093
TRISS probability of survival, n (%)										
50% ≤	1947/2209 (88.1)	2437/2708 (90.0)	2465/2766 (89.1)	6112/6674 (91.6)	7271/8020 (90.7)	9010/9788 (92.1)	11674/12544 (93.1)	13833/14907 (92.8)	15895/17124 (92.8)	13309/14093 (94.4)
<50%	262/2209 (11.9)	271/2708 (10.0)	301/2766 (10.9)	562/6674 (8.4)	749/8020 (9.3)	778/9788 (7.9)	870/12544 (6.9)	1074/14907 (7.2)	1229/17124 (7.2)	784/14093 (5.6)

\*Private car, unknown and others.

†Free falls from high place like a building.

‡Falls from stair or slope and so on.

§Train, machine, falling object, explosion, compression and others.

ISS, Injury Severity Scores; RTS, Revised Trauma Score; TRISS, Trauma and Injury Severity Score.

**Relationship to previous studies**

Our results are in accordance with several observational studies that reported trends in in-hospital mortality for patients with trauma.<sup>8-13</sup> A retrospective cohort study from Israel reported that in-hospital mortality of patients with ISS scores of 16 or higher decreased significantly from 22% in 2000 to 16% in 2010 and that the OR for in-hospital mortality in 2010 compared with 2000 was 0.53, confirming a downward trend. The authors mentioned that establishment of a trauma response system brought a significant decrease in in-hospital mortality although any single factor that explained the reduction was not identified.<sup>8</sup> A retrospective study performed in New South Wales, Australia used multivariable logistic regression analyses to show that the OR for in-hospital mortality of patients with an ISS score higher than 15 in 2007 compared with 2003 was 0.71 and that the OR for in-hospital mortality was significantly lower among patients receiving definitive care at level I trauma centres compared with regional trauma centres.<sup>9</sup> Also, in Japan, a retrospective observational study reported that adjusted in-hospital trauma mortality was significantly lower in patients from 2009 to 2011 than from 2004 to 2006 (OR 0.64 in 2009–2011 compared with 2004–2006) possibly due to establishment of trauma education.<sup>10</sup> In previous studies, it was mentioned that changes in the trauma response system, such as establishment of trauma education in the hospitals where patients were transported, contributed to the decrease in in-hospital mortality. However, it is difficult to explain reductions of in-hospital mortality with any single factor.

Our results showed changes in the injury mechanisms within the blunt trauma category. Traffic accidents had a tendency to decrease, and falls down/falls at same level showed a yearly increase in trend. A previous study reported that the in-hospital mortality OR of fall victims were lower than that of motor vehicle crash (MVC) victims in a multivariable logistic analysis (OR 0.60 in fall, reference in MVC).<sup>14</sup> Changes in the injury mechanisms within the blunt trauma category might contribute to reductions of in-hospital mortality in our study. Therefore, we conducted an injury mechanism-stratified Mantel-extension test for blunt trauma and investigated the relationship between year and in-hospital mortality among patients with blunt trauma. However, in-hospital mortality for patients with blunt trauma significantly decreased each year after stratifying for injury mechanism, and we noticed that this stratification did not affect trends in in-hospital mortality with respect to this test. In addition, our study investigated changes in emergency transportation methods and CT imaging methods because previous studies reported that helicopter transportation and usage of whole-body CT were associated with improvement in mortality among severe patients with trauma.<sup>15-18</sup> Emergency transportation by helicopter and usage of whole-body CT tended to increase over the study decade, and we noted that helicopter transportation had lower ORs for in-hospital mortality compared

**Table 2** Examination, treatment and hospitalisation of patients with trauma by years

Year	2004, n=2209	2005, n=2708	2006, n=2766	2007, n=6674	2008, n=8020	2009, n=9788	2010, n=12544	2011, n=14907	2012, n=17124	2013 n=14093
Whole-body CT, n (%)										
All patients	48/2209 (2.2)	95/2708 (3.5)	59/2766 (2.1)	439/6674 (6.6)	791/8020 (9.9)	1237/9788 (12.6)	1765/12544 (14.1)	2101/14907 (14.1)	2815/17124 (16.4)	2465/14093 (17.5)
Patients with ISS>16	35/1066 (3.3)	62/1187 (5.2)	42/1237 (3.4)	264/2861 (9.2)	443/3381 (13.1)	679/3872 (17.5)	900/4886 (18.4)	1120/5734 (19.5)	1411/6295 (22.4)	1158/4707 (24.6)
IVR, n (%)										
Abdomen	77/2209 (3.5)	93/2708 (3.4)	82/2766 (3.0)	224/6674 (3.4)	312/8020 (3.9)	481/9788 (4.1)	481/12544 (3.8)	538/14907 (3.6)	716/17124 (4.2)	627/14093 (4.5)
Pelvis	97/2209 (4.4)	104/2708 (3.8)	96/2766 (3.5)	212/6674 (3.2)	259/8020 (3.2)	366/9788 (3.7)	401/12544 (3.2)	425/14907 (2.9)	601/17124 (3.5)	534/14093 (3.8)
Hospitalisation location, n (%)										
Emergency and critical care centre (ICU)	1819/2138 (85.1)	2070/2597 (79.7)	2042/2660 (76.8)	4543/6320 (71.9)	5527/7405 (74.6)	6478/9301 (69.6)	8109/11994 (67.6)	9160/14269 (64.2)	10358/16342 (63.4)	8037/13446 (59.8)
General ward	307/2138 (14.9)	527/2597 (20.3)	618/2660 (23.2)	1777/6320 (28.1)	1878/7405 (25.4)	2823/9301 (30.4)	3885/12106 (32.1)	5109/14269 (35.8)	5984/16342 (36.6)	5409/13446 (40.2)
Length-of-hospital stay (days)	14 (5-35)*	15 (5-34)†	15 (6-33)‡	16 (6-35)§	16 (6-34)¶	17 (6-34)**	17 (6-34)††	18 (6-34)‡‡	17 (7-33)§§	16 (6-31)¶¶
Disposition at discharge, n (%)										
Home	886/1775 (49.9)	1242/2313 (53.7)	1208/2385 (50.6)	2786/5889 (47.3)	3240/7057 (45.9)	4111/8759 (46.9)	5515/11429 (48.3)	6631/13630 (48.7)	7820/15701 (49.8)	6758/13135 (51.5)
Hospital transfer	886/1775 (49.9)	1056/2313 (45.7)	1167/2385 (48.9)	3043/5889 (51.7)	3769/7057 (53.4)	4508/8759 (51.5)	5731/11429 (50.1)	6737/13630 (49.4)	7662/15701 (48.8)	6205/13135 (47.2)
Others*	3/1775 (0.2)	15/2313 (0.6)	10/2385 (0.5)	60/5889 (1.0)	48/7057 (0.7)	140/8759 (1.6)	183/11429 (1.6)	262/13630 (1.9)	219/15701 (1.4)	172/13135 (1.3)

\*Other places except for homes and hospitals.

†Missing data=85.

‡Missing data=57.

§Missing data=61.

¶Missing data=110.

\*\*Missing data=141.


††Missing data=146.

‡‡Missing data=246.

§§Missing data=222.

¶¶Missing data=170.

ISS, Injury Severity Score; IVR, interventional radiology.



**Figure 2** Ten-year in-hospital mortality trends for patients with trauma. ISS, Injury Severity Scores; TRISS Ps, Trauma and Injury Severity Score probability of survival.

with ambulance transportation among patients with ISS scores of 16 or higher. Whole-body CT was also associated with in-hospital mortality decreases among patients with

ISS scores of 16 or higher. Even after adjusting for helicopter transportation and usage of whole-body CT, ORs for in-hospital mortality of patients with trauma with ISS

**Table 3** Multivariable logistic regression analysis for in-hospital mortality of patients with trauma

	All patients, n=90833		Patients with ISS >16, n=35226		Patients with TRISS probability of survival ≥0.5, n=83953	
	OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value
Age	1.04 (1.04 to 1.04)	<0.001	1.04 (1.03 to 1.04)	<0.001	1.05 (1.05 to 1.05)	<0.001
Gender						
Male	1.24 (1.16 to 1.33)	<0.001	1.13 (1.04 to 1.21)	0.002	1.42 (1.31 to 1.54)	<0.001
Mechanism of injury						
Penetrating	1.00	–	1.00	–	1.00	–
Blunt	1.82 (1.46 to 2.27)	<0.001	1.26 (0.93 to 1.72)	0.14	1.36 (1.01 to 1.84)	0.04
ISS	1.06 (1.06 to 1.06)	<0.001	1.05 (1.04 to 1.05)	<0.001	1.09 (1.08 to 1.09)	<0.001
GCS	0.72 (0.71 to 0.72)	<0.001	0.73 (0.73 to 0.74)	<0.001	0.72 (0.71 to 0.73)	<0.001
Systolic blood pressure	0.99 (0.99 to 0.99)	<0.001	0.99 (0.99 to 0.99)	<0.001	0.99 (0.99 to 1.00)	<0.001
Respiratory rate	1.01 (1.00 to 1.01)	0.003	1.00 (1.00 to 1.01)	0.06	1.03 (1.03 to 1.04)	<0.001
Year						
2004	1.58 (1.32 to 1.91)	<0.001	1.55 (1.26 to 1.91)	<0.001	1.60 (1.28 to 1.99)	<0.001
2005	1.53 (1.27 to 1.82)	<0.001	1.56 (1.28 to 1.91)	<0.001	1.38 (1.12 to 1.71)	<0.001
2006	1.25 (1.04 to 1.49)	0.01	1.24 (1.01 to 1.52)	0.04	1.04 (0.83 to 1.31)	0.41
2007	1.08 (0.94 to 1.23)	0.42	1.09 (0.93 to 1.28)	0.49	1.02 (0.87 to 1.21)	0.33
2008	1.07 (0.94 to 1.21)	0.48	1.11 (0.96 to 1.29)	0.28	0.95 (0.81 to 1.11)	0.83
2009	1.00	–	1.00	–	1.00	–
2010	0.90 (0.80 to 1.02)	0.002	0.95 (0.83 to 1.09)	0.03	0.86 (0.75 to 0.99)	0.03
2011	0.79 (0.70 to 0.89)	<0.001	0.84 (0.73 to 0.96)	<0.001	0.76 (0.66 to 0.87)	<0.001
2012	0.79 (0.70 to 0.88)	<0.001	0.81 (0.71 to 0.92)	<0.001	0.74 (0.64 to 0.84)	<0.001
2013	0.69 (0.61 to 0.79)	<0.001	0.70 (0.60 to 0.81)	<0.001	0.63 (0.54 to 0.73)	<0.001

GCS, Glasgow Coma Scale; ISS, Injury Severity Score.



**Table 4** Multivariable logistic regression analysis for in-hospital mortality of patients with trauma with ISS >16 (n=34764)

	OR (95% CI)	P value
Age	1.04 (1.03 to 1.04)	<0.001
Gender		
Male	1.14 (1.05 to 1.23)	0.001
Mechanism of injury		
Penetrating	1.00	–
Blunt	1.27 (0.93 to 1.73)	0.13
Transportation		
Ambulance	1.00	
Ambulance with doctors	0.86 (0.73 to 1.01)	0.90
Helicopter	0.70 (0.63 to 0.79)	0.02
Others*	0.88 (0.64 to 1.20)	0.83
ISS	1.05 (1.04 to 1.05)	<0.001
GCS	0.73 (0.73 to 0.74)	<0.001
Systolic blood pressure	0.99 (0.99 to 0.99)	<0.001
Respiratory rate	1.00 (1.00 to 1.01)	0.06
Whole-body CT	0.91 (0.83 to 0.99)	0.04
Year		
2004	1.50 (1.22 to 1.85)	<0.001
2005	1.53 (1.25 to 1.88)	<0.001
2006	1.20 (0.98 to 1.48)	0.08
2007	1.06 (0.90 to 1.24)	0.78
2008	1.11 (0.95 to 1.29)	0.25
2009	1.00	–
2010	0.95 (0.82 to 1.09)	0.05
2011	0.85 (0.74 to 0.97)	<0.001
2012	0.82 (0.72 to 0.93)	<0.001
2013	0.71 (0.61 to 0.82)	<0.001

\*Private car, unknown and others.

GCS, Glasgow Coma Scale; ISS, Injury Severity Score.

scores of 16 or higher significantly decreased each year like primary outcome.

### Possible explanation and implications

Various factors that influence outcomes for patients with trauma have been reported during the past decade: changes in trauma systems and emergency transportation by helicopter in prehospital settings<sup>15 16 19 20</sup>; usage of whole-body CT in examinations<sup>17 18 21</sup>; tranexamic acid and damage control resuscitation including blood product ratio and massive transfusion protocols in treatment<sup>22–26</sup>; Advanced Trauma Life Support (ATLS) education<sup>10 27–29</sup> and region/facility differentials.<sup>30 31</sup> However, it is controversial as to whether or not these factors improve outcomes for patients with trauma. Therefore, improvement of in-hospital mortality in our study was considered to be due to a complex of changes in multiple factors.

Additionally, social policy might affect in-hospital mortality. During our study period, laws requiring seat-belt usage in rear seats and strengthened penalties for drunken driving were established.<sup>32</sup> Also, the improvement of motor vehicle engineering and safety devices might affect in-hospital mortality. These changes have mainly affected the prevalence of certain trauma types and their severity (namely, MVC), as seen in our results, and such decreases in trauma severity which cannot be reflected in ISS and RTS might additionally contribute to changes in in-hospital mortality.

Finally, we saw an increase in the mean age of patients with trauma from 47 to 64. This seems to be supported by the well-reported demographic shift in Japan to an elderly society. As older adults are most likely to become victims of falls down or falls at same level (which we found to increase over the study period), our results may reflect the need for new guidelines for trauma care that focus specifically on the challenges of this trauma type. However, we controlled for age and other variables in our analyses, and still found the downward trend to be statistically significant. Age may continue to be a concern, however, and should be considered as an important variable in future studies.

Taken together, our results indicate a need to continuously monitor available long-term patient data in order to create detailed snapshots of trauma prognoses. A deeper understanding of the effect that complex variables such as education, means of transportation, age, injury type, and treatment type have on patient outcomes will give a solid foundation on which to build and refine future guidelines.

### Limitations

Our study had several limitations. First, there was selection bias because all Japanese hospitals that treat trauma do not participate in the JTDB. Also, the number of participating hospitals differed across the study period. The severity of patients with trauma might differ by participating hospitals. However, most hospitals were tertiary emergency medical centres in Japan, where tertiary emergency medical centres defined by government. Therefore, participants in each year would be from similar environment. In addition, in-hospital mortality of patients with trauma significantly decreased each year after adjusting for severity. Second, our study excluded a large number of patients with missing data (n=31 650). It might affect our outcome. However, the results of Cochran-Armitage testing were similar between study population without missing data and it included any patients who had missing data. Furthermore, we could not investigate regional and facility differences. However, our study was conducted using a nationwide database which had a large sample size to bolster our statistical analyses.

### CONCLUSION

Our study has shown that in-hospital mortality for patients with trauma in Japan significantly decreased over a

10-year period (2004–2013) after adjusting for age and severity. The composite factors including improvement of medical care and services may contribute to reductions of inhospital mortality for patients with trauma, although further study is needed.

**Acknowledgements** The authors like to acknowledge Thomas D. Mayers and Bryan Mathis for his English-language manuscript revision.

**Contributors** IN conceived and designed this study, conducted data processing, analysed the data and wrote the manuscript. TA contributed to the conception and design of this study and drafted the manuscript. MU and NT contributed to design and critical revision of the manuscript. DS contributed to the acquisition of the data, study design and critical revision of the manuscript. All authors read and approved the final manuscript.

**Funding** This study was supported by the Ministry of Health, Labour and Welfare (H27-seisaku-senryaku-012).

**Competing interests** None declared.

**Patient consent** Detail has been removed from this case description/these case descriptions to ensure anonymity. The editors and reviewers have seen the detailed information available and are satisfied that the information backs up the case the authors are making.

**Ethics approval** The study protocol was reviewed and approved by the Ethical Committee of Yokohama City Minato Red Cross Hospital.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

## REFERENCES

1. Ministry of Health, Labour, and Welfare. Retrieved 27 Jun 2017. <http://www.mhlw.go.jp/toukei/list/81-1a.html> (In Japanese).
2. Ministry of Health, Labour, and Welfare. Retrieved 27 June 2017. <http://www.mhlw.go.jp/toukei/saikin/hw/jinkou/tokusyuu/furyo10/01.html> (In Japanese).
3. JATEC. Retrieved 27 Jun 2017. [http://www.jtcr-jatec.org/index\\_jatec.html](http://www.jtcr-jatec.org/index_jatec.html) (In Japanese).
4. Japan Trauma Data Bank. Retrieved 27 Jun 2017. <http://www.jtcr-jatec.org/traumabank/index.htm> (In Japanese).
5. Cabinet Office, Government of Japan. Retrieved 27 Jun 2017. [http://www8.cao.go.jp/kourei/whitepaper/w-2015/html/gaiyou/s1\\_1.html](http://www8.cao.go.jp/kourei/whitepaper/w-2015/html/gaiyou/s1_1.html) (In Japanese).
6. Sammy I, Lecky F, Sutton A, *et al.* Factors affecting mortality in older trauma patients-A systematic review and meta-analysis. *Injury* 2016;47:1170–83.
7. Trauma registry. *Trauma registry review board of the Japanese Association for the Surgery of trauma*. Tokyo: Herusu Shuppan, 2013. (In Japanese).
8. Siman-Tov M, Radomislensky I, Peleg K. Reduction in trauma mortality in Israel during the last decade (2000-2010): the impact of changes in the trauma system. *Injury* 2013;44:1448–52.
9. Curtis KA, Mitchell RJ, Chong SS, *et al.* Injury trends and mortality in adult patients with major trauma in New South Wales. *Med J Aust* 2012;197:233–7.
10. Hondo K, Shiraishi A, Fujie S, *et al.* In-hospital trauma mortality has decreased in Japan possibly due to trauma education. *J Am Coll Surg* 2013;217:850–7.

11. Di Saverio S, Gambale G, Coccolini F, *et al.* Changes in the outcomes of severe trauma patients from 15-year experience in a Western European trauma ICU of Emilia Romagna region (1996-2010). A population cross-sectional survey study. *Langenbecks Arch Surg* 2014;399:109–26.
12. Ruchholtz S, Lefering R, Paffrath T, *et al.* Reduction in mortality of severely injured patients in Germany. *Dtsch Arztebl Int* 2008;105:225–31.
13. Moore L, Turgeon AF, Lauzier F, *et al.* Evolution of patient outcomes over 14 years in a mature, inclusive Canadian trauma system. *World J Surg* 2015;39:1397–405.
14. Haider AH, Chang DC, Haut ER, *et al.* Mechanism of injury predicts patient mortality and impairment after blunt trauma. *J Surg Res* 2009;153:138–42.
15. Galvagno SM, Haut ER, Zafar SN, *et al.* Association between helicopter vs ground emergency medical services and survival for adults with major trauma. *JAMA* 2012;307:1602–10.
16. Abe T, Takahashi O, Saitoh D, *et al.* Association between helicopter with physician versus ground emergency medical services and survival of adults with major trauma in Japan. *Crit Care* 2014;18:R146.
17. Caputo ND, Stahmer C, Lim G, *et al.* Whole-body computed tomographic scanning leads to better survival as opposed to selective scanning in trauma patients: a systematic review and meta-analysis. *J Trauma Acute Care Surg* 2014;77:534–9.
18. Jiang L, Ma Y, Jiang S, *et al.* Comparison of whole-body computed tomography vs selective radiological imaging on outcomes in major trauma patients: a meta-analysis. *Scand J Trauma Resusc Emerg Med* 2014;22:54.
19. Mckee JL, Roberts DJ, van Wijngaarden-Stephens MH, *et al.* The right treatment at the right time in the right place: a population-based, before-and-after study of outcomes associated with implementation of an all-inclusive trauma system in a large Canadian province. *Ann Surg* 2015;261:558–64.
20. Twijnstra MJ, Moons KG, Simmermacher RK, *et al.* Regional trauma system reduces mortality and changes admission rates: a before and after study. *Ann Surg* 2010;251:339–43.
21. Sierink JC, Treskes K, Edwards MJ, *et al.* Immediate total-body CT scanning versus conventional imaging and selective CT scanning in patients with severe trauma (REACT-2): a randomised controlled trial. *Lancet* 2016;388:673–83.
22. Shakur H, Roberts I, Bautista R, *et al.* Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet* 2010;376:23–32.
23. Cannon JW, Khan MA, Raja AS, *et al.* Damage control resuscitation in patients with severe traumatic hemorrhage: A practice management guideline from the Eastern Association for the Surgery of Trauma. *J Trauma Acute Care Surg* 2017;82:605–17.
24. Zink KA, Sambasivan CN, Holcomb JB, *et al.* A high ratio of plasma and platelets to packed red blood cells in the first 6 hours of massive transfusion improves outcomes in a large multicenter study. *Am J Surg* 2009;197:565–70. discussion.
25. Mitra B, O'Reilly G, Cameron PA, *et al.* Effectiveness of massive transfusion protocols on mortality in trauma: a systematic review and meta-analysis. *ANZ J Surg* 2013;83:918–23.
26. Valle EJ, Allen CJ, Van Haren RM, *et al.* Do all trauma patients benefit from tranexamic acid? *J Trauma Acute Care Surg* 2014;76:1373–8.
27. Navarro S, Montmany S, Rebasa P, *et al.* Impact of ATLS training on preventable and potentially preventable deaths. *World J Surg* 2014;38:2273–8.
28. Wang P, Li NP, Gu YF, *et al.* Comparison of severe trauma care effect before and after advanced trauma life support training. *Chin J Traumatol* 2010;13:341–4.
29. Mohammad A, Branicki F, Abu-Zidan FM. Educational and clinical impact of Advanced Trauma Life Support (ATLS) courses: a systematic review. *World J Surg* 2014;38:322–9.
30. Jarman MP, Castillo RC, Carlini AR, *et al.* Rural risk: Geographic disparities in trauma mortality. *Surgery* 2016;160:1551–9.
31. Moore L, Evans D, Hameed SM, *et al.* Mortality in Canadian Trauma Systems: a Multicenter Cohort Study. *Ann Surg* 2017;265:212–217.
32. Road Traffic Low. Retrieved 27 Jun 2017. <http://law.e-gov.go.jp/htmldata/S35/S35HO105.html> (In Japanese).