



Degerud, E., Ystrom, E., Tambs, K., Ariansen, I., Morland, J., Magnus, P., ... Næss, O. (2018). The interplay between cognitive ability, alcohol consumption, and health characteristics. *Psychological Medicine*, 48(12), 2011-2022. <https://doi.org/10.1017/S0033291717003543>, <https://doi.org/10.1017/S0033291717003543>

Peer reviewed version

Link to published version (if available):

[10.1017/S0033291717003543](https://doi.org/10.1017/S0033291717003543)

[10.1017/S0033291717003543](https://doi.org/10.1017/S0033291717003543)

[Link to publication record in Explore Bristol Research](#)

PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via CUP at <https://www.cambridge.org/core/journals/psychological-medicine/article/interplay-between-cognitive-ability-alcohol-consumption-and-health-characteristics/CDF09DD4679686E78027AEF372D2A4C2>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:

<http://www.bristol.ac.uk/pure/about/ebr-terms>

Title page

Title: The interplay between cognitive ability, alcohol consumption, and health characteristics.

Authors: E Degerud, E Ystrøm, K Tambs, I Ariansen, J Mørland, P Magnus, G Davey Smith, Ø Næss.

Norwegian Institute of Public Health, P.O Box 4404, N-0403, Oslo, Norway, ED EY KT IA JM PM ØN. Institute of Clinical Medicine, University of Oslo, P.O box 1171, Blindern, 0318, Oslo, Norway, JM. MRC Integrative Epidemiology Unit, School of Social and Community Medicine, University of Bristol, Senate House, Tyndall Avenue, Bristol, BS8 1TH, UK, GDS. Institute of Health and Society, University of Oslo, P.O box 1130, Blindern, 0318, Oslo, Norway, ØN.

Corresponding author: Eirik Degerud, Norwegian Institute of Public Health, P.O. Box 4404, Oslo N-0403, Norway; eide@fhi.no

Keywords (MESH-terms): Drinking behaviour; alcohol drinking; cardiovascular diseases; behaviour and behaviour mechanisms.

Funding: Norges Forskningsråd (ES530767)

Current word count: 5512. Abstract 245.

Current reference count: 61

Abstract

Background: Higher cognitive ability is associated with favourable health characteristics. The relation between ability and alcohol consumption, and their interplay with other health characteristics, is unclear. We aimed to assess the relationship between cognitive ability and alcohol consumption and to assess whether alcohol consumption relate differently to health characteristics across strata of ability.

Methods: For 63,120 Norwegian males, data on cognitive ability in early adulthood were linked to midlife data on alcohol consumption frequency (times per month, 0-30) and other health characteristics, including cardiovascular risk factors and mental distress. Relations were assessed using linear regression and reported as unstandardized beta coefficients (95% CI).

Results: The mean \pm SD frequency of total alcohol consumption in the sample was 4.0 \pm 3.8 times per month. In the low, medium and high group of ability, the frequencies were 3.0 \pm 3.3, 3.7 \pm 3.5, and 4.7 \pm 4.1, respectively. In the full sample, alcohol consumption was associated with physical activity, heart rate, fat mass, smoking, and mental distress. Most notably, each additional day of consumption was associated with a 0.54% (0.44, 0.64) and 0.14% (0.09, 0.18) increase in the probability of current smoking and mental distress, respectively. In each strata of ability (low, medium, high), estimates were 0.87% (0.57, 1.17), 0.48% (0.31, 0.66) and 0.49% (0.36, 0.62) for current smoking, and 0.44% (0.28, 0.60), 0.10% (0.02, 0.18), and 0.09% (0.03, 0.15) for mental distress, respectively.

Conclusions: Participants with low cognitive ability drink less frequently, but in this group, more frequent alcohol consumption is more strongly associated with adverse health characteristics.

Introduction

The terms cognitive ability and intelligence are used interchangeably to describe general mental capability that, among other things, involves the ability to reason, plan, solve problems, think abstractly, comprehend complex ideas, learn quickly and learn from experience (Gottfredson, 1997). Poorer performance on tests of cognitive ability earlier in life tends to be associated with adverse risk factors of cardiovascular diseases (CVDs) later in life (Batty *et al*, 2007; Clouston *et al*, 2015; Marmot & Kivimaki, 2009; Richards *et al*, 2010; Taylor *et al*, 2003). More consistent is an association with mental distress (Gale *et al*, 2009; Hatch *et al*, 2007). The overrepresentation of adverse health characteristics at lower levels of cognitive ability is a source of health inequality, and underlines the importance of social policies ensuring that everyone has the possibility to obtain their cognitive potential (Gottfredson, 1997; Tucker-Drob & Bates, 2016).

Alcohol consumption is a complex behaviour in the sense that peoples' drinking patterns differ widely. It is an even more complex risk factor, because these drinking patterns associate differentially with health risk. Chronic and episodic heavy drinking, as well as light-to-moderate drinking, increases the risk of hundreds of diseases and conditions spanning both physical and mental health (Management of Substance Abuse Team., 2014; Rehm *et al*, 2009). Light-to-moderate consumption, on the other hand, is also associated with a reduced risk of ischemic heart disease (IHD) (Roerecke & Rehm, 2012; Ronksley *et al*, 2011). Because IHD is the leading cause of death in many countries, this association may propagate into a net beneficial association with all-cause mortality in certain age groups (Knott *et al*, 2015). The beneficial association could result from unaccounted bias in the design of observational studies, such as residual confounding, and few studies have accounted for the potential role of cognitive ability, which influence on CVD has been observed to encompass and surpass that of traditional CVD risk factors (Batty *et al*, 2009).

To address this, we study the interplay between cognitive ability, alcohol consumption, and different health characteristics, including cardiovascular risk factors and mental distress. The first aim of the study is to assess whether cognitive ability in early adulthood may influence midlife alcohol consumption. The second aim is to

assess whether the relationships between alcohol consumption and cardiovascular risk factors and mental distress differ by cognitive ability. We hypothesised that we would find differences that could contribute to the understanding of the alcohol harm paradox, which is a manifestation of social inequalities in health, where individuals with low socioeconomic position, or in this study low cognitive ability, experience more alcohol-related harm from the same level of exposure (Beard *et al*, 2016).

Methods

Study population

The source population was Norwegian men born after 1944 and before 1961 who participated in cardiovascular health surveys in midlife. From this population we selected two samples depending on the type of data available on alcohol consumption phenotypes. The first and main sample was selected from the National Health Screening Service's Age 40 Program (1994-1999) and selected surveys within the Cohort of Norway (1994-1999). Participants were required to have data available on alcohol consumption frequency or alcohol abstaining, cognitive ability from the mandatory evaluation for military conscription as well as data on other relevant covariates from the health surveys and additional covariates obtained by linkages to national health registries. The second sample was selected from the Finnmark III study (1987-1988, part of the Counties Study) and selected surveys within the Cohort of Norway (2000-2003). The statistical analyses performed in the second sample did not require exclusion of individuals with missing covariate data from health surveys and national registries.

Cognitive ability

Most Norwegian males were summoned for mandatory evaluation for military conscription, of which ~90% attended, mostly (95%) between their 18th and 21st birthday (Sundet *et al*, 2008). Reasons for non-summoning or non-attendance included chronic diseases, physical and mental disability, addiction problems, criminal records, or employment abroad or at sea. At the evaluation, the young males were required to complete three cognitive tests in which performance on each test was scored from 0 to 11. The test-retest correlation has been shown to be reliable (0.84 for arithmetic, 0.90 for word similarities, and 0.70 for progressive matrices, respectively). Intercorrelations between the different tests have been shown to range from 0.53 to 0.64 (Sundet *et al*, 1988). The three scores were standardised according to a reference population and then, with equal weight on each test score, combined into a stanine score (range 1-9, mean=5, SD=2) which provided a measure of general cognitive ability. The stanine score was used to categorise participants into low (1-4), medium (5-6), and high (7-9) ability. The stanine scores were entered in the Norwegian Armed Forces Personnel Data Base and obtained for this study by

data linkage. Because the test battery was re-standardised in 1963 and again in 1980 (mean scores lowered on average with 1.02 stanine points (Sundet *et al*, 2004)), the study population was limited to subjects for which stanine scores could be compared, which included males born after 1944 and before 1961.

Covariates

The health surveys provided self-reported measurements of alcohol consumption, current smoking (yes or no), level of leisure time physical activity (range 1-4, in which 1 reflect a sedentary lifestyle and 4 a very active lifestyle), history of diabetes (any), history of CVD (angina pectoris, myocardial infarction or stroke), family history of IHD, and mental distress. The measure of leisure time physical activity reflected either a general question (with 4 categories), the combination of two questions on the number of hours of hard and light activity, respectively, or all questions. Objective measurements was obtained for systolic blood pressure (mmHg), heart rate (beats per minute) and body mass index (BMI, kg/m²). Non-fasting biochemical measurements in blood samples (mmol/l) was obtained for serum triglycerides, total cholesterol, and high-density lipoprotein (HDL) cholesterol. Marital status (divorced/separated versus married/never married/widower) was obtained in conjunction with the health survey invitations (Population registry of Norway). The highest level of obtained education (range 1-8, in which 1 equals primary school and 8 equals a completed research education such as a master or doctoral degree, corresponding to the International Classification of Education 1997 categories) was obtained until 2011 from the National Educational Database.

Mental distress

The health surveys provided subjective measurement of mental health by the use of the Cohort of Norway Mental Health Index (CONOR-MHI) (Bramness *et al*, 2010; Sjøgaard *et al*, 2003). The index is based on 7 questions assessing whether the person in the past two weeks felt “nervous and unsettled”, “troubled by anxiety”, “secure and calm”, “irritable”, “happy and optimistic”, “sad/depressed”, or “lonely”. The questions have 4 possible answers: “no”, “a little”, “quite a bit”, and “very”, which are scored from 1-4 (scores for the questions “secure and calm” and “happy and

optimistic” are reversed). The index is constructed from the sum (7-28), in which a high score indicates mental distress. Subjects with two or more missing values were excluded, while a single missing value was replaced by the sample mean value for that question. The CONOR-MHI has been validated as a tool for assessing mental distress in epidemiological research by comparison to previously validated instruments of depression and anxiety (Søgaard *et al*, 2003), and the suggested cut-off (mean score >2.15) was applied to define current mental distress in this study.

Alcohol consumption

From the survey questions, we identified current abstainers and current drinkers from the question, “*Are you a total abstainer from alcohol (yes or no)*”. Among current drinkers, we further derived their frequency of total alcohol consumption (0-30 times) from the question, “*How many times per month do you usually drink alcohol?*” This variable was used continuously and categorically (less than monthly, 1-3, 4-6, 7-15, or 16-30 times per month). We derived data on the consumption frequency of different type of alcoholic beverages (0-50 glasses), including beer, wine and spirits, from the question “*How many glasses of beer/wine/spirits do you usually drink in the course of two weeks?*” Very high values were recoded to a maximum of 50. From the question, “*Approximately how often during the past year have you consumed alcohol corresponding to at least 5 small (35 cl) bottles of beer, a bottle of wine, or a quarter of a bottle of spirits?*”, we defined a binge drinker, or heavy episode drinker, as someone reporting a frequency of at least once a month, and in some cases at least once per week. Irregular monthly heavy drinking has been shown to cancel out any cardio-protective effects of moderate alcohol consumption (Roerecke & Rehm, 2010).

Statistical analysis

Analyses were performed in *RStudio*, version 1.0.44 (RStudio-Team, 2015) with additional use of the packages *Hmisc* and *ggplot2* (Harrell, 2013; Wickham, 2009). We do not impose a specific threshold to define statistical significance and have not performed any correction for the number of comparisons made. For both cognitive ability and for alcohol consumption, we presented the distribution of covariates at

mean values (standard deviations) or as counts (percentages). Group differences were assessed using analysis of variance and the chi-squared test, as appropriate.

The first aim was to assess whether cognitive ability influence alcohol consumption. To address this aim, we regressed alcohol consumption on ability (categorical variable) and age, and in a second model also on obtained education. Two methods of regression were applied depending on the outcome. For dichotomous outcomes, which included a comparison of current abstainers to seldom consumers (less than once per month) and binge drinkers to non-binge drinkers, we used ordinary least squares linear regression models to obtain unstandardized beta coefficients with 95% CIs. The analysis was repeated for binge drinking defined by a threshold of at least once per month and at least once per week.

For continuous outcomes, which included discrete counts of total alcohol consumption frequency (0-30) and the number of glasses (0-50) consumed in the course of two weeks on average for beer, wine, and spirits, we used negative binomial regression to obtain incidence rate ratios (IRRs) with 95% CIs. This was necessary because the continuous variables had greater variance than mean, which could result in over-dispersion and inaccurate estimation when using linear regression. The influence of ability on these outcomes were also visualised as smoothed means with 95% confidence intervals.

The second aim was to assess whether the relationship between alcohol consumption and different health characteristics differed by cognitive ability, or in other words, whether ability modified the relationships. The characteristics included dichotomous variables (yes/no) for current smoking, family history of IHD, divorced/separated, and current mental distress (CONOR-MHI >2.15), and continuous variables for BMI, systolic blood pressure, serum triglycerides, and serum HDL-cholesterol. We addressed the aim by regressing the health characteristics on alcohol consumption, age, and ability using ordinary least squares regression models to obtain unstandardized beta coefficients with 95% CIs. Two alcohol exposure variables were fitted in separate models, including a dichotomous variable comparing current abstainers to seldom consumers, and for a continuous variable of

total alcohol consumption frequency in which current abstainers were excluded. The analyses were performed in the full sample and in strata of ability. . Lastly, we fitted models that included an interaction term between alcohol and ability to test for effect modification on a multiplicative scale. The interaction term allows the linear slope (marginal effect of alcohol consumption) to vary by ability, and we report the unstandardized beta coefficient of the term with 95% CI and p-value.

Results

Participants

From 88,576 potentially eligible males, we included 63,120 current abstainers and current drinkers to the first and main study sample (see Supplementary Figure 1). There were only minor differences between the complete cases and the larger groups of excluded participants. Participants (n=16,220) with no entry in the military data base and thus ineligible for the study were slightly older on average than complete cases (41.1 and 43.5 years), but otherwise similar in terms of alcohol consumption (4.0 and 4.1 times per month) and obtained education (4.2 and 4.0 on a scale from 1 to 8), respectively. Participants excluded for missing values on ability (n=6932) were comparable in age (41.2 years), but had slightly lower values for alcohol consumption (3.7 times per month) and education (3.8).

From 12,196 males, we included 5425 current drinkers with data on heavy episode drinking (binge drinking) to the second sample. There were more individuals with missing values on alcohol consumption (n=2495) in the second sample relative to the first and main sample. They were on average of similar age as the complete cases (42.9 and 43.0 years), but had obtained 1 point lower education (4.3 and 3.4), and among the 632 with a value on ability (only 940 had an entry in the military data base), the mean stanine score was 1 point lower than for complete cases (6.2 and 5.2), respectively.

Descriptive statistics

Table 1 presents descriptive statistics for cognitive ability and for total alcohol consumption frequency in the main sample (n=63,120). Overall, estimates tended to be less favourable in terms of cardiovascular risk factors, divorced/separated, and mental distress for the subjects with low ability (n=10,986), more favourable for those with high ability (n=26,455), and in between for those with medium ability (n=25,679). Differences between categories of alcohol consumption were less consistent, with the exception of HDL-cholesterol, current smoking and divorces/separations, which appeared to follow a linear positive trend with more frequent consumption.

Cognitive ability and alcohol consumption

The percentage currently abstaining from alcohol was 5.5% overall and 6.2%, 4.7%, and 6.0% in groups of ability (low, medium, high), respectively. In comparison to the low group, the probability (95% CI) of abstaining in the medium and high groups was -1.45% (-1.96, -0.95) and -0.22% (-0.73, 0.29), respectively. Figure 1A depicts the relation of ability with total alcohol consumption frequency. Average frequency per month was 4.0 ± 3.8 overall and 3.0 ± 3.3 , 3.7 ± 3.5 , and 4.7 ± 4.1 within each group (low, medium, high). In comparison to the low group, the IRR (95% CI) of consumption in the medium and high groups was 1.23 (1.20, 1.25) and 1.53 (1.50, 1.57) when adjusted for age, and 1.20 (1.17, 1.23) and 1.43 (1.39, 1.46) when further adjusted for obtained education, respectively.

Figure 1B depicts the relation of ability with the consumption of different types of beverages. The number of glasses consumed in the course of two weeks were 3.7 ± 4.9 for beer, 1.8 ± 2.9 for wine, and 1.5 ± 2.8 for spirits overall. Within each group of ability (low, medium, high), the number of glasses were 3.5 ± 5.1 , 3.7 ± 4.8 , and 3.8 ± 4.9 for beer, 1.0 ± 2.4 , 1.5 ± 2.6 , and 2.3 ± 3.4 for wine, and 1.9 ± 3.4 , 1.6 ± 2.8 , 1.3 ± 2.5 for spirits, respectively. In comparison to the low group, the age adjusted IRR (95% CI) of consumption in the medium and high group were 1.06 (1.03, 1.10) and 1.10 (1.06, 1.13) for beer, 1.49 (1.43, 1.56) and 2.19 (2.09, 2.28) for wine, and 0.88 (0.85, 0.92) and 0.70 (0.67, 0.73) for spirits, respectively. When further adjusted for education, IRR were 1.10 (1.06, 1.13) and 1.15 (1.11, 1.18) for beer, 1.37 (1.31, 1.43) and 1.74 (1.67, 1.83) for wine, and 0.96 (0.92, 1.00) and 0.86 (0.82, 0.90) for spirits, respectively.

In the study population with data on heavy episodic drinking (n=5425), the percentage reporting a frequency of at least once per month was 27.8%, while 8.3% reported a frequency of at least once per week. The corresponding percentages in each group of ability (low, medium, high) were 22.9%, 27.2%, and 30.1%, respectively, and 6.5%, 8.0%, and 9.2%. In comparison to the low group, the age-adjusted probability (95% CI) of bingeing at least once per month in the medium and high group was 6.3% (2.7, 10.0) and 9.8% (6.2, 13.3), respectively, and 7.4% (3.6,

11.1) and 11.0% (6.9, 15.1) when further adjusted for education. The corresponding values for binge drinking at least once per week were 1.4% (-0.9, 3.7) and 2.9% (0.6, 5.1) when adjusted for age, and 2.1% (-0.0, 4.5) and 3.8% (1.2, 6.3) when further adjusted for education, respectively.

Interplay between cognitive ability, alcohol consumption, and health characteristics

Figure 2 provides a visual presentation of the relation between total alcohol consumption frequency (n=59,655) and health characteristics within each strata of cognitive ability, including cardiovascular risk factors and mental distress. There is less data and more uncertainty at very frequent intakes, as indicated by the wider confidence intervals. At a glance, the curves indicate linear relationships with BMI, triglycerides, HDL-cholesterol, and the probability of current smoking, being divorced/separated, and current mental distress, and non-linearity for physical activity and heart rate. The relation between consumption and systolic blood pressure and the probability of a family history of IHD appear less consistent in the different strata.

Table 2 provides the linear relations between alcohol consumption and health characteristics overall and within each strata of ability. More frequent alcohol consumption was associated with an increase in the probability of smoking, mental distress and divorce/separation, serum HDL-cholesterol and physical activity, and a decrease in heart rate, BMI and serum triglycerides. There was no association with systolic blood pressure. The slopes in the medium and high strata of ability differed (effect modification) from the low stratum for smoking, BMI, physical activity, heart rate, and mental distress. The increase in current smoking and mental distress with more frequent consumption, as well as the decrease in BMI, was more pronounced in the low stratum. In contrast to the two upper strata, there was no increase in physical activity or a decrease in heart rate in the low stratum.

Supplementary Table 1 shows the relative comparison of health characteristics between current abstainers (n=3465) and seldom consumers (n=8922) overall and in strata of ability. Current abstaining was associated with lower systolic blood pressure, heart rate, probability of smoking and divorce/separation, and higher probability of mental distress. Associations in the

medium and high strata of ability differed consistently from the low stratum for some characteristics. Current abstainers in the medium and high strata, but not in the low stratum, had lower systolic blood pressure and heart rate than seldom consumers, while current abstainers in the low strata, but not in the medium and high strata, had a higher probability of mental distress.

Discussion

Principle findings

Among Norwegian men, higher cognitive ability in early adulthood was associated with more frequent alcohol consumption in midlife. Higher wine consumption seems to account for this difference, and to some degree higher beer consumption, while consumption of spirits was inversely associated with ability. Heavy drinking episodes, or binge drinking, was positively associated with ability in a smaller separate sample. Aside from alcohol, the individuals with lower ability were more highly exposed to adverse health characteristics such as smoking and mental distress. . More frequent alcohol consumption associated with higher levels of physical activity and lower heart rate, but also more tobacco smoking and mental distress. However, the associations were different depending on ability. The association with smoking and mental distress was more pronounced among participants with low ability, and the favourable association with physical activity and heart rate was confined to participants with higher ability.

Cognitive ability and alcohol consumption

The hypothesis that cognitive ability could influence alcohol consumption, and not only the other way around, was stated decades ago (Parker & Noble, 1977). Our findings support the hypothesis and suggest that the relation for Norwegian men is positive. The bulk of previous studies on the topic also support a positive relation in other countries (Batty *et al*, 2008; Cheng & Furnham, 2013; Clouston *et al*, 2015; Corley *et al*, 2011; Fleming *et al*, 1982; Hatch *et al*, 2007; Hunt *et al*, 1984; Johnson *et al*, 2009; Kanazawa & Hellberg, 2010; Latvala *et al*, 2014; Mortensen *et al*, 2005). However, the related studies are very heterogeneous in terms of how and when cognitive ability and alcohol consumption were measured. A few studies observed no relation (Kubicka *et al*, 2001; Stautz *et al*, 2016; Wennberg *et al*, 2002; Windle & Blane, 1989), and some observed inverse or non-linear relationships (Batty *et al*, 2006; Muller *et al*, 2013; Sjolund *et al*, 2015).

Cognitive ability and socioeconomic position are closely related. In adolescence and adulthood, individuals are selected on ability into different

socioeconomic environments, such as educational and career trajectories, where social status, income, health consciousness, risk factors, and social support may differ (Ariansen *et al*, 2015; Deary, 2013). It is also within this environment that people tend to find their partner, subjecting cognitive ability to assortative mating, and causes it to correlate strongly between partners (Plomin & Deary, 2015). Cognitive ability is also very heritable, thus when highly educated and intelligent partners reproduce, assortative mating contributes to a positive relation between the offspring's genetic propensity for cognitive ability and the early life socioeconomic position (Plomin & Deary, 2015). Alcohol consumption has consistently been observed to be more frequent among socioeconomically advantaged groups (Management of Substance Abuse Team., 2014), and this relationship could overlap with the one we observed for ability. However, the influence of ability on total alcohol consumption was only slightly attenuated when we adjusted for the potential mediating effect of obtained education, which is an indicator of socioeconomic position.

The relation between cognitive ability and the consumption of different alcoholic beverages was strongest for wine, which was also observed in Danish men (Mortensen *et al*, 2005) and Scottish women and men (Corley *et al*, 2011). Wine is not produced in Norway, retail is confined to alcohol monopolies, and alcohol taxes are particularly high on a relative scale (Osterberg, 2011). Socioeconomically advantaged groups may be more inclined to purchase wine for culinary purposes, and afford to do so, which further support heritability and selection as possible mechanisms. The inverse association with spirit consumption is opposite of the findings in the Scottish study (Corley *et al*, 2011). Spirits are, similar to wine, only available for sale through the monopolies, but in contrast has a history of home production in Norway. Interestingly, when further adjusted for education, the influence of ability on wine consumption remained strong, but was notably attenuated. The influence on higher beer consumption increased in contrast, while the influence on lower spirit consumption was attenuated towards null. This suggests that education mediates in part the influence of ability on wine, and that this role was masked when the relationship was analysed on total consumption level where different types of beverages were not taken into account.

Cognitive ability may help individuals understand and appreciate the risks and benefits associated with different behaviours, and by that virtue influence cardiovascular health (Clouston *et al*, 2015; Marmot & Kivimaki, 2009). Drawing rational and healthy decisions when all risks are known, however, also involve aspects not assessed by the measurements which constitutes general cognitive ability (Toplak *et al*, 2010). While the adverse health effects of tobacco are clear, the effects of alcohol are complex, and when applying this perspective, it is not intuitive in which direction cognitive ability may influence consumption. Fifty-eight percent of adult Norwegians believe that a glass of wine is healthy (Norwegian Directorate of Health, 2012), perhaps because of biased media coverage or the tendency to prefer information that confirms existing or wishful beliefs (Nickerson, 1998). It is plausible that some Norwegians consume wine for perceived health effects, but we did not have information on the relation between cognitive ability, the perception of the healthiness of alcohol and alcoholic beverages, and alcohol consumption. Different type of alcoholic beverages are consumed for pleasure, but there are aspects of wine that might be best appreciated by those with knowledge and skill to taste them. This would again requires interest, training, and financial resources, and preferably an environment that shares the interest, which might be a reason why we observe that both ability and education has the strongest influence on wine consumption.

We found a positive relation between cognitive ability and binge drinking. The relation did not change to a large degree when further adjusted for education, and if any resulted in a stronger association. This finding was unexpected, given that the known harmful effects of binge drinking should be an incentive for health conscious groups to avoid the behaviour. It is also in sharp contrast to the inverse relation observed in the limited number of previous studies on the subject, including a prospective Scottish study using hangovers as a proxy for binge drinking (Batty *et al*, 2006), a prospective study from Great Britain (Cheng & Furnham, 2013), and a cross-sectional study of Swedish military conscripts (Sjolund *et al*, 2015). Although unexpected, comparable studies of the relation between ability in childhood and illegal drug use in adulthood are also mixed (White *et al*, 2012). To explain why ability should cause alcohol and drug use, one study emphasised the co-variation between higher ability and personality traits associated with alcohol and drug use (Hakulinen *et al*, 2015; White & Batty, 2011). Given that both cognitive ability and these personality traits are heritable (Devlin *et al*, 1997; Polderman *et al*, 2015; Ystrom *et*

al, 2011), common genetic factors with pleiotropic effects could explain why subjects with higher ability more frequently binge drink. Another possibility is that the amount of alcohol consumed per episode is different depending on ability, meaning that despite a higher frequency, each episode is less heavy among those with higher ability.

Interplay between cognitive ability, alcohol consumption and health characteristics

We observed that individuals with low cognitive ability were more highly exposed to cardiovascular risk factors and mental distress. Traditional risk factors are accounted for in studies assessing the relation between alcohol consumption and the risk of cardiovascular events, but mental distress is seldom measured. The consistent differences for measured risk factors indicate that ability might be associated with other risk factors, including those that are subtle, less influential and difficult to measure individually, but which together could constitute an important confounding effect. For example, a Danish study neatly revealed that wine buyers, which is consumed more frequently with higher ability, more often purchase healthy food items than beer buyers (Johansen *et al*, 2006). The measurement and inclusion of ability would thus be required to account for these subtle differences.

We also observed that the associations between alcohol consumption and health characteristics were different depending on by ability, or in other words, that cognitive ability modifies the relation between consumption, risk factors, and mental distress. This implicates ability and alcohol consumption in a more complex interplay with other health characteristics and the risk of cardiovascular events. It further suggests that the relationship between alcohol consumption and the risk of cardiovascular events, and potentially other alcohol-related outcomes, is confounded differently in subjects with low and high ability, and that a combined analysis of these subjects may fail to incorporate this information. It should be underlined that the co-variation did not diverge in opposing directions for any of the health characteristics, which would be the case if alcohol consumption were associated with less smoking in the high strata of ability and more smoking in the low strata.

In light of the cross-sectional data, the interplay may be interpreted from different perspectives. Subjects with low ability could be less able to incorporate alcohol into a healthy lifestyle, as indicated by the lack of co-variation with physical activity and heart rate, and more susceptible or less resistant to the disinhibitory or spillover effects of alcohol, as indicated by the stronger association with more smoking and mental distress. Consequently, more frequent alcohol consumption could worsen the existing tendency of adverse health characteristics to accumulate at lower ability. Cognition is involved in how humans regulate emotions (Ochsner & Gross, 2005; Ochsner *et al*, 2012), and subjects with higher ability are more often born and selected into more advantaged socioeconomic environments where the level of social support and health awareness is probably stronger. Another possible interpretation is that the measurement of cognitive ability in this study successfully differentiated the participants according to how adequately they were able to deal with mental distress, in which the steeper increase in alcohol consumption among subjects with low cognitive ability is a consequence of this difference, not a cause.

Current abstainers consistently smoked less than seldom consumers in all strata of cognitive ability, and in the medium and high strata, current abstaining was also associated with lower systolic blood pressure and heart rate. In contrast, mental distress was more frequent among current abstainers in the medium stratum, and even more so in the low stratum, but no association was observed in the high stratum. These findings indicate that in the medium and high strata of cognitive ability, current abstainers are healthier than seldom consumers are, while differences in the low strata are inconsistent.

Methodological considerations

The study population was limited to Norwegian men who attended both the mandatory evaluation for military conscription and a health survey in midlife, which is likely a somewhat healthier sample than the general population. Participants without an entry in the military database did not differ from complete cases, but participants with missing values on alcohol tended to have lower education and cognitive ability. There were few missing values on alcohol consumption frequency in the main sample, but more missing on binge drinking in the second sample, suggesting that

participants with lower ability were somewhat underrepresented in analyses of binge drinking. As it is not possible to elucidate the distribution pattern of binge drinking among those with missing values, we assume that it was comparable to complete cases. It is questionable to which extent the findings apply to women, as they tend to drink less than men do. The relationships observed in this study are context specific and could differ when this changes, such as in sub-populations or other countries.

Alcohol consumption may vary over time and be difficult to recall, leading to measurement errors when using self-reporting. The large sample size provides robustness to account for random errors, but not systematic errors, and cognitive ability could be a source of systematic error. However, if we consider serum HDL-cholesterol concentrations to be a biomarker of alcohol consumption (Brien *et al*, 2011), we observed that the linear increase in HDL-cholesterol concentration with increasing alcohol consumption was comparable in strata of ability (Table 2). When comparing the group consuming alcohol over 15 times per month to the group consuming alcohol 7-15 times within each strata of ability (Table 1), HDL-cholesterol differed between 0.07 and 0.08 mmol/l. This represented an increase of about 25 g alcohol per day (Brien *et al*, 2011), and indicated no particular increase in the low ability group in which alcohol consumption associated more strongly with adverse health characteristics. Therefore, we regard it as unlikely that differential relationships between alcohol and other health characteristics were spurious results of this potential bias.

Cognitive ability was measured in early adulthood, which is evidence to support that it could influence midlife alcohol consumption. However, we are not able to rule out the possibility of reverse causality, in which mental distress or alcohol consumption in childhood and early adolescence could have influenced the development of cognitive ability or the performance on the tests. We limited our study to estimate the overall association with cognitive ability and the potential mediating effect of obtained education, but in contrast to some previous studies, did not attempt to elucidate whether cognitive ability was mediating an effect of childhood socioeconomic position.

We have based inferences on the existence of an overall linear association, which may be somewhat rigid and misleading, as there were indications for non-linearity for physical activity and heart rate. There could be subjects combining moderate alcohol consumption with a healthy lifestyle at both high and low levels of cognitive ability, as well as subjects combining heavy drinking with an unhealthy lifestyle, but the distribution of these subjects may differ by ability. This could also be the reason why we do not observe an association with systolic blood pressure, despite evidence for a causal effect of alcohol on this cardiovascular risk factor (Chen *et al*, 2008).

The current study had a large size, a clear temporal separation where ability was measured prior to alcohol consumption, it measured different types of alcoholic beverages as well as binge drinking, and incorporated the potential mediating role of education. It therefore conveys an important contribution to existing literature on the influence of ability on alcohol consumption. It also implicates ability in the development of social inequalities in health, linking together studies showing that individuals with low cognitive ability are more highly exposed to risk factors and mental distress, and studies showing that socioeconomically disadvantaged groups experience more harmful effects of alcohol from a given level of exposure.

Conclusions

Among Norwegian men, higher early adulthood cognitive ability was associated with more frequent alcohol consumption and binge drinking in midlife. More frequent alcohol consumption associated with higher levels of physical activity and lower heart rate, but also more tobacco smoking and mental distress. While the association with smoking and mental distress was more pronounced among participants with low ability, the favourable association with physical activity and heart rate was confined to participants with higher ability. These findings show that ability is involved in how a given level of alcohol consumption relates to major risk factors and mental health, and thereby implicates ability in the development of the alcohol harm paradox and social inequalities in health.

References

- Ariansen I, Mortensen L, Igland J, Tell GS, Tambs K, Graff-Iversen S, Strand BH, Naess O** (2015). The educational gradient in coronary heart disease: the association with cognition in a cohort of 57,279 male conscripts. *Journal of Epidemiology and Community Health* **69**, 322-9.
- Batty GD, Deary IJ, Macintyre S** (2006). Childhood IQ and life course socioeconomic position in relation to alcohol induced hangovers in adulthood: the Aberdeen children of the 1950s study. *Journal of Epidemiology and Community Health* **60**, 872-4.
- Batty GD, Deary IJ, Schoon I, Emslie C, Hunt K, Gale CR** (2008). Childhood mental ability and adult alcohol intake and alcohol problems: the 1970 British cohort study. *American Journal of Public Health* **98**, 2237-43.
- Batty GD, Deary IJ, Schoon I, Gale CR** (2007). Mental ability across childhood in relation to risk factors for premature mortality in adult life: the 1970 British Cohort Study. *Journal of Epidemiology and Community Health* **61**, 997-1003.
- Batty GD, Shipley MJ, Dundas R, Macintyre S, Der G, Mortensen LH, Deary IJ** (2009). Does IQ explain socio-economic differentials in total and cardiovascular disease mortality? Comparison with the explanatory power of traditional cardiovascular disease risk factors in the Vietnam Experience Study. *European Heart Journal* **30**, 1903-9.
- Beard E, Brown J, West R, Angus C, Brennan A, Holmes J, Kaner E, Meier P, Michie S** (2016). Deconstructing the Alcohol Harm Paradox: A Population Based Survey of Adults in England. *PLoS One* **11**, e0160666.
- Bramness JG, Walby FA, Hjellvik V, Selmer R, Tverdal A** (2010). Self-reported mental health and its gender differences as a predictor of suicide in the middle-aged. *American Journal of Epidemiology* **172**, 160-6.
- Brien SE, Ronksley PE, Turner BJ, Mukamal KJ, Ghali WA** (2011). Effect of alcohol consumption on biological markers associated with risk of coronary heart disease: systematic review and meta-analysis of interventional studies. *BMJ* **342**, d636.
- Chen L, Smith GD, Harbord RM, Lewis SJ** (2008). Alcohol intake and blood pressure: a systematic review implementing a Mendelian randomization approach. *PLoS Medicine* **5**, e52.
- Cheng H, Furnham A** (2013). Correlates of adult binge drinking: evidence from a British cohort. *PLoS One* **8**, e78838.
- Clouston SA, Richards M, Cadar D, Hofer SM** (2015). Educational Inequalities in Health Behaviors at Midlife: Is There a Role for Early-life Cognition? *Journal of Health and Social Behavior* **56**, 323-40.

- Corley J, Jia X, Brett CE, Gow AJ, Starr JM, Kyle JA, McNeill G, Deary IJ** (2011). Alcohol intake and cognitive abilities in old age: the Lothian Birth Cohort 1936 study. *Neuropsychology* **25**, 166-75.
- Deary IJ** (2013). Intelligence. *Current Biology* **23**, R673-6.
- Devlin B, Daniels M, Roeder K** (1997). The heritability of IQ. *Nature* **388**, 468-71.
- Fleming JP, Kellam SG, Brown CH** (1982). Early predictors of age at first use of alcohol, marijuana, and cigarettes. *Drug and Alcohol Dependence* **9**, 285-303.
- Gale CR, Hatch SL, Batty GD, Deary IJ** (2009). Intelligence in childhood and risk of psychological distress in adulthood: The 1958 National Child Development Survey and the 1970 British Cohort Study. *Intelligence* **37**, 592-599.
- Gottfredson LS** (1997). Mainstream science on intelligence: An editorial with 52 signatories, history, and bibliography (Reprinted from The Wall Street Journal, 1994). *Intelligence* **24**, 13-23.
- Hakulinen C, Elovainio M, Batty GD, Virtanen M, Kivimaki M, Jokela M** (2015). Personality and alcohol consumption: Pooled analysis of 72,949 adults from eight cohort studies. *Drug and Alcohol Dependence* **151**, 110-4.
- Harrell FE** (2013). R-package 'Hmisc'.
- Hatch SL, Jones PB, Kuh D, Hardy R, Wadsworth ME, Richards M** (2007). Childhood cognitive ability and adult mental health in the British 1946 birth cohort. *Social Science and Medicine* **64**, 2285-96.
- Hunt E, Coggan P, Berg C, Tianwattanatana P** (1984). The effects of social drinking practices on cognitive performance. *Department of psychology, University of Washington, unpublished manuscript, 1984, described in Parson, 1986.*
- Johansen D, Friis K, Skovenborg E, Gronbaek M** (2006). Food buying habits of people who buy wine or beer: cross sectional study. *BMJ* **332**, 519-22.
- Johnson W, Hicks BM, McGue M, Iacono WG** (2009). How Intelligence and Education Contribute to Substance Use: Hints from the Minnesota Twin Family Study. *Intelligence* **37**, 613-624.
- Kanazawa S, Hellberg JEEU** (2010). Intelligence and Substance Use. *Review of General Psychology* **14**, 382-396.
- Knott CS, Coombs N, Stamatakis E, Biddulph JP** (2015). All cause mortality and the case for age specific alcohol consumption guidelines: pooled analyses of up to 10 population based cohorts. *BMJ* **350**, h384.
- Kubicka L, Matejcek Z, Dytrych Z, Roth Z** (2001). IQ and personality traits assessed in childhood as predictors of drinking and smoking behaviour in middle-aged adults: a 24-year follow-up study. *Addiction* **96**, 1615-28.

- Latvala A, Rose RJ, Pulkkinen L, Dick DM, Kaprio J** (2014). Childhood verbal development and drinking behaviors from adolescence to young adulthood: a discordant twin-pair analysis. *Alcoholism, Clinical and Experimental Research* **38**, 457-65.
- Management of Substance Abuse Team.** (2014). Global status report on alcohol and health 2014 ed. World Health Organization: Geneva, Switzerland.
- Marmot M, Kivimaki M** (2009). Social inequalities in mortality: a problem of cognitive function? *European Heart Journal* **30**, 1819-20.
- Mortensen LH, Sorensen TI, Gronbaek M** (2005). Intelligence in relation to later beverage preference and alcohol intake. *Addiction* **100**, 1445-52.
- Muller M, Kowalewski R, Metzler S, Stettbacher A, Rossler W, Vetter S** (2013). Associations between IQ and alcohol consumption in a population of young males: a large database analysis. *Social Psychiatry and Psychiatric Epidemiology* **48**, 1993-2005.
- Nickerson R** (1998). Confirmation Bias: A Ubiquitous Phenomenon in Many Guises. *Review of General Psychology* **2**, 175-220.
- Norwegian Directorate of Health** (2012). Unpublished survey of the general population, performed in conjunction with a campaign named "Hvite uker".
- Ochsner KN, Gross JJ** (2005). The cognitive control of emotion. *Trends in Cognitive Sciences* **9**, 242-9.
- Ochsner KN, Silvers JA, Buhle JT** (2012). Functional imaging studies of emotion regulation: a synthetic review and evolving model of the cognitive control of emotion. *Annals of the New York Academy of Sciences* **1251**, E1-24.
- Osterberg EL** (2011). Alcohol tax changes and the use of alcohol in Europe. *Drug Alcohol Rev* **30**, 124-9.
- Parker ES, Noble EP** (1977). Alcohol consumption and cognitive functioning in social drinkers. *Journal of Studies on Alcohol* **38**, 1224-32.
- Plomin R, Deary IJ** (2015). Genetics and intelligence differences: five special findings. *Molecular Psychiatry* **20**, 98-108.
- Polderman TJ, Benyamin B, de Leeuw CA, Sullivan PF, van Bochoven A, Visscher PM, Posthuma D** (2015). Meta-analysis of the heritability of human traits based on fifty years of twin studies. *Nature Genetics* **47**, 702-9.
- Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J** (2009). Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet* **373**, 2223-33.
- Richards M, Stephen A, Mishra G** (2010). Health returns to cognitive capital in the British 1946 birth cohort. *Longitudinal and Life Course Studies* **1**, 16.

- Roerecke M, Rehm J** (2010). Irregular heavy drinking occasions and risk of ischemic heart disease: a systematic review and meta-analysis. *American Journal of Epidemiology* **171**, 633-44.
- Roerecke M, Rehm J** (2012). The cardioprotective association of average alcohol consumption and ischaemic heart disease: a systematic review and meta-analysis. *Addiction* **107**, 1246-60.
- Ronksley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA** (2011). Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *BMJ* **342**, d671.
- RStudio-Team** (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA.
- Sjolund S, Hemmingsson T, Allebeck P** (2015). IQ and level of alcohol consumption-findings from a national survey of Swedish conscripts. *Alcoholism, Clinical and Experimental Research* **39**, 548-55.
- Stautz K, Pechey R, Couturier DL, Deary IJ, Marteau TM** (2016). Do Executive Function and Impulsivity Predict Adolescent Health Behaviour after Accounting for Intelligence? Findings from the ALSPAC Cohort. *PloS One* **11**, e0160512.
- Sundet JM, Barlaug DG, Torjussen TM** (2004). The end of the Flynn effect? A study of secular trends in mean intelligence test scores of Norwegian conscripts during half a century. *Intelligence* **32**, 349-362.
- Sundet JM, Eriksen W, Tambs K** (2008). Intelligence Correlations Between Brothers Decrease With Increasing Age Difference Evidence for Shared Environmental Effects in Young Adults. *Psychological Science* **19**, 843-847.
- Sundet JM, Tambs K, Magnus P, Berg K** (1988). On the Question of Secular Trends in the Heritability of Intelligence-Test Scores - a Study of Norwegian Twins. *Intelligence* **12**, 47-59.
- Søgaard A, Bjelland I, Tell GS, Røysamb E** (2003). A comparison of the CONOR Mental Health Index to the HSCL-10 and HADS. *Norsk Epidemiologi* **13**, 279-284.
- Taylor MD, Hart CL, Davey Smith G, Starr JM, Hole DJ, Whalley LJ, Wilson V, Deary IJ** (2003). Childhood mental ability and smoking cessation in adulthood: prospective observational study linking the Scottish Mental Survey 1932 and the Midspan studies. *Journal of Epidemiology and Community Health* **57**, 464-5.
- Toplak ME, Sorge GB, Benoit A, West RF, Stanovich KE** (2010). Decision-making and cognitive abilities: A review of associations between Iowa Gambling Task performance, executive functions, and intelligence. *Clinical Psychology Review* **30**, 562-81.
- Tucker-Drob EM, Bates TC** (2016). Large Cross-National Differences in Gene x Socioeconomic Status Interaction on Intelligence. *Psychological Science* **27**, 138-49.

- Wennberg P, Andersson T, Bohman M** (2002). Psychosocial characteristics at age 10; differentiating between adult alcohol use pathways: a prospective longitudinal study. *Addictive Behaviors* **27**, 115-30.
- White J, Batty GD** (2011). Intelligence across childhood in relation to illegal drug use in adulthood: 1970 British Cohort Study. *Journal of Epidemiology and Community Health* **66**, 767-74.
- White JW, Gale CR, Batty GD** (2012). Intelligence quotient in childhood and the risk of illegal drug use in middle-age: the 1958 National Child Development Survey. *Annals of Epidemiology* **22**, 654-7.
- Wickham H** (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
- Windle M, Blane HT** (1989). Cognitive ability and drinking behavior in a national sample of young adults. *Alcoholism, Clinical and Experimental Research* **13**, 43-8.
- Ystrom E, Reichborn-Kjennerud T, Aggen SH, Kendler KS** (2011). Alcohol dependence in men: reliability and heritability. *Alcoholism, Clinical and Experimental Research* **35**, 1716-22.

Table 1

Descriptive statistics for 63,120 adult males according to cognitive ability and alcohol consumption.

	Cognitive ability	All participants (n=63,120)	Current abstainer (n=3465)	Current drinkers: Total alcohol consumption frequency per month (n=59,655)					p-value
				<1 time (n=8922)	1-3 times (n=22,775)	4-6 times (n=17,369)	7-15 times (n=9752)	>15 times (n=837)	
Age	1-4	41.13 (1.92)	41.09 (1.86)	41.08 (1.86)	41.05 (1.91)	41.24 (2.00)	41.33 (1.89)	41.46 (2.05)	<0.001
	5-6	41.07 (1.91)	41.09 (1.72)	41.02 (1.87)	40.93 (1.93)	41.16 (1.93)	41.30 (1.86)	41.66 (2.14)	<0.001
	7-9	41.16 (1.93)	41.14 (1.81)	41.10 (1.92)	40.98 (1.96)	41.20 (1.92)	41.37 (1.93)	41.70 (1.94)	<0.001
	p-value		0.468	0.582	0.177	0.641	0.177	0.357	
Education (1-8)	1-4	3.05 (1.01)	3.01 (1.05)	2.97 (0.94)	3.07 (1.00)	3.06 (1.03)	3.19 (1.09)	3.05 (1.04)	<0.001
	5-6	3.81 (1.27)	3.84 (1.36)	3.70 (1.23)	3.77 (1.23)	3.85 (1.27)	3.95 (1.38)	3.98 (1.40)	<0.001
	7-9	5.07 (1.52)	5.13 (1.55)	4.85 (1.51)	4.93 (1.51)	5.10 (1.51)	5.33 (1.51)	5.20 (1.63)	<0.001
	p-value		<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	
Physical activity (1-4)	1-4	2.09 (1.00)	1.99 (1.03)	2.04 (1.04)	2.12 (1.01)	2.10 (0.97)	2.13 (0.96)	1.85 (0.97)	0.012
	5-6	2.21 (0.98)	2.17 (1.00)	2.17 (1.00)	2.20 (0.98)	2.23 (0.97)	2.24 (0.97)	2.27 (1.02)	<0.001
	7-9	2.28 (0.93)	2.18 (0.91)	2.21 (0.97)	2.27 (0.94)	2.30 (0.92)	2.33 (0.92)	2.29 (0.93)	<0.001
	p-value		<0.001	<0.001	<0.001	<0.001	<0.001	0.001	
Body mass index (kg/m ²)	1-4	26.57 (3.63)	26.85 (4.26)	26.83 (3.93)	26.60 (3.56)	26.34 (3.34)	26.26 (3.43)	26.20 (3.22)	<0.001
	5-6	26.28 (3.32)	26.32 (3.67)	26.35 (3.50)	26.35 (3.35)	26.24 (3.19)	26.10 (3.14)	25.91 (3.30)	<0.001
	7-9	26.03 (3.18)	25.97 (3.42)	26.23 (3.56)	26.10 (3.24)	26.03 (3.08)	25.86 (2.9)	25.68 (3.18)	<0.001
	p-value		<0.001	<0.001	<0.001	<0.001	<0.001	0.120	
Systolic blood pressure (mm Hg)	1-4	133.3 (13.6)	133.3 (14.7)	133.4 (13.2)	133.1 (13.4)	133.4 (13.7)	133.7 (14.4)	134.8 (15.1)	0.383
	5-6	132.6 (13.2)	131.1 (12.8)	133.0 (13.1)	132.5 (13.3)	132.7 (13.2)	132.8 (13.4)	133.2 (13.7)	0.032
	7-9	131.8 (13.2)	130.3 (13.1)	132.3 (13.7)	131.7 (13.3)	131.9 (13.1)	131.9 (13.0)	131.9 (13.3)	0.034
	p-value		<0.001	0.004	<0.001	<0.001	<0.001	0.037	
Heart rate (beats per minute)	1-4	73.0 (12.2)	74.2 (13.4)	73.7 (12.3)	72.6 (11.8)	72.4 (12.0)	73.4 (13.1)	74.2 (12.2)	0.006
	5-6	71.1 (12.0)	70.8 (12.3)	72.2 (12.0)	71.2 (12.0)	70.9 (11.9)	70.5 (11.9)	71.8 (12.3)	<0.001
	7-9	69.6 (11.9)	69.4 (11.8)	70.8 (12.4)	69.9 (11.7)	69.1 (11.8)	69.1 (12.1)	70.2 (11.4)	<0.001
	p-value		<0.001	<0.001	<0.001	<0.001	<0.001	0.001	
Triglycerides (mmol/l)	1-4	2.25 (1.51)	2.36 (1.78)	2.30 (1.47)	2.25 (1.52)	2.18 (1.36)	2.26 (1.75)	2.06 (1.73)	0.004
	5-6	2.16 (1.43)	2.09 (1.31)	2.21 (1.47)	2.17 (1.42)	2.15 (1.45)	2.10 (1.45)	2.13 (1.47)	0.020
	7-9	2.06 (1.30)	2.10 (1.31)	2.13 (1.5)	2.08 (1.28)	2.04 (1.27)	2.02 (1.27)	1.93 (1.15)	<0.001
	p-value		<0.001	<0.001	<0.001	<0.001	<0.001	0.120	
Total cholesterol (mmol/l)	1-4	5.86 (1.09)	5.86 (1.10)	5.86 (1.11)	5.85 (1.09)	5.88 (1.07)	5.89 (1.05)	5.76 (1.16)	0.519
	5-6	5.78 (1.05)	5.58 (1.02)	5.75 (1.06)	5.77 (1.05)	5.80 (1.03)	5.84 (1.04)	5.87 (1.10)	<0.001
	7-9	5.69 (1.02)	5.59 (1.02)	5.66 (1.06)	5.65 (1.02)	5.71 (1.01)	5.76 (1.02)	5.79 (1.05)	<0.001
	p-value		<0.001	<0.001	<0.001	<0.001	<0.001	0.821	
HDL-cholesterol (mmol/l)	1-4	1.18 (0.31)	1.13 (0.30)	1.14 (0.29)	1.17 (0.30)	1.21 (0.32)	1.26 (0.31)	1.33 (0.48)	<0.001
	5-6	1.20 (0.31)	1.13 (0.28)	1.14 (0.29)	1.17 (0.29)	1.23 (0.32)	1.26 (0.32)	1.34 (0.40)	<0.001
	7-9	1.21 (0.31)	1.14 (0.28)	1.15 (0.29)	1.18 (0.29)	1.22 (0.31)	1.27 (0.32)	1.34 (0.34)	<0.001
	p-value		0.530	0.298	0.277	0.185	0.045	0.998	

Family history of IHD	1-4	4630 (42.1%)	274 (40.3%)	921 (41.6%)	1949 (43.3%)	1101 (42.2%)	353 (39.9%)	32 (33.0%)	0.104
	5-6	10,640 (40.2%)	495 (39.5%)	1548 (39.7%)	4055 (40.1%)	2989 (40.5%)	1468 (41.3%)	85 (32.4%)	0.096
	7-9	9884 (38.5%)	588 (38.4%)	1081 (38.5%)	3087 (37.8%)	2926 (39.7%)	2003 (37.7%)	199 (41.6%)	0.090
	<i>p-value</i>		0.664	0.093	<0.001	0.081	0.003	0.028	
Diabetes	1-4	104 (0.90%)	12 (1.80%)	28 (1.30%)	39 (0.90%)	19 (0.70%)	5 (0.60%)	1 (1.00%)	0.069
	5-6	179 (0.70%)	13 (1.00%)	38 (1.00%)	69 (0.70%)	40 (0.50%)	18 (0.50%)	1 (0.40%)	0.043
	7-9	152 (0.60%)	9 (0.60%)	39 (1.40%)	41 (0.50%)	40 (0.50%)	20 (0.40%)	3 (0.60%)	<0.001
	<i>p-value</i>		0.034	0.270	0.045	0.511	0.560	0.771	
Current smoker	1-4	5398 (49.1%)	211 (31.0%)	1018 (45.9%)	2264 (50.4%)	1364 (52.2%)	483 (54.6%)	58 (59.8%)	<0.001
	5-6	10,152 (38.4%)	225 (18.0%)	1348 (34.6%)	4007 (39.7%)	3011 (40.8%)	1443 (40.6%)	118 (45.0%)	<0.001
	7-9	6400 (24.9%)	113 (7.40%)	600 (21.4%)	2069 (25.3%)	1991 (27.0%)	1457 (27.4%)	170 (35.6%)	<0.001
	<i>p-value</i>		<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	
Divorced or separated	1-4	1314 (12.0%)	52 (7.60%)	226 (10.2%)	570 (12.7%)	324 (12.4%)	120 (13.6%)	22 (22.7%)	<0.001
	5-6	2822 (10.7%)	85 (6.80%)	309 (7.90%)	1095 (10.8%)	849 (11.5%)	442 (12.4%)	42 (16.0%)	<0.001
	7-9	2152 (8.40%)	41 (2.70%)	167 (6.00%)	704 (8.60%)	624 (8.50%)	547 (10.3%)	69 (14.4%)	<0.001
	<i>p-value</i>		<0.001	<0.001	<0.001	<0.001	0.001	0.128	
Mental distress	1-4	774 (7.60%)	77 (12.3%)	157 (7.70%)	256 (6.10%)	181 (7.50%)	89 (10.8%)	14 (16.3%)	<0.001
	5-6	1309 (5.10%)	82 (6.90%)	214 (5.80%)	449 (4.60%)	349 (4.90%)	190 (5.50%)	25 (10.0%)	<0.001
	7-9	949 (3.80%)	71 (4.70%)	121 (4.40%)	290 (3.60%)	229 (3.20%)	199 (3.80%)	39 (8.30%)	<0.001
	<i>p-value</i>		<0.001	<0.001	<0.001	<0.001	<0.001	0.067	

Presented as mean (SD) and counts (%). Groups of cognitive ability were defined as stanine scores from 1-4 (n=10,986), 5-6 (n=26,455), and 7-9 (n=25,679). Mental distress was defined as a CONOR-mental health index ≥ 2.15 . Group differences were assessed with analysis of variance and the chi-squared test and included both current abstainers and current drinkers. Family history of ischemic heart disease (IHD).

Table 2

Linear relation between alcohol consumption and different health characteristics among current drinkers (n=59,655 adult males) in the full sample and in strata of cognitive ability.

Outcome	Cognitive ability ^b	Change in outcome per unit increase in the frequency of total alcohol consumption per month (range 0-30) ^a	Test for effect modification
		Unstandardized linear beta coefficient (95% CI) ^c	Unstandardized beta coefficient (95% CI), p-value ^d
Current smoker	All	0.54% (0.44, 0.64)	Reference group -0.38% (-0.71, 0.06), p=0.02 -0.39% (-0.71, 0.08), p=0.01
	1-4	0.87% (0.57, 1.17)	
	5-6	0.48% (0.31, 0.66)	
	7-9	0.49% (0.36, 0.62)	
Body mass index, kg/m ²	All	-0.03 (-0.04, -0.02)	Reference group 0.03 (0.00, 0.05), p=0.02 0.03 (0.01, 0.05), p=0.02
	1-4	-0.06 (-0.08, -0.03)	
	5-6	-0.03 (-0.04, -0.02)	
	7-9	-0.03 (-0.04, -0.02)	
Physical activity, mean 1-4	All	0.005 (0.003, 0.007)	Reference group 0.007 (0.000, 0.014), p=0.04 0.006 (-0.000, 0.012), p=0.07
	1-4	-0.001 (-0.007, 0.005)	
	5-6	0.007 (0.003, 0.010)	
	7-9	0.005 (0.002, 0.008)	
Systolic blood pressure, mm Hg	All	0.01 (-0.02, 0.04)	Reference group -0.02 (-0.11, 0.07) p=0.65 -0.04 (-0.13, 0.05) p=0.41
	1-4	0.04 (-0.05, 0.12)	
	5-6	0.01 (-0.04, 0.06)	
	7-9	0.00 (-0.04, 0.04)	
Heart rate, beats per minute	All	-0.07 (-0.09, -0.04)	Reference group -0.09 (-0.17, -0.01), p=0.04 -0.08 (-0.16, -0.00), p=0.04
	1-4	0.01 (-0.07, 0.08)	
	5-6	-0.08 (-0.12, -0.04)	
	7-9	-0.07 (-0.11, -0.04)	
Triglycerides, mmol/l	All	-0.009 (-0.012, -0.006)	Reference group -0.003 (-0.012, 0.007), p=0.59 -0.002 (-0.011, 0.007) p=0.69
	1-4	-0.007 (-0.017, 0.001)	
	5-6	-0.010 (-0.015, -0.005)	
	7-9	-0.009 (-0.013, -0.005)	
HDL-cholesterol, mmol/l	All	0.011 (0.011, 0.012)	Reference group -0.000 (-0.002, 0.002), p=0.92 -0.001 (-0.003, 0.001), p=0.44
	1-4	0.012 (0.010, 0.014)	
	5-6	0.012 (0.011, 0.013)	
	7-9	0.011 (0.010, 0.012)	
Family history of IHD	All	-0.04% (-0.15, 0.07)	Reference group 0.26% (-0.08, 0.59), p=0.14 0.33% (0.00, 0.66), p=0.04
	1-4	-0.31% (-0.60, -0.01)	
	5-6	-0.05% (-0.22, 0.13)	
	7-9	0.04% (-0.11, 0.19)	
Divorced or separated	All	0.36% (0.30, 0.43)	Reference group -0.02% (-0.23, 0.19), p=0.87 -0.10% (-0.30, 0.10), p=0.32
	1-4	0.42% (0.22, 0.61)	
	5-6	0.41% (0.30, 0.52)	
	7-9	0.32% (0.23, 0.41)	
Mental distress	All	0.14% (0.09, 0.18)	Reference group -0.33% (-0.49, -0.18), p<0.01 -0.35% (-0.50, -0.20), p<0.01
	1-4	0.44% (0.28, 0.60)	
	5-6	0.10% (0.02, 0.18)	
	7-9	0.09% (0.03, 0.15)	

^a Models adjusted for age and cognitive ability (if not used as a stratifying variable)

^b Cognitive ability strata derived from a stanine score (range 1-9).

^c Coefficients for dichotomous outcomes are multiplied by 100 and should be interpreted as the percentage point change in the probability of the outcome per one-unit increase in alcohol consumption frequency.

^d Test for effect modification on multiplicative scale, reflecting whether the linear slope in the medium (5-6) and high (7-9) group of ability differ from the slope in the low group (1-4), respectively.




Figure 1. Cognitive ability and alcohol consumption among 59,655 adult male current drinkers. Smoothed means with 95% confidence intervals for the relation of cognitive ability measured in early adulthood with midlife (A) total alcohol consumption frequency (n=59,655) and (B) consumption of different alcoholic beverages, including wine (n=55,432), beer (n=57,900) and spirits (n=56,095).




Figure 2. The crude relationship between total alcohol consumption, cardiovascular risk factors, and mental distress, within strata of cognitive ability for 59,655 adult male current drinkers. Regression lines are smoothed means with 95% confidence intervals. The probabilities of current smoking, mental distress, a family history of ischemic heart disease (IHD), and being divorced or separated range from 0 to 1.

Supplementary table 1

Difference in health characteristics between seldom consumers (n=8922) and current abstainers (n=3465) in the full sample and within strata of cognitive ability.

Response variable	Cognitive ability ^b	Current abstaining versus consumption less than once a month ^a	Test for effect modification
		Unstandardized linear beta coefficient (95% CI) ^c	Unstandardized beta coefficient (95% CI), p-value ^d
Current smoker	All	-15.3% (-16.9, -13.5)	Reference group -1.67% (-6.32, 2.98), p=0.48 0.90% (-3.70, 5.52), p=0.70
	1-4	-14.9% (-19.1, -10.7)	
	5-6	-16.6% (-19.5, -13.7)	
	7-9	-14.0% (-16.3, -11.7)	
Body mass index, kg/m ²	All	-0.12 (-0.26, 0.03)	Reference group -0.06 (-0.45, 0.33), p=0.77 -0.29 (-0.68, 0.09), p=0.14
	1-4	0.03 (-0.32, 0.37)	
	5-6	-0.03 (-0.26, 0.19)	
	7-9	-0.27 (-0.48, -0.05)	
Physical activity, mean 1-4	All	-0.02 (-0.06, 0.02)	Reference group 0.05 (-0.06, 0.15), p=0.38 0.01 (-0.09, 0.12), p=0.81
	1-4	-0.05 (-0.14, 0.04)	
	5-6	-0.00 (-0.06, 0.06)	
	7-9	-0.04 (-0.09, 0.02)	
Systolic blood pressure, mm Hg	All	-1.58 (-2.11, -1.05)	Reference group -1.76 (-3.19, -0.34), p=0.02 -1.92 (-3.34, -0.51), p=0.01
	1-4	-0.13 (-1.29, 1.04)	
	5-6	-1.88 (-2.71, -1.05)	
	7-9	-2.04 (-2.89, -1.20)	
Heart rate, beats per minute	All	-0.99 (-1.47, -0.51)	Reference group -1.90 (-3.20, -0.59), p<0.01 -1.93 (-3.22, -0.63), p<0.01
	1-4	0.52 (-0.56, 1.59)	
	5-6	-1.38 (-2.15, -0.61)	
	7-9	-1.41 (-2.16, -0.65)	
Triglycerides, mmol/l	All	-0.048 (-0.106, 0.009)	Reference group -0.18 (-0.34, -0.03), p=0.02 -0.10 (-0.25, 0.05), p=0.22
	1-4	0.060 (-0.073, 0.193)	
	5-6	-0.121 (-0.212, -0.030)	
	7-9	-0.035 (-0.125, 0.054)	
HDL-cholesterol, mmol/l	All	-0.011 (-0.023, 0.002)	Reference group 0.003 (-0.030, 0.036), p=0.85 0.000 (-0.033, 0.033), p=0.99
	1-4	-0.012 (-0.039, 0.015)	
	5-6	-0.009 (-0.028, 0.011)	
	7-9	-0.012 (-0.031, 0.007)	
Family history of IHD	All	-0.47% (-2.40, 1.46)	Reference group 1.00% (-4.22, 6.22), p=0.71 1.07% (-4.11, 6.25), p=0.69
	1-4	-1.28% (-5.51, 2.95)	
	5-6	-0.29% (-3.40, 2.82)	
	7-9	-0.22% (-3.25, 2.81)	
Divorced or separated	All	-2.31% (-3.32, -1.30)	Reference group 1.40% (-1.34, 4.14), p=0.32 -0.73% (-3.45, 1.98), p=0.60
	1-4	-2.55% (-5.08, -0.02)	
	5-6	-1.17% (-2.86, 0.52)	
	7-9	-3.29% (-4.62, -1.96)	
Mental distress	All	1.46% (0.50, 2.43)	Reference group -3.52% (-6.17, -0.86), p=0.01 -4.27% (-6.90, -1.65), p<0.01
	1-4	4.58% (2.04, 7.12)	
	5-6	1.05% (-0.51, 2.61)	
	7-9	-0.29% (-1.02, 1.61)	

^a Models adjusted for age and cognitive ability (if not used as a stratifying variable)

^b Cognitive ability strata derived from a stanine score (range 1-9).

^c Coefficients for dichotomous outcomes are multiplied by 100 and should be interpreted as the percentage point (%) change in the probability of the outcome per one-unit increase in alcohol consumption frequency.

^d Test for effect modification on multiplicative scale, reflecting whether the difference between seldom consumers and current abstainers in the medium (5-6) and high (7-9) groups of ability differ from the difference in the low group (1-4), respectively.

Supplementary figure 1

The flow of participants from the source surveys to the study population groups.

