

# Early micro-rheological consequences of single fraction total body low-dose photon irradiation in mice

Kornelia Szluha<sup>a,b</sup>, Kornelia Lazanyi<sup>a</sup>, Andrea Furka<sup>a</sup>, Ferenc Kiss<sup>c</sup>, Imre Szabo<sup>a</sup>, Eva Pintye<sup>a</sup>, Iren Miko<sup>c</sup> and Norbert Nemeth<sup>c,\*</sup>

<sup>a</sup>Department of Radiotherapy, Medical and Health Science Center, University of Debrecen, Debrecen, Hungary,

<sup>b</sup>Department of Diagnostic Radiology and Oncotherapy, Semmelweis University, Budapest, Hungary

<sup>c</sup>Department of Operative Techniques and Surgical Research, Institute of Surgery, Medical and Health Science Center, University of Debrecen, Debrecen, Hungary

**Abstract.** Despite of the studies on widespread biological effects of irradiation, surprisingly only little number of papers can be found dealing with its *in vivo* hemorheological impact. Furthermore, other studies suggested that low-dose irradiation might differ from high-dose in more than linear ways. On Balb/c Jackson female adult mice hematological and hemorheological impacts of total body irradiation were investigated 1 hour following 0.002, 0.005, 0.01, 0.02, 0.05 and 0.1 Gy dose irradiation. In case of 0.01 Gy further groups were analyzed 30 minutes, 2, 4, 6, 24 and 48 h after irradiation. According to the results, it seems that the dose-dependent changes of blood micro-rheological parameters are not linear. The irradiation dose of 0.01 Gy acted as a point of 'inflection', because by this dose we found the most expressed changes in hematological parameters, as well as in red blood cell aggregation, deformability and osmoscan data. The time-dependent changes showed progressive decrease in pH, rise in lactate concentration, further decrease in erythrocyte aggregation index and deformability, with moderate shifting of the optimal osmolality point and modulation in membrane stability. As conclusion, low-dose total body irradiation may cause micro-rheological changes, being non-linearly correlated with the irradiation dose.

Keywords: Red blood cell deformability, red blood cell aggregation, osmotic gradient ektacytometry, low-dose irradiation, mice

## 1. Introduction

Although medical use of radioactive agents is older than a century, it is still an unknown territory in more than a few details. It is widely known, that the ionising effect of radiotherapy causes biological alteration due to its energy being absorbed in tissues. This effect is quantifiable by the means of linear energy transfer (LET) [7]. Irradiations are divided into two main groups based on of their LET value. High LET values belong to alpha radiation, neutrons and protons, while low values are characteristic of X-ray, gamma-, beta-, photon and electron radiation [11, 16, 23]. Present article concentrates on the biological effect of irradiation of high LET.

With the spread of modern imaging technologies (CT, PET-CT, etc.) it is inevitable to explore the radiation biological effects of low-dose irradiation. However, literature data are limited in this area.

\*Corresponding author: Norbert Nemeth, M.D., Ph.D., Department of Operative Techniques and Surgical Research, Medical and Health Science Centre, University of Debrecen, H-4032 Debrecen, Nagyterdei krt. 98., Hungary. Tel./Fax: +36 52 416 915; E-mail: nemeth@med.unideb.hu.

The biological and neuroimmunological effects of low-dose irradiation are traditionally only estimated deductively from those of high dose, however, recent studies suggest, that low-dose irradiation may differ from high-dose in more than linear ways [6, 8, 11, 27, 48]. Dauer and co-workers even state that low-dose radiation research should be holistic and empirically founded for models, defining the shape of the dose-response relationship to be developed [14].

Despite of the studies on widespread biological effects of irradiation, surprisingly only little number of papers can be found dealing with its *in vivo* hemorheological impact. However, rheological link to radiotherapy seems to be an important aspect. Clinical studies demonstrated the beneficial effects of pentoxifylline (with improving effect on red blood cell deformability) in radiotherapy of breast and lung cancer [20, 36] and experimentally in rhabdomyosarcoma [50] among others. Other clinical hemorheological studies showed such evidences, for example in connection of high-dose brachytherapy in uterine cancer published by von Tempelhoff and co-workers [46]. Some papers deal with the effect of irradiation on red blood cell rheology, mostly in relation of blood storage [19, 38, 45], and sporadically can be found follow-up animal studies [e.g. 49].

Red blood cells' micro-rheological properties, such as deformability and aggregation are determined by numerous factors (cell shape, volume, surface-to-volume ratio, inner viscosity, membrane viscosity, composition of cell surface glycocalyx, fibrinogen concentration, lactate concentration, pH, osmolarity, oxygenation), and play important role in characterization, determination of blood flow characteristic at various levels of the circulation [1, 4, 10, 21, 22, 24, 32, 33, 35, 37, 39, 43, 47].

The effects of irradiation on red blood cells (e.g., change in volume, morphology and membrane integrity, induced oxidative damage, potassium efflux, sodium influx) [7, 17, 19, 30, 31, 38], depending on its dosage and fractionating, thus may have micro-rheological impact, also affecting tissue microcirculation. However, there is still a lack of *in vivo* animal studies in this field.

In this study we hypothesized that low-dose photon irradiation, under 0.1 Gray (Gy), may have detectable micro-rheological effects in mice. We also hypothesized that this impact is not linearly correlated with the dosage and may change in time.

## 2. Materials and methods

### 2.1. Experimental animals and groups

The experiments were approved and registered by the University of Debrecen Committee of Animal Research (registration Nr.: 16/2011 UD CAR), in accordance with the Hungarian Animal Protection Act (Law XVIII/1998).

Sixty-five Balb/c Jackson female mice (bodyweight:  $20.67 \pm 1.68$  g; age: 98–105 days) were subjected into two experimental series: in the *Experimental series I* the effects of various doses of irradiation were investigated, in the *Experimental series II* time-dependent effect of one chosen dose of irradiation was analyzed. The mice were randomly divided into the following groups.

#### 2.1.1. Experimental series I

Control group ( $n = 5$ ): no irradiation was made, but all the circumstances of housing, transport, time staying in the irradiation cage were similar to the groups with real irradiation.

Groups of various irradiation doses: 0.002 Gy ( $n = 5$ ), 0.005 Gy ( $n = 5$ ), 0.01 Gy ( $n = 5$ ), 0.02 Gy ( $n = 5$ ), 0.05 Gy ( $n = 5$ ) and 0.1 Gy ( $n = 5$ ). The samplings were performed under general anesthesia exactly 1 hour after irradiation.




Fig. 1. Schematic drawing of the positioner for homogenous total-body irradiation of mice.

### 2.1.2. Experimental series II

After the irradiation with 0.01 Gy further time points of investigations were set at 30 minutes ( $n=5$ ), 2 hours ( $n=5$ ), 4 hours ( $n=5$ ), 6 hours ( $n=5$ ), 24 hours ( $n=5$ ) and 48 hours ( $n=5$ ), respectively.

### 2.2. Technique of irradiation

In a pilot study, we analyzed the body weight-volume relations in the same mice strain. The body volume/body weight ratio was found to be a mean of 1.13 (volume =  $22.67 \pm 0.81 \text{ cm}^3$ , weight =  $20 \pm 1.26 \text{ g}$ ; ratio =  $1.13 \pm 0.03$ ,  $r^2 = 0.9375$ ;  $n = 6$ ). The irradiation dose fractions were calculated accordingly.

Since the mice were irradiated *in vivo* we had to prepare a special positioner. This device also had a kind of homogenizing effect. Since the mice are relatively small, compared to the size of the irradiation field, the positioner was embedded into a paraffin frame that ensured almost homogeneous dose distribution. Nineteen positioners were placed on a tray that also had a dosimeter on it and were irradiated with the same fraction (Fig. 1). This way, not only the *ex ante* fine-tuning of dosing, but the *ex post* controlling became possible.

Besides creating the special positioner we had to find the suitable technology that can radiate extraordinarily small doses such as 0.002 Gray. Since linear accelerators are not capable of such a fine-tuning and low dose, we decided to use Chisobalt 2B-75 device (Chirama, Czechoslovakia). Chisobalt irradiates with  $^{60}\text{Co}$  isotope ( $^{60}\text{Co}$   $T_{1/2} = 5.27$  years) at a dose rate of 0.6789 Gy/min, with 1.25 MeV (SSD 100 cm, field size  $21 \times 21 \text{ cm}$ ).

Although  $^{60}\text{Co}$  is capable of lower dose, than linear accelerators, shielding of the research objects was also necessary in order to produce the appropriate dose distribution. The thickness of the absorber plastic-sheet was 60 mm. When using Chisobalt the application of PTW Unidos Power Chamber Electrometer (PTW, Germany), a special device became indispensable in order to adjust and measure the exact dose.

### 2.3. Anesthesia and sampling protocol

Thiopental<sup>®</sup> was used intraperitoneally in 62.5 mg/kg dosage for general anesthesia, under which median thoraco-laparotomy was performed for blood sampling and tissue biopsies. For blood sampling cardiac puncture was made (26 G needle), the samples were taken into tubes with  $\text{K}_3\text{-EDTA}$  (1.5 mg/ml) as anticoagulant. Laboratory measurements were completed shortly after sampling, all within 1-2 hours.

After blood sampling biopsies were taken from the heart, lung, liver, spleen, kidney, small bowel and femur for histomorphological examinations. In this paper the laboratory investigations are reported.

## 2.2. Laboratory measurements

### 2.2.1. Blood pH and lactate concentration

An ABL555 blood gas analyzer automate (Radiometer Copenhagen, Denmark) was used to determine blood pH and lactate concentration (mmol/l). Following blood sampling, the tests were immediately performed (within 1-2 minutes), without direct contact with air.

### 2.2.2. Hematological parameters

To determine the general hematological parameters a Sysmex F-800 semi-automated microcell counter (TOA Medical Electronics Co., Japan) was used. Measurements could be carried out within 10-15 minutes after sampling. In this study hemoglobin concentration (Hgb [g/dl]), hematocrit (Hct [%]), mean corpuscular volume (MCV [fl]), mean corpuscular hemoglobin content (MCH [pg]), mean corpuscular hemoglobin concentration (MCHC [g/dl]) and red cell distribution width (RDW-CV% [%]) were analyzed.

### 2.2.3. Red blood cell deformability

A LoRRca MaxSis Osmoscan device (Mechatronics BV, The Netherlands) was used to measure red blood cell elongation index in the function of shear stress, osmotic gradient ektacytometry parameters and performing membrane stability test.

For *regular red blood cell deformability tests* 5  $\mu$ l blood sample was required, which was gently mixed in 1 ml of isotonic polyvinyl-pyrrolidone solution (360 kDa PVP in normal phosphate buffered saline; viscosity = 27 mPa.s, osmolarity = 290–300 mOsm/kg; pH  $\sim$  7.3). The suspension was injected into the bob-cup system of the device, carefully watching not to leave any air bubbles in the measuring chamber. During the measurements, the device generated shear stress (SS) range from 0.3 to 30 Pa, while the laser diffraction pattern was analyzed at set shear stress levels to calculate the elongation index (EI) values. The EI is equal to  $(L - W) / (L + W)$ , where L is the length and W is the width of the diffractogram. EI increases with red blood cell deformability [2, 18]. The tests were carried out at constant temperature of 37°C. Since data reduction methods of EI-SS curves are useful tools for comparison [3, 5, 44], Lineweaver-Burk analyses were performed, calculating the maximal elongation index ( $EI_{max}$ ) and the shear stress at half  $EI_{max}$  ( $SS_{1/2}$  [Pa]) values, according to the following formula:  $1/EI = SS_{1/2}/EI_{max} \times 1/SS + 1/EI_{max}$  [3].

For the *osmotic gradient ektacytometry (osmoscan) measurements* 250  $\mu$ l blood was needed, which was placed and gently mixed in 5 ml PVP solution, having the same parameters described above. During the measurements, a constant shear stress of 30 Pa was applied to the sample, which is continuously aspirated into the measurement chamber together with changing amount of 0 and 500 mOsmol/kg PVP solutions. The EI values were continuously registered along the osmolarity scale [13, 18]. In this study we analyzed the peak of the elongation index-osmolarity curve, comparing the maximal elongation index values (EI values at the peak of the curve) and the osmolarity at maximal EI ('optimal' osmolarity).

In the *membrane stability test*, we used a continuous shear stress of 100 Pa for 300 seconds. Before and after the mechanical shearing the device measures regular red blood cell deformability tests. We used this function on blood samples of the *Experimental series II*.




Fig. 2. Changes of blood pH (A) and lactate concentration [mmol/l] (B) 1 hour after various doses of irradiation. means  $\pm$  S.D.; \* $p < 0.05$  vs. Control.

#### 2.2.4. Red blood cell aggregation

Based on light-transmittance method, a Myrenne MA-1 erythrocyte aggregometer (Myrenne GmbH, Germany) was used to determine aggregation index values M (at shear rate of  $0 \text{ s}^{-1}$ ) and M1 (at shear rate of  $3 \text{ s}^{-1}$ ) 5 or 10 seconds after disaggregation. The indices (M 5 s, M1 5 s, M 10 s, M1 10 s) increase with enhanced red blood cell aggregation [2, 18]. The measurements could be carried out within 20–30 minutes after sampling.

#### 2.3. Statistical analyses

Data are expressed as means and standard deviations (S.D.). For comparison of groups with various doses of irradiation, *t*-test or Mann-Whitney rank sum test were used, according to the data distribution. Comparing the changes in time, one way ANOVA on ranks with Bonferroni's method was used. A *p* value of  $< 0.05$  was considered as statistically significant.

### 3. Results

#### 3.1. Effects of various low-dose irradiation (0.002, 0.005, 0.01, 0.02, 0.05 and 0.1 Gy)

##### 3.1.1. Blood pH and lactate concentration

Changes in blood pH or lactate concentration did not reflect obvious relationship to dose-dependency of the irradiation (Fig. 2).

After irradiation at 0.005 and 0.05 Gy we measured significantly lower pH values compared to the Control ( $p = 0.009$  and  $p = 0.016$ , respectively). Lactate concentration decreased in all irradiated groups in various manners. Compared to the Control group significant decrease was found in groups irradiated by 0.002 Gy ( $p = 0.006$ ), 0.005 Gy ( $p = 0.001$ ), 0.02 Gy ( $p < 0.001$ ) and 0.1 Gy ( $p = 0.012$ ).

##### 3.1.2. Hematological data

Selected hematological parameters are summarized in Table 1.

Hematocrit (Hct [%]) showed slight decrease after irradiation at all dose. The most expressed decrease was observed at the 0.01 Gy ( $p = 0.01$ ) and 0.1 Gy doses. If the values were compared to the lowest one, to the 0.01 Gy data, the differences were significant at 0.002 Gy ( $p = 0.01$ ), 0.005 Gy ( $p = 0.023$ ), 0.02 Gy ( $p < 0.001$ ) and 0.05 Gy ( $p = 0.007$ ).

Mean corpuscular volume (MCV [fl]) moderately decreased after 0.002 Gy ( $p = 0.003$  vs 0.01 Gy), 0.005 Gy ( $p = 0.031$  vs Control;  $p < 0.001$  vs 0.01 Gy) and 0.05 Gy irradiation ( $p < 0.001$  vs 0.01 Gy).

Table 1  
Erythrocyte-related hematological, deformability and osmoscan parameters in blood samples taken from mice 1 hour after various dose of irradiation

Variable	Control	0.002 Gy	0.005 Gy	0.01 Gy	0.02 Gy	0.05 Gy	0.1 Gy
Hct [%]	44.52 ± 5.29	44.39 ± 3.12 #	42.58 ± 1.38 #	37.4 ± 4.95*	44.9 ± 2.34 #	43.42 ± 3.3 #	40.53 ± 7.19
Hgb [g/dl]	11.2 ± 0.81	11.81 ± 0.41	12.04 ± 0.2	10.53 ± 1.47	10.89 ± 0.54	11.64 ± 0.76	11.56 ± 1.46
MCV [fl]	57.33 ± 5.35	54.47 ± 5.79 #	51.47 ± 1.44*	61.03 ± 1.77+	59.27 ± 1.31	53.78 ± 2.68 #	60.56 ± 1.91
RDW-CV% [%]	14 ± 0.22	13.71 ± 1.18	13.25 ± 0.34	14.1 ± 0.45	14.23 ± 0.29	14.07 ± 0.57	13.86 ± 0.31
MCH [pg]	14.45 ± 0.91	14.46 ± 0.65	14.57 ± 0.43 #	15.55 ± 1.28	14.37 ± 0.2 #	14.3 ± 0.4 #	16.75 ± 1.99
MCHC [g/dl]	25.45 ± 3.34	26.68 ± 1.82	28.3 ± 0.67	28.15 ± 4.99	24.26 ± 0.47	26.66 ± 1.37	27.52 ± 3.44
EI <sub>max</sub>	0.544 ± 0.01	0.538 ± 0.04	0.548 ± 0.02	0.539 ± 0.02	0.557 ± 0.02	0.547 ± 0.01	0.543 ± 0.02
SS <sub>1/2</sub> [Pa]	1.87 ± 0.3	2.55 ± 0.49*	2.76 ± 0.3*	1.77 ± 0.46	1.85 ± 0.32	1.78 ± 0.12	1.88 ± 0.36
Maximal EI	0.444 ± 0.04	0.487 ± 0.03	0.496 ± 0.01*	0.445 ± 0.08	0.5 ± 0.004*	0.5 ± 0.02*	0.478 ± 0.04
Osmolarity at max. EI [mOsm/kg]	312 ± 20.55	326.2 ± 11.23 #	326.6 ± 9.44 #	293.2 ± 9.98	293.2 ± 7.85	313 ± 6	321 ± 11.66 #

means ± S.D. EI<sub>max</sub> = calculated maximal elongation index (EI); SS<sub>1/2</sub> = calculated shear stress (SS) at half EI<sub>max</sub> by Lineweaver-Burk analysis; Maximal EI = maximal EI values measured in osmoscan function. \* $p < 0.05$  vs. Control; #  $p < 0.05$  vs. 0.01 Gy; +  $p = 0.053$  vs. Control.




Fig. 3. Changes of aggregation index M 5 s (A), M 10 s (B), M1 5 s (C) and M1 10 s (D) 1 hour after various doses of irradiation. means  $\pm$  S.D.; \* $p < 0.05$  vs. Control.

MCV values slightly increased in samples taken from groups with irradiation of 0.01 Gy ( $p = 0.053$ ), 0.02 Gy and 0.1 Gy.

Mean corpuscular hemoglobin content (MCH [pg]) showed moderate increase in groups of 0.01 Gy and 0.1 Gy. The other parameters, such as RDW-CVR% or MCHC did not show significant differences.

### 3.1.3. Red blood cell deformability

Comparative parameters of regular red blood cell deformability and osmotic gradient ektacytometry (osmoscan) measurements are showed in Table 1.

Calculated parameters from the elongation index (EI) - shear stress (SS [Pa]) curves, such as  $EI_{max}$  and  $SS_{1/2}$  [Pa] did not show obvious relation to the irradiation dose. We did not find any significant difference in  $EI_{max}$ , however,  $SS_{1/2}$  values were significantly increased compared to the Control only in samples from the groups which were irradiated with 0.002 Gy ( $p < 0.001$ ) or 0.005 Gy ( $p < 0.001$ ).

During osmoscan measurement, we could observe that the maximal elongation index values rather increased in all irradiated groups, but in the smallest manner at 0.01 Gy. The differences versus Control were significant in cases of 0.005 Gy ( $p = 0.034$ ), 0.02 Gy ( $p = 0.016$ ) and 0.05 Gy ( $p = 0.046$ ). However, the osmolarity values at the maximal EI (at the peak of EI-osmolarity curves) were the highest in case of 0.002 Gy ( $p < 0.001$  vs. 0.01 Gy), 0.005 Gy ( $p < 0.001$  vs. 0.01 Gy) and 0.1 Gy ( $p = 0.008$  vs. 0.01 Gy).

### 3.1.4. Red blood cell aggregation

Investigating the aggregation index M and M1 at 5 and 10 seconds, we could observe, that up to 0.01 Gy, the values tended to decrease (even till zero values), than rather increased values were seen, however, all being lower than the Control data. The 0.01 Gy dose seemed like a site of an 'inflexion' point (Fig. 3).




Fig. 4. Changes of blood pH (A) and lactate concentration [mmol/l] (B) in the function of time after 0.01 Gy dose of irradiation. means  $\pm$  S.D.; \* $p < 0.05$  vs. Control; # $p < 0.05$  vs. 1 h.

M index of 5 seconds (M 5 s) showed significantly lower values at 0.002 Gy ( $p < 0.001$ ), 0.01 Gy ( $p < 0.001$ ), 0.02 Gy ( $p < 0.001$ ), 0.05 Gy ( $p < 0.001$ ) and 0.1 Gy ( $p < 0.001$ ), compared to Control (Fig. 3A).

M index of 10 seconds (M 10 s) also showed significantly lower values at 0.002 Gy ( $p < 0.001$ ), 0.005 Gy ( $p < 0.001$ ), 0.02 Gy ( $p < 0.001$ ), 0.05 Gy ( $p < 0.001$ ) and 0.1 Gy ( $p < 0.001$ ), compared to Control. At 0.01 Gy all measured values were zero (Fig. 3B).

M1 index of 5 seconds (M 5 s) showed the same tendency as M values did. Compared to Control significantly lower values were measured at 0.002 Gy ( $p = 0.001$ ), 0.005 Gy ( $p < 0.001$ ), 0.01 Gy ( $p < 0.001$ ), 0.02 Gy ( $p < 0.001$ ), 0.05 Gy ( $p < 0.001$ ) and 0.1 Gy ( $p < 0.001$ ) (Fig. 3C).

M1 index of 10 seconds (M 10 s) showed significant difference at 0.005 Gy ( $p = 0.002$ ), 0.02 Gy ( $p = 0.003$ ), 0.05 Gy ( $p = 0.002$ ) and 0.1 Gy ( $p = 0.003$ ), compared to Control. Here at 0.01 Gy all measured values were zero, too (Fig. 3D).

### 3.2. Changes in the function of time after irradiation with 0.01 Gy

#### 3.2.1. Blood pH and lactate concentration

In general, blood pH values showed a continuous declining, while lactate concentration expressed an increasing tendency in the function of time, after irradiation with 0.01 Gy (Fig. 4).

Compared to Control, blood pH values were significantly lower 30 minutes ( $p = 0.017$ ), 2 hours ( $p = 0.009$ ), 4 hours ( $p = 0.002$ ), 24 hours ( $p < 0.001$ ) and 48 hours ( $p = 0.001$ ) after irradiation. Values at 4, 24 and 48 hours were significantly lower versus the 1-hour data, too ( $p = 0.006$ ,  $p = 0.005$  and  $p = 0.004$ , respectively).

Lactate concentration continuously elevated, reaching the highest levels from the 4<sup>th</sup> hour until the end of the follow-up period. Compared to the Control, we found significant differences at the 24th hour ( $p = 0.008$ ) and by the 48th hour ( $p = 0.007$ ). These 1 and 2-day values were significantly higher compared to the 1-hour data, too ( $p = 0.004$  and  $p = 0.032$ ), together with the 4-hour values ( $p = 0.043$ ).

#### 3.2.2. Hematological data

Selected, erythrocyte-related hematological parameters are summarized in Table 2. Hematocrit (Hct [%]) did not show important changes until the 4th hour (except for the 1-hour samples), and over 6 hours the values started to be elevated, being significantly different versus Control (6 h:  $p = 0.044$ , 24 h:  $p = 0.008$  and 48 h:  $p = 0.012$ ) and compared to the 1-hour samples, too ( $p < 0.001$  for all). Hemoglobin concentration values (Hgb [g (dl)]) slightly increased, while mean corpuscular volume (MCV [fl]) showed



216 significant or almost significant decrease compared to Control (30-min:  $p = 0.024$ , 2 h:  $p = 0.055$ , 4 h:  
217  $p = 0.059$ , 6 h  $p = 0.069$ ). These values were highly significant versus the data obtained from the 1-hour  
218 samples ( $p < 0.001$  for all). The other parameters, such as RDW-CV%, MCH and MCHC did not show  
219 important changes.

### 220 3.2.3. Red blood cell deformability

221 Comparative data of regular deformability tests and osmotic gradient ektacytometry measurements are  
222 presented in Table 2.

223 Except for the data from the 1-hour samples,  $EI_{max}$  values showed decrease compared to Control,  
224 reaching significant level at 4 hours ( $p = 0.022$ ), 6 hours ( $p = 0.015$ ), 24 hours ( $p = 0.014$ ) and 48 hours  
225 ( $p = 0.002$ ). Compared to the 1-hour values,  $EI_{max}$  values were significantly lowered at 4 hours ( $p = 0.05$ ),  
226 24 hours ( $p = 0.039$ ) and 48 hours ( $p = 0.026$ ). In turn,  $SS_{1/2}$  values were higher versus Control (except  
227 for the 1-hour data), showing significant difference at 30 minutes (versus Control:  $p = 0.03$ ; versus 1 h:  
228  $p = 0.024$ ), 2 hours (versus Control:  $p < 0.001$ ; versus 1 h:  $p < 0.001$ ) and 4 hours (versus Control:  $p = 0.001$ ;  
229 versus 1 h:  $p = 0.002$ ).

230 During osmotic gradient ektacytometry measurement, the maximal EI values were increased compared  
231 to the Control, being significant at 30 minutes ( $p = 0.032$ ), 2 hours ( $p = 0.03$ ), 6 hours ( $p = 0.008$ ) and the  
232 same for 48 hours ( $p = 0.008$ ). The osmolarity values at the maximal EI were increased versus Control (at  
233 6 h:  $p = 0.023$ ; 48 h:  $p = 0.003$ ) and versus the 1-hour data (30 min:  $p = 0.008$ ; 2 h:  $p = 0.001$ ; 4 h:  $p = 0.026$ ;  
234 6 h:  $p < 0.001$ ; 24 h:  $p = 0.004$ ; 48 h:  $p < 0.001$ ).

235 Investigating membrane stability, we used shear stress of 100 Pa for 5 minutes. The elongation index (EI)  
236 - shear stress (SS) curves of the deformability measurements completed before and after the mechanical  
237 shearing of samples, taken at various time points after irradiation, and results are plotted on Fig. 6.  
238 Comparative data on EI-SS curves are summarized in Table 3. As expected, after the given mechanical  
239 stress, deformability of red blood cells decreased, but in different manner in the samples taken at various  
240 times after irradiation. The smallest difference was found at the 2-hour, 4-hour and 6-hour samples. Table 4  
241 shows the ratios of the values measured after versus before mechanical stress. In the function of shear  
242 stress, values over 1 Pa showed similar magnitude. Data at 3 Pa were selected for statistical comparison.  
243 Compared to the 30-minute data, values were significantly higher at 2 hours ( $p < 0.001$ ), 4 hours ( $p = 0.004$ )  
244 and 6 hours ( $p = 0.042$ ), when the before-after differences were the smallest in the original EI-SS curves,  
245 too. Values at 6 hours showed difference compared to the 2-hour values, too ( $p = 0.039$ ).

### 246 3.2.4. Red blood cell aggregation

247 In the following-up study part, a moderate decreasing tendency was observed in all the four aggregation  
248 index parameters, except for the values from the 1-hour samples (Fig. 5).

249 Compared to the Control group, M index of 5 seconds (M 5 s) showed significantly lower values only  
250 at 6 hours ( $p < 0.001$ ), 24 hours ( $p = 0.002$ ) and 48 hours ( $p < 0.001$ ). While compared to the 1-hour data,  
251 values were significantly different at 30 minutes ( $p < 0.001$ ), 2 hours ( $p < 0.001$ ), 4 hours ( $p < 0.001$ ), 6  
252 hours ( $p = 0.002$ ) and 24 hours ( $p < 0.001$ ) (Fig. 5A).

253 M index of 10 seconds (M 10 s) showed significantly lower values only at 6 hours ( $p < 0.001$ ) and 24  
254 hours ( $p < 0.001$ ), compared to Control (Fig. 5B).

255 Compared to the Control groups, all M1 index of 5 seconds (M 5 s) values were significantly lower  
256 over the follow-up period ( $p < 0.001$  for all, except for the 24-hour:  $p = 0.002$ ). Compared to the 1-hour  
257 data, the differences were significant at 30 minutes ( $p < 0.001$ ), 2 hours ( $p < 0.001$ ), 4 hours ( $p = 0.003$ ),  
258 6 hours ( $p = 0.002$ ) and 24 hours ( $p < 0.001$ ) (Fig. 5C).

Table 2  
Time-dependent changes of erythrocyte-related hematological, deformability and osmoscan parameters in blood samples taken from mice irradiated with 0.01 Gy  
(Control and 1 h values are transferred from Table 1)

Variable	Control	30 min	1 h	2 h	4 h	6 h	24 h	48 h
Hct [%]	44.52 ± 5.29	44.36 ± 2.67 #	37.4 ± 4.95*	44.27 ± 4.95 #	44.44 ± 3.23 #	49.42 ± 4.25*#	50.92 ± 4.25*#	52.01 ± 6.6*#
Hgb [g/dl]	11.2 ± 0.81	12.08 ± 0.48*#	10.53 ± 1.47	12.16 ± 0.27*#	12.33 ± 0.82*#	13.46 ± 0.5*#	13.45 ± 0.42*#	13.44 ± 0.42*#
MCV [fl]	57.33 ± 5.35	52.89 ± 1.98*#	61.03 ± 1.77	51.82 ± 1.14 #	51.75 ± 1.01 #	53.3 ± 3.78 #	55.1 ± 3.85 #	54.8 ± 3.95 #
RDW-CV% [%]	14 ± 0.22	14.21 ± 0.79	14.1 ± 0.45	13.64 ± 0.34	13.79 ± 0.34	14.71 ± 1.05	14.26 ± 0.66	14.46 ± 0.38
MCH [pg]	14.45 ± 0.91	14.36 ± 0.51	15.55 ± 1.28	14.15 ± 0.52	14.36 ± 0.27	14.56 ± 0.491	14.73 ± 0.78	14.22 ± 0.59
MCHC [g/dl]	25.45 ± 3.34	27.31 ± 1.71	28.15 ± 4.99	27.5 ± 0.86	27.76 ± 0.52	27.43 ± 2.41	26.81 ± 2.17	26.13 ± 2.55
EI <sub>max</sub>	0.544 ± 0.01	0.502 ± 0.02	0.539 ± 0.02	0.527 ± 0.04	0.494 ± 0.06*#	0.508 ± 0.03*	0.492 ± 0.05*#	0.51 ± 0.02*#
SS <sub>1/2</sub> [Pa]	1.87 ± 0.3	2.28 ± 0.46*#	1.77 ± 0.46	2.73 ± 0.49*#	2.72 ± 0.62*#	2.12 ± 0.6	2.1 ± 0.49	2.01 ± 0.56
Maximal EI	0.444 ± 0.04	0.497 ± 0.005*	0.445 ± 0.08	0.504 ± 0.007*	0.465 ± 0.04	0.504 ± 0.002*	0.471 ± 0.04	0.505 ± 0.002*
Osmolarity at max. EI [mOsm/kg]	312 ± 20.55	334.8 ± 8.64 #	293.2 ± 9.98	332.5 ± 12.07 #	326.3 ± 22.81#	346 ± 17.5*#	333 ± 19.35 #	360.2 ± 14.7*#

means ± S.D. EI<sub>max</sub> = calculated maximal elongation index (EI); SS<sub>1/2</sub> = calculated shear stress (SS) at half EI<sub>max</sub> by Lineweaver-Burk analysis; Maximal EI = maximal EI values measured in osmoscan function. \**p* < 0.05 vs. Control; # *p* < 0.05 vs. 1 h.

Table 3

Changes of red blood cell deformability parameters during membrane stability test, presenting values before and after the mechanical stress (shearing at 100 Pa for 300 sec)

Variable		30 min	2 h	4 h	6 h	24 h	48 h
EI at 3 Pa	before	0.238 ± 0.01	0.277 ± 0.02	0.242 ± 0.03	0.32 ± 0.02	0.311 ± 0.02	0.33 ± 0.01
	after	0.137 ± 0.01*	0.23 ± 0.01*	0.211 ± 0.04*	0.237 ± 0.02*	0.202 ± 0.02*	0.223 ± 0.02*
EI <sub>max</sub>	before	0.493 ± 0.02	0.494 ± 0.02	0.489 ± 0.02	0.535 ± 0.05	0.499 ± 0.03	0.502 ± 0.01
	after	0.407 ± 0.01*	0.438 ± 0.04*	0.408 ± 0.06*	0.442 ± 0.02*	0.402 ± 0.02*	0.434 ± 0.03*
SS <sub>1/2</sub> [Pa]	before	1.82 ± 0.34	2.43 ± 0.24	2.85 ± 1.04	1.34 ± 0.66	1.58 ± 0.46	1.32 ± 0.29
	after	7.03 ± 1.01*	4.13 ± 0.57*	4.18 ± 1.59*	5.79 ± 1.24*	6.73 ± 1.3*	6.61 ± 1.31*

means ± S.D. EI = elongation index; EI<sub>max</sub> = calculated maximal EI; SS<sub>1/2</sub> = calculated shear stress (SS) at half EI<sub>max</sub> by Lineweaver-Burk analysis \**p* < 0.05 vs. before.




Fig. 5. Changes of aggregation index M 5 s (A), M 10 s (B), M1 5 s (C) and M1 10 s (D) in the function of time after 0.01 Gy dose of irradiation. means ± S.D.; \**p* < 0.05 vs. Control; #*p* < 0.05 vs. 1 h.

259 M1 index of 10 seconds (M 10 s) showed significantly lower values versus Control at 24 hours  
 260 (*p* = 0.018) (Fig. 5D).

#### 261 4. Discussion and conclusions

262 The anti-inflammatory and sterilisation effects of irradiation are well known and clinically widespread  
 263 for a long time [7]. It is used in case of traumas, viral inflammations, leukaemia and even in case of




Fig. 6. Elongation index (EI) - shear stress (SS) curves of the deformability measurements completed before (white) and after (black) the mechanical shearing (100 Pa for 5 minutes) of samples taken at various time points (A: 30 minutes, B: 2 hours, C: 4 hours, D: 6 hours, E: 24 hours, F: 48 hours) after irradiation at 0.01 Gy. means  $\pm$  S.D.; \* $p < 0.05$  vs. before.

264 transfused blood to prevent graft versus host disease [7, 9, 30, 31, 38]. Nevertheless, the optimisation of  
 265 dosing is in its inauguration stage.

266 Selim et al. analyzed the effects of 25, 50 and 100 Gy gamma-rays on stored red blood cells using  
 267 alpha-lipoic acid, in order to help preserving cell viability and structural integrity of the cell membrane,  
 268 reducing the effect of oxidative damage [15, 40, 41]. The dose of 25 Gy was found to be safe related to  
 269 erythrocyte integrity. However, this dose may cause significantly increased mean cell volume, as showed

Table 4

Ratios (after/before) calculated from elongation index values at different shear stress values measured before and after mechanical shearing at 100 Pa for 300 sec

Timepoint	Shear stress [Pa]						
	0.95	1.69	3	5.33	9.49	16.87	30
30 min	0.72 ± 0.04	0.57 ± 0.03	0.63 ± 0.07	0.69 ± 0.06	0.79 ± 0.03	0.83 ± 0.03	0.88 ± 0.02
2 h	0.91 ± 0.13	0.8 ± 0.07	0.83 ± 0.04*	0.87 ± 0.01	0.89 ± 0.03	0.9 ± 0.02	0.91 ± 0.03
4 h	0.85 ± 0.1	0.78 ± 0.07	0.86 ± 0.11*	0.86 ± 0.08	0.87 ± 0.06	0.89 ± 0.07	0.89 ± 0.07
6 h	0.7 ± 0.11	0.65 ± 0.12	0.74 ± 0.07*#	0.79 ± 0.09	0.82 ± 0.09	0.86 ± 0.04	0.92 ± 0.04
24 h	0.71 ± 0.14	0.57 ± 0.04	0.65 ± 0.04	0.71 ± 0.05	0.75 ± 0.04	0.81 ± 0.03	0.87 ± 0.03
48 h	0.75 ± 0.07	0.62 ± 0.04	0.67 ± 0.07	0.74 ± 0.06	0.81 ± 0.05	0.86 ± 0.05	0.92 ± 0.05

means ± S.D. \* $p < 0.05$  vs. 30 min, #  $p < 0.05$  vs. 2 h.

by Kim et al. [19]. Cicha et al. also demonstrated that erythrocyte deformability impairs in exposure to a 35 Gy gamma irradiation [12].

Mihaescu et al. used 20 Gy in mice to investigate the prothrombotic impact in the microcirculation of the large bowel, which was partly mediated by P-selectin and P-selectin glycoprotein ligand-1 (PSGL-1) [28]. Inhibition or lack of P-selectin or PSGL-1 may reduce the thrombotic events in blood vessels [29].

Micke et al. investigated the effects of low-dose irradiation on neutrophil granulocytes of human donors [27], on the other hand, Reverberi, Govoni and Verenini reported the effects on human red blood cells [38]. Serhatlioglu et al. concentrated on the long term effects of recurrent low-dose irradiations on CD4+ T-lymphocyte ratio and serum total IgA, IgG, IgM and C3, C4 levels with the help of volunteering radiologists and those who have never been directly or indirectly irradiated so far [42].

However, immediate (short-term) effects of irradiation of blood of living organisms are near to nonexistent. Maks et al. analyzed the white blood cell counts of mice after proton and gamma irradiation up to 2 Gy by dose rates of 0.5 Gy/min or 0.5 Gy/h. They found a dose-dependent decrease of total leukocyte count [23]. Bogdandi et al. used as low as 0.01-0.1 Gy dose total-body irradiation in mice, demonstrating dose-dependent differences up to 2 Gy in apoptotic processes in various immune cells [8]. Still, *in vitro* research data on low-dose photon irradiation of the blood of living organism are abundant.

Present paper endeavours to narrow this gap in knowledge by supplying data on short-term red blood cell alteration of living mice after low-dose photon irradiation.

According to the results, it seems that the dose-dependent changes of blood micro-rheological parameters are not linear. The irradiation dose of 0.01 Gy acted as a point of 'inflexion', because by this dose we found the most expressed changes in the hematological parameters, as well as in the red blood cell aggregation, deformability and osmoscan data. This was the reason why we have chosen this dose for the second part of the study, where the time-dependent changes were analyzed further. In the literature, we did not find other study using these very low doses of irradiation together investigation of micro-rheological parameters.

Wen et al. also used  $^{60}\text{Co}$  irradiation, but in higher dose, for studying hemorheological changes in an animal model. They found that 7 Gy total-body irradiation caused long-lasting decrease of hematocrit (over 10 days, and slow normalization over 30 days) in rabbits, accompanied by decreased deformability index [49].

Red blood cell deformability is determined by cell volume, surface-volume ratio, cell morphology, inner viscosity (hemoglobin content), and membrane viscosity [1, 2, 24, 35]. Red blood cell aggregation

is determined or influenced also by the cell-morphology, but by the composition of the surface glycocalyx, as well as the plasmatic factors, such as the fibrinogen concentration together with the shearing forces [4, 35, 37]. Important influencing factors are the oxygenation, the temperature, the pH and the osmolarity of the micro-environment of the red blood cells [1, 4, 33, 34, 37]. Irradiation of various origins may interact with these factors at numerous points, by influencing the cell volume, the membrane integrity by oxidative damage and structural changes [9, 12, 17, 26, 30, 38]. Since red blood cell deformability plays important role in determining blood flow and microcirculation [1, 10, 22, 43], these changes may cause further alteration in tissue perfusion.

However, the background, dose- and time-dependency of the direction of the changes, like improving or impairing micro-rheological variables are still unclear. Does decreased aggregation mean a rheological improvement after irradiation? Or does it reflect such changes in cellular factors that may reduce the aggregation ability but together with a relatively stable deformability and with shifting of the optimal osmolarity (osmolarity at maximal EI)? Too composite questions so far.

Using laser irradiation (632.8 and 532 nm) Mi et al. reported reduced blood viscosity, and red blood cell deformability could be improved in human blood [25]. They hypothesized that the 'mobilization' of membrane-linked hemoglobin might cause this enhancing effect, which has been demonstrated in their later porcine study [26].

It seems that together with micro-rheological changes, irradiation may cause a complex impact on the blood organ with labyrinthine interactions.

Therefore, it would be also interesting to see more exact evidences for a kind of inflexion point of changes in the function of dose and time. For revealing the background and find evidences, further studies are needed in the future using hemorheological, hematological, hemostaseological, immunological and histomorphological as well as ultrastructural complex investigations.

## Acknowledgments

Authors are grateful to the technical and laboratory staff of the Department of Radiotherapy and the Department of Operative Techniques and Surgical Research. Special thanks to Krisztina Gazso, biologist and Mihaly Simon, physicist (Department of Radiotherapy), to Zoltan Klarik, M.D. and Eniko Toth, M.D. (Department of Operative Techniques and Surgical Research) for their assistance.

Grants: Baross Gabor Programme of National Research and Technology Office (OMFB-00411/2010); and Janos Bolyai Research Scholarship of the Hungarian Academy of Sciences (2010-2013, N. Nemeth).

The authors comply with the Ethical Guidelines for Publication in *Clinical Hemorheology and Microcirculation* as published on the IOS Press website and in Volume 44, 2010, pp. 1-2 of this journal.

## References

- [1] O.K. Baskurt, Mechanisms of blood rheology alterations, in: *Handbook of Hemorheology and Hemodynamics*, O.K. Baskurt, M.R. Hardeman, M.W. Rampling and H.J. Meiselman, eds., IOS Press, Amsterdam, The Netherlands, 2007, pp. 170–190.
- [2] O.K. Baskurt, M. Boynard, G.C. Cokelet, P. Connes, B.M. Cooke, S. Forconi, M.R. Hardeman, F. Jung, F. Liao, H.J. Meiselman, G. Nash, N. Nemeth, B. Neu, B. Sandhagen, S. Shin, G. Thurston and J.L. Wautier, International Expert Panel for Standardization of Hemorheological Methods, New guidelines for hemorheological laboratory techniques, *Clin Hemorheol Microcirc* **42** (2009), 75–97.

- 341 [3] O.K. Baskurt, M.R. Hardeman, M. Uyuklu, P. Ulker, M. Cengiz, N. Nemeth, S. Shin, T. Alexy and H.J. Meiselman,  
342 Parameterization of red blood cell elongation index – shear stress curves obtained by ektacytometry, *Scand J Clin Lab*  
343 *Invest* **69** (2009), 777–788.
- 344 [4] O. Baskurt, B. Neu and H.J. Meiselman, Determinants of red blood cell aggregation, in: *Red Blood Cell Aggregation*, CRC  
345 Press, Boca Raton, USA, 2012. pp. 9–29.
- 346 [5] O.K. Baskurt and H.J. Meiselman, Data reduction methods for ektacytometry in clinical hemorheology, *Clin Hemorheol*  
347 *Microcirc* (2012), DOI: 10.3233/CH-2012-1616
- 348 [6] I. Beli, G. Koukourakis, K. Platoni, M. Tolia, N. Kelekis, J. Kouvaris, C. Syrigos, K. Mystakidou, C. Varveris and V.  
349 Kouloulias, Hypofractionated radiotherapy in non small cell lung cancer: A review of the current literature, *Rev Recent*  
350 *Clin Trials* **5** (2010), 103–111.
- 351 [7] S.M. Bentzen, Quantitative clinical radiobiology, *Acta Oncol* **32** (1993), 259–275.
- 352 [8] E.N. Bogdandi, A. Balogh, N. Felgyinszki, T. Szatmari, E. Persa, G. Hildebrandt, G. Safrany and K. Lumniczky, Effects  
353 of low-dose radiation on the immune system of mice after total-body irradiation, *Radiat Res* **174** (2010), 480–489.
- 354 [9] L.N. Button, W.C. DeWolf, P.E. Newburger, M.S. Jacobson and S.V. Keyv, The effects of irradiation on blood components,  
355 *Transfusion* **21** (1981), 419–426.
- 356 [10] S. Chien, Red cell deformability and its relevance to blood flow, *Annu Rev Physiol* **49** (1987), 177–192.
- 357 [11] Y. Chen, X. Chen, Y. Li, H. Zhang, Y. Xie, X. Zhang, H. Ren, Y. Wang, S. Liao, M. He, J. Ren, J. Zhang, X. Zhou, T.  
358 Zheng, B.A. Kilfoy and Y. Zhang, Early effects of low dose C ion or x-ray irradiation on peripheral blood lymphocytes of  
359 patients with alimentary tract cancer, *Dose Response* **9** (2011), 356–368.
- 360 [12] I. Cicha, Y. Suzuki, N. Tateishi, M. Shiba, M. Muraoka, K. Tadokoro and N. Maeda, Gamma-ray-irradiated red blood cells  
361 stored in mannitol-adenine-phosphate medium: Rheological evaluation and susceptibility to oxidative stress, *Vox Sang* **79**  
362 (2000), 75–82.
- 363 [13] M.R. Clark, N. Mohandas and S.B. Shohet, Osmotic gradient ektacytometry: Comprehensive characterization of red cell  
364 volume and surface maintenance, *Blood* **61** (1983), 899–910.
- 365 [14] L.T. Dauer, A.L. Brooks, D.G. Hoel, W.F. Morgan, D. Stram and P. Tran, Review and evaluation of updated research on  
366 the health effects associated with low-dose ionising radiation, *Radiat Prot Dosimetry* **140** (2010), 103–136.
- 367 [15] O.S. Desouky, N.S. Selim, E.M. Elbakrawy and R.A. Rezk, Impact evaluation of  $\alpha$ -lipoic acid in gamma-irradiated  
368 erythrocytes, *Radiat Phys Chem* **80** (2011), 446–452.
- 369 [16] D.S. Gridley, T.L. Freeman, A.Y. Makinde, A.J. Wroe, X. Luo-Owen, J. Tian, X.W. Mao, S. Rightnar, A.R. Kennedy, J.M.  
370 Slater and M.J. Pecaut, Comparison of proton and electron radiation effects on biological responses in liver, spleen and  
371 blood, *Int J Radiat Biol* **87** (2011), 1173–1181.
- 372 [17] J. Hirayama, H. Abe, H. Azuma and H. Ikeda, Leakage of potassium from red blood cells following gamma ray irradiation  
373 in the presence of dipyridamole, trolox, human plasma or mannitol, *Biol Pharm Bull* **28** (2005), 1318–1320.
- 374 [18] M.R. Hardeman, P.T. Goedhart and S. Shin, Methods in hemorheology, in: *Handbook of Hemorheology and Hemodynamics*,  
375 O.K. Baskurt, M.R. Hardeman, M.W. Rampling and H.J. Meiselman, eds., IOS Press, Amsterdam, The Netherlands, 2007,  
376 pp. 242–266.
- 377 [19] Y.K. Kim, E.H. Kwon, D.H. Kim, D.I. Won, S. Shin and J.S. Suh, Susceptibility of oxidative stress on red blood cells  
378 exposed to gamma rays: Hemorheological evaluation, *Clin Hemorheol Microcirc* **40** (2008), 315–324.
- 379 [20] H.C. Kwon, S.K. Kim, W.K. Chung, M.J. Cho, J.S. Kim, J.S. Kim, S.R. Moon, W.Y. Park, S.J. Ahn, Y.K. Oh, H.G. Yun and  
380 B.S. Na, Effect of pentoxifylline on radiation response of non-small cell lung cancer: A phase III randomized multicenter  
381 trial, *Radiother Oncol* **56** (2000), 175–179.
- 382 [21] J.P. Lopes de Almeida, T. Freitas-Santos and C. Saldanha, Erythrocyte deformability dependence on band 3 protein in an  
383 *in-vitro* model of hyperfibrinogenemia, *Clin Hemorheol Microcirc* **50** (2012), 213–219.
- 384 [22] J.P. Lopes de Almeida, S. Oliveira and C. Saldanha, Erythrocyte as a biological sensor, *Clin Hemorheol Microcirc* **51**  
385 (2012), 1–20.
- 386 [23] C.J. Maks, X.S. Wan, J.H. Ware, A.L. Romero-Weaver, J.K. Sanzari, J.M. Wilson, S. Rightnar, A.J. Wroe, P. Koss, D.S.  
387 Gridley, J.M. Slater and A.R. Kennedy, Analysis of white blood cell counts in mice after gamma- or proton-radiation  
388 exposure, *Radiat Res* **176** (2011), 170–176.
- 389 [24] H.J. Meiselman, Morphological determinants of red blood cell deformability, *Scand J Clin Lab Invest* **41** Suppl 156 (1981),  
390 27–34.
- 391 [25] X.Q. Mi, J.Y. Chen, Y. Cen, Z.J. Liang and L.W. Zhou, A comparative study of 632.8 and 532nm laser irradiation on some  
392 rheological factors in human blood *in vitro*, *J Photochem Photobiol B* **74** (2004), 7–12.

- [26] X.Q. Mi, J.Y. Chen and L.W. Zhou, Effect of low laser irradiation on disconnecting the membrane-attached hemoglobin from erythrocyte membrane, *J Photochem Photobiol B-Biol* **83** (2006), 146–150.
- [27] O. Micke, A. Haidenberger, T. Auer, S. Egger, M.H. Seegenschmiedt, P. Hengster and A.F. DeVries, Irradiation causes biphasic neutrophilic granulocyte phagocytic function, *Strahlenther Onkol* **181** (2005), 313–318.
- [28] A. Mihaescu, C. Thornberg, S. Santen, S. Mattsson, B. Jeppsson and H. Thorlacius, Radiation-induced platelet-endothelial cell interactions are mediated by P-selectin and P-selectin glycoprotein ligand-1 in the colonic microcirculation, *Surgery* **151** (2012), 606–611.
- [29] K. Miszti-Blasius, I.B. Debreceni, S. Felszeghy, B. Dezsó and J. Kappelmayer, Lack of P-selectin glycoprotein ligand-1 protects mice from thrombosis after collagen/epinephrine challenge, *Thromb Res* **127** (2011), 228–234.
- [30] G. Moroff and N.L.C. Luban, The influence of gamma radiation on red cell and platelet properties, *Transfus Sci* **15** (1994), 141–148.
- [31] G. Moroff, S. Holme, J.P. AuBuchon, W.A. Heaton, J.D. Sweeney and L.I. Friedman, Viability and *in vitro* properties of AS-1 red cells after gamma irradiation, *Transfusion* **39** (1999), 128–134.
- [32] A.V. Muravyov and I.A. Tikhomirova, Role molecular signaling pathways in changes of red blood cell deformability, *Clin Hemorheol Microcirc* (2012), DOI: 10.3233/CH-2012-1575
- [33] N. Nemeth, I. Miko, A. Furka, F. Kiss, I. Furka, A. Koller and M. Szilasi, Concerning the importance of changes in hemorheological parameters caused by acid-base and blood gas alterations in experimental surgical models, *Clin Hemorheol Microcirc* **51** (2012), 43–50.
- [34] N. Nemeth, F. Kiss, Z. Klarik and I. Miko, Comparative osmotic gradient ektacytometry data on inter-species differences of experimental animals, *Clin Hemorheol Microcirc* (2012), DOI: 10.3233/CH-2012-1620
- [35] S. de Oliveira and C. Saldanha, An overview about erythrocyte membrane, *Clin Hemorheol Microcirc* **44** (2010), 63–74.
- [36] B. Ozturk, I. Egehan, S. Atavci and M. Kitapci, Pentoxifylline in prevention of radiation-induced lung toxicity in patients with breast and lung cancer: A double-blind randomized trial, *Int J Radiat Oncol Biol Phys* **58** (2004), 213–219.
- [37] W.H. Reinhart, Peculiar red cell shapes: Fahraeus Lecture 2011, *Clin Hemorheol Microcirc* **49** (2011), 11–27.
- [38] R. Reverberi, M. Govoni and M. Verenini, Deformability and viability of irradiated red cells, *Ann Ist Super Sanita* **43** (2007), 176–185.
- [39] P. Ruef, M. Ehrhard, D. Frommhold, L. Koch, B. Fritzsching and J. Poeschl, Lipid A decreases human erythrocytes deformability by increasing intracellular Ca (2+): Effects of verapamil, staurosporine and the rho-kinase inhibitor Y-27632, *Clin Hemorheol Microcirc* **49** (2011), 315–322.
- [40] N.S. Selim, O.S. Desouky, E.M. Elbakrawy and R.A. Rezk, Electrical behavior of stored erythrocytes after exposure to gamma radiation and the role of  $\alpha$ -lipoic acid as radioprotector, *Appl Radiat Isot* **68** (2010), 1018–1024.
- [41] N.S. Selim, O.S. Desouky, N.M. Ismail and A.Z. Dakrory, Spectroscopic analysis of irradiated erythrocytes, *Radiat Phys Chem* **80** (2011), 1337–1342.
- [42] S. Serhatlioglu, E. Ogur, A.T. Ozan, F. Gursu, A. Godekmerdan and A. Ayar, Biochemical and immunological effects of ionizing radiation in radiology staff members, *Tani Girisim Radyol* **10** (2004), 97–102.
- [43] K.H. Son, C.B. Ahn, S.H. Lee, I. Noh, E. Shin, M. So, H. Kim, K. Sun and K.T. Kim, Changes in RBC deformability and oxygen-delivering ability in cold blood cardioplegia, *Clin Hemorheol Microcirc* **48** (2011), 223–229.
- [44] H. Stadnick, R. Onell, J.P. Acker and J.L. Holovati, Eadie-Hofstee analysis of red blood cell deformability, *Clin Hemorheol Microcirc* **47** (2011), 229–239.
- [45] Y. Suzuki, N. Tateishi, I. Cicha, M. Shiba, M. Muraoka, K. Tadokoro and N. Maeda, Decreased deformability of the X-ray-irradiated red blood cells stored in mannitol-adenine-phosphate medium, *Clin Hemorheol Microcirc* **22** (2000), 131–141.
- [46] G.F. von Tempelhoff, L. Heilmann, K. Pollow and G. Hommel, Monitoring of rheologic variables during postoperative high-dose brachytherapy for uterine cancer, *Clin Appl Thromb Hemost* **10** (2004), 239–348.
- [47] I.A. Tikhomirova, A.O. Oslyakova and S.G. Mikhailova, Microcirculation and blood rheology in patients with cerebrovascular disorders, *Clin Hemorheol Microcirc* **49** (2011), 295–305.
- [48] H.D. Thames, Jr, H.R. Withers, L.J. Peters and G.H. Fletcher, Changes in early and late radiation responses with altered dose fractionation: Implications for dose-survival relationships, *Int J Radiat Oncol Biol Phys* **8** (1982), 219–226. JJr.
- [49] Z. Wen, L. Xie, Z. Yan, W. Yao, K. Chen, W. Ka and S. Chien, Effect of  $^{60}\text{Co}$  irradiation on characteristics of hemorheology in rabbits, *Clin Hemorheol Microcirc* **25** (2001), 75–81.
- [50] F. Zywiets, L. Böhm, C. Sagowski and W. Kehrl, Pentoxifylline enhances tumor oxygenation and radiosensitivity in rat rhabdomyosarcomas during continuous hyperfractionated irradiation, *Strahlenther Onkol* **180** (2004), 306–314.