



Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s00431-012-1925-z>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

1	Article Title	Relation between biomarkers and clinical severity in patients with Smith–Lemli–Opitz syndrome	
2	Article Sub- Title		
3	Article Copyright - Year	Springer-Verlag Berlin Heidelberg 2013 (This will be the copyright line in the final PDF)	
4	Journal Name	European Journal of Pediatrics	
5	Corresponding Author	Family Name	Oláh
6		Particle	
7		Given Name	Anna V.
8		Suffix	
9		Organization	University of Debrecen
10		Division	Department of Laboratory Medicine, Medical and Health Science Center
11		Address	98 Nagyerdei kt., Debrecen 4032, Hungary
12		e-mail	olaha@med.unideb.hu
13	Author	Family Name	Szabó
14		Particle	
15		Given Name	Gabriella P.
16		Suffix	
17		Organization	University of Debrecen
18		Division	Department of Pediatrics
19		Address	Debrecen, Hungary
20		e-mail	
21	Author	Family Name	Varga
22		Particle	
23		Given Name	József
24		Suffix	
25		Organization	University of Debrecen
26		Division	Department of Nuclear Medicine, Medical and Health Science Center
27		Address	Debrecen, Hungary
28		e-mail	
29	Author	Family Name	Balogh

30		Particle	
31		Given Name	Lídia
32		Suffix	
33		Organization	Semmelweis University
34		Division	1st Department of Pediatrics
35		Address	Budapest, Hungary
36		e-mail	
<hr/>			
37		Family Name	Csábi
38		Particle	
39		Given Name	Györgyi
40	Author	Suffix	
41		Organization	University of Pécs
42		Division	Department of Pediatrics
43		Address	Pecs, Hungary
44		e-mail	
<hr/>			
45		Family Name	Csákváry
46		Particle	
47		Given Name	Violetta
48	Author	Suffix	
49		Organization	Markusovszky County Hospital
50		Division	
51		Address	Szombathely, Hungary
52		e-mail	
<hr/>			
53		Family Name	Erwa
54		Particle	
55		Given Name	Wolfgang
56	Author	Suffix	
57		Organization	LKH Medical University
58		Division	
59		Address	Graz, Austria
60		e-mail	
<hr/>			
61		Family Name	Balogh
62		Particle	
63		Given Name	István
64	Author	Suffix	
65		Organization	University of Debrecen
66		Division	Department of Laboratory Medicine, Medical and Health Science Center

67		Address	98 Nagyerdei kt., Debrecen 4032, Hungary
68		e-mail	
69		Received	28 October 2012
70	Schedule	Revised	19 December 2012
71		Accepted	23 December 2012
72	Abstract	<p>Smith–Lemli–Opitz syndrome (SLOS), a multiple congenital anomaly with severe mental retardation, is caused by decreased activity of 7-dehydrocholesterol reductase. Fifteen Hungarian patients were diagnosed with SLOS on the basis of clinical symptoms, serum cholesterol, 7-dehydrocholesterol, and molecular genetic testing. Their age at the time of diagnosis in mild SLOS ($n=4$, clinical score <20) was 0.5–18 years, cholesterol was 2.37 ± 0.8 mmol/L, and 7DHC was 0.38 ± 0.14 mmol/L. In the group of typical SLOS ($n=7$, score 20–50), the diagnosis was set up earlier (age of 0.1–7 years); t-cholesterol was 1.47 ± 0.7 mmol/L, and 7DHC was 0.53 ± 0.20 mmol/L. Patients with severe SLOS ($n=4$, clinical score >50) died as newborns and had the lowest t-cholesterol (0.66 ± 0.27 mmol/L), and 7DHC was 0.47 ± 0.14 mmol/L. Correlation coefficient with clinical severity was 0.74 for initial t-cholesterol and 0.669 for Cho/7DHC. Statistically significant difference was between the initial t-cholesterol of mild and severe SLOS ($p=0.01$), and between the Cho/7DHC ratios of groups ($p=0.004$). In severe SLOS, the percentage of α-lipoprotein was significantly lower than in typical ($p=0.003$) and mild SLOS ($p=0.004$). Although serum albumin, total bilirubin, and hemostasis parameters remained in the reference range during cholesterol supplementation ($n=10$) combined with statin therapy ($n=9$), increase of aspartate aminotransferase and alanine aminotransferase in 50 % of the patients probably refers to a reversible alteration of liver function; therefore, statin therapy was suspended. <i>Conclusion:</i> life expectancy is fundamentally determined by the initial t-cholesterol, but dehydrocholesterol and α-lipoprotein have prognostic value. Accumulation of hepatotoxic DHC may inhibit the synthesis of α-lipoproteins, decreasing the reverse cholesterol transport. During statin therapy, we suggest monitoring of lipid parameters and liver function.</p>	
73	Keywords separated by ' - '	7-Dehydrocholesterol - Cholesterol - Lipoprotein electrophoresis - Liver function - Smith–Lemli–Opitz syndrome - Statin	
74	Foot note information		

Relation between biomarkers and clinical severity in patients with Smith–Lemli–Opitz syndrome

Anna V. Oláh · Gabriella P. Szabó · József Varga · Lidia Balogh · Györgyi Csábi · Violetta Csákváry · Wolfgang Erwa · István Balogh

Received: 28 October 2012 / Revised: 19 December 2012 / Accepted: 23 December 2012
© Springer-Verlag Berlin Heidelberg 2013

Abstract Smith–Lemli–Opitz syndrome (SLOS), a multiple congenital anomaly with severe mental retardation, is caused by decreased activity of 7-dehydrocholesterol reductase. Fifteen Hungarian patients were diagnosed with SLOS on the basis of clinical symptoms, serum cholesterol, 7-dehydrocholesterol, and molecular genetic testing. Their age at the time of diagnosis in mild SLOS ($n=4$, clinical score <20) was 0.5–18 years, cholesterol was 2.37 ± 0.8 mmol/L, and 7DHC was 0.38 ± 0.14 mmol/L. In the group of typical SLOS ($n=7$, score 20–50), the diagnosis was set up earlier (age of 0.1–7 years); t-cholesterol was 1.47 ± 0.7 mmol/L, and 7DHC was 0.53 ± 0.20 mmol/L. Patients with severe SLOS ($n=4$, clinical

score >50) died as newborns and had the lowest t-cholesterol (0.66 ± 0.27 mmol/L), and 7DHC was 0.47 ± 0.14 mmol/L. Correlation coefficient with clinical severity was 0.74 for initial t-cholesterol and 0.669 for Cho/7DHC. Statistically significant difference was between the initial t-cholesterol of mild and severe SLOS ($p=0.01$), and between the Cho/7DHC ratios of groups ($p=0.004$). In severe SLOS, the percentage of α -lipoprotein was significantly lower than in typical ($p=0.003$) and mild SLOS ($p=0.004$). Although serum albumin, total bilirubin, and hemostasis parameters remained in the reference range during cholesterol supplementation ($n=10$) combined with statin therapy ($n=9$), increase of aspartate aminotransferase and alanine aminotransferase in 50 % of the patients probably refers to a reversible alteration of liver function; therefore, statin therapy was suspended. *Conclusion:* life expectancy is fundamentally determined by the initial t-cholesterol, but dehydrocholesterol and α -lipoprotein have prognostic value. Accumulation of hepatotoxic DHC may inhibit the synthesis of α -lipoproteins, decreasing the reverse cholesterol transport. During statin therapy, we suggest monitoring of lipid parameters and liver function.

27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

A. V. Oláh (✉) · I. Balogh
Department of Laboratory Medicine, Medical and Health Science Center, University of Debrecen, 98 Nagyterdei krt., Debrecen 4032, Hungary
e-mail: olaha@med.unideb.hu

G. P. Szabó
Department of Pediatrics, University of Debrecen, Debrecen, Hungary

J. Varga
Department of Nuclear Medicine, Medical and Health Science Center, University of Debrecen, Debrecen, Hungary

L. Balogh
1st Department of Pediatrics, Semmelweis University, Budapest, Hungary

G. Csábi
Department of Pediatrics, University of Pécs, Pécs, Hungary

V. Csákváry
Markusovszky County Hospital, Szombathely, Hungary

W. Erwa
LKH Medical University, Graz, Austria

Keywords 7-Dehydrocholesterol · Cholesterol · Lipoprotein electrophoresis · Liver function · Smith–Lemli–Opitz syndrome · Statin

48 Q3
49
50

List of abbreviations

ALT	Alanine aminotransferase	54
AST	Aspartate aminotransferase	56
ALP	Alkaline phosphatase	58
t-Cho	Total cholesterol	60
CK	Creatine kinase	63
7-DHC	7-Dehydrocholesterol	64

51
52 Q4
53
54
55
56
57
58
59
60
61
62
63
64

66 DHCR7 7-Dehydrocholesterol reductase
 68 HDL-C High-density cholesterol
 70 KM Kilomicron
 73 GGT gamma-Glutamyltransferase
 74 LDH Lactate dehydrogenase
 76 SLOS Smith–Lemli–Opitz syndrome
 78

syndrome in our first patient was identified by L. Kozak 123
 and his coworkers [31]. Very recently, we published a 124
 paper [1] on the genetic background of the Hungarian 125
 patients with SLOS. Similar to the finding of Porter 126
 [21], we did not find a strong connection between the 127
 genotype and phenotype. Therefore, the relation between 128
 biomarkers and clinical feature was investigated in di- 129
 agnostic and therapeutic aspects. 130

79 **Introduction**

80 The Smith–Lemli–Opitz syndrome (OMIM 270400), an au-
 81 toosomal recessive, severe developmental disorder with multi-
 82 ple congenital anomalies, is caused by a defect of cholesterol
 83 biosynthesis [3, 12, 20–22]. The syndrome was first reported
 84 by Smith et al. in 1964 and was characterized by a dysmorphic
 85 face, microcephaly, hypospadiasis, and severe growth retard-
 86 ation [26]. The cause of Smith–Lemli–Opitz syndrome
 87 (SLOS) is the defective function of the 7-dehydrocholesterol
 88 reductase (DHCR7) enzyme which catalyzes the last step of
 89 cholesterol biosynthesis [7, 32]. This enzyme is responsible
 90 for the transformation of 7-dehydrocholesterol to cholesterol.
 91 Cholesterol is an important component of the cell membrane,
 92 mitochondrial membrane, and myelin formation in the brain,
 93 spinal cord, and peripheral nervous system. Cholesterol acts
 94 also as a precursor for bile acids and steroid hormones, and
 95 plays an important role in the embryonic hedgehog signaling
 96 mechanism during embryogenesis as well [13]. For phenotypic
 97 characterization, the modified Bialer scoring system of
 98 Kelley and Hennekam has been used which weight embryo-
 99 logically separate organ systems equally [12, 13].

100 Calculation of clinical severity scores is based on evalu-
 101 ation of anatomical abnormalities in ten embryologically
 102 separated organs (brain, oral region, eye, heart, kidney, liver,
 103 lung, bowel, and genitals) [13]. In 1998, the human *DHCR7*
 104 gene was cloned by three different groups [4, 35, 36]. Over
 105 140 different mutations in the *DHCR7* gene have been
 106 published to date [39, 40 and Human Gene Mutation Data-
 107 base]. For the treatment of the disease, during the past
 108 20 years, different therapeutic approaches were applied such
 109 as cholesterol substitution, with or without bile acids, and
 110 simvastatin that decreases the level of 7DHC and enhances
 111 the residual activity of the DHCR7, reportedly with im-
 112 provement in both biochemical parameters and clinical
 113 symptoms [7, 8, 18, 32]. Recently, the efficiency of choles-
 114 terol supplementation has been debated [5, 25], and statins
 115 cannot be considered as a safety approach in each SLO
 116 patient because of potential side effects [28].

117 In the past, the diagnosis of SLO syndrome was estab-
 118 lished mainly on the basis of characteristic phenotypic fea-
 119 tures, including severe mental and somatic retardation. The
 120 wide range of gene mutations in SLO syndrome and altera-
 121 tion of biochemical parameters have been described in
 122 different populations [2, 27]. A new mutation of SLO

Patients and methods 131

132 During the last decade, 15 patients (age of 0.1–18 years, 132
 eight males and seven females) were diagnosed with SLO 133
 syndrome in Hungary, the first case in 2002 [30]. After 134
 observation of clinical symptoms, their diagnosis was 135
 proved by serum 7DHC level. Anatomical abnormalities 136
 of ten embryologically separated organs (brain, oral re- 137
 gion, acral, eye, heart, kidney, liver, lung, bowel, and 138
 agenitals) have been scored [13]. On the base of clinical 139
 severity scores, patients were assigned to three groups: 140
 patients with mild SLOS were defined by a score below 141
 20, typical SLOS means 20–50 scores, and a score above 142
 50 means a severe type of SLO syndrome [12, 13]. The 143
 patients enrolled into the mild group are still alive ($n=4$; 144
 age when the diagnosis was set up, 0.5–18 years). In the 145
 typical SLOS group ($n=7$; age at diagnosis, 0.1–7 years), 146
 two children lived less than 2 years of age; five children 147
 are still alive. All patients with severe SLOS died in the 148
 newborn period ($n=4$, age < 2 months). After setting up 149
 the diagnosis, all patients received cholesterol substitution 150
 ($n=10$, Cholesterol Module, 50–250 mg/kg/day, Nutricia; 151
 no 18.012). It was completed with statin therapy in nine 152
 patients. Dosage of the statin was 0.2–0.4 mg/kg/day, and 153
 clinical state and efficiency of therapy were monitored by 154
 regular clinical checkup in 3- to 6-month periods includ- 155
 ing the evaluation of clinical condition, anthropometric 156
 parameters, serum cholesterol, 7DHC level, liver enzymes 157
 (aspartate aminotransferase (AST), alanine aminotransfer- 158
 ase (ALT), lactate dehydrogenase (LDH), alkaline phos- 159
 phatase, gamma-glutamyltransferase (GGT)), and creatine 160
 kinase (CK) activity. Statin therapy was suspended in five 161
 patients because of the side effects. 162

163 Rapid determination of 7DHC in serum was performed
 164 by the modified UV spectrophotometric method of Honda et
 165 al. [6] and was compared with the gas chromatography–
 166 mass spectrometry (GC/MS) method described by Kelly
 167 [10]. Serum samples of the patients were stored frozen at
 168 –20 °C until use (~2 weeks). A total of 200 µL serum was
 169 extracted with 1.6 mL of *c*-hexane/*i*-propanol mixture (3:1).
 170 Humatrol normal serum (Human, Magdeburg, Germany)
 171 was used as negative control. For calibration, 200 µL ali-
 172 quot of negative control serum and 100 µL of 400 mg/L

173 7DHC stock solution (c-hexane/i-propanol, 3:1) were
 174 mixed, and then 1.5 mL extracting solution was added to
 175 it. All samples were covered and centrifuged at $400\times g$ for
 176 5 min. The absorbance of clear supernatant was measured at
 177 285 nm. For reagent, blank c-hexane/i-propanol, 3:1, was
 178 used. The between-run and within-run coefficients of varia-
 179 tion were $<10\%$. The detection limit of this method is about
 180 10 mg/L (or 5 mg/L when using 400 μL of serum), and it is
 181 linear in the range of 10–400 mg/L. The t-cholesterol/7-
 182 dehydrocholesterol (7-DHC) ratio was calculated in the
 183 same unit (in milligram per liter). These 7-DHC concentra-
 184 tions were compared to those obtained by a published GC/
 185 MS method [10]. When we have compared the UV method
 186 with GC in ten samples (LKH Graz, Austria), the same
 187 samples proved to be positive although the UV method
 188 resulted in lower 7-DHC values. Serum total cholesterol
 189 level was determined by a routinely used enzymatic color-
 190 imetric assay (Modular, Roche Ltd, Mannheim, Germany).
 Q5 191 Therefore, we compared total cholesterol (cholesterol oxi-
 192 dase (CHOD)–peroxidase (POD)) results with LC-MS, and
 193 UV method of 7DHC with LC-MS (7+8DHC) in eight
 194 patients. The correlation coefficient was 0.962 between the
 195 two cholesterol methods and 0.9477 between (7+8DHC) and
 196 7DHC results. The proportion of alpha-, beta-lipoproteins,
 197 and kilomicron fractions was analyzed by agarose gel electro-
 198 phoresis (Hydragel 15, Sebia, AL Instruments, Lisses,
 199 France). Enzyme activities were determined in serum by IFCC
 200 (AST, ALT, GGT, CK) or optimized UV kinetic method
 201 (LDH) on a Modular P800 analyzer (Roche Ltd, Mannheim).
 202 Serum total and conjugated bilirubin was determined by col-
 203 orimetric assay and cholesterol by enzymatic colorimetric
 204 method (CHOD–POD) on the same analyzer.

205 Statistical comparison of cholesterol, 7DHC, and α -
 206 lipoprotein levels among the three groups was carried out
 207 by Kruskal–Wallis test. For the cholesterol and α -lipoprotein
 208 levels that showed Gaussian distribution, the Bonferroni test
 209 was applied for pairwise comparisons.

210 Results

211 The age of patients when the diagnosis was set up was in wide
 212 range (0.5–18 years); in the mild-type SLOS group ($n=4$, clinical
 213 score <20), the mean level of serum cholesterol was $2.37\pm$
 214 0.8 mmol/L and the mean of 7DHC was 0.38 ± 0.14 mmol/L
 215 (147 ± 55 mg/L). In the group of typical SLOS, diagnosis was set
 216 up earlier (age of 0.1–7 years) ($n=7$; clinical score, 20–50); the
 217 mean of serum cholesterol level was 1.47 ± 0.7 mmol/L, and
 218 7DHC was 0.53 ± 0.20 mmol/L (202 ± 77 mg/L).

219 Those patients who died as newborns (at the age of less than
 220 2 months) were enrolled into the severe SLOS group ($n=4$,
 221 clinical score >50). Their 7DHC level (0.47 ± 0.14 mmol/L;
 222 181 ± 52 mg/L) was similar to the typical SLO group with great

223 scatter, but their cholesterol level (0.66 ± 0.27 mmol/L) was
 224 significantly lower than in mild SLOS (2.37 ± 0.8 mmol/L). In
 225 spite of the limited number of patients, our data refer to the
 226 prognostic value of initial cholesterol level regarding the life
 227 expectance.

228 Clinical severity scores, genotypes, and initial lipid param-
 229 eters are listed in Table 1. Correlation between initial serum
 230 cholesterol and clinical scores is shown in Fig. 1 ($n=15$;
 231 regression line, $r=0.74$). The initial Cho/7DHC ratio showed
 232 a similar weak inverse relationship with clinical scores
 233 ($r=0.669$). Statistical evaluation of the three SLOS groups
 234 showed significant difference in the initial cholesterol levels
 235 of the mild and severe SLOS groups (Bonferroni test, $p=0.01$;
 236 Fig. 2a). A significant difference could be observed between
 237 the Cho/7DHC ratios of the groups as well (Kruskal–Wallis
 238 test, $p=0.004$; Fig. 2b).

239 Lipoprotein gel electrophoresis detected decreased per-
 240 centage of α -lipoprotein in severe SLOS ($7\pm 5\%$) compared
 241 to the age-matched control group $25.4\pm 1.6\%$ ($n=5$; age, 0–
 242 3 years) without lipid disorder or to the typical ($31.6\pm 9\%$)
 243 and mild SLOS ($33\pm 6\%$). Bonferroni test proved that the
 244 ratio of α -lipoprotein in the severe SLOS group was signifi-
 245 cantly lower than in the typical ($p=0.003$) and mild SLOS
 246 group ($p=0.004$); see Fig. 2c. It might be clinically relevant
 247 that alpha lipoproteins are hardly detectable by gel electro-
 248 phoresis in severe SLOS (Fig. 3a), while the distribution of
 249 lipoproteins is generally normal in mild cases (Fig. 3b).

250 Our findings suggest that the initial level of serum cho-
 251 lesterol fundamentally determines the severity and life ex-
 252 pectancy in SLOS, and the ratio of Cho/7DHC and α -
 253 lipoprotein has additional prognostic value.

254 Liver function was monitored in those patients who sur-
 255 vived the age of 1 year ($n=10$). LDH activity was elevated
 256 in one patient, and creatine kinase was high in another
 257 patient—both of them were treated by simvastatin. We have
 258 observed more cases of elevated AST and ALT activities in
 259 typical SLOS ($n=4/5$; age, 6 ± 5.1 years; score, >20) than in
 260 mild cases ($n=1/5$; age, 5.1 ± 1 years; score, <20) during
 261 cholesterol supplementation ($n=10$) combined with statin
 262 therapy ($n=9$). The transaminase activities were twice as
 263 much in typical SLOS (AST, 50 ± 29 U/L; ALT, 47 ± 25 U/L)
 264 than in mild type (AST, 23 ± 7 U/L; ALT, 21 ± 21 U/L).
 265 Besides the similar age in the two groups, we have to notice
 266 that the duration of therapy was longer (2.9 ± 2.6 years) in
 267 the typical group compared to the mild SLOS (0.8 ± 1 year).
 268 Although serum albumin (36.8 – 47.3 g/L), total bilirubin,
 269 and the hemostasis parameters remained in the reference
 270 range, the increase of AST and ALT in 50 % of the patients
 271 probably refers to a reversible alteration of liver function.
 272 Increased sensibility of liver in SLOS may be the conse-
 273 quence of higher DHC level. When statin treatment (sim-
 274 vastatin, atorvastatin) was suspended in the affected five
 275 patients, their liver enzyme activities returned to the

Table 1 Initial individual clinical data, genotypes, and lipid levels in patients with SLOS ($n=15$)

Patient	Sex	Age at diagnosis (years)	Age (years)	Genotype	Severity score	Clinical severity type	7DHC (mmol/L)	7DHC (mg/L)	Cho (mmol/L)	Cho (mg/L)	Cho/7DHC (mg/mg)	Lipoprotein		KM (%)
												Alpha (%)	Beta (%)	
t1.1				Reference ranges →	Mild <20, severe >50		<0.00038	<0.15	<1 year, 1.3–4.9; >1 year, 2.8–5.2	– ^a	>10,000	24–27 ^b	73–76 ^b	0 ^b
t1.2														
t1.3														
t1.4														
t1.5	F	6	10	c.452G>A, c.740 C>T	10	M	0.34	130	3.47	1343	10.30	32	68	0
t1.6	F	18	18	c.452 G>A, ?	15	M	0.22	87	2.44	944	10.90	42	58	0
t1.7	F	0.5	2	c.1295A>G; c.1328G>A	15	M	0.56	217	2.00	774	3.56	30	70	0
t1.8	M	0.6	1	c.976G>T, c.452G>A	20	M	0.41	156	1.57	608	3.89	28	72	0
t1.9	M	7	15	c.964-1G>C, c.1097G>T	25	T	0.26	102	2.77	1072	10.50	38	62	0
t1.10	F	0.1	1 ^c	c.976G>T, c.374 A>G	40	T	0.53	205	2.10	813	3.96	25	75	0
t1.11	M	0.1	1	c.452G>A, c.1295 A>G	20	T	0.71	274	1.47	569	2.07	41	59	0
t1.12	M	0.1	3	IVS8-1G>C, c.1190C>T	40	T	0.66	253	1.08	418	1.65	32	64	4
t1.13	M	0.2	8	IVS8-1G>C, c.1190C>T	50	T	0.78	302	1.40	542	1.79	28	72	0
t1.14	M	0.1	2	c.730G>A, c.976G>T	40	T	0.40	155	0.72	279	1.80	16	84	0
t1.15	M	0.1	1 ^c	c.326T>C, c.452G>A	30	T	0.33	126	0.77	298	2.36	41	59	0
t1.16	F	0.1	Newborn ^c	IVS8-1G>C homozygote	55	S	0.28	109	0.31	120	1.10	12	88	0
t1.17	F	0.1	<0.2 ^c	IVS8-1G>C, c.1190 C>T	55	S	0.45	174	0.58	224	1.29	2	98	0
t1.18	F	0.1	Newborn ^c	c.725G>A; c.452 G>A	55	S	0.56	215	0.89	344	1.60	8	92	0
t1.19	M	0.1	Newborn ^c	n.d.	55	S	0.58	224	0.86	333	1.49			0

KM kilomicrograms

^a Reference ranges for Cho <1 year, 503–1,896 mg/L; >1 year, 1,084–2,012 mg/L

^b Percent of lipoproteins in the age-matched healthy control group (0–3 years, $n=5$)

^c Died as newborn or toddler

^d Patients 8 and 9 are siblings




Fig. 1 Relation between initial cholesterol level and clinical severity in SLO syndrome ($n=15$). Initial cholesterol fundamentally determines the severity of SLOS and life expectancy

276 reference range—a typical case is demonstrated on Fig. 4.
 277 Cholesterol monotherapy was enough to improve total cho-
 278 lesterol with 0.5–1 mmol/L in three patients.

279 **Discussion**

280 The incidence of Smith–Lemli–Opitz syndrome is rang-
 281 ing from 1:20,000 to 1:60,000 [11, 17]. Our northern
 282 neighboring countries, in the Czech Republic and in the
 283 Slovak population, the incidence is even higher: from
 284 1:10,000 to 1:20,000 [2]. We could correctly identify
 285 15 patients by the rapid determination of 7DHC in serum
 286 which was performed with UV spectrophotometry that
 287 showed good correlation with the GC/MS method [6,
 288 10]. It is important to mention that UV spectrophotom-
 289 etry is an easy-to-use method but less sensitive compared
 290 to GC–MS, and there is a difference between the two
 291 methods (2–39 %) as Honda and Batta also described
 292 [5]. Our findings are in accordance with the recently
 293 emerged and accepted opinion that the initial value of
 294 serum cholesterol fundamentally determines the severity,
 295 development, and life expectancy of SLOS [25]. The
 296 cholesterol/7DHC ratio has additional prognostic value
 297 in the classification of SLOS.

298 The decreased ratio of α -lipoproteins detected in se-
 299 vere SLO compared to the other types of SLO or to the
 300 age-matched control group may be the consequence of
 301 cholesterol biosynthesis disorder. On the other hand, the
 302 impaired liver function which can be observed in typical
 303 and severe SLO because of accumulated toxic dehydro-
 304 cholesterol and other sterol metabolites may further in-
 305 hibit the synthesis of lipoproteins. As the reactivity of
 306 7DHC with oxygene radicals increased [42], a high
 307 blood level of 7DHC in severe SLO phenotype tends to





Fig. 2 a Distribution of initial serum cholesterol level in severe, typical, and mild clinical types of SLOS (box and whiskers). Bonferroni test showed significant difference between the initial cholesterol levels of mild and severe SLOS groups ($p=0.01$). b Distribution of initial serum cholesterol/7DHC ratio in severe, typical, and mild clinical types of SLOS. There was significant difference between the Cho/7DHC ratios of the patient groups (Kruskal–Wallis test: $p=0.004$). c Ratio of initial serum α -lipoprotein in severe, typical, and mild types of SLOS. Bonferroni test proved that the ratio of α -lipoprotein in severe SLOS group was significantly lower than in the typical ($p=0.003$) and mild SLOS group ($p=0.004$)

308 accelerate lipid peroxidation causing further damage of
 309 proteins and antioxidant enzymes on the surface of high-
 310 density cholesterol. Because of an extremely low
 311 α -lipoprotein level in severe SLOS, we suppose that cho-
 312 lesterol reverse transport will slow down causing a great-
 313 er extent of cholesterol deficiency. Although the number
 314 of our patients is limited and results cannot be evaluated
 315 properly statistically, our long-run clinical experiences,
 316 e.g., poor life expectancy, seem to be in accordance with
 317 this hypothesis.

318 Treatment with cholesterol with or without bile acids can
 319 improve the sterol abnormalities observed in patients with
 320 SLO syndrome [18, 34]. Introduction of statins in the treat-
 321 ment of SLO patients is based on the fact that inhibition of
 322 HMG-CoA reductase results in a decrease in the precursors

Fig. 3 **a** Alpha lipoproteins are hardly detectable in severe SLOS (patient 13, Sebia gel electrophoresis). **b** Percentage of lipoprotein fractions may be normal in mild-type SLOS (patient 1, Sebia gel electrophoresis)



such as 7DHC or 8DHC [8]. Moreover, in in vitro human fibroblast culturing in a cholesterol-deficient medium supplemented with statin, an upregulation of the DHCR7 activity was detected [24]. The phenomenon that cholesterol substitution in combination with simvastatin treatment decreases the level of the abnormally high 7DHC as well as increases the cholesterol level recently is debated. In accordance with Starck, during statin therapy, we observed significant liver function impairment in SLOS which emphasizes the vulnerability of patients with limited liver detoxication capacity, and that needs special attention in therapeutic approach [28]. We agree with Starck et al. that simvastatin treatment in SLOS cannot be considered as a safe approach in each case. When hepatotoxic effect is detected, modification of therapy (e.g., cholesterol supplementation without statin) may be considered which can increase the cholesterol

and reestablish the liver function [28]. According to Haas' opinion, the mechanism of these therapeutic approaches is different: the level of cholesterol can be elevated by supplementation, and statins may reduce the DHC level which increases the cholesterol/7DHC ratio [5].

Determination of the lipid parameters in different categories of SLO syndrome is essential, because initial lipid levels have prognostic value. Monitoring of lipids and liver function helps to evaluate the efficiency of cholesterol supplementation and detects the side effect of statin therapy.

Although it is generally accepted that SLO syndrome has wide genetic variability, and genotype and phenotype are not in close connection, the blood level of cholesterol precursor 7DHC and clinical severity depend on mutation types [3, 5, 12, 41]. Therefore, biochemical markers still have a significant role besides the phenotype in setting up the diagnosis, prognosis, and later in the follow-up.

The early diagnosis is the precondition of the effective therapy although the individual results are different with strong limitation. Prenatal diagnosis with biochemical methods and molecular genetic test are available if the disease-causing mutation(s) in the family is known [14, 15, 20, 23]. Traditional biochemical screening of the substrates (7DHC and 8DHC) in serum is not reliable for detection of carrier status because there is an overlap between the ranges of serum concentrations of cholesterol and 7DHC in carriers and noncarriers; therefore, the biochemical testing of fibroblasts (7DHC) or the molecular genetic analysis of the disease-causing mutations in the family is recommended [24].

Based on the identification of family-specific gene mutations in the affected families, the prenatal genetic examination was introduced in Hungary in 2009. A web page has been set up to provide detailed information about the diagnostic and therapeutic possibilities (www.smithlempiopitz.hu).




Fig. 4 A typical case of statin intolerance in patient 8 during cholesterol supplementation combined with simvastatin therapy. When the liver function and clinical condition were impaired, statin therapy was finished, and transaminase activities returned to the normal range. The arrow shows the duration of statin therapy

Q10 323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339

340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376

Acknowledgments The authors are grateful to László Szabó (Markusovszky County Hospital, Szombathely, Hungary), Ákos Szekeres (Semmelweis University, Budapest), and Ilona György for discussions, and Nóra Holhós, Erika Dzsudzsák, Emilia Szabó, and Szilvia Fülöp for their technical support. The work was supported by the TÁMOP 4.2.1./B-09/1/KONV-2010-0007 project. The project is implemented through the New Hungary Development Plan, co-financed by the European Social Fund.

Conflict of interest The authors stated that there are no conflicts of interest regarding the publication of this article. Research granting played no role in the study design, interpretation of data, or writing of the report.

References

1. Balogh I, Koczok K, Szabó GP, Török O, Hadzsiev K, Csábi G, Balogh L, Dzsudzsák E, Ajzner É, Szabó L, Csákváry V, Oláh AV (2012) Mutational spectrum of Smith–Lemli–Opitz syndrome patients in Hungary. *Mol Syndromol* 3:215–222

2. Bzduch V (2000) Incidence of Smith–Lemli–Opitz syndrome in Slovakia. *Am J Med Genet* 90:260

3. Cunnif C, Kratz LE, Moser A, Natowicz MR, Kelley RI (1997) Clinical and biochemical spectrum of patients with RSH/Smith–Lemli–Opitz syndrome and abnormal cholesterol metabolism. *Am J Med Genet* 68:263–269

4. Fitzky BU, Witsch-Baumgartner M, Erdel M, Lee JN, Paik YK, Glossmann H, Utermann G, Moebius FF (1998) Mutations in the delta 7-sterol reductase gene in patient with the Smith–Lemli–Opitz syndrome. *Proc Natl Acad Sci USA* 95:8181–8186

5. Haas D, Garbade SF, Vohwinkel C, Muschol N, Trefz FK, Penzien JM, Zschocke J, Hoffmann GF, Burgard P (2007) Effect of cholesterol and simvastatin treatment in patient with Smith–Lemli–Opitz syndrome. *J Inherit Met Dis* 30:375–387

6. Honda A, Batta AK, Salen G, Tint GS, Chen T, Shefer S (1997) Screening for abnormal cholesterol biosynthesis in the Smith–Lemli–Opitz syndrome. *Am J Med Genet* 68:288–293

7. Irons M, Elias ER, Salen G, Tint GS, Batta AK (1993) Defective cholesterol biosynthesis in Smith–Lemli–Opitz syndrome. *Lancet* 341:1414

8. Jira PE, Wevers RA, de Jong J, Rubio-Gozalbo E, Janssen-Zijlstra FS, van Heyst AF, Sengers RC, Smeitink JA (2000) Simvastatin. A new therapeutic approach for Smith–Lemli–Opitz syndrome. *J Lipid Res* 41:1339–1346

9. Jira PE, Wanders RJ, Smeitink JA, De Jong J, Wevers RA, Oostheim W, Tuerlings JH, Hennekam RC, Sengers RC, Waterham HR (2001) Novel mutations in the 7-dehydrocholesterol reductase gene of 13 patients with Smith–Lemli–Opitz syndrome. *Ann Hum Genet* 65:229–236

10. Kelley RI (1995) Diagnosis of Smith–Lemli–Opitz syndrome by gas chromatography/mass spectrophotometry of 7-dehydrocholesterol in plasma, amniotic fluid and cultured skin fibroblasts. *Clin Chim Acta* 236:45–58

11. Kelley RI (1997) Editorial: a new face for an old syndrome. *Am J Med Genet* 65:251–256

12. Kelley RI, Hennekam RCM (2001) Smith–Lemli–Opitz syndrome. In: Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Kinzler KW, Vogelstein B (eds) *The metabolic and molecular bases of inherited disease*. Mc Graw-Hill, New York, pp 6183–6201

13. Kelley RI, Hennekam RCM (2000) The Smith–Lemli–Opitz syndrome. *J Med Genet* 37:321–335

14. Kratz LE, Kelley RI (1999) Prenatal diagnosis of the RSH/Smith–Lemli–Opitz syndrome. *Am J Med Genet* 82:376–381

15. Mills K, Mandel H, Montemagno R, Soothill P, Gershoni-Baruch R, Clayton PT (1996) First trimester prenatal diagnosis of Smith–Lemli–Opitz syndrome (7-dehydrocholesterol reductase deficiency). *Pediatr Res* 39:816–819

16. Moebius FF, Fitzky BU, Lee JN, Paik YK, Glossmann H (1998) Molecular cloning and expression of the human Δ7-sterol reductase. *Proc Natl Acad Sci* 95:1899–1902

17. Nowaczyk MJM, McCaughey D, Whelan DT, Porter FD (2001) Incidence of Smith–Lemli–Opitz syndrome in Ontario, Canada. *Am J Med Genet* 102:18–20

18. Nwokoro NA, Mulvihill JJ (1997) Cholesterol and bile acid replacement therapy in children and adults with Smith–Lemli–Opitz (SLO/RSH) syndrome. *Am J Med Genet* 68:315–321

19. Opitz JM (1999) RHD (so-called Smith–Lemli–Opitz) syndrome. *Curr Opin Pediatr* 11:353–362

20. Palomaki GE, Bradley LA, Knight GJ, Craig WY, Haddow JE (2002) Assigning risk for Smith–Lemli–Opitz syndrome as part of 2nd trimester screening for Down’s syndrome. *J Med Screen* 9:43–44

21. Porter FD (2008) Smith–Lemli–Opitz syndrome: pathogenesis, diagnosis and management. *Eur J Human Genet* 16:535–541

22. Ryan AK, Bartlett K, Clayton P, Eaton S, Mills L, Donnai D, Winter RM, Burn J (1998) Smith–Lemli–Opitz syndrome: a variable clinical and biochemical phenotype. *J Med Genet* 35:558–565

23. Sharp P, Haan E, Fletcher JM, Khong TY, Carey WF (1997) First-trimester diagnosis of Smith–Lemli–Opitz syndrome. *Prenat Diagn* 17:355–361

24. Shefer S, Salen G, Honda A, Batta A, Hauser S, Tint GS, Honda M, Chen T, Holick MF, Nguyen LB (1997) Rapid identification of Smith–Lemli–Opitz syndrome homozygotes and heterozygotes (carriers) by measurement of deficient 7-dehydrocholesterol-delta 7-reductase activity in fibroblasts. *Metabolism* 46:844–850

25. Sikora DM, Ruggiero M, Petit-Kekel K, Merckens LS, Connor WE, Steiner RD (2004) Cholesterol supplementation does not improve developmental progress in Smith–Lemli–Opitz syndrome. *J Pediatr* 144:783–791

26. Smith DW, Lemli L, Opitz JM (1964) A newly recognized syndrome of multiple congenital anomalies. *J Pediatr* 64:210–217

27. Stanek AJ, Ciara E, Malunovic EM, Korniszewski L, Piekutowska-Abramczuk D, Popowska E, Krajewska-Walasek M (2008) Mild Smith–Lemli–Opitz syndrome. *Eur J Med Gen* 51:124–140

28. Starck L, Lovgren-Sandblom A, Bjorkhem I (2002) Simvastatin treatment in the SLO syndrome: a safe approach? *Am J Med Genet* 113:183–189

29. Starck L, Lovgren-Sandblom A, Bjorkhem I (2002) Cholesterol treatment forever? The first Scandinavian trial of cholesterol supplementation in the cholesterol-synthesis defect Smith–Lemli–Opitz syndrome. *J Intern Med* 252:314–321

30. Szabó L, Erwa W, Soroncz M (2002) Metabolic background of the Smith–Lemli–Opitz (genito-facio-digital) syndrome. *Gyermekgyógyászat* 53:695–702

31. Szabó GP, Oláh AV, Kozák L, Balogh E, Nagy A, Blahakova I, Oláh É (2010) A patient with Smith–Lemli–Opitz syndrome: novel mutation of the *DHCR7* gene and effects of therapy with simvastatin and cholesterol supplement. *Eur J Pediatr* 169:121–123

32. Tierney E, Conley SK, Goodwin H, Porter FD (2010) Analysis of short-term behavioral effects of dietary cholesterol supplementation in Smith–Lemli–Opitz syndrome. *Am J Med Genet* 152A:91–95

33. Tint GS, Salen G, Batta AK, Shefer S, Irons M, Elias ER, Abuelo DN, Johnson VP, Lambert M, Lutz R (1995) Correlation of severity and outcome with plasma sterol levels in variants of the Smith–Lemli–Opitz syndrome. *J Pediatr* 127:82–87

- 506 34. Ullrich K, Koch HG, Meschede D (1996) Smith–Lemli–Opitz
507 syndrome; treatment with cholesterol and bile acids. *Neuropediat-*
508 *rics* 27:111–112
- 509 35. Wassif CA, Maslen C, Kachiele-Linjewile S (1998) Mutation in
510 the human sterol delta-7 reductase gene at 11q12-13 cause Smith–
511 Lemli–Opitz syndrome. *Am J Hum Genet* 63:55–62
- 512 36. Waterham HR, Wijburg FA, Hennekam RC (1998) Smith–Lemli–
513 Opitz syndrome is caused by mutations in the 7-dehydrocholesterol
514 reductase gene. *Am J Hum Genet* 63:329–338
- 515 37. Waterham HR, Oostheim W, Romeijn GJ (2000) Incidence and
516 molecular mechanism of aberrant splicing owing to a G→C splice
517 acceptor site mutation causing Smith–Lemli–Opitz syndrome. *J*
518 *Med Genet* 37:387–389
- 519 38. Waterham HR (2002) Inherited disorders of cholesterol biosynthe-
520 sis. *Clin Genet* 61:393–403
- 521 39. Wittsch-Baumgartner M, Loffler J, Utermann G (2001) Mutations
522 in the human DHCR7 gene. *Hum Mutat* 17:172–182
- 523 40. Wittsch-Baumgartner M, Loffler J, Menzel HJ (2001) Frequency
524 gradients of DHCR7 mutations in patient with Smith–Lemli–Opitz
525 syndrome in Europe: evidence for different origins of common
526 mutations. *Eur J Hum Genet* 9:45–50
- 527 41. Wittsch-Baumgartner M, Fitzky BU, Ogorelkova M, Kraft HG,
528 Moebius FF, Glossmann H, Seedorf U, Gillessen-Kaesbach G,
529 Hoffmann GF, Clayton P, Kelley RI, Utermann G (2000) Muta-
530 tional spectrum in the 7-sterol reductase gene and genotype-
531 phenotype correlation in 84 patients with Smith–Lemli–Opitz syn-
532 drome. *Am J Hum Genet* 66:402–412
- 533 42. Xu L, Davis TA, Porter NA (2009) Rate constants for peroxidation
534 of polyunsaturated fatty acids and sterols in solution and in lip-
535 osomes. *J Am Chem Soc* 131:13037–13044

UNCORRECTED PROOF

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Please check if the affiliations are correctly presented.
- Q2. This phrase was changed to "on the basis of." Please check if appropriate.
- Q3. Please check if the keywords were correctly captured.
- Q4. The abbreviations "ALP", "t-CHO", "HDL-C", and "KM" were used only once in the text and were deleted. Please consider deleting the abbreviations in the list as well.
- Q5. The acronym "CHOD-POD" was expanded as "cholesterol oxidase–peroxidase". Please check if appropriate.
- Q6. Please check if Table 4 entries are correctly presented.
- Q7. Please check if the changes in the sentence "Our findings suggest that the initial level of serum cholesterol..." are appropriate.
- Q8. This sentence was changed to "When the liver function and clinical condition were impaired..." Please check if appropriate.
- Q9. Please check if the changes in the sentence "Although the number of our patients is limited..." are appropriate.
- Q10. Please check if the changes in the sentence "Moreover, in in vitro human fibroblast culturing..." are appropriate.
- Q11. References 9, 16, 19, 29, 33, 37 and 38 were not cited anywhere in the text. Please provide citations. Alternatively, delete the items from the list.