

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

#### Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

# Metadata of the article that will be visualized in OnlineFirst

ArticleTitle	Killing Rates for Caspofungin Against <i>Candida albicans</i> After Brief and Continuous Caspofungin Exposure in the Presence and Absence of Serum	
Article Sub-Title		
Article CopyRight	Springer Science+Business Media Dordrecht (This will be the copyright line in the final PDF)	
Journal Name	Mycopathologia	
Corresponding Author	Family Name	<b>Majoros</b>
	Particle	
	Given Name	<b>László</b>
	Suffix	
	Division	Department of Medical Microbiology
	Organization	University of Debrecen
	Address	Nagyerdei krt. 98, Debrecen, 4032, Hungary
	Email	major@med.unideb.hu
Author	Family Name	<b>Kovács</b>
	Particle	
	Given Name	<b>Renátó</b>
	Suffix	
	Division	Department of Medical Microbiology
	Organization	University of Debrecen
	Address	Nagyerdei krt. 98, Debrecen, 4032, Hungary
	Email	
Author	Family Name	<b>Gesztelyi</b>
	Particle	
	Given Name	<b>Rudolf</b>
	Suffix	
	Division	Department of Pharmacology and Pharmacodynamics
	Organization	University of Debrecen
	Address	Debrecen, Hungary
	Email	
Author	Family Name	<b>Perlin</b>
	Particle	
	Given Name	<b>David S.</b>
	Suffix	
	Division	Public Health Research Institute
	Organization	New Jersey Medical School-Rutgers
	Address	Newark, NJ, USA
	Email	
Author	Family Name	<b>Kardos</b>
	Particle	
	Given Name	<b>Gábor</b>
	Suffix	

Division Department of Medical Microbiology  
Organization University of Debrecen  
Address Nagyerdei krt. 98, Debrecen, 4032, Hungary  
Email

---

Author Family Name **Domán**  
Particle  
Given Name **Marianna**  
Suffix  
Division Department of Medical Microbiology  
Organization University of Debrecen  
Address Nagyerdei krt. 98, Debrecen, 4032, Hungary  
Email

---

Author Family Name **Berényi**  
Particle  
Given Name **Réka**  
Suffix  
Division Department of Medical Microbiology  
Organization University of Debrecen  
Address Nagyerdei krt. 98, Debrecen, 4032, Hungary  
Email

---

Schedule Received 12 June 2014  
Revised  
Accepted 1 August 2014

---

Abstract It was previously demonstrated that brief ( $\leq 1$  h) exposures to echinocandins are as effective to kill *Candida albicans* cells as continuous 24-h exposure. However, killing rates after continuous and short (1 h) echinocandin exposures to *C. albicans* have not yet been evaluated in RPMI-1640 with and without 50 % serum. We evaluated four echinocandin susceptible *C. albicans* bloodstream isolates, ATCC 10231 type strain and an echinocandin-resistant isolate (DPL20, FKS F645P). Caspofungin MICs, time-kill and postantifungal effect (PAFE) tests were performed in RPMI-1640 with and without 50 % serum. Killing rates (*k* values) in time-kill and PAFE experiments were determined for each strain and concentration. In time-kill experiments, colony count decreases were isolate- and concentration-dependent at 0.25, 1, 4, 8, 16 and 32 mg/L in RPMI-1640, but concentration-independent at 1, 4, 8, 16 and 32 mg/L in 50 % serum. One-hour caspofungin exposure at 4, 16 and 32 mg/L resulted in CFU decreases comparable with the results obtained in time-kill experiments in RPMI-1640, but 50 % serum at 4, 16 and 32 mg/L allowed growth of all isolates (*k* values were negative) ( $P < 0.05$ – $0.001$ ). PAFE in 50 % serum decreased markedly at 4, 16 and 32 mg/L. Killing rates remained high and concentration-independent in 50 % serum in case of continuous but not in case of brief caspofungin exposure. As only a short growth inhibition without killing was observed in 50 % serum, clinical relevance of caspofungin PAFE in vivo is questionable.

---

Keywords (separated by '-') Echinocandins - Serum-based susceptibility testing - Postantifungal effect

---

Footnote Information

---

# Killing Rates for Caspofungin Against *Candida albicans* After Brief and Continuous Caspofungin Exposure in the Presence and Absence of Serum

Renátó Kovács · Rudolf Gesztelyi · David S. Perlin ·  
Gábor Kardos · Marianna Domán · Réka Berényi ·  
László Majoros

Received: 12 June 2014 / Accepted: 1 August 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** It was previously demonstrated that brief ( $\leq 1$  h) exposures to echinocandins are as effective to kill *Candida albicans* cells as continuous 24-h exposure. However, killing rates after continuous and short (1 h) echinocandin exposures to *C. albicans* have not yet been evaluated in RPMI-1640 with and without 50 % serum. We evaluated four echinocandin susceptible *C. albicans* bloodstream isolates, ATCC 10231 type strain and an echinocandin-resistant isolate (DPL20, FKS F645P). Caspofungin MICs, time-kill and postantifungal effect (PAFE) tests were performed in RPMI-1640 with and without 50 % serum. Killing rates ( $k$  values) in time-kill and PAFE experiments were determined for each strain and concentration. In time-kill experiments, colony count decreases were isolate- and concentration-dependent at 0.25, 1, 4, 8, 16 and 32 mg/L in RPMI-1640, but concentration-independent at 1, 4, 8, 16 and 32 mg/L in 50 % serum. One-hour caspofungin exposure at 4, 16 and 32 mg/L

resulted in CFU decreases comparable with the results obtained in time-kill experiments in RPMI-1640, but 50 % serum at 4, 16 and 32 mg/L allowed growth of all isolates ( $k$  values were negative) ( $P < 0.05$ – $0.001$ ). PAFE in 50 % serum decreased markedly at 4, 16 and 32 mg/L. Killing rates remained high and concentration-independent in 50 % serum in case of continuous but not in case of brief caspofungin exposure. As only a short growth inhibition without killing was observed in 50 % serum, clinical relevance of caspofungin PAFE in vivo is questionable.

**Keywords** Echinocandins · Serum-based susceptibility testing · Postantifungal effect

## Introduction

Echinocandins (caspofungin, micafungin and anidulafungin) are reported to exhibit concentration-dependent fungicidal or fungistatic activity against the majority of *Candida* species. Additionally, echinocandins exert prolonged postantifungal effect (PAFE) after short (1 h) echinocandin exposure against many *Candida* species, including *C. parapsilosis* [1–4]. Recent in vitro findings with RPMI-1640 suggest that a very short ( $\leq 1$  h) exposure to caspofungin kills *Candida* cells as effectively as a continuous 24-h exposure [5]. As echinocandins are highly protein-bound ( $\geq 96.5$  %) agents (i.e., serum fundamentally

R. Kovács · G. Kardos · M. Domán · R. Berényi ·  
L. Majoros (✉)  
Department of Medical Microbiology, University of  
Debrecen, Nagyerdei krt. 98, 4032 Debrecen, Hungary  
e-mail: major@med.unideb.hu

R. Gesztelyi  
Department of Pharmacology and Pharmacodynamics,  
University of Debrecen, Debrecen, Hungary

D. S. Perlin  
Public Health Research Institute, New Jersey Medical  
School-Rutgers, Newark, NJ, USA

56 influences killing activities of echinocandins) [6–8],  
 57 we compared killing rates produced by short (1 h) and  
 58 continuous (24 h) caspofungin exposure as well as  
 59 PAFE in RPMI-1640 and 50 % serum against *C.*  
 60 *albicans*.

## 61 Materials and Methods

### 62 Isolates

63 All *C. albicans* originated from blood samples  
 64 (Table 1) and were identified as described previously  
 65 [9]. ATCC 10231 and an echinocandin-resistant  
 66 isolate (DPL20, FKS F645P) were also included in  
 67 the study.

### 68 MIC Determination

69 Caspofungin (Sigma, Budapest, Hungary) MICs in  
 70 RPMI-1640 with and without 50 % human serum  
 71 (from a human male, type AB, Sigma, Budapest,  
 72 Hungary) were determined using the CLSI broth  
 73 macrodilution method [6–8, 10]. Caspofungin final  
 74 concentration ranged between 0.015 and 32 mg/L.  
 75 MICs were read after 24 h using the partial inhibition  
 76 criterion [10].

### 77 Postantifungal Effect and Time-Kill Curves

78 PAFE was measured in both media simultaneously. As  
 79 in our preliminary experiments 5-, 10- and 30-min  
 80 exposures to 0.5, 1 or 2 mg/L caspofungin did  
 81 not produce measurable PAFEs in 50 % serum

**Fig. 1** Time-kill plots of caspofungin against four *Candida* ▶  
*albicans* isolates (averages ± standard deviation) in RPMI-  
 1640 (a) and 50 % serum (d) against ATCC 10231 type strain in  
 RPMI-1640 (b) and 50 % serum (e), and against the echino-  
 candin-resistant *C. albicans* DPL20 in RPMI-1640 (c) and 50 %  
 serum (f). Clinical isolates and type strain were exposed to 0.25,  
 1, 4, 8, 16 and 32 mg/L, while DPL20 isolate was exposed to 4,  
 8, 16 and 32 mg/L of caspofungin for 24 h (continuous  
 caspofungin exposure)







(1–16 × MIC in 50 % serum), we used caspofungin 82  
 at 4, 16 and 32 mg/L concentrations with a 60-min 83  
 exposure time (brief caspofungin exposure) [3–5]. As 84  
 the maximum administrable daily 150–200 mg ca- 85  
 spofungin doses produce 30.4–40.6 mg/L geometric 86  
 mean of peak concentrations in humans [11], the 87  
 highest caspofungin concentration used in this study 88  
 was 32 mg/L [6]. 89

The starting inocula in PAFE experiments were 90  
 1–5 × 10<sup>5</sup> cells/ml [3–5]. After 1 h, the cells were 91  
 collected by centrifugation at 1.500×g for 10 min and 92  
 were washed three times with sterile saline, resus- 93  
 pended in 10 ml drug-free warm RPMI-1640 with and 94  
 without 50 % human serum. Samples (100 µl) were 95  
 removed at 0, 4, 8, 12 and 24 h, serially diluted 96  
 tenfold, plated (4 × 30 µl) onto Sabouraud dextrose 97  
 agar and incubated at 35 °C for 48 h [3–5]. 98

For time-kill assays (continuous 24-h caspofungin 99  
 exposure), test solutions were not centrifuged or 100  
 washed. We used 0.25, 1, 4, 8, 16 and 32 mg/L 101  
 caspofungin. In case of isolate DPL20, we used 4, 8, 102  
 16 and 32 mg/L caspofungin. Otherwise, the method 103  
 was the same as described for PAFE [3–5]. All 104  
 experiments were performed at least twice, and means 105  
 of data are presented. 106

**Table 1** *Candida albicans* isolates, MICs of caspofungin and the effect caspofungin in time-kill studies in RPMI-1640 (RPMI) and RPMI-1640 supplemented with 50 % serum (50 % serum)


Isolates	MIC (mg/L)	Effect in time-kill studies at concentrations shown (mg/L)		
		RPMI	50 % Serum	50 % Serum
183	0.03	0.25	≥0.03 fungistatic	≥0.25 fungistatic
3666	0.03	0.125	≥0.03 fungistatic	≥0.25 fungistatic
12132	0.015	0.25	≥0.03 fungistatic	≥0.25 fungistatic
10920	0.03	0.125	≥0.03 fungistatic	≥0.25 fungistatic
ATCC 10231	0.03	0.5	≥0.03 fungistatic	≥0.5 fungistatic
DPL20	4	>32	≥16 fungistatic	No effect



## 107 Data Analysis

108 Killing kinetics was analyzed mathematically, as  
 109 described previously [6, 12]. An exponential equation  
 110 was fitted to the mean data at each time point:  
 111  $N_t = N_0 \times e^{-kt}$ , where  $N_t$  is the number of viable  
 112 yeasts at time  $t$ ,  $N_0$  is the number of viable yeasts in the

113 initial inoculum,  $k$  is the killing rate and  $t$  is the  
 114 incubation time. Negative  $k$  values indicate growth,  
 115 and positive  $k$  values indicate killing. The goodness of  
 116 fit for each isolate was assessed by the  $r^2$  value ( $>0.8$ )  
 117 [6, 12]. PAFE was defined as the difference between  
 118 the time required for control and test isolates to grow 1  
 119  $\log_{10}$  following drug removal [3–5].



**Fig. 2** Postantifungal effect curves of caspofungin against *Candida albicans* isolate 183 in RPMI-1640 (a) and 50 % serum (c), and against *C. albicans* isolate 3666 in RPMI-1640 (b) and

50 % serum (d). Isolates were exposed to 4, 16 and 32 mg/L of caspofungin for 1 h (brief caspofungin exposure)

120 One-way ANOVA with Tukey's posttesting was  
 121 used to analyze differences in killing rates by  
 122 concentrations in either RPMI-1640 or 50 % serum  
 123 [6]. Paired *T* test was used to compare the effect of the  
 124 medium as well as the killing and growth rates in time-  
 125 kill and PAFE experiments at the same drug  
 126 concentration.

## Results

### Susceptibility

MIC values are presented in Table 1. Paradoxical  
 growth was not observed. Clinical isolates and 10231  
 ATCC strain were susceptible to caspofungin

127

128

129

130

131

**Table 2** Postantifungal effect (PAFE) of caspofungin against *Candida albicans* isolates at 4, 16 and 32 mg/L in RPMI-1640 (RPMI) and RPMI-1640-50 % serum (50 % serum)







Isolate number	Medium	PAFE in hours		
		4 (mg/L)	16 (mg/L)	32 (mg/L)
183	RPMI-1640	>19.34	>19.34	14.81
	50 % Serum	2.27	10.14	>18.79
3666	RPMI-1640	7.40	>19.23	>19.23
	50 % Serum	0	0.79	1.01
10920	RPMI-1640	>19.05	>19.05	>19.05
	50 % Serum	0	0.24	0.24
12132	RPMI-1640	14.21	13.84	7.65
	50 % Serum	0.60	0.95	3.46
ATCC 10231	RPMI-1640	4.89	>19.88	>19.88
	50 % Serum	0.09	0.34	1.39
DPL20	RPMI-1640	0	0	0
	50 % Serum	0	0	0

**Table 3** Maximum log changes in log CFU/mL compared to starting inoculum in time-kill and postantifungal (PAFE) studies in RPMI-1640 and RPMI-1640-50 % serum (50 % serum)

Isolate number	Media	Maximum log decreases in CFU in time-killing and PAFE experiments at the indicated caspofungin concentration					
		4 mg/L		16 mg/L		32 mg/L	
		Time kill	PAFE	Time kill	PAFE	Time kill	PAFE
183	RPMI-1640	-1.1	-0.53	-1.18	-0.42	-0.75	-0.48
	50 % serum	-1.06	+0.37	-0.9	-0.08	-0.9	-0.04
3666	RPMI-1640	-1.38	-0.66	-0.54	-1.6	-0.34	-2.08
	50 % serum	-1.40	+0.13	-1.12	+0.12	-2.68	-0.12
10920	RPMI-1640	-1.23	-1.23	-1.22	-1.48	-0.69	-1.48
	50 % serum	-0.69	+0.49	-1.46	+0.38	-1.43	+0.64
12132	RPMI-1640	-0.44	-0.30	-0.19	-0.27	-0.23	-0.62
	50 % serum	-1.56	+0.67	-1.71	+0.66	-1.86	+0.27
ATCC 10231	RPMI-1640	-1.56	-1.48	-0.55	-2.08	-1.65	-1.78
	50 % serum	-1.18	+0.26	-1.38	+0.09	-1.95	+0.08
DPL20	RPMI-1640	-0.39	+0.61	-1.19	+0.78	-1.00	+0.37
	50 % serum	+0.81	+0.84	+0.69	+0.57	-0.05	+0.71



Author Proof



◀ **Fig. 3** Killing rates of caspofungin and the corresponding fitted regression lines (*dashed lines*) in time-kill experiments against four *C. albicans* isolates (averages  $\pm$  standard deviation) in RPMI-1640 (**a**) and 50 % serum (**d**), against *C. albicans* DPL20 in RPMI-1640 (**b**) and 50 % serum (**e**), and against ATCC 10231 type strain in RPMI-1640 (**c**) and 50 % serum (**f**). Positive and negative *k* values indicate the decrease and increase, respectively, in viable cell numbers

132 according to the revised CLSI break points in RPMI-  
133 1640 [13]. As expected, the DPL20 isolate with a  
134 prominent *fk*s mutation was resistant to caspofungin  
135 [13]. MIC values were 4- to 16-fold higher in the  
136 presence of 50 % serum.

### 137 Time-Kill Experiments

138 In time-kill experiments, caspofungin was fungistatic  
139 at  $\geq 1-2 \times$  MIC in both media against the clinical  
140 isolates as well as against the 10231 ATCC strain  
141 ( $<99.9$  % reduction in viable cell count compared to  
142 the starting inoculum) (Table 1; Fig. 1a, b, d, e).

143 Against the resistant strain DPL20, caspofungin  
144 produced a weak fungistatic effect in RPMI-1640  
145 (Fig. 1C). In 50 % serum, the killing curves were  
146 generally similar to control; at 32 mg/L, a negligible  
147 reduction was observed after 4 h, but later the killing  
148 curves again became similar to control (Fig. 1f).

### 149 Postantifungal Effect

150 The time required for control (drug-free) isolates to  
151 grow 1 log<sub>10</sub> was similar in both media (4.12–4.95 h in  
152 RPMI-1640 and 4.69–5.17 h in 50 % serum). PAFE  
153 plots for isolates 183 and 3666 in RPMI-1640 and in  
154 50 % serum are shown in Fig. 2. In RPMI-1640,  
155 clinical isolates and the ATCC 10231 strain showed  
156 the inhibition of re-growth at 4, 16 and 32 mg/L for  
157 4.89 to  $>19.34$ , 13.84 to  $>19.88$  and 7.65 to  $>19.88$  h,  
158 respectively (Table 2). PAFE in 50 % serum  
159 decreased markedly at 4, 16 and 32 mg/L concentra-  
160 tions (Table 2; Fig. 2). Most isolates showed growth  
161 in 50 % serum; colony count decreases occurred only  
162 in cases of isolates 183 at 16 and 32 mg/L (Fig. 2c)  
163 and 3666 at 32 mg/L (Fig. 2d) and only after 4 h and  
164 were negligible (Table 3). In case of DPL20, isolate  
165 PAFE was never observed regardless of media  
166 (Table 2).

### Comparison of Colony Count Changes in Time-Kill and Postantifungal Effect Experiments 167 168

169 Maximum colony count changes compared to the  
170 starting inocula in time-kill and PAFE experiments at  
171 4, 16 and 32 mg/L are presented in Table 3.

172 Comparing the different media at the same con-  
173 centrations in killing experiments, the CFU decrease  
174 was generally higher in 50 % serum than in RPMI-  
175 1640 (Table 3). Contrastingly, the CFU decrease in  
176 PAFE experiments was significantly higher in RPMI-  
177 1640 than in 50 % serum with all isolates and  
178 concentrations ( $P < 0.05-0.001$ ) (Table 3).

179 Comparing the colony counts reductions at the  
180 same concentrations in PAFE and time-kill experi-  
181 ments, we noticed comparable or sometimes higher  
182 reductions (in cases of 10920 and 3666 isolates) in  
183 PAFEs than that seen with continuous 24-h exposure  
184 in RPMI-1640 (Table 3). However, 50 % serum  
185 significantly decreased the PAFE killing for all tested  
186 isolates ( $P < 0.01-0.001$ ) (Table 3; Fig. 4a–e).

### Killing Rates in Time-Kill Experiments 187

188 In time-kill experiments, killing activity of caspofun-  
189 gin was significantly weaker at 16–32 mg/L than at  
190 0.25, 1, 4 and 8 mg/L ( $P < 0.05-0.001$ ) in RPMI-  
191 1640 (mini-paradoxical effect) (Fig. 3a). However,  
192 killing rates at 1–32 mg/L were concentration-inde-  
193 pendent in 50 % serum against the susceptible isolates  
194 (Fig. 3d). Similar effect was noticed in case of the  
195 strain ATCC 10231 (Fig. 3c, f). In case of isolate,  
196 DPL20 *k* values were negative (indicating the growth  
197 instead of killing) regardless of media (Fig. 3b, e).






### Killing Rates in Postantifungal Effect Experiments 198

199 In PAFE experiments, killing rates for clinical isolates  
200 and the ATCC strain in RPMI-1640 were isolate- and  
201 concentration-dependent (*k* values from  $-0.111$  to  
202  $+1.019$  1/h), while in 50 % serum, the *k* values  
203 showed markedly narrower range (from  $-0.017$  to  $-$   
204  $0.185$  1/h) (Fig. 4a–e).

### Discussion 205

206 This study is the first in which killing rates in short and  
207 continuous caspofungin exposures to *C. albicans* were

Author Proof



**Fig. 4** Killing rates of caspofungin and the corresponding fitted regression lines (*dashed lines*) in PAFE experiments against 183 (a), 10920 (b), 3666 (d) and 12132 (e) *C. albicans* isolates in RPMI-1640 and 50 % serum, and against *C. albicans* ATCC 10231 type strain (c) in RPMI-1640 and 50 % serum. Positive and negative  $k$  values indicate the decrease and increase, respectively, in viable cell numbers. For panels C and D, the scale of the y axis was broken for better visualization of regression lines

208 compared head to head in RPMI-1640 with and  
209 without 50 % serum. In agreement with previous  
210 results [3–5], the CFU decreases in time-kill and  
211 PAFE experiments were similar in RPMI-1640;  
212 however, at 16 and 32 mg/L, killing rates decreased.  
213 Decreased killing rates in time-kill experiments at 16  
214 and 32 mg/L can be explained by the adaptive and  
215 compensatory response to high caspofungin concen-  
216 trations in the fungal cells that limit killing effect and  
217 allow for the growth [14]. Addition of 50 % serum  
218 significantly decreased killing rates at 4, 16 and  
219 32 mg/L in the PAFE experiments as compared to  
220 RPMI-1640, while the killing rates in time-kill  
221 experiments (continuous exposure) at  $\geq 1$  mg/L con-  
222 centrations remained high and concentration-indepen-  
223 dent (i.e., killing rate reached its maximum at 1 mg/L).  
224 These findings are in accordance with our previous  
225 results, where killing rates of *C. krusei* and *C.*  
226 *inconspicua* did not differ significantly in 50 % serum  
227 at effective concentrations [6]. Moreover, 1-h expo-  
228 sure of *C. albicans* to 0.5, 1 and 2 mg/L of caspofungin  
229 in 50 % serum is not long enough to produce any  
230 growth inhibition, as opposed to what found with  
231 RPMI-1640 alone.

232 Louie et al. [15] demonstrated that tissues serve as  
233 drug reservoirs from which the drug is released slowly,  
234 explaining that serum caspofungin half-life tripled  
235 when both serum and tissues half-life were taken into  
236 account in the terminal half-life calculation. They  
237 concluded that the primary tissue reservoir rather than  
238 PAFE was responsible for the excellent in vivo  
239 activity of caspofungin [15]. Their results are in line  
240 with the present study, as 1-h exposure to caspofungin  
241 produced negative  $k$  values (growth) and significantly  
242 decreased PAFEs in 50 % serum.

243 PAFE is frequently regarded as a contributor to  
244 clinical efficacy of echinocandins [1–4, 13]. It is  
245 defined as prolonged growth inhibition following  
246 limited (generally 1 h) in vitro drug exposure; how-  
247 ever, it must be noted that this is not equal to killing, as

248 slower growth may also be regarded as prominent  
249 PAFE. While killing may directly lead to eradication,  
250 growth, even slower growth, of fungi still carries a risk  
251 of persistent infection and fungal re-growth. These  
252 facts should be taken into consideration when trans-  
253 lating PAFE as a contributor to clinical efficacy.  
254 Present results strongly suggest that PAFE is lost in the  
255 presence of 50 % serum, even though marked PAFE is  
256 detected in RPMI-1640 after 5-min exposure [5]. The  
257 negligible PAFEs found in 50 % serum indicate that  
258 prolonged in vitro PAFEs (in RPMI-1640), frequently  
259 interpreted as a contributor to better clinical efficacy  
260 [1–4], may be less important in vivo (better mimicked  
261 by the serum-containing medium) at least against *C.*  
262 *albicans*. Whether these also apply for non-albicans  
263 species is to be answered by further studies.

264 In summary, continuous but not brief caspofungin  
265 exposure produced measurable killing rates against *C.*  
266 *albicans* clinical isolates in killing studies in the  
267 presence of 50 % serum. PAFE after brief exposure to  
268 caspofungin (and probably to other echinocandins),  
269 even when marked, may play a limited role in the  
270 excellent clinical efficacy of echinocandins.

271 **Acknowledgments** R. Kovács was supported by the TÁMOP  
272 4.2.4. A/2-11-1-2012-0001 “National excellence Program-  
273 Elaborating and operating an inland student and researcher  
274 personal support system.” The project was subsidized by  
275 European Union and co-financed by the European Social Fund.

276 **Conflict of interest** L. Majoros received conference travel  
277 grants from MSD, Astellas and Pfizer.  
278

## 279 References

- 280 1. Andes D, Diekema DJ, Pfaller MA, Bohrmuller J, Marchillo  
281 K, Lepak A. In vivo comparison of the pharmacodynamic  
282 targets for echinocandin drugs against *Candida* species.  
283 Antimicrob Agents Chemother. 2010;54:2497–506.
- 284 2. Clancy CJ, Huang H, Cheng S, Derendorf H, Nguyen MH.  
285 Characterizing the effects of caspofungin on *Candida*  
286 *albicans*, *Candida parapsilosis*, and *Candida glabrata* iso-  
287 lates by simultaneous time-kill and postantifungal-effect  
288 experiments. Antimicrob Agents Chemother. 2006;50:  
289 2569–72.
- 290 3. Nguyen KT, Ta P, Hoang BT, Cheng S, Hao B, Nguyen  
291 MH, et al. Characterising the post-antifungal effects of  
292 micafungin against *Candida albicans*, *Candida glabrata*,  
293 *Candida parapsilosis* and *Candida krusei* isolates. Int J  
294 Antimicrob Agents. 2010;35:80–4.
- 295 4. Nguyen KT, Ta P, Hoang BT, Cheng S, Hao B, Nguyen  
296 MH, et al. Anidulafungin is fungicidal and exerts a variety

- 297 of postantifungal effects against *Candida albicans*, *C.*  
 298 *glabrata*, *C. parapsilosis* and *C. krusei* isolates. Antimicrob  
 299 Agents Chemother. 2009;53:3347–52.
- 300 5. Shields RK, Nguyen MH, Du C, Press E, Cheng S, Clancy  
 301 CJ. Paradoxical effect of caspofungin against *Candida*  
 302 bloodstream isolates is mediated by multiple pathways, but  
 303 eliminated in human serum. Antimicrob Agents Chemother.  
 304 2011;55:2641–7.
- 305 6. Kovács R, Gesztelyi R, Berényi R, Domán M, Kardos G,  
 306 Juhász B, et al. Killing rates exerted by caspofungin in 50%  
 307 serum and its correlation with in vivo efficacy in a neutro-  
 308 penic murine model against *Candida krusei* and *Candida*  
 309 *inconspicua*. J Med Microbiol. 2014;63:186–94.
- 310 7. Földi R, Kovács R, Gesztelyi R, Kardos G, Berényi R, Juhász  
 311 B, et al. Comparison of in vitro and in vivo efficacy of ca-  
 312 spofungin against *Candida parapsilosis*, *C. orthopsilosis*, *C.*  
 313 *metapsilosis* and *C. albicans*. Mycopathol. 2012;174:311–8.
- 314 8. Nasar A, Ryan L, Frei CR, Cota JM, Wiederhold NP.  
 315 Influence of serum and albumin on echinocandin in vitro  
 316 potency and pharmacodynamics. Curr Fungal Infect Rep.  
 317 2013;7:89–95.
- 318 9. Szabó Z, Tóth B, Kovács M, Kardos G, Maráz A, Rozgonyi  
 319 F, et al. Evaluation of the new Micronaut-Candida system  
 320 compared to the API ID32C method for yeast identification.  
 321 J Clin Microbiol. 2008;46:1824–5.
- 322 10. Clinical and Laboratory Standards Institute. Reference  
 323 method for broth dilution antifungal susceptibility testing of  
 yeasts. Approved standard, 3rd ed. M27-A3. Clinical and  
 Laboratory Standards Institute, Wayne, PA (2008).
- 324 11. Cornely OA, Vehreschild JJ, Vehreschild MJ, Würthwein  
 325 G, Arenz D, Schwartz S, et al. Phase II dose escalation study  
 326 of caspofungin for invasive *Aspergillosis*. Antimicrob  
 327 Agents Chemother. 2011;55:5798–803.
- 328 12. Cantón E, Pemán J, Valentin A, Espinel-Ingroff A, Gober-  
 329 nado M. In vitro activities of echinocandins against *Candida*  
 330 *krusei* determined by three methods: MIC and minimal  
 331 fungicidal concentration measurements and time-kill stud-  
 332 ies. Antimicrob Agents Chemother. 2009;53:3108–11.
- 333 13. Pfaller MA, Diekema DJ, Andes D, Arendrup MC, Brown  
 334 SD, Lockhart SR, et al. Clinical breakpoints for the echino-  
 335 candins and *Candida* revisited: integration of molecular,  
 336 clinical, and microbiological data to arrive at species-specific  
 337 interpretive criteria. Drug Resist Updat. 2011;14:164–76.
- 338 14. Walker LA, Gow NA, Munro CA. Elevated chitin content  
 339 reduces the susceptibility of *Candida* species to caspofun-  
 340 gin. Antimicrob Agents Chemother. 2013;57:146–54.
- 341 15. Louie A, Deziel M, Liu W, Drusano MF, Gumbo T, Drusano  
 342 GL. Pharmacodynamics of caspofungin in a murine model  
 343 of systemic candidiasis: importance of persistence of ca-  
 344 spofungin in tissues to understanding drug activity. Anti-  
 345 microb Agents Chemother. 2005;49:5058–68.
- 346  
347  
348

UNCORRECTED