

Body size changes in ground beetle assemblages – a reanalysis of Braun *et al.* (2004)'s data

GÁBOR L. LÖVEI¹ and TIBOR MAGURA² ¹Department of Integrated Pest Management, Danish Institute of Agricultural Sciences, Flakkebjerg Research Centre, Denmark, ²Hortobágy National Park Directorate, Debrecen, POB. 216, H-4002, Hungary

Abstract. 1. Data in Braun *et al.*'s (2004) recent paper on size trends in ground beetles (Carabidae) are re-analysed, after re-classifying the species according to feeding categories.

2. When ground beetles are classified as predatory species, mixed feeders or herbivores, and evaluated separately, the size trends indicate group-specific differences. Most of these do not support the efficiency-specialisation hypothesis.

3. Taxonomic groups rarely fit into one ecological category and ecological analyses should explicitly consider this hurdle.

Key words. Body size, Carabidae, disturbance, feeding type, life history, pollution.

Introduction

Size has a significant, substantial impact on ecological interactions (Peters, 1983). It modulates resource use, the outcome of species interactions (Peters, 1983), and more indirectly, the periods of activity, habitat suitability, and numerous other parameters. The study of size relationships and their role in shaping interactions is an active research area in ecology (Peters, 1983; Brown, 1995).

Changes in either the size in individual species, or the size distribution of the species in a habitat are also parameters potentially indicating environmental stress (McGeoch, 1998). Recently, Braun *et al.* (2004) used this approach to analyse species size trends in ground beetles (Carabidae) at an abandoned fertiliser factory in Germany. Using the mean size (estimated biovolume) of beetles present, they found that the average species size decreased over time (16 years), in parallel with a decrease in habitat contamination caused by the former fertiliser factory. Based on these results, they speculated about the validity of two conflicting hypotheses of resource use distribution within the family of ground beetles (Carabidae). In Braun *et al.*'s study (2004) ground beetles were used as a taxonomic (and predatory) group; no consideration was made of the various feeding strategies that are found within the group.

In the present contribution the results by Braun *et al.* (2004) are reconsidered and their data were subjected to an evaluation

that considered feeding habits of the species. The main motivation for this was that an important assumption, the uniform feeding type of ground beetles, is not correct. Neither all ground beetles, nor even the majority of them can be considered predatory. In the European fauna, there are many species that are herbivores, or mixed feeders (Lövei and Sunderland, 1996). A final point is that one of the hypotheses evaluated by Braun *et al.* (2004) and attributed to Blake *et al.* (1994), in fact was suggested earlier by Szyszko (1983).

Materials and methods

Data presented in Braun *et al.* (2004) were re-analysed after resorting their species list into three adult feeding categories: predatory species, mixed feeders, and herbivores. This categorisation was based on work by Larochelle (1990), Hürka (1996), and Lindroth (1985, 1986). We then re-analysed size change trends using the same methods as Braun *et al.* (2004) but chose a different graphical representation.

Size trends were analysed using ANOVA, followed by Tukey tests if significant differences were found. Calculations were done using the SPSS statistical software (SPSS, 2000).

Results

Size trends of all species

Considering all carabids together (regardless of their feeding categories), the mean size in the 90s was significantly smaller

Correspondence: Gabor L. Lövei, Department of Integrated Pest Management, Danish Institute of Agricultural Sciences, Flakkebjerg Research Centre, DK-4200 Slagelse, Denmark. E-mail: gabor.lovei@agrsci.dk

than that in the 80s for all the studied areas (Fig. 1 and Table 1; see also fig. 3 and table 2 in Braun *et al.*, 2004). This picture changed when the species in the different feeding categories were considered separately.

Analysis by feeding categories

Carnivore species. In the severely polluted area, body size was smallest in 1991, significantly smaller than either in 1980 and 1996 (but not 1990). The other three body size values did not significantly differ from each other, although 1981 was smaller than either 1980 or 1996 (Fig. 1 and Table 1).

In the moderately polluted area, there is a decreasing trend, but this consists only of a drop from the first year's higher value to a lower level in the following year. Ten years later, this level is somewhat lower, now becoming significantly smaller than the corresponding value for 1980 (but not the more variable 1981). This level shows virtually no difference 5 years later (Fig. 1 and Table 1).

In the lightly polluted area, there are no significant time trends. The mean size in 1980 was somewhat larger than in the other 3 years (Fig. 1 and Table 1).

Herbivore species. There were few herbivorous species. The size trend in the severely polluted area showed a zigzag pattern, with some significant differences in the middle years. The initial and final years did not differ. The same appeared in the other two areas, at a somewhat lower level (Fig. 1 and Table 1).

Mixed feeder species. The largest number of species were classified as mixed feeders, and, consequently, this group should contribute considerably to the overall trend. In fact, the trends shown by the mixed feeders were similar to that of 'all species'. This group shows no significant size difference among the dif-

ferently polluted areas in 1980 or 1981. Severely and moderately polluted areas seem to support assemblages that became progressively smaller, more in the severely polluted than in the moderately polluted area. There was no decreasing trend in the lightly polluted area, except a smaller assemblage in 1991 (Fig. 1 and Table 1).

Discussion

This re-analysis further developed the conclusions of Braun *et al.* (2004). Most of the support for the role of the increasing specialisation hypothesis in the organisation of ground beetle assemblages disappears when the feeding behaviour of the species is taken into account.

Our analysis underlines the notion that taxonomic units rarely correspond to ecological units, and it is rarely justified to consider them as such. Considering the mean size of ground beetles in general may give rise to a statistical artefact. This is due to the oversimplified classification of the family as predators. Considering natural history is essential for the correct ecological interpretation of patterns in nature. Carabids can conveniently be collected by a single method as most of the species, at least in the northern temperate region (but not in the tropics, see Erwin, 1979), have surface-active adults. Traditionally, carabids have been considered predators, but even Thiele (1977) stressed that this can be a misleading simplification. Results published since the early 70s fully support this suggestion (Lövei and Sunderland, 1996; Toft and Bilde, 2002). Consequently, the approach followed by Braun *et al.* (2004) is of limited value, and in need of further development. The different feeding types of ground beetles should be considered separately and it was demonstrated that this indeed changed the time-size relationship. It is ecologically more valid to consider the changes in size and the assumed resource partitioning with at least




Fig. 1. Size trends (biovolume, mm^3) of ground beetle species captured in three areas under different pollution levels near a fertiliser factory, Germany, in 4 years between 1980 and 1996. Data from Braun *et al.* (2004).

Table 1. Results of the one-way ANOVA for the ground beetle species collected in the severely, moderately, and lightly polluted areas in 1980, 1981, 1991, and 1996, near a fertiliser factory, Germany. Data from Braun *et al.* (2004)

Source	SS	d.f.	MS	F	P	Tukey-test
All species, severely polluted area						
Year	79.855	3	26.618	93.023	< 0.001	1980 = 1981 > 1991 > 1996
Error	1092.223	3817	0.286			
Total	11937.437	3821				
All species, moderately polluted area						
Year	7.279	3	2.426	15.992	< 0.001	1980 = 1981 = 1991 > 1996
Error	71.312	470	0.152			
Total	1647.199	474				
All species, lightly polluted area						
Year	4.588	3	1.529	9.386	< 0.001	1980 > 1991; 1980 > 1996
Error	118.786	729	0.163			
Total	2311.356	733				
Carnivore species, severely polluted area						
Year	68.275	3	22.758	56.302	< 0.001	1980 > 1991; 1996 > 1991
Error	191.194	473	0.404			
Total	794.861	477				
Carnivore species, moderately polluted area						
Year	4.069	3	1.356	3.976	0.009	1980 > 1991; 1980 > 1996
Error	48.440	142	0.341			
Total	521.016	146				
Carnivore species, lightly polluted area						
Year	0.942	3	0.314	1.465	0.224	
Error	66.863	312	0.214			
Total	956.640	316				
Herbivore species, severely polluted area						
Year	4.155	3	1.385	24.174	< 0.001	1996 > 1991
Error	50.995	890	0.057			
Total	2497.278	894				
Herbivore species, moderately polluted area						
Year	4.923	3	1.641	30.136	< 0.001	1991 > 1996
Error	4.847	89	0.054			
Total	249.327	93				
Herbivore species, lightly polluted area						
Year	2.430	3	0.810	12.819	< 0.001	not performed*
Error	4.423	70	0.063			
Total	175.991	74				
Mixed feeding species, severely polluted area						
Year	51.720	3	17.240	72.583	< 0.001	1980 > 1991; 1980 > 1996, 1981 > 1996, 1991 > 1996
Error	580.979	2446	0.238			
Total	8645.298	2450				
Mixed feeding species, moderately polluted area						
Year	1.447	3	0.482	9.555	< 0.001	1980 = 1981 = 1991 > 1996
Error	11.713	232	0.050			
Total	876.856	236				
Mixed feeding species, lightly polluted area						
Year	3.370	3	1.123	9.220	< 0.001	1980 = 1981 = 1996 > 1991
Error	41.419	340	0.122			
Total	1178.725	344				

*Too few individuals in 1980 and 1981.

the different feeding types separately. The assumptions of feeding categories are crucial, and future studies refining these categories (for example, the extent of carrion feeding in carabids, which currently is not well understood, Lövei and Sunderland, 1996) might lead to still different results.

When seeking explanations for assemblage organisation, single-group studies can lead to spurious results. In the case of carabids, spiders and ants should be considered, as they have a significant impact on ground beetles (Thiele, 1977; Lövei and Sunderland, 1996).

A further, modulating influence on assemblage organisation is the permanence of species in a habitat. Species presence is often taken as an indication that the species can find all its needs and survive in the habitat. The presence of non-resident species, termed tourists, transient or occasional species in ecological communities, however, is frequent (Magurran, 2003). Such species do not have sufficiently tight links to the habitat where found, and even though their ratio can be high (Novotny and Missa, 2000), they are not legitimate members of the ecological community. Considering all species captured as integral components of the ecosystem is debatable. Species presence is not necessarily permanent, and community organisation could be better understood if permanent and transient species are identified and separated (Magurran and Henderson, 2003). The latter may have little explanatory power in the search for assembly and resource distribution patterns (Magurran, 2003). Size trends could be different also in this case if we could separate permanent from temporary species in the present case but this is not possible due to the non-continuous sampling.

Braun *et al.* (2004) termed the hypothesis that increasing disturbance causes the average size of ground beetles to decrease in a habitat as 'Blake *et al.* hypothesis'. The same idea, although formulated a little differently, was suggested in 1983 by Jan Szyszko, who studied the possibility of characterising pine forest regeneration by using an index of 'mean individual biomass' (Szyszko, 1983). His hypothesis, while formulated as 'decreasing disturbance allows greater average body size', is essentially the same as in Blake *et al.* (1994). Therefore, in accordance with scientific priority, we suggest that this hypothesis should be termed the 'Szyszko-hypothesis', or 'increasing average size hypothesis'.

Acknowledgements

We thank Drs A. Báldi, B. Hawkins, J. Kotze, B. Tóthmérész, and three anonymous reviewers for comments on earlier drafts of the manuscript.

References

- Blake, S., Foster, G.N., Eyre, M.D. and Luff, M.L. (1994) Effects of habitat type and grassland management practices on the body size distribution of carabid beetles. *Pedobiologia*, **38**, 502–512.
- Braun, S.D., Jones, T.H. and Perner, J. (2004) Shifting average body size during regeneration after pollution – a case study using ground beetle assemblages. *Ecological Entomology*, **29**, 543–554.
- Brown, J.H. (1995) *Macroecology*. University of Chicago Press, Chicago, Illinois.
- Cleveland, W.C. (1994) *The Elements of Graphing Data*. Hobart Press, Summit, New Jersey.
- Erwin, T.L. (1979) Thoughts on the evolutionary history of ground beetles: hypotheses generated from comparative faunal analysis of lowland forest sites in temperate and tropical regions. *Carabid Beetles: Their Evolution, Natural History, and Classification* (ed. by T. L. Erwin, G. E. Ball, D. R. Whitehead and A. L. Halpern), pp. 539–592. Junk Publishers, The Hague.
- Hürka, K. (1996) *Carabidae of the Czech and Slovak Republics*. Kabourek, Zlin, Czech Republic.
- Larochelle, A. (1990) The food of carabid beetles. *Fabreeriesupplement*, **5**, 1–132.
- Lindroth, C.H. (1985) *The Carabidae (Coleoptera) of Fennoscandia and Denmark*. Fauna Entomologica Scandinavica, **15**, part 1. E.J. Brill, Leiden, The Netherlands.
- Lindroth, C.H. (1986) *The Carabidae (Coleoptera) of Fennoscandia and Denmark*. Fauna Entomologica Scandinavica, **15**, part 2. E.J. Brill, Leiden, The Netherlands.
- Lövei, G.L. and Sunderland, K.D. (1996) Ecology and behavior of ground beetles (Coleoptera: Carabidae). *Annual Review of Entomology*, **41**, 231–256.
- Magurran, A.E. (2003) *Measuring Biological Diversity*. Oxford University Press, Oxford.
- Magurran, A.E. and Henderson, P.A. (2003) Explaining the excess of rare species in natural species abundance distributions. *Nature*, **422**, 714–716.
- McGeoch, M.A. (1998) The selection, testing, and application of terrestrial insects as bioindicators. *Biological Reviews*, **73**, 181–201.
- Novotny, V. and Missa, O. (2000) Local versus regional species richness in tropical insects: one lowland site compared with the island of New Guinea. *Ecological Entomology*, **25**, 445–451.
- Peters, R.H. (1983) *The Ecological Implications of Body Size*. Cambridge University Press, Cambridge.
- SPSS Inc. (2000) *SPSS 10.0.7 for Windows*. SPSS Inc., Chicago, Illinois.
- Szyszko, J. (1983) State of Carabidae (Col.) fauna in fresh pine forest and tentative valorisation of this environment. SGGW-AG Monographs No. 28. Warsaw Agricultural University Press, Warsaw, Poland.
- Thiele, H.-U. (1977) *Carabid Beetles in Their Environments*. Springer, Berlin.
- Toft, S. and Bilde, T. (2002) Carabid diets and food value. *The Agroecology of Carabid Beetles* (ed. by J. M. Holland), pp. 81–110. Intercept, Andover, U.K.
- Tufte, E.R. (2003) *The Visual Display of Quantitative Information*, 2nd edn. Graphics Press, Cheshire, Connecticut.

Accepted ?????? 2005

Author Query Form

Journal: Ecological Entomology

Article: een_794.xml

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication. Many thanks for your assistance.

Query No.	Query	Remark
1	Au: Please supply accepted date	

MARKED PROOF

Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Stet
Insert in text the matter indicated in the margin	⤴	New matter followed by ⤴
Delete	⤵ through matter to be deleted	⤵
Delete and close up	⤵ through matter to be deleted	⤵
Substitute character or substitute part of one or more word(s)	/ through letter or ⤵ through word	New letter or new word
Change to italics	— under matter to be changed	ƒ
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	= under matter to be changed	=
Change to bold type	~ under matter to be changed	~
Change to bold italic	≡ under matter to be changed	≡
Change to lower case	Encircle matter to be changed	⊖
Change italic to upright type	(As above)	⤴
Insert 'superior' character	/ through character or ⤴ where required	⤴ under character e.g. ⤴
Insert 'inferior' character	(As above)	⤵ over character e.g. ⤵
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	⤴ and/or ⤵
Insert double quotation marks	(As above)	⤴ and/or ⤵
Insert hyphen	(As above)	⊖
Start new paragraph	⤴	⤴
No new paragraph	⤵	⤵
Transpose	⤴	⤴
Close up	linking Ⓞ letters	Ⓞ
Insert space between letters	⤴ between letters affected	#
Insert space between words	⤴ between words affected	#
Reduce space between letters	⤴ between letters affected	⤴
Reduce space between words	⤴ between words affected	⤴