

SCIENTIFIC REPORTS



OPEN

Characterization of the *aodA*, *dnmA*, *mnSOD* and *pimA* genes in *Aspergillus nidulans*

Éva Leiter¹, Hee-Soo Park², Nak-Jung Kwon², Kap-Hoon Han^{2,3}, Tamás Emri¹, Viktor Oláh⁴, Ilona Mészáros⁴, Beatrix Dienes⁵, János Vincze⁵, László Csernoch⁵, Jae-Hyuk Yu² & István Pócsi¹

Received: 12 October 2015

Accepted: 05 January 2016

Published: 05 February 2016

Mitochondria play key roles in cellular energy generation and lifespan of most eukaryotes. To understand the functions of four nuclear-encoded genes predicted to be related to the maintenance of mitochondrial morphology and function in *Aspergillus nidulans*, systematic characterization was carried out. The deletion and overexpression mutants of *aodA*, *dnmA*, *mnSOD* and *pimA* encoding alternative oxidase, dynamin related protein, manganese superoxide dismutase and Lon protease, respectively, were generated and examined for their growth, stress tolerances, respiration, autolysis, cell death, sterigmatocystin production, hyphal morphology and size, and mitochondrial superoxide production as well as development. Overall, genetic manipulation of these genes had less effect on cellular physiology and ageing in *A. nidulans* than that of their homologs in another fungus *Podospora anserina* with a well-characterized senescence. The observed interspecific phenotypic differences can be explained by the dissimilar intrinsic stabilities of the mitochondrial genomes in *A. nidulans* and *P. anserina*. Furthermore, the marginally altered phenotypes observed in *A. nidulans* mutants indicate the presence of effective compensatory mechanisms for the complex networks of mitochondrial defense and quality control. Importantly, these findings can be useful for developing novel platforms for heterologous protein production, or on new biocontrol and bioremediation technologies based on *Aspergillus* species.

Fungal ageing and longevity are highly dependent on mitochondrial integrity and functions in both the yeast *Saccharomyces cerevisiae* and the filamentous fungus *Podospora anserina*^{1–4}. The age-dependent instability and rearrangements of mitochondrial DNA (mtDNA) result in cease of the expression of mtDNA-encoded genes, the subsequent impairment of basically important mitochondrial functions like respiration and, finally, in the definite lifespan and senescence phenomenon typical of *P. anserina* cultures^{3–8}. Because any failure in the maintenance of mitochondrial functions may result in the early onset of programmed cell death the elements of mitochondrial quality control system including the elimination of reactive oxygen species (ROS), damaged biomolecules and even impaired mitochondria (mitophagy) are intensively studied but still not fully understood in fungal biology^{4,7,8}.

In *P. anserina*, disturbances in the supply and maintenance of respiratory chain proteins may lead to deleterious consequences and, hence, a strong connection between respiration and longevity should exist^{6,9,10}. Paradoxically at first glance, the cytochrome C oxidase (COX) deficient mutant strains of *P. anserina* showed increased lifespan^{6,10,11} although *P. anserina* is an obligate aerobic fungus and, consequently, the loss of COX should be lethal. However, the fungus has the capability to induce an alternative oxidase (AOX) pathway to compensate for the impairment of the COX protein^{6,10,11}.

Under normal physiological conditions, ROS are eliminated by an arsenal of antioxidative enzymes in fungal cells including superoxide dismutases, which convert superoxide to H₂O₂. Although these enzymes affect the mitochondrial ROS levels, mitochondrial stability, subsequent ageing and lifespan beneficially^{12,13} the

¹Department of Biotechnology and Microbiology, Faculty of Science and Technology, University of Debrecen, Debrecen, Hungary. ²Departments of Bacteriology and Genetics, The University of Wisconsin-Madison, Wisconsin, USA. ³Department of Pharmaceutical Engineering, Woosuk University, Wanju, Republic of Korea. ⁴Department of Botany, Faculty of Science and Technology, University of Debrecen, Debrecen, Hungary. ⁵Department of Physiology, Faculty of Medicine, University of Debrecen, Debrecen, Hungary. Correspondence and requests for materials should be addressed to L.C. (email: csl@edu.unideb.hu) or J.-H.Y. (email: jyu1@wisc.edu) or I.P. (email: pocsi.istvan@science.unideb.hu)

overexpression of manganese superoxide dismutase encoded by *PaSod3* unexpectedly led to a reduced lifespan in *P. anserina*, and the deletion of the same gene increased the paraquat sensitivity but did not affect the peroxide tolerance and the lifespan of the fungus¹⁴. These data warn us that the roles of ROS in lifespan and ageing may be controversial under certain conditions^{4,14,15}.

Mitochondrial dynamics (changes in the frequency of fission and fusion phenotypes) as part of the cellular quality control is an indispensable part of the cellular adaptation to rapidly changing environmental conditions including stress^{4,7,16}. The fission of mitochondria is controlled by a dynamin-related protein, which is found in all eukaryotes including yeasts and filamentous fungi^{7,17,18}. In *P. anserina*, mutations in the dynamin-related protein 1 encoding gene *PaDnm1* resulted in filamentous-like mitochondria and prolonged the lifespan of the fungus, *i.e.* some mycelia grew up to 1,000 days¹⁸. On the other hand, overexpression of *PaDnm1* increased disintegration of mitochondria even in young cultures¹⁸.

There is one more strategy to preserve the quality of mitochondrial proteins and, therefore, to increase the longevity of the mitochondria in fungi through the induction of the damaged and excess protein removal mechanisms. In these organelles, the soluble serine protease LON plays a central role in controlling degradation of the impaired proteins^{4,7}. In *P. anserina*, the yeast LON ortholog *PaLon* protein is expressed at higher level in senescent cultures than in juvenile cultures. Overexpression of *PaLon* led to elevated resistance against external hydrogen-peroxide compared to a control strain, and increased amount of *PaLon* prolonged the lifespan of the cultures¹⁵. Cytochrome c oxidase deficient long-lived mutants of *P. anserina* was also characterized by increased LON activities¹⁰ and Adam *et al.*¹⁹ reported a decreased lifespan for the *PaLon* gene deletion mutant grown under physiological conditions and also demonstrated the contribution of PaLon protease to ascospore germination and sexual reproduction.

In this study, we characterize a group of nuclear-encoded genes that are thought to be important for mitochondrial functions in the model organism *Aspergillus nidulans*, a model fungus to many Aspergilli with outstanding industrial or biomedical importance²⁰. The group of selected genes included homologs of functionally characterized fungal AOX alternative oxidase, DNM1 dynamin-related protein 1, MnSOD superoxide dismutase and the LON serine protease genes^{14,21,22}. Notably, a more recent study by Li *et al.*²³ managed to link oxidative stress, mitochondrial dysfunction and subsequent apoptotic cell death to the spontaneous degeneration of *A. nidulans* maintained on artificial culture media, which adds a strong argument supporting the applicability of this study.

Results

A series of *A. nidulans* mutants were generated through the deletion and overexpression of nuclear-encoded genes that were shown to be directly involved in the maintenance of the integrity and function of mitochondria in other fungi, and a number of key physiological, morphological and developmental phenotypes of the mutants were analyzed.

Growth and stress challenges on nutrient agar plates. Various stress sensitivities of the mutant and control strains were tested on minimal nitrate medium (MNM) agar stress plates²⁴. In general, fungal growth and stress sensitivities were hardly affected by the deletions and overexpressions of the mitochondrial function and morphology related genes and the phenotypic changes were rather sporadic (Fig. 1 and Supplementary Fig. S1). For example, the $\Delta dnmA$ mutant showed a reduced growth (approximately 40% decrease in colony diameter) even without any stress treatment in comparison to the control strain and the deletion of *mnSOD* displayed a remarkably high sensitivity to menadione exposure (Fig. 1B and Supplementary Fig. S1). The inactivation of *pimA* brought about decreased tolerance meanwhile overexpression of the same gene resulted in an increased tolerance to heavy metal stress initiated by CdCl₂ (Fig. 1B and Supplementary Fig. S1). Inactivation of *aodA* did not influence the oxidative stress sensitivity of the fungus in accordance with previous findings reported by Suzuki *et al.*²⁵ (2012). Quite surprisingly, only the $\Delta mnSOD$ mutant showed significantly increased sensitivity to PAF, an anti-fungal protein eliciting apoptotic cell death in sensitive fungi²⁶. When employed at a concentration of 200 µg/ml PAF reduced the growth of the control and $\Delta mnSOD$ strains with approximately 60 and 75%, respectively (Fig. 1C and Supplementary Fig. S1).

Growth and oxidative stress defense in submerged culture. Growth of the mutants was also characterized in submerged cultures by recording gains in the dry cell mass (DCM) (Supplementary Table S1). Complex media were inoculated with conidiospores and, after 16 h incubation, mycelia were washed to MNM medium supplemented with 2% glucose and also to glucose-free minimal medium. Although significant delays in biomass gains were observed in glucose supplemented cultures of the $\Delta aodA$, *dnmAOE* (“OE” indicates “overexpression” mutant) and *mnSODOE* strains and, to a lesser extent, with the $\Delta dnmA$ and $\Delta pimA$ gene deletion strains after 4 h incubation these differences disappeared when the incubation times were increased up to 10 h (Supplementary Table S1). It is noteworthy that the biomass values of the strains were comparable after 24 h carbon starvation in glucose-free culture medium independently of the genotypes with the exception of the $\Delta aodA$ strain where no significant autolytic loss of biomass was recorded. In addition, all strains reached similar biomass gains when they were transferred to MNM supplemented with 1% glucose, which indicates similar survival rates of the carbon-starving control and mutant strains (Supplementary Table S1).

To test the reactive species (RS) productions of the strains tested, mycelia grown in submerged cultures were stained by the fluorescent dye 2',7'-dichlorofluorescein diacetate (Tables 1 and 2). DCF (2',7'-dichlorofluorescein) production, which is proportional to RS generation^{24,27}, was higher in glucose-supplemented cultures of the $\Delta aodA$ mutant incubated for 4 h but was lower when the fungus was incubated in glucose-free minimal medium for 24 h. Both the deletion and overexpression of *pimA* increased DCF levels in glucose-supplemented submerged cultures when compared to the control. It is noteworthy that similar tendency was also observed when




Figure 1. Comparison of the growth, stress sensitivities, cell death and autolysis of the control and mutant strains. Part A: Growth of control (THS30.3) and mutant strains on MNM agar plates. Part B: Selected stress sensitivity experiments with the control and the $\Delta mnSOD$, $\Delta pimA$ and *pimAOE* strains. Part C: Growth inhibiting effect of the apoptosis-inducing antifungal protein PAF on the $\Delta mnSOD$ mutant in comparison to the control strain. Part D: Proteinase activities as autolysis markers^{29,30} recorded in carbon-starving (24 h) submerged cultures of the THS30.3 control and $\Delta aodA$ and *aodAOE* mutants. In proteinase assays, azocasein was used as substrate, and mean \pm SD proteinase activity values calculated from three independent experiments are presented. The statistically significant difference calculated for the $\Delta aodA$ strain by the Student's t test is marked with asterisks: *** $p < 0.1\%$. Further phenotypes were not detected as shown in Supplementary Figs S1 and S3.

Strains	DCF	GPx	GR	Catalase	SOD
Control (THS30.3)	0.76 ± 0.08	3.63 ± 0.06	4.2 ± 0.1	0.94 ± 0.02	0.082 ± 0.009
$\Delta aodA$	3.40 ± 0.05***	2.6 ± 0.1***	4.6 ± 0.2*	2.3 ± 0.1***	0.14 ± 0.02**
<i>aodAOE</i>	0.9 ± 0.3	4.19 ± 0.06***	4.7 ± 0.3*	0.8 ± 0.1*	0.09 ± 0.01
$\Delta dnmA$	0.8 ± 0.2	5.8 ± 0.2***	7.6 ± 0.3***	1.7 ± 0.1***	0.130 ± 0.009***
<i>dnmAOE</i>	0.59 ± 0.07*	2.78 ± 0.03***	4.0 ± 0.1	1.3 ± 0.1***	0.078 ± 0.006
$\Delta mnSOD$	0.5 ± 0.1*	3.62 ± 0.03	4.6 ± 0.1*	1.2 ± 0.1*	0.063 ± 0.004*
<i>mnSODOE</i>	0.6 ± 0.1*	3.6 ± 0.1	4.6 ± 0.2*	0.68 ± 0.03*	0.15 ± 0.03*
$\Delta pimA$	1.8 ± 0.4**	4.7 ± 0.1***	4.7 ± 0.2**	1.49 ± 0.05***	0.088 ± 0.008
<i>pimAOE</i>	2.4 ± 0.3***	3.67 ± 0.06	4.6 ± 0.2*	1.1 ± 0.1	0.074 ± 0.003

Table 1. Changes in the specific DCF production and in the specific GPx, GR, catalase and SOD activities^a in growing cultures. ^aSpecific enzyme activities are expressed as mkat/kg protein (GPx and GR) or kat/kg protein (catalase). Specific SOD activities are given as mU/kg protein⁵⁹. Specific DCF productions are expressed as mmol/kg DCM. Specific DCF productions as well as specific enzyme activity values are presented as mean ± SD calculated from three independent experiments. **p* < 5%, ***p* < 1% and ****p* < 0.1%. *p* values were calculated using the Student's t-test.

Strains	DCF	ST production	GPx	GR	Catalase	SOD
Control (THS30.3)	0.18 ± 0.02	0.14 ± 0.02	0.62 ± 0.07	2.3 ± 0.2	2.9 ± 0.1	0.209 ± 0.009
$\Delta aodA$	0.10 ± 0.01***	0.030 ± 0.004***	0.71 ± 0.01	2.39 ± 0.04	1.6 ± 0.2***	0.167 ± 0.009***
<i>aodAOE</i>	0.18 ± 0.01	0.026 ± 0.003***	0.33 ± 0.06***	1.3 ± 0.1***	1.4 ± 0.2***	0.10 ± 0.01***
$\Delta dnmA$	0.28 ± 0.04*	0.024 ± 0.003***	0.57 ± 0.07	2.1 ± 0.1	3.2 ± 0.1	0.20 ± 0.01
<i>dnmAOE</i>	0.12 ± 0.01**	0.15 ± 0.02	0.36 ± 0.04***	1.67 ± 0.04***	1.8 ± 0.1***	0.13 ± 0.01***
$\Delta mnSOD$	0.15 ± 0.01	0.18 ± 0.02*	0.59 ± 0.01	2.38 ± 0.06	2.7 ± 0.1	0.132 ± 0.005***
<i>mnSODOE</i>	0.10 ± 0.01***	0.2 ± 0.02**	0.41 ± 0.01***	2.0 ± 0.1	1.0 ± 0.2***	0.138 ± 0.008***
$\Delta pimA$	0.6 ± 0.1***	0.020 ± 0.002***	0.59 ± 0.01	1.9 ± 0.4	4.7 ± 0.2***	0.24 ± 0.03
<i>pimAOE</i>	0.27 ± 0.01***	0.35 ± 0.04***	0.24 ± 0.03***	1.03 ± 0.01***	1.4 ± 0.1***	0.12 ± 0.012***

Table 2. Changes in the specific DCF and ST productions and in the specific GPx, GR, catalase and SOD activities^a in carbon-starving ageing cultures. ^aSpecific enzyme activities are expressed as mkat/kg protein (GPx and GR) or kat/kg protein (catalase). Specific SOD activities are given as mU/kg protein⁵⁹. Specific DCF and ST productions are expressed as mmol/kg DCM and mg/g DCM, respectively. Specific DCF and ST productions as well as specific enzyme activity values are presented as mean ± SD calculated from three independent experiments. **p* < 5%, ***p* < 1% and ****p* < 0.1%. *p* values were calculated using the Student's t-test.

the $\Delta pimA$ strain was incubated in glucose-free medium for 24 h. The overexpression of *mnSOD* also decreased the intracellular RS levels considerably in both glucose supplemented and glucose-free cultures. Although further minor phenotypes appeared in $\Delta mnSOD$ (in the presence of 2% glucose) and $\Delta dnmA$ (in the absence of glucose) cultures these changes were significant only at *p* < 5% levels (Tables 1 and 2).

To screen the oxidative stress defense systems of the mutants, a selection of antioxidant enzyme activities, including glutathione peroxidase (GPx), glutathione reductase (GR), catalase and superoxide dismutase (SOD) specific activities, were measured (Tables 1 and 2). In cultures supplemented with 2% glucose, the deletion of alternative oxidase increased the specific catalase, SOD and GR activities, while the GPx activity was lower than that found in the control strain. In the *aodAOE* mutant, GPx and GR levels increased significantly. The deletion of *dnmA* elevated the specific activities of all tested enzymes meanwhile the GPx activity decreased and, surprisingly, the catalase activity concomitantly increased significantly in the *dnmAOE* strain. Manipulation of MnSOD production affected the GR, catalase and, as expected, the SOD activities (Table 1). Deletion of *mnSOD* raised GR and catalase activities while overproduction of MnSOD lowered the activity of catalase but increased the SOD activity. Interestingly, high specific GPx, GR and catalase activities were found in the $\Delta pimA$ strain (Table 1).

In carbon-starving ageing cultures of the $\Delta aodA$ strain, both the specific catalase and SOD activities decreased meanwhile the *aodAOE* mutant displayed reduced GPx, GR, catalase and SOD activity levels. The overexpression of *dnmA* mitigated all enzyme activities tested but the deletion of *dnmA* did not influence the antioxidative enzyme activities at all. The overexpression of *mnSOD* resulted in lower GPx, catalase as well as SOD activities. In the *pimAOE* mutant, low specific GPx, GR, catalase and SOD activities were measured, and the specific catalase activities in the $\Delta pimA$ strain were of higher than those recorded in the control strain (Table 2).

Production of sterigmatocystin and autolysis-related proteinase. As oxidative stress response has been connected to secondary metabolite production²⁸, we determined the sterigmatocystin (ST) yields²⁴ in carbon-starving submerged cultures of the mutants (Table 2, Supplementary Fig. S2). Importantly, the overexpression of *aodA* as well as the deletion of *aodA*, *dnmA* and *pimA* decreased ST productions meanwhile mycotoxin production increased considerably (more than twofold) in the *pimAOE* strain and, to a lesser extent, in the $\Delta mnSOD$ and *mnSODOE* strains (Table 2; Supplementary Fig. S2).




Figure 2. Respiration rates. Changes in total (■), KCN-sensitive (▨) alternative oxidase (AOX; □) and KCN + SHAM-resistant respirations (▩) ($\mu\text{mol}/\text{min}/\text{g DCM}$) recorded either in young cultures supplemented with 2% glucose (Part A) or in old, carbon starving cultures (Part B). KCN-sensitive and AOX-dependent respiration were calculated by subtraction of KCN-resistant from total respiration and of KCN-SHAM from KCN respiration, respectively. In Part A, the addition of KCN even stimulated the respiration of the *aodAOE* strain and, hence, we got a virtual negative value for the KCN-sensitive respiration in this case. This paradoxical behavior can be explained with the considerably increased capacity of the AOX-dependent respiratory pathway similar to that observed before in the *P. anserina* AOX overexpression strain by Lorin *et al.*⁶. Mean \pm SD values calculated from three independent experiments are presented. Statistically significant differences compared to the control determined by the Student's t test are marked with asterisks: * $p < 5\%$, ** $p < 1\%$, *** $p < 0.1\%$. In all respiration rate assays, the THS30.3 strain was used as the control strain.

We also determined the autolytic processes of the carbon-starving (16 + 24 h) submerged cultures by measuring proteinase activity accumulating in aging *A. nidulans* cultures²⁹. Among our mutants, only the deletion of *aodA* alternative oxidase gene decreased significantly the specific autolytic proteinase activity which prevented any autolytic loss of dry cell mass (DCM) (Fig. 1D; Supplementary Fig. S3; Supplementary Table S1).

Respiration. Mitochondria are the central organelles of respiration and, therefore, total, cytochrome c-dependent KCN-sensitive, alternative oxidase (AOX)-dependent [KCN-resistant, SHAM (salicylhydroxamic acid)-sensitive and remaining (KCN + SHAM resistant, *i.e.* resistant to the concomitant addition of KCN and SHAM) respirations were determined.

As shown in Fig. 2, all types of respirations were typically lower in carbon-starving cultures but these differences were highly unequal, *e.g.* 52.3, 27.1, 63.7 and 84% lower values were recorded for the total, cytochrome c-dependent, AOX-dependent and remaining respirations, respectively, when carbon-starving and glucose supplemented cultures of the THS30.3 control strain were compared (Fig. 2A,B). Very similar tendencies in total and KCN-resistant respirations were detected previously in carbon-starving *A. nidulans* cultures³⁰.

In this study, in cultures supplemented with 2% glucose, manipulation of AOX production resulted in significant changes in the cyanide-resistant respiration. The deletion of *aodA* decreased while the overexpression of the same gene increased the cyanide-resistant as well as the KCN + SHAM-resistant (remaining) respirations of the fungus (Fig. 2A). It is noteworthy that the total respiration of the $\Delta aodA$ strain decreased in the presence of glucose meanwhile the respiration of the *aodAOE* strain even increased paradoxically resulting in a negative value KCN-sensitive respiration when the culture was supplemented with KCN (Fig. 2A). In carbon-starving cultures,

similar tendencies were observed with some differences; meanwhile the total and the cytochrome c-dependent respirations were not affected in $\Delta aodA$ cultures they were considerably reduced in the *aodAOE* strain (Fig. 2B).

Furthermore, the deletion of *dnmA* significantly increased the total and the KCN-resistant AOX-dependent and decreased the remaining respirations of the cultures grown in the presence of 2% glucose (Fig. 2A). In ageing cultures, the gene deletion decreased considerably the total and the KCN-sensitive cytochrome c-dependent respirations (Fig. 2B). The KCN + SHAM resistant remaining respiration was significantly lower in both the $\Delta dnmA$ and the *dnmA*AOE strains in the presence of glucose (Fig. 2A). In both the $\Delta mnSOD$ and *mnSODOE* strains, the KCN + SHAM resistant respiration decreased in comparison to the control strain (Fig. 2A). In ageing cultures of the $\Delta mnSOD$ and *mnSODOE* strains, the KCN + SHAM resistant respiration remained higher than that found in control cultures (Fig. 2B). Importantly, both the deletion and overexpression of *mnSOD* decreased the total and the KCN-sensitive cytochrome c-dependent respirations in ageing cultures (Fig. 2B) and a lower AOX-dependent respiration was also detected in the $\Delta mnSOD$ strain. The overexpression of *pimA* decreased the total, the KCN-sensitive cytochrome c-dependent and the KCN + SHAM resistant remaining respirations in the presence of glucose while the AOX-dependent respiration was increased and the remaining activity was decreased by $\Delta pimA$ mutation (Fig. 2A). In ageing cultures, the total and the cytochrome c-dependent respirations of the $\Delta pimA$ strain decreased meanwhile the respiration of the *pimAOE* strain was decreased only a lower level of significance ($p < 5\%$) when compared to that of the THS30.3 control strain (Fig. 2B).

Mitochondrial morphology. We studied mitochondrial morphology, superoxide production and hyphal diameter in 20–24 h cultures of *A. nidulans* mycelia using MitoTracker Green, dihydroethidium and Calcofluor White stainings, respectively^{23,31} (Fig. 3A–C, Supplementary Fig. S4). Examining the volumetric ratio of the mitochondria inside the selected region of interest, we found significant differences between the control and the mutant strains only in two cases. Namely, the deletion of *dnmA* and the overexpression of *pimA* increased the volumetric ratio of mitochondria in the second hyphal segments from the apices (Fig. 3D; Supplementary Fig. S4). Importantly, meanwhile the numbers of these organelles did not change significantly in these mutants the sizes of mitochondria expanded considerably (Fig. 3E; Supplementary Fig. S4). It is worth noting that no significant differences were found in superoxide productions visualized by dihydroethidium staining irrespective of changes in the volumetric ratio and size of mitochondria (Supplementary Fig. S4; Supplementary Table S4) although these organelles are the main source of ROS and also the main targets of ROS-induced organelle damages^{1,4,7,23}. Surprisingly, overexpression of *aodA*, *dnmA* and *pimA* but not that of *mnSOD* resulted in significantly thinner hyphae (with approximately 30–50% reductions in hyphal diameters) in comparison to the control strain (Supplementary Fig. S4E).

Cleistothecia formation and asexual sporulation, viability of conidiospores. We also quantified fruiting body formation and conidiospore production in all mutants^{24,32}. Both the deletion and overexpression of *aodA* increased the number of cleistothecia meanwhile the relative numbers of fruiting bodies were less in the *pimAOE* and *mnSODOE* strains than those found in control cultures (Fig. 4A). The maturation of cleistothecia has also been accelerated in the $\Delta aodA$ and *aodAOE* mutants, and completely matured cleistothecia were observed after 4 d incubation in the *aodAOE* strain ($175 \pm 32 \text{ cm}^{-2}$ compared to the control $2.2 \pm 0.1 \text{ cm}^{-2}$; Supplementary Fig. S6). Surprisingly, the $\Delta pimA$ mutant was unable to produce ascospores even after 12 d incubation although the number of cleistothecia was within the normal range. Considering asexual sporulation, the deletion of each gene tested resulted in considerably reduced conidiospore formation. The relative numbers of conidiospores were also slightly reduced by overexpression of *dnmA* and *pimA* (Fig. 4B).

We also tested the viability of the conidiospores under different stress conditions^{32,33}. Incubation of spores of the $\Delta mnSOD$ mutant at 50 °C for 10 min resulted in an appr. 50% reduction in viability meanwhile the asexual spores of the control and other genetically modified strains were not affected by thermal stress at all (Fig. 4C and Supplementary Fig. S5). Short-term storage as well as long-term storage (3, 6 and 12 d) of the conidiospores at 4 °C reduced drastically the viability of the $\Delta mnSOD$, and also decreased significantly the viabilities of the $\Delta aodA$, $\Delta dnmA$ and $\Delta pimA$ asexual spores (Fig. 4D).

Discussion

Mitochondria are the powerhouses of the cells in the overwhelming majority of eukaryotes including fungi³⁴. As a major source of endogenous ROS in aerobic organisms, mitochondria are continuously subjected to oxidative damages. Thus, not surprisingly, the ROS-elicited decay of mitochondrial functions is considered as a leading cause for cell ageing processes and, ultimately, the onset of programmed cell death^{1,4}. Although mitochondrial ageing and senescence is an intensively studied area in fungal biology, our current knowledge in this important field is limited mostly to two species, the baker's yeast *S. cerevisiae* and the filamentous fungus *P. anserina*^{1–4}.

In this study, we aimed at the functional investigation of four nuclear-encoded *A. nidulans* genes, whose homologs are associated with mitochondrial function and morphology, and have been found to influence ageing and longevity in *P. anserina*^{14,21,22}. The *aodA*, *dnmA*, *mnSOD* and *pimA* genes predicted to encode alternative oxidase, dynamin-related protein, manganese superoxide dismutase and LON protease, respectively, were genetically manipulated, and a number of physiological, morphological and developmental features of the mutants were examined. Altogether 82 phenotypes were observable, which confirmed the importance of mitochondrial proteins in the maintenance of cellular and mitochondrial morphology and functions as well as in sexual and asexual developments (Fig. 5; Supplementary Note S1). Importantly, a decreased production and viability of conidiospores were observed commonly in all gene deletion mutants (Fig. 4; Supplementary Table S4), suggesting that these nucleus-encoded genes play a role in sporogenesis and the integrity of conidia. On the other hand, many phenotypes were sporadic and randomly distributed among the gene deletion and overexpression strains (Supplementary Table S4).




Figure 3. Visualization and characterization of mitochondria. Panel A: Three-channel confocal Z-stack images were taken where the green channel shows mitochondria (Mt) stained by MitoTracker Green (a), the red channel visualizes the superoxide (O₂^{•-}) indicator dihydroethidium (b), while the blue channel (chitin staining by Calcofluor White) shows structure of the hypha and the boundaries for each segment (c)^{23,60}. Panel B: Visualizing mitochondria-identifying these cell organelles. (d) Mitochondria stained by MitoTracker Green. (e) 2nd wavelet level of the 2D SWT of the same frame. (f) Contours of ROI (solid yellow line) and background (dashed yellow line) selected manually and mitochondria (solid white line) as provided by the automatic segmentation. (g) Identified mitochondria (cyan colored areas) superimposed to the denoised image. Panel C: Comparing mitochondrial morphology in the second hyphal segment of the THS30.3 control strain (h–j), $\Delta dnmA$ (k–m) and *pimAOE* strains (n–p). Markings on subpanels (i, l and o) correspond to those on subpanel (f). Markings on subpanels (j, m and p) correspond to those on subpanel (g). Scale bar: 10 μm. Panels D and E: Comparison of the volumetric ratio and size of mitochondria in the THS30.3 control, and the $\Delta dnmA$ and *pimAOE* mutant strains. Mean ± SD values calculated from four independent experiments are presented. Statistically significant differences determined by the Student's t test are marked with asterisks: **p < 1%, ***p < 0.1%. No further phenotypes concerning mitochondrial morphology were observed (Supplementary Fig. S4).

We then compared the phenotypes of relevant gene deletion and overexpression strains in *P. anserina*^{6,14,15,18,19,21} and *A. nidulans* as summarized in Table 3. Up to 26 features were chosen and the respective




Figure 4. Cleistothecia and conidiospore productions and viability of conidia. Part A: Cleistothecia productions observed after 5 (■) and 8 d (▨) incubations. Both the deletion and overexpression of *aodA* stimulated meanwhile the overexpression of *mnSOD* and of *pimA* hindered the formation of fruiting bodies. Photographs show 8 d cleistothecia of the control (THS30.3) and $\Delta pimA$ strains. Note that cleistothecia of the $\Delta pimA$ strain were sterile. (Scale bars: 200 μ m). Part B: Conidiospore productions. All gene deletion strains and the *pimA*OE overexpression strain produced less conidiospores than the control strain. Part C: Heat stress sensitivity of the $\Delta mnSOD$ conidia. Conidia without heat treatment (50 °C for 10 min) were used as control³². As shown in Supplementary Fig. S5, only conidia of the $\Delta mnSOD$ strain were sensitive to heat stress. Part D: Decreasing asexual spore viabilities during storage at 4 °C. Symbols represent the following strains: ◆, THS30.3 (control strain), ■, $\Delta aodA$, ▲, $\Delta dnmA$, ●, $\Delta mnSOD$ and ■, $\Delta pimA$. The spore viabilities of all gene deletion strains were significantly lower ($p < 5\%$ at 6 and 12 d incubation times) than that of the THS30.3 control strain. Note that no significant decreases in the conidiospore viabilities were recorded in the gene overexpression strains and, hence, these data are not shown here for clarity. In Parts A–C, mean \pm SD values calculated from three independent experiments are presented. Significant differences determined by the Student's t test are marked with asterisks: ** $p < 1\%$, *** $p < 0.1\%$. In Part D, per cent decreases in the spore viabilities (mean \pm SD values calculated from three independent experiments) are presented.

phenotypes were placed in one of the following four groups: (i) the same or similar phenotypes were described in the relevant *P. anserina* and the *A. nidulans* mutants, (ii) phenotype was observed in *P. anserina* but not in the corresponding *A. nidulans* mutant, (iii) phenotype was observed in *A. nidulans* but not in the corresponding *P. anserina* mutant and (iv) opposite phenotypes were found in the *P. anserina* and *A. nidulans* mutants (Table 3). The distribution of the features between these groups (6:12:7:1) showed that approximately the half (46.1%) of the phenotypes reported in the *P. anserina* strains were not observable in the corresponding *A. nidulans* mutants (Table 3). These findings clearly indicated that genetic modifications of these four genes have less impact on the




Figure 5. Summary of functions of AodA, DnmA, MnSOD and PimA in *A. nidulans*. Major functions revealed by the modulation of mitochondrial morphology and functions *via* the deletion and overexpression of the genes *aodA*, *dnmA*, *mnSOD* and *pimA* are summarized. Schematic mitochondrial, hyphal, cleistotheceal and conidiophore structures are shown. In the dividing mitochondria, hypothetical protein structures and shapes are presented, which are based on homologous proteins characterized in various species (© Ibolya Pócsi; for further details see Supplementary Note S1).

cellular physiology and lifespan of *A. nidulans* compared to *P. anserina*^{6,14,15,18,19,21} (Table 3). Furthermore, eight phenotypes (30.8%) were either present only in *A. nidulans* or opposite to that recorded in *P. anserina* (Table 3), shedding light on further differences in their roles between the two ascomycetous filamentous fungi. These suggest a clear limit in using *P. anserina* as a model for predicting mitochondrial functions, life-span and longevity. This implies that the evolutionary divergence between *P. anserina* and other fungal species of interest²⁰ makes the application of *Podospora*-based models on ageing and senescence^{10,11,22} quite difficult when important questions like strain degeneration processes and the control of lifespan are addressed in industrially and biomedically important filamentous fungi like the Aspergilli²³.

Because only nuclear-encoded genes thought to have significant impacts on mitochondrial physiology and morphology were selected for genetic studies in *A. nidulans*, and their orthologs have been functionally characterized in *P. anserina*^{6,14,15,18,19,21}, one may raise the important question that how the evolutionary distance between these two species influenced the unevenly emerging phenotypes of the mutants (Table 3). One option could be the divergent evolution of the mitochondrial gene contents through distinct species-specific gene transfer events to the nucleus³⁵. However, the *P. anserina* mitochondrial genome harbors the same protein-coding potential except for the ATPase 9 gene as found before in *Neurospora crassa* and *A. nidulans*³⁶. Therefore, we should consider another option, the significant differences in the size, organization, stability and intraspecies polymorphism of the

Parameters tested	<i>Podospora anserina</i> strains							
	<i>aox::HPH</i> ⁶	<i>GPD-AOX</i> ⁶	<i>PaDnm1::ble</i> ¹⁸	<i>PaDnm1_Ex</i> ²¹	$\Delta PaSOD3$ ¹⁴	<i>PaSOD3_OEx</i> ¹⁴	$\Delta PaLON$ ¹⁹	<i>PaLON_Oex</i> ¹⁵
Growth in surface culture	0	0	- (on cornmeal medium) \blacklozenge	n.d.	0	- \blacksquare	0	0
Oxidative stress tolerance	0	0	- (to H ₂ O ₂) \blacksquare	n.d.	- (to paraquat) \blacklozenge	- (to paraquat and H ₂ O ₂) \blacksquare	- (to H ₂ O ₂) \blacksquare	+ (to H ₂ O ₂) \blacksquare
RS production	0 \blacktriangle	+ \blacksquare	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Cell death	0	0	- (resistant to etoposide) \blacksquare	+ (on PASM +1 % glucose) \blacksquare	0 \blacktriangle	+ \blacksquare	+ \blacksquare	- \blacksquare
Total respiration rate	0 \blacktriangle	0	n.d.	n.d.	n.d.	n.d.	n.d.	+ \bullet
Alternative respiration rate	no AOX protein \blacklozenge	+ \blacklozenge	n.d.	n.d.	n.d.	n.d.	n.d.	0
Mitochondrial morphology	n.d.	n.d.	filamentous structure (but not in old culture) \blacklozenge	fragmented mitochondria \blacksquare	n.d.	n.d.	n.d.	0 \blacktriangle
Sexual development	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	less ascospores \blacklozenge	n.d.
Number of fruiting bodies	0 (female fertility) \blacktriangle	0 (female fertility) \blacktriangle	0 (female fertility)	n.d.	n.d.	n.d.	0 (female fertility)	0 (female fertility) \blacktriangle
Parameters tested	<i>Aspergillus nidulans</i> strains							
	$\Delta aodA$	<i>aodAOE</i>	$\Delta dnmA$	<i>dnmAOE</i>	$\Delta mnSOD$	<i>mnSODOE</i>	$\Delta pimA$	<i>pimAOE</i>
Growth in surface culture	0	0	- \blacklozenge	0	0	0 \blacksquare	0	0
Oxidative stress tolerance	0	0	0 \blacksquare	0	- (to MSB) \blacklozenge	0 \blacksquare	0 \blacksquare	0 \blacksquare
RS production [§]	+ \blacktriangle	0 \blacksquare	0	-	-	-	+	+
Cell death	0	0	0 \blacksquare	0 \blacksquare	- (to PAF) \blacktriangle	0 \blacksquare	0 \blacksquare	0 \blacksquare
Total respiration rate [§]	- \blacktriangle	0	+	0	0	0	0	- \bullet
Alternative respiration rate [§]	- \blacklozenge	+ \blacklozenge	+	0	0	0	+	0
Mitochondrial morphology	0	0	higher volumetric ratio and number \blacklozenge	0 \blacksquare	0	0	0	higher volumetric ratio and number \blacktriangle
Sexual development	0	0	0	0	0	0	no ascospores \blacklozenge	0
Number of fruiting bodies	+ \blacktriangle	+ \blacktriangle	0	0	0	-	0	- \blacktriangle

Table 3. Comparison of the phenotypes of relevant *Podospora anserina* and *Aspergillus nidulans* mutant strains. [§]Reactive species (RS) productions^{24,27} and respiration rates²⁹ measured in growing cultures were taken into consideration. \blacklozenge The same or similar phenotypes were described in the relevant *P. anserina* and the *A. nidulans* mutants. \blacksquare Phenotype was observed in *P. anserina* but not in the appropriate *A. nidulans* mutant. \blacktriangle Phenotype was observed in *A. nidulans* but not in the appropriate *P. anserina* mutant. \bullet Opposite phenotypes in the relevant *P. anserina* and the *A. nidulans* mutants. 0 stands for no alteration meanwhile + and - indicate positive or negative alterations in comparison to appropriate controls; n.d.-not determined.

P. anserina and *A. nidulans* mtDNAs^{5,36-43}. We speculate that the primary reason for the phenotypic differences between *P. anserina*^{6,14,15,18,19} and *A. nidulans* would be due to the intrinsic instability of the *P. anserina* mtDNA.

Importantly, Li *et al.*²³ found a causal connection between the spontaneous formation of fluffy sectors and mitochondrial dysfunctions in *A. nidulans* cultures maintained on potato dextrose agar. Although a linkage of culture degeneration to programmed cell death was demonstrated, the fluffy sector culture could be maintained further without any growth arrest or any sign of *P. anserina*-like senescence probably owing to compensatory mechanisms like the induction of stress defence elements including heat shock proteins, anti-apoptotic factors and DNA repair proteins²³.

As mentioned above, these interesting phenotypes have also shed light on the remarkable importance of mitochondria in the developmental processes in filamentous fungi, especially concerning conidiogenesis and the preservation of the viability of conidia under storage at 4 °C (Fig. 4; Table 3). Further basic questions can include the elucidation of how the various upstream and downstream regulatory elements controlling development, stress, and metabolism in *A. nidulans*, such as ‘velvet’ proteins^{44,45}, SskA and SrrA response regulators^{32,33}, SakA mitogen-activated protein kinase⁴⁶ and AtfA transcription factor^{47,48} contribute to the quality control of mitochondria^{4,7}, and understanding how changes in the quality of mitochondrial functions will be sensed and how these signals are channeled towards regulatory elements by specific signaling pathways^{49,50}.

It is worth noting that some observations presented in this paper also seem to be of great value from the biotechnologist’s point of view. For example, the deletion of *aodA* resulted in reduced production of proteinase and ST in *A. nidulans* (Table 2, Fig. 1D; Supplementary Figs S2 and S3), which may be suitable for heterologous protein production⁵¹. The *A. nidulans* $\Delta aodA$ mutant maintains satisfactory antioxidative defence even in the presence of endogenous oxidative stress (Supplementary Fig. S1), further suggesting the utility of novel $\Delta aodA$ -based *Aspergillus* industrial platforms for heterologous protein production. Interestingly, ST production was also decreased significantly *via* deletions of *dnmA* and *pimA* as well as through the overexpression of *aodA* (Table 2).

It is also remarkable that the overexpression of *pimA* resulted in the emergence of a new, heavy metal (Cd²⁺) tolerant *A. nidulans* strain, *pimAOE* (Fig. 1B, Supplementary Fig. S1), which could be beneficial when *Aspergillus*-based environmental technologies are elaborated⁵². Importantly, both asexual and sexual developments responded to the manipulation of the selected genes in many cases, and formation of cleistothecia was positively affected by either deletion or overexpression of *aodA* (Fig. 4, Supplementary Fig. S6). These observations on asexual and sexual sporulations may also be important when the potential industrial applications of genetically engineered *Aspergillus* strains are considered⁵³.

Future studies are definitely needed to answer the intriguing questions whether these observations are limited to *A. nidulans* or are applicable and exploitable in other *Aspergillus* as well, *e.g.* in the biological control of toxigenic fungi²⁸, in the bioremediation of heavy metal polluted areas⁵² in combating *Aspergillus* strain degenerations²³.

To sum it up, the manipulation of four nuclear-encoded genes thought to be important for mitochondrial functions shed light on a number of physiological, morphological and developmental roles played by these genes in *A. nidulans* (Fig. 5; Supplementary Note S1). The remarkable differences observed between the phenotypes of the relevant *A. nidulans* and *P. anserina* mutants (Table 3; Supplementary Table 3) suggest that *A. nidulans*-based platforms should also be considered in future studies aiming at mitochondria-related phenomena like ageing and programmed cell death, complementarily to the well-established *P. anserina*-based and *S. cerevisiae*-based platforms^{1–4}.

Methods

Culture media, construction of gene deletion and overexpression strains. For cultivation of *A. nidulans* strains, a standard complete medium containing 0.5% yeast extract²⁹ and minimal nitrate medium (MNM) were used with appropriate nutritional supplements⁵⁴.

Target genes in the *A. nidulans* genome were identified and confirmed by based on annotation data in the Broad Institute *Aspergillus* Comparative Database as well as by using *S. cerevisiae* and *P. anserina* homologues in protein vs. protein queries in NCBI BLAST (blastp; http://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastp&PAGE_TYPE=BlastSearch&LINK_LOC=blasthome) and in the *P. anserina* Genome Project website (blastp; <http://podospora.igmors.u-psud.fr/blast.php>). The selected target genes with locus IDs and homologies to relevant *P. anserina* genes were *aodA* (encoding mitochondrial alternative oxidase, AN2099.2²⁵, Expect value: 4e-123 when compared to *P. anserina* Pa_3_1710 coding for alternative oxidase, which is identical to PaAox⁶), *dnmA* (hypothetical protein gene, AN8874.2, Expect value: 0.0 to Pa_1_12670 coding for dynamin-related protein 1 DNMI involved in mitochondrial fission, which is identical to PaDnm1¹⁸), *mnSOD* (putative manganese superoxide dismutase gene, AN5577.2, Expect value: 1E-102 to *P. anserina* Pa_5_1740 encoding superoxide dismutase, which is identical to PdSod3⁵⁵) and *pimA* (hypothetical protein gene, AN6193.2, Expect value: 0.0 to Pa_3_4170 encoding putative Lon protease, mitochondrial precursor, which is identical to PaLON¹⁵). To construct deletion mutants, the double-joint PCR (DJ-PCR) method of Yu *et al.*⁵⁶ was used with primers listed in Supplementary Table S2. The amplified deletion cassettes were used to transform RJMP1.59 strain using the Vinoflow FCE lysing enzyme (Novo Nordisk)⁵⁷. Single copy transformants were selected after Southern blot analysis and crossed with rRAW16 to get prototroph strains. All progenies of the independent crosses were proved to be single copy deleted mutants by Southern analyses⁵⁶. The genotypes of the studied *A. nidulans* strains are summarized in Supplementary Table S3.

To generate overexpression (OE) mutants, individual ORFs were amplified with the primers presented in Supplementary Table S2. The amplicons were digested with restriction enzymes as indicated in Supplementary Table S2, and ligated between the *niiA* promoter and the *trpC* terminator in pHS11 (Park and Yu, unpublished data). The final plasmid containing ¾ *pyroA* gene was expected to have single copy integration at the *pyroA* locus, which together with the overexpression of each gene was confirmed by PCR and Northern blot analyses, respectively⁵⁸.

In all assays, the prototrophic THS30.3 strain (for genotype see Supplementary Table S3) was used as the control strain.

Stress sensitivity assays, apoptotic cell death of mutants, conidiospore heat stress-sensitivity and viability tests.

To determine the stress sensitivity of mutants, the agar plate assays of Hagiwara *et al.*³² were used with slight modifications. Freshly grown (6 days) conidia (10⁵ suspended in 5 µl aliquots of physiological saline –0.01% Tween 80) were spotted on MNM agar plates in Petri dishes⁵⁴ containing the following

stress generating agents: oxidative stress: 6.0 mM H₂O₂, 0.08 mM menadione sodium bisulphite (MSB), 2.0 mM diamide, 0.08 mM tBOOH, osmotic stress: 1.5 M KCl, 1.5 M NaCl, 2.0 M sorbitol, cell wall stress: 75 µg/ml CongoRed, heavy metal stress: 300 µM CdCl₂. Stress plates were incubated for 5 days at 37 °C²⁴.

The small molecular mass antifungal protein produced by *Penicillium chrysogenum* (PAF) causes apoptosis-like cell death in sensitive filamentous fungi like *A. nidulans*²⁶. In these assays, PAF supplemented MNM agar prepared in 12-well tissue culture plates were point-inoculated with 2×10^3 conidia, and were incubated at 37 °C for 72 h²⁶.

Determination of vitality and physiological parameters in submerged cultures of *A. nidulans*. *A. nidulans* strains were pre-grown in Erlenmeyer flasks (500 ml) containing 100 ml complete medium (pH 6.5), where culture media were inoculated always with 5×10^7 spores and incubated for 16 h at 37 °C and at 3.7 Hz shaking frequency. Following that, mycelia were collected by filtration on sintered glass, washed with sterile distilled water and transferred into MNM (100 ml aliquots) containing either 2% or no glucose, and were incubated at 37 °C and 3.7 Hz shaking frequency for 4, 10 (with extra glucose) or 24 h (without any extra glucose). Growth of the mutants was characterized by increases in the dry cell mass (DCM). To determine the vitalities of the carbon-starving cultures of the tested strains, the cultures were washed to minimal medium containing 1% glucose and increases in the dry cell mass (DCM) were recorded²⁴.

The intracellular reactive species (RS) levels were characterized by the formation of 2',7'-dichlorofluorescein (DCF) from 2',7'-dichlorofluorescein diacetate. RS includes all reactive oxygen species (ROS) and all reactive nitrogen species, which are able to oxidize 2',7'-dichlorofluorescein to DCF^{24,27}. At the incubation times tested, 10 µmol/ml 2',7'-dichlorofluorescein diacetate was added to 20 ml aliquots of the cultures, and after incubating further for 1 h in 100 ml culture flasks, the mycelia were harvested by centrifugation. The production of DCF was determined spectrofluorimetrically^{24,59}.

Changes in the specific activities of certain antioxidant enzymes were also recorded in separate experiments. In these cases, cell-free extracts were prepared by X-pressing and centrifugation⁵⁹. Specific catalase, glutathione peroxidase (GPx), glutathione reductase (GR), superoxide dismutase (SOD) activities were measured as before⁵⁹.

Respiration was measured in an Oxigraph O₂ electrode (Hansatech, UK) at 37 °C. The cytochrome c-dependent pathway and alternative oxidase respiration was inhibited with 1.0 mM KCN and 0.25 mM SHAM, respectively²⁹.

Proteinase activities of carbon-starving (24 h starvation) cultures were measured using azocasein as substrate³⁰.

Prior to secondary metabolite analysis, mycelial mass collected from submerged cultures were lyophilized and resuspended in 70% v/v acetone (15 mg freeze-dried mycelium in 200 µl solvent)²⁴. Following that, secondary metabolites in 25 µl aliquots of extracts were analyzed on silica gel TLC plates using sterigmatocystin (ST) standard.

Morphological studies on mitochondria. Fungal conidia (10^3 to 10^4 ml⁻¹) were grown in complete medium³⁰ on coverslips for 20–24 h and washed in MNM before loading with 30 nM MitoTracker Green for 30 min, 2.5 nM dihydroethidium for 20 min and 2.5 nM Calcofluor White (CFW) for 5 min to visualize mitochondria, intracellular superoxide radicals and chitin component of the cell wall, respectively. Then the coverslips were mounted on the microscope chamber and images were obtained using a laser scanning microscope LSM 510 Meta (Zeiss, Jena, Germany)^{23,31}.

Three-channel confocal Z-stack images were taken where the green channel showed mitochondria stained by MitoTracker Green, the red channel visualized the superoxide indicator dihydroethidium, while the blue channel (Calcofluor White staining) showed structure of the hypha and the boundaries for each segment (Fig. 3A).

In order to remove high-frequency noise, 2D stationary wavelet transform (SWT) based denoising was performed⁶⁰ with delta parameter of 0.7. A region of interest (ROI) was manually marked on the Z-frame showing most of the hypha in focus respecting the boundaries of a single hyphal segment (Fig. 3B–f; dashed yellow line). A cell-free background (Fig. 3B–f; dashed white line) was also selected and the average intensity in the region was subtracted from all intensity values on the frame. We selected the region between the first and second septa behind the tip. Since the dye used for the green channel stained mitochondria, every pixel within the ROI either belonged to a mitochondrion (high intensity pixels) or not (low intensity pixels). Segmentation was done on the W2 level of the 2D SWT of the analyzed frame (Fig. 3B–e) using a noise level related threshold value optimized for every record. Actual threshold values varied between 0.08–0.20 and pixels above this threshold were marked as mitochondria (Fig. 3B–g). The volumetric ratio of mitochondria within the hyphal segment is given as the number of high intensity pixels compared to the total number of pixels within the selected ROI. Contiguous areas of high intensity pixels were marked as mitochondria (Fig. 3B–f; solid white lines) and their sizes were calculated based on the number of pixels. The average of three consecutive Z-stacks was taken into account. The diameter of the hypha was measured in the blue channel (Calcofluor White staining) by the LSM Image Browser (Zeiss, Jena, Germany) program using the segments where the mitochondrial structure was examined.

Sexual and asexual developments. To induce cleistothecium formation, 6 d old conidia were plated in top agar at 1×10^5 conidia/plate density and were incubated at 37 °C. After 24 h incubation, plates were sealed with Parafilm and samples were taken with a cork borer every day between 3–12 d of incubation, and cleistothecia/cm² were counted under a dissection microscope and cleistothecia/cm² values were calculated⁴⁶.

The conidiospore forming capabilities of the *A. nidulans* strains were determined as published by Vargas-Pérez *et al.*³³. Briefly, conidia (10^5) of the mutant and control strains were spotted onto MNM agar plates as described above, and were incubated and were allowed to sporulate at 37 °C for 5 days. Conidia were harvested by washing,

counted in a Burkner chamber and spore numbers were expressed as number/cm² of colony surface. The areas of the colony surfaces were calculated using photographs.

To test the heat sensitivity of asexual spores, conidia were harvested from 6 days old colonies and suspended in physiological saline—0.01% Tween 80²⁴. Conidia in 10⁵/ml concentration were incubated at 50 °C for 10 min and, following that, were diluted and spread on MNM agar plates. The numbers of colonies representing successfully germinated conidia were counted after incubation for 2 days at 37 °C. Conidia without any heat treatment were used as reference. For viability test, conidia were stored at 4 °C and germination rates were determined after 3, 6 and 12 d storages as described above³².

Statistical analysis of experimental data. All experiments were performed in three independent sets with the exception of the mitochondrial volumetric ratio and size determinations where four independent experiments were carried out. Mean±SD values are presented, statistical significances were calculated using Student's t-test, and *p*-values less than 0.05 were considered as statistically significant.

References

- Osiewacz, H. D. & Scheckhuber, C. Q. Impact of ROS on ageing of two fungal model systems: *Saccharomyces cerevisiae* and *Podospora anserina*. *Free Rad Res* **40**, 1350–1358 (2006).
- Ugidos, A., Nyström, T. & Caballero, A. Perspectives on the mitochondrial etiology of replicative aging in yeast. *Exp Gerontol* **45**, 512–515 (2010).
- Osiewacz, H. D. Mitochondrial quality control in aging and lifespan control of the fungal aging model *Podospora anserina*. *Biochem Soc Trans* **39**, 1488–1492 (2011).
- Osiewacz, H. D. & Bernhardt, D. Mitochondrial quality control: impact on aging and life span—a mini-review. *Gerontology* **59**, 413–420 (2013).
- Begel, O., Boulay, J., Albert, B., Dufour, E. & Sainsard-Chanet, A. Mitochondrial group II introns, cytochrome c oxidase, and senescence in *Podospora anserina*. *Mol Cell Biol* **19**, 4093–4100 (1999).
- Lorin, S. *et al.* Overexpression of the alternative oxidase restores senescence and fertility in a long-lived respiration-deficient mutant of *Podospora anserina*. *Mol Microbiol* **42**, 1259–1267 (2001).
- Bernhardt, D., Hamann, A. & Osiewacz, H. D. The role of mitochondria in fungal aging. *Curr. Opin. Microbiol.* **22**, 1–7 (2014).
- Bernhardt, D., Müller, M., Reichert, A. S. & Osiewacz, H. D. Simultaneous impairment of mitochondrial fission and fusion reduces mitophagy and shortens replicative lifespan. *Sci. Rep.* **5**, 7885 (2015).
- Sellem, C. H., Marsy, S., Boivin, A., Lemaire, C. & Sainsard-Chanet, A. A mutation in the gene encoding cytochrome c1 leads to a decreased ROS content and to a long-lived phenotype in the filamentous fungus *Podospora anserina*. *Fungal Genet Biol* **44**, 648–658 (2007).
- Scheckhuber, C. Q. *et al.* Alternative oxidase dependent respiration leads to an increased mitochondrial content in two long-lived mutants of the aging model *Podospora anserina*. *PLoS One* **6**, e16620 (2011).
- Gredilla, R., Grief, J. & Osiewacz, H. D. Mitochondrial free radical generation and lifespan control in the fungal ageing model *Podospora anserina*. *Exp Gerontol* **41**, 439–447 (2006).
- Longo, V. D., Gralla, E. B. & Valentine, J. S. Superoxide dismutase activity is essential for stationary phase survival in *Saccharomyces cerevisiae*. Mitochondrial production of toxic oxygen species *in vivo*. *J Biol Chem* **271**, 12275–12280 (1996).
- Harris, N. *et al.* Mnsod overexpression extends the yeast chronological (G(0)) life span but acts independently of Sir2p histone deacetylase to shorten the replicative life span of dividing cells. *Free Radic Biol Med* **34**, 1599–1606 (2003).
- Zintel, S., Schwitalla, D., Luce, K., Hamann, A. & Osiewacz, H. D. Increasing mitochondrial superoxide dismutase abundance leads to impairments in protein quality control and ROS scavenging systems and to lifespan shortening. *Exp Gerontol* **45**, 525–532 (2010).
- Luce, K. & Osiewacz, H. D. Increasing organismal healthspan by enhancing mitochondrial protein quality control. *Nat Cell Biol* **11**, 852–858 (2009).
- Westermann, B. Mitochondrial fusion and fission in cell life and death. *Nat Rev Mol Cell Biol* **11**, 872–884 (2010).
- Otsuga, D. *et al.* The dynamin-related GTPase, Dnm1p, controls mitochondrial morphology in yeast. *J Cell Biol* **143**, 333–349 (1998).
- Scheckhuber, C. Q. *et al.* Reducing mitochondrial fission results in increased life span and fitness of two fungal ageing models. *Nat Cell Biol* **9**, 99–105 (2007).
- Adam, C. *et al.* Biological roles of the *Podospora anserina* mitochondrial Lon protease and the importance of its N-domain. *PLoS One* **7**, e38138 (2012).
- Wang, H., Xu, Z., Gao, L. & Hao, B. A fungal phylogeny based on 82 complete genomes using the composition vector method. *BMC Evol Biol* **9**, 195 (2009).
- Scheckhuber, C. Q., Rödel, E. & Wüsthube, J. Regulation of mitochondrial dynamics—characterization of fusion and fission genes in the ascomycete *Podospora anserina*. *Biotechnol J* **6**, 781–790 (2008).
- Scheckhuber, C. Q., Mitterbauer, R. & Osiewacz, H. D. Molecular basis of and interference into degenerative processes in fungi: potential relevance for improving biotechnological performance of microorganisms. *Appl Microbiol Biotechnol* **85**, 27–35 (2009).
- Li, L. *et al.* Linkage of oxidative stress and mitochondrial dysfunctions to spontaneous culture degeneration in *Aspergillus nidulans*. *Mol Cell Proteomics* **13**, 449–461 (2014).
- Yin, W. B. *et al.* bZIP transcription factors affecting secondary metabolism, sexual development and stress responses in *Aspergillus nidulans*. *Microbiology-UK* **159**, 77–88 (2013).
- Suzuki, Y., Murray, S. L., Wong, K. H., Davis, M. A. & Hynes, M. J. Reprogramming of carbon metabolism by the transcriptional activators AcuK and AcuM in *Aspergillus nidulans*. *Mol Microbiol* **84**, 942–964 (2012).
- Hegedűs, N. *et al.* The small molecular mass antifungal protein of *Penicillium chrysogenum*—a mechanism of action oriented review. *J Basic Microbiol* **51**, 561–571 (2011).
- Halliwell, B. & Gutteridge, J. M. C. *Free Radicals in Biology and Medicine* Ch. 5, 268–330 (Oxford University Press, 2007).
- Reverberi, M., Ricelli, A., Zjalic, S., Fabbri, A. A. & Fanelli, C. Natural functions of mycotoxins and control of their biosynthesis in fungi. *Appl. Microbiol. Biotechnol.* **87**, 899–911.
- Emri, T., Molnár, Z., Pusztahelyi, T. & Pócsi, I. Physiological and morphological changes in autolyzing *Aspergillus nidulans* cultures. *Folia Microbiol* **49**, 277–284 (2004).
- Szilágyi, M. *et al.* Extracellular proteinase formation in carbon starving *Aspergillus nidulans* cultures—physiological function and regulation. *J. Basic Microbiol.* **51**, 625–634 (2011).
- Lam, Y. T., Aung-Htut, M. T., Lim, Y. L., Yang, H. & Dawes, I. W. Changes in reactive oxygen species begin early during replicative aging of *Saccharomyces cerevisiae* cells. *Free Rad Biol Med* **50**, 963–970 (2011).
- Hagiwara, D. *et al.* The SskA and SrrA response regulators are implicated in oxidative stress responses of hyphae and asexual spores in the phosphorelay signaling network of *Aspergillus nidulans*. *Biosci Biotechnol Biochem* **71**, 1003–1014 (2007).

33. Vargas-Pérez, I., Sánchez, O., Kawasaki, L., Georgellis, D. & Aguirre, J. Response regulators SrrA and SskA are central components of a phosphorelay system involved in stress signal transduction and asexual sporulation in *Aspergillus nidulans*. *Eukaryot Cell* **6**, 1570–1583 (2007).
34. Gray, M. W., Burger, G. & Lang, B. F. Mitochondrial evolution. *Science* **283**, 1476–1481 (1999).
35. Adams, K. L. & Palmer, J. D. Evolution of mitochondrial gene content: gene loss and transfer to the nucleus. *Mol Phylogenet Evol* **29**, 380–395 (2003).
36. Cummings, D. J., McNally, K. L., Domenico, J. M. & Matsuura, E. T. The complete DNA sequence of the mitochondrial genome of *Podospora anserina*. *Curr Genet* **17**, 375–402 (1990).
37. Brown, T. A., Waring, R. B., Scaccocchio, C. & Davies, R. W. The *Aspergillus nidulans* mitochondrial genome. *Curr Genet* **9**, 113–117 (1985).
38. Joardar, V. *et al.* Sequencing of mitochondrial genomes of nine *Aspergillus* and *Penicillium* species identifies mobile introns and accessory genes as main sources of genome size variability. *BMC Genomics* **13**, 698 (2012).
39. Croft, J. H. Genetic variation and evolution in *Aspergillus* in *Evolutionary Biology of the Fungi* (eds. Rayner, A. D. M., Brasier, C. M. & Moore, D.) 311–323 (Cambridge University Press, 1987).
40. Hamari, Z., Juhász, A. & Kevei, F. Role of mobile introns in mitochondrial genome diversity of fungi (a mini review). *Acta Microbiol Immunol Hung* **49**, 331–335 (2002).
41. Van Diepeningen, A. D. *et al.* Mitochondrial recombination increases with age in *Podospora anserina*. *Mech Ageing Dev* **131**, 315–322 (2010).
42. Maas, M. F., de Boer, H. J., Debets, A. J. & Hoekstra, R. F. The mitochondrial plasmid pAL2-1 reduces calorie restriction mediated life span extension in the filamentous fungus *Podospora anserina*. *Fungal Genet Biol* **41**, 865–871 (2004).
43. Lorin, S., Dufour, E. & Sinsard-Chanet, A. Mitochondrial metabolism and ageing in the filamentous fungus *Podospora anserina*. *Biochim Biophys Acta* **1757**, 604–610 (2016).
44. Bayram, O. & Braus, G. H. Coordination of secondary metabolism and development in fungi: the velvet family of regulatory proteins. *FEMS Microbiol Rev* **36**, 1–24 (2012).
45. Park, H. S. & Yu, J. H. Genetic control of asexual sporulation in filamentous fungi. *Curr Opin Microbiol* **15**, 669–677 (2012).
46. Kawasaki, L., Sánchez, O., Shiozaki, K. & Aguirre, J. SakA MAP kinase is involved in stress signal transduction, sexual development and spore viability in *Aspergillus nidulans*. *Mol Microbiol* **45**, 1153–1163 (2002).
47. Hagiwara, D., Asano, Y., Yamashino, T. & Mizuno, T. Characterization of bZip-type transcription factor AtfA with reference to stress responses of conidia of *Aspergillus nidulans*. *Biosci Biotechnol Biochem* **72**, 2756–2760 (2008).
48. Lara-Rojas, F., Sánchez, O., Kawasaki, L. & Aguirre, J. *Aspergillus nidulans* transcription factor AtfA interacts with the MAPK SakA to regulate general stress responses, development and spore functions. *Mol Microbiol* **80**, 436–454 (2011).
49. Jaimes-Arroyo, R. *et al.* The SrkA kinase is part of the SakA MAPK interactome and regulates stress responses and development in *Aspergillus nidulans*. *Eukaryot Cell* **14**, 495–510 (2015).
50. Pusztahelyi, T. & Pócsi, I. Functions, cooperation and interplays of the vegetative growth signaling pathways in the aspergilli. *J Mycol* **2013**, Article ID 832521 (2013).
51. van den Hombergh, J. P., van de Vondervoort, P. J., Fraissinet-Tachet, L. & Visser, J. *Aspergillus* as a host for heterologous protein production: the problem of proteases. *Trends Biotechnol* **15**, 256–263 (1997).
52. Chakraborty, S., Mukherjee, A., Khuda-Bukhsh, A. R. & Das, T. K. Cadmium-induced oxidative stress tolerance in cadmium resistant *Aspergillus foetidus*: its possible role in cadmium bioremediation. *Ecotoxicol Environ Saf* **106**, 46–53 (2014).
53. Ishi, K., Maruyama, J., Juvvadi, P. R., Nakajima, H. & Kitamoto, K. Visualizing nuclear migration during conidiophore development in *Aspergillus nidulans* and *Aspergillus oryzae*: multinucleation of conidia occurs through direct migration of plural nuclei from phialides and confers greater viability and early germination in *Aspergillus oryzae*. *Biosci Biotechnol Biochem* **69**, 747–754 (2005).
54. Barratt, R. W., Johnson, G. B. & Ogata, W. N. Wild-type and mutant stocks of *Aspergillus nidulans*. *Genetics* **52**, 233–246 (1965).
55. Grimm, C., Böhl, L. & Osiewacz, H. D. Overexpression of Pa_1_10620 encoding a mitochondrial *Podospora anserina* protein with homology to superoxide dismutases and ribosomal proteins leads to lifespan extension. *Curr Genet* **61**, 73–86 (2015).
56. Yu, J. H. *et al.* Double-joint PCR: a PCR-based molecular tool for gene manipulations in filamentous fungi. *Fungal Genet Biol* **41**, 973–981 (2004).
57. Szczytyk, E. *et al.* Fusion PCR and gene targeting in *Aspergillus nidulans*. *Nat Protoc* **1**, 3111–3120 (2006).
58. Kwon, N. J., Shin, K. S. & Yu, J. H. Characterization of the developmental regulator FlbE in *Aspergillus fumigatus* and *Aspergillus nidulans*. *Fungal Genet Biol* **47**, 981–993 (2010).
59. Emri, T., Pócsi, I. & Szentirmai, A. Analysis of the oxidative stress response of *Penicillium chrysogenum* to menadione. *Free Radic Res* **30**, 125–132 (1999).
60. Szabó, L. Z., Vincze, J., Csernoch, L. & Szentesi, P. Improved spark and ember detection using stationary wavelet transforms. *J Theor Biol* **264**, 1279–1292 (2010).

Acknowledgements

The authors thank Mrs. Lászlóné Gábor Tóth and Ms. Melinda Takács for their valuable contributions to the experimental work, Drs. Sándor Kéki and Lajos Nagy for discussing and validating analytical data, and Ms. I. Pócsi for drawing Fig. 5. É.L. was a recipient of a Fulbright Research Fellowship to the UW-Madison. This work was primarily supported by the Hungarian Scientific Research Fund (OTKA K100464 for IP) and the SROP-4.2.2.B-15/1/KONV-2015-0001 project in Debrecen, which has been supported by the European Union, co-financed by the European Social Fund. The work was also supported by the National Science Foundation (IOS0950850 to JHY) and the Intelligent Synthetic Biology Center of Global Frontier Projects (2011-0031955 for JHY and NRF-2015M3A6A8065838 for KHH), funded by the Ministry of Education, Science and Technology, Republic of Korea.

Author Contributions

The study was performed at various locations and was conducted by the following senior researchers: genetic manipulations: J.H.Y., University of Wisconsin, Madison; confocal microscopy studies: L.Cs., University of Debrecen; physiological characterization of the strains with the exception of respiration measurements: I.P., University of Debrecen; respiration measurements: I.M., University of Debrecen. I.P. conceived the study, É.L., N.J.K., T.E., I.M., K.H.H., B.D., L.Cs., J.H.Y. and I.P. designed the experiments, É.L., H.P., V.O., B.D. and J.V. performed the experiments, and all authors took part in the evaluation and discussion of experimental data. The manuscript was written by I.P. with the contribution of É.L., B.D., L.Cs. and J.H.Y.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Leiter, É. *et al.* Characterization of the *aodA*, *dnmA*, *mnSOD* and *pimA* genes in *Aspergillus nidulans*. *Sci. Rep.* **6**, 20523; doi: 10.1038/srep20523 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>