

An acrylated isocyanonaphthalene based solvatochromic click reagent: Optical and biolabeling properties and quantum chemical modeling

Miklós Nagy¹, Dávid Rácz¹, Zsolt László Nagy¹, Tibor Nagy¹, Péter Pál Fehér², Mihály Purgel^{2,3}, Miklós Zsuga¹, Sándor Kéki^{1*}

¹*Department of Applied Chemistry, ²Department of Physical Chemistry, ³MTA-DE Homogeneous Catalysis and Reaction Mechanisms Research Group, University of Debrecen, H-4032 Debrecen, Hungary*

* Corresponding author: keki.sandor@science.unideb.hu, fax: +36 52 518662; H-4032 Debrecen, HUNGARY

Abstract

The optical and biolabeling properties of a novel molecule 1-(2-acryloyloxy-3-chloro-prop-1-yl)-amino-5-isocyanonaphthalene (ACAIN) is reported. In addition to being a real solvatochromic fluorophore it reacts quantitatively and rapidly with simple thiols in a thiol-ene click reaction. DFT calculations revealed a dark nonfluorescent state of ACAIN due to a close energy triplet state where electron transition can happen between the acrylic double bond and the aromatic core through an intramolecular hydrogen bond between the NH and C=O moieties. The hydrothiolation reaction is accompanied by a 1.5-19 fold increase in fluorescence intensity depending on the solvent used owing to the saturation of the acrylic group. The quantum yield and reactivity of the molecules were found to be largely dependent on the substituent of the acryl moiety.

The biolabeling properties were investigated in detail by fluorometry and ESI mass spectrometry using cysteine, KAC as a simple tripeptide and BSA as a model protein.

Keywords: Fluorescence, Click reaction, Solvatochromism, Density-functional calculations, Bovine serum albumin

Introduction

Over the past few decades, interest in the thiol-Michael addition reaction has increased dramatically, since it proceeds with high yield, stereoselectivity, rate, and is thermodynamically favored. [1-4] Thiol-ene additions can be extremely useful in the detection, localization, qualification and quantitation of intracellular thiols such as cysteine, glutathione or the free cysteine units of complex peptides. [5] The development of fluorescent probes that react through catalyzed (by base or nucleophile) Michael additions resulting in anti-Markovnikov addition products [6,7] has been in the focus in the last decades. The application of the reactive fluorophore is more favorable when the base molecule is non/less fluorescent and the thiol addition is accompanied by a substantial increase in fluorescent intensity. Such fluorophores include pyrazolines [5] BODIPY [8], α,β -unsaturated ketones [9] and nitroolefin-based coumarin [10].

Solvatochromic dyes are important members of smart materials. The wavelength (color) and intensity of their emitted fluorescent light is affected by the environment, particularly by its polarity. This feature can be applied advantageously in molecular biology, especially as biolabeling dyes for fluorescence microscopy, where the different parts of biomolecules can be easily differentiated from each other by the various colors of the emitted fluorescent light caused by the polarity changes around the fluorophores. This phenomenon is also suitable for the determination of the structure of biomacromolecules (eg. peptides, or locate binding sites of enzymes), or even for following their interactions by observing the local changes.[14]

Acryl(ate)s and methacryl(ate)s are easily reacted with thiols and can also be easily attached to different solvatochromic fluorophores. A good example is Acrylodan, a Prodan derivative bearing a reactive acryl moiety, which has a high affinity to thiols, therefore selectively labels proteins with free thiol groups [15,16] and has many important applications in bioanalytics [17-20]. After reacting with the $-SH$ group of biomolecules the conjugation of the acryloyl group with the close aromatic ring in Acrylodan will cease and fluorescence intensity increases dramatically. This phenomenon is useful in differentiating the unreacted dye from the labelled biomolecules and can be used in the construction of molecular switches. This intensity increase after the reaction does not necessarily require conjugation, it can also happen by the help of inter- or intramolecular H-bonding [21-24].


In addition to biolabeling thiol-ene reactions can be preferable routes for the synthesis of various types of polymers, such as end functionalized polymers, dendrimers, furthermore several controlled

radical polymerization techniques have also been developed based on this technique [11-13]. When a reactive fluorophore is used in the construction of smart networks, larger amounts are needed and consequently price and availability becomes serious issue. Most of these fluorophores are of complex structure or based on a pricey and hard to prepare backbone as is the case with Acrylodan. Therefore, there is a constant need for click reagents that are easily prepared from cheap and common starting materials using simple reactions, while the resulting fluorophore is solvatochromic, reacts completely with thiols, and has a dark and highly fluorescent state.

In this paper we report the design and preparation of a reactive solvatochromic dye (ACAIN) (and its methylated and chlorinated derivatives) by the modification of our recently developed molecule, 1-amino-5-isocyanonaphthalene (ICAN) [25] with only easily available reagents such as epichlorohydrin and acryloyl chloride. Quantum chemical calculations showed that when the acryloyl moiety is attached to the NH₂ group through an ethyl spacer unit, an intramolecular H-bond can form between the NH hydrogen and the carbonyl oxygen of the acryl group. This H-bonding will result in a dark state owing to a HOMO-LUMO electron transfer between the acrylic double bond and the naphthalene core and is largely solvent dependent. The acryl group reacts with thiols through a click reaction rapidly, followed by a significant increase of the emitted fluorescent light intensity because of the saturation of the double bond. ACAIN showed high affinity to thiol groups on a wide range of free thiol-containing molecules without the use of any catalyst. The kinetics of its biolabeling properties were investigated in detail using cysteine, lysyl-alanyl-cysteine (KAC), and bovine serum albumin (BSA) by fluorometry and ESI mass spectrometry. Based on density functional (DFT) calculations a model was developed for the description of the unique fluorescent behavior that can be useful in the development of fluorescent molecular switches. The resulting solvatochromic, highly fluorescent bioconjugate products can be particularly useful in pharmaceutical chemistry and fluorescence microscopy.

Results and Discussion

The acryl group cannot be attached directly to the amino group of ICAN without disrupting its electron donating properties and as a consequence flawing the solvatochromic properties of the molecule. The attachment should be carried out using a spacer, favorably a short chain hydrocarbon. Epichlorohydrin offers an easy alternative because after the opening of the epoxy ring with the amine a secondary hydroxyl group is formed which in turn is easily reacted with acryloyl chloride as is presented in Scheme 1. The reaction was also carried out using methacryloyl chloride and 2-choloroacrylic acid to investigate the change in the optical properties when an electron donating (Me) and an electron withdrawing (Cl) group is present on the acryl moiety.



Scheme 1. Synthesis of 1-(3-chloro-2-hydroxyprop-1-yl)-amino-5-isocyanonaphthalene (**2**, **CHAIN**), 1-(2-acryloyloxy-3-chloro-prop-1-yl)-amino-5-isocyanonaphthalene (**3**, **ACAIN**), 1-N-(2-methacryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (**4**, **MACAIN**) and (2-chloroacryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (**5**, **CACAIN**).

To test the solvatochromic properties of ACAIN and its derivatives UV-vis and steady-state fluorescence measurements were carried out. It should be noted, however, that this paper focuses on ACAIN, therefore its properties are presented in the main article, while the corresponding data for MACAIN and CACAIN are presented in the supporting information.

The UV-Vis spectrum of ACAIN recorded in THF shows a diffuse band attributed to the HOMO-LUMO transition between $\lambda=300$ nm and 400 nm with a peak value of 366 nm (spectrum No. 1. in Figure 1.).




Figure 1. The normalized absorption (1) and emission spectra of ACAIN recorded in hexane (2), THF (3) and water (4). 1 was recorded in THF.

The peak position showed only a slight variation between $\lambda_{\max}=356\text{-}373$ nm depending on the solvent used. The extinction coefficient corresponding to this peak varies between $\epsilon=5700$ Mcm^{-1} in water and $\epsilon=9100$ Mcm^{-1} in dichloromethane. The double peak structure of the band is most probably due to the superposition of the NH vibrational transition onto the absorption spectrum. It should be noted that the experimental UV-Vis spectra are in very good agreement with the calculated spectra (Figure S22 in the Supporting Information (SI)). Emission spectra were recorded in solvents of different polarity. The results are summarized in Table 1. for ACAIN and for MACAIN and CACAIN in TableS1(a)-S1(b) in the SI.

Table 1. Excitation (λ_{ex}), emission (λ_{em}) maxima, Stokes shift ($\Delta\nu$), molar absorbance (ϵ) and quantum yield (Φ_{F}) of ACAIN in different solvents

Solvent	λ_{ex} (nm)	λ_{em} (nm)	Stokes shift (cm^{-1})	ϵ (M^{-1})	Φ_{F} (%)
Hexane	356	423	4449	8500	3
Cyclohexane	359	423	4214	7040	8
Toluene	362	439	4845	8100	5
Dichloromethane	362	456	5694	9100	7
i-Propanol	367	477	6284	6450	26
Tetrahydrofuran	366	460	5583	7950	18
Chloroform	363	455	5570	7200	10
Ethyl acetate	366	459	5536	8000	19
Dioxane	364	456	5543	7640	15
Acetone	368	472	5987	6800	35
Methanol	367	485	6629	7740	18
Acetonitrile	364	479	6596	7500	25
Pyridin	369	479	6223	7110	9
Dimethyl formamide	370	482	6280	7280	46
Dimethyl sulfoxide	373	490	6401	6800	63
Water	362	502	7704	5700	3

As can be seen from the data of Table 1. ACAIN has visible emission in every solvent investigated. Characteristically for solvatochromic behavior the Stokes shifts increase with solvent polarity, that is bathochromic shift of the emitted light with increasing solvent polarity was found. The lowest emission wavelength was observed in hexane with $\lambda_{\text{em,max}}=423$ nm, the largest in water $\lambda_{\text{em,max}}=502$ nm and that of the THF solution can be found in between them at $\lambda_{\text{em,max}}=460$ nm as is presented in Figure 1. Surprisingly low quantum yields were observed in nonpolar solvents for all three compounds compared to the values of the starting ICAN molecule, which has the strongest emission in such solvents. According to the data of Table S1 the introduction of an electron donating methyl group to the 2 position of the acryl moiety of ACAIN increases the quantum yields in almost every solvent by ~10-20 %. On the other hand, the introduction of the electron withdrawing chlorine atom to the same position drastically lowers the QY values, that is the QYs do not exceed 10% even in polar solvents in the case of CACAIN. The explanation for these phenomena will be presented later.

The solvent-dependent behavior of ACAIN was quantitatively described by the most recent Catalán model according to Equation 1.

$$Y = Y_0 + aSA + bSB + sSP + tSdP, \quad (1)$$

where Y_0 is the property of the substance of interest (e.g., emission maximum and Stokes shift) in the absence of solvent, for example, in the gas phase. SA is the quantitative empirical measure of the ability of bulk solvent to act as a hydrogen-bond donor towards a solute. SB is the quantitative empirical measure of the ability of a bulk solvent to act as a hydrogen-bond acceptor or electron-pair donor towards a solute, forming a solute-to-solvent hydrogen bond or a solvent-to-solute coordinative bond, respectively. SP and SdP are the solvent polarizability and dipolarity parameters, respectively, determined using reference dye molecules. a, b, s and t are the corresponding coefficients and their inclusion in the equation indicates the dependence of the property under investigation upon the respective solvent parameter.

The Catalán coefficients for the emission wavenumber at the maximum ($\bar{\nu}_{em,max}$) and the Stokes shifts ($\Delta\bar{\nu}$) were obtained by multilinear regression analysis and are summarized in Table 2. The corresponding SA, SB, SP and SdP values of the solvents along with the plots of the measured values of $\bar{\nu}_{em,max}$ and $\Delta\bar{\nu}$ versus their calculated values according to Eq. 1 for ACAIN (and its derivatives) are presented in the Supporting Information as Table S4 and Figure S8. According to the data ACAIN, MACAIN and CACAIN can be considered as real solvatochromic dyes.

Table 2. Solvent-independent correlation coefficients a_{SA} , b_{SB} , s_{SP} and t_{SdP} of the Catalán parameters SA, SB, SP and SdP, respectively, solute property of the reference system $(\bar{\nu}_{em,max})_0$, correlation coefficient (r) and number of solvents (n) calculated by multiregression analysis for the solvatochromism of CHAIN and ACAIN.

Compound	$\bar{\nu}_{em,max,0}$	a_{SA}	b_{SB}	s_{SP}	t_{SdP}	r	n
		(cm ⁻¹)					
CHAIN	29300±1139	-2715±480	-2746±588	-5918±1384	-2675±1064	0.968	14
ACAIN	29113±1252	-2349±635	-2037±775	-5356±1714	-3062±1376	0.947	15

MS (**8**, **9**) methods. The reaction conditions are summarized in Table 3 except for **8** and **9** which will be dealt in detail later.

Table 3. The reaction conditions of the hydrothiolation reaction of **ACAIN** with different thiols. The conversion values are given for **ACAIN** and all reactions were carried out at 25 °C.

Thiol No.	[Thiol]	[ACAIN]	solvent	time	conversion
4	0.106 M	0.053 M	d ₆ -DMSO	<5 min	100%
5	0.318 M	0.159 M	d ₆ -DMSO	<5 min	100%
6	0.079 M	0.053 M	d ₆ -DMSO:D ₂ O (1:1)	<5 min	100%
7	0.190 M	0.095 M	DMSO	1 h	46 %

A number of solvents were examined but acceptable conversion was observed only in DMSO and water. The hydrothiolation reaction of the acryl group usually requires an amine catalyst such as triethylamine (TEA). However, in the case of propanethiol and thiophenol an almost instantaneous reaction was observed in d₆-DMSO at room temperature after the addition of the thiols in twofold molar amount relative to ACAIN. As can be seen in the ¹H NMR spectra presented in Figure 2 after 5 minutes reaction time the olefin proton signals of the acryl group completely disappeared. The formation of the addition product was confirmed by ESI-MS measurements and by the appearance of the S-CH₂ signals at 2.62 and 2.41 ppm. It can also be seen from the spectra that the thiol addition does not affect the electronic structure of the naphthalene core, only the signal of the NH proton shifts slightly downfield from 5.39 to 5.33 ppm.

Unlike the ¹H NMR spectrum, the adduct formation largely affects the fluorescence behavior of ACAIN. It was noted earlier that ACAIN has low quantum yields especially in nonpolar solvents. In contrast, the ACAIN-propanethiol adduct showed intense fluorescence in almost all solvents as can be seen in Figure 3. This intensity increase was not limited to the ACAIN-propanethiol adduct but could be observed in the case of every other thiols (**5**, **6**, **7**, **8**, **9**), as well.



Figure 3. Visible fluorescence of ACAIN (top) and ACAIN-propanethiol adduct (bottom) in solvents in order of increasing polarity excited at $\lambda=365$ nm. All solutions have the same concentration: 25.4 μ M

The fluorescence properties of ACAIN and its PrSH adduct are compiled in Table 4. The most dramatic increase in fluorescence intensity was observed in the less polar solvents hexane, toluene and dichloromethane where the increase was ~13-19 fold, while in the most polar solvents except water it was only 1.5-2.5 fold. The emission maxima and the shape of the emission peak were virtually unchanged. The extinction coefficients of the thiol adducts were also found to be very close to those of ACAIN within a ~10% error. Based on these observations it is surmised that the acryl group gives rise to a non-radiative path from the excited state to the ground state. By saturating the double bond this non-radiative way is eliminated and the quantum efficiencies increase considerably. The explanation of this phenomenon will be given in the quantum chemical part of the paper.

The reactivity of MACAIN and CACAIN was also investigated in the presence of propanethiol. Using the same experimental conditions as with ACAIN no reaction of PrSH with MACAIN could be observed, while CACAIN reacted approximately 100 times faster than ACAIN in DMSO as is presented in FigS10. Unfortunately the increased reactivity of CACAIN is accompanied by lower hydrolytic stability, namely CACAIN was found to hydrolyze in water even at pH=7. It should be noted, however, that no significant hydrolysis could be detected in the case of ACAIN in aqueous buffers applied during the kinetic measurements at pH=7, pH=8 and pH=9. In addition, ACAIN has very low quantum yield in water, a common feature among many solvatochromic fluorophores. In polar protic solvents such as water the energy gap that separates the S_0 and S_1 states is reduced

leading to a marked increase in non-radioactive decay and results in a decrease in the fluorescence quantum yield [14]. The low quantum yield may be favorable in applications such as biolabeling.

Table 4. The fluorescence intensities and emission maxima of **ACAIN** and its adduct with propanethiol in different solvents at 20 °C. All solutions have the same concentration of 2.54 μ M. The intensity ratios were calculated relative to the intensities of the pure **ACAIN** solutions.

Solvent	Pl. Intensity		λ (em) nm		Intensity ratio
	ACAIN	ACAIN+PrSH	ACAIN	ACAIN+PrSH	
Hexane	189	2836	423	424	15.0
Toluene	160	3011	439	438	18.8
Dichloromethane	237	3210	456	457	13.5
i-Propanol	838	3226	477	481	3.8
Tetrahydrofurane	760	2831	460	458	3.7
Ethyl-acetate	713	2808	459	460	3.9
Dioxane	648	3824	456	450	5.9
Acetone	1277	2536	472	470	2.0
Methanol	674	2268	485	485	3.4
Piridine	634	932	479	479	1.5
Acetonitrile	1132	2598	479	480	2.3
Dimethyl formamide	2183	3352	482	484	1.5
Dimethyl sulfoxide	1977	3241	490	490	1.6
Water	85	305	502	494	3.6

Kinetic Investigation of the reaction of ACAIN with biomolecules

The reaction between ACAIN and different compounds containing thiol (-SH) groups were followed spectrofluorometrically. The fluorescence intensity *versus* time curves for the reaction of ACAIN with cysteine, KAC and BSA at different pH are shown in Fig. 4.

Figure 4

As seen in Fig. 4. the fluorescence intensities increase with time constantly until plateau values have been reached indicating the completion of the reaction, i.e., covalent binding of all ACAIN molecules to the target compounds through acryl-thiol reaction. Moreover, in the case of BSA as can be seen in Fig. 4.c., there are large increases in the fluorescence intensity at the very beginning

of the reaction followed by much slower increases in the fluorescence intensities. The initial, very sharp and large increase in the fluorescence intensities, which occur instantly upon adding of BSA to the ACAIN solution, can be ascribed to the fast non-covalent binding of ACAIN molecules to the BSA as was previously demonstrated for Acrylodan and BSA by Mims et al. [16]. ACAIN itself reveals poor fluorescence in water due to the low quantum yield but becomes brightly fluorescent on non-covalent binding to BSA ascribed to the change of the micro-environment of ACAIN molecules. The change of the environment of ACAIN molecules results in a considerable increase in quantum yield and blue shifted emission maximum from 502 nm to 462 nm. The further, relatively slow increase in the fluorescence intensities are due to the covalent binding of ACAIN molecules to the BSA via acryl-thiol reaction involving the free thiol group of Cys-34 unit of BSA. It should be noted, that the second slow increase in fluorescence intensity was completely missing when the experiment was repeated using **1** as the fluorophore (FigureS11 in the SI).




Figure 4. The fluorescence intensity versus time curves for the reaction of ACAIN with cysteine (a) ($\lambda_{em}=505$ nm), KAC (b) ($\lambda_{em}=505$ nm) and BSA (c) ($\lambda_{em}=460$ nm). Experimental conditions: $[cystein]_o = [KAC]_o = [BSA]_o = 50 \mu M$, $[ACAIN]_o = 6.3 \mu M$, $T = 20 \text{ }^\circ C$.

To evaluate the kinetic curves presented in Fig. 4., we can assume a second-order kinetics for the reaction of the fluorophore (F) with the substrate (S) according to eq. 2.



In these kinetic experiments large excess of molar concentrations of S to those of F were applied, i.e., $[S]_0 \gg [F]_0$, to obtain pseudo first-order kinetics. Using the pseudo first-order kinetics eq. 3 can be derived for the variation of the fluorescence intensity with time (the derivation of eq. 3 can be found in the Supporting Information)

$$I(t) = I_0 + \Delta I(1 - e^{-k_{\text{obs}}t}) \quad (3)$$

where $\Delta I = I_\infty - I_0$, and I_0 and I_∞ are the initial and final fluorescence intensities, respectively.

Eq. 3. was fitted to the experimental fluorescence intensity *versus* time curves to determine the values of k_{obs} . According to the results, k_{obs} values increased with the pH, suggesting that the deprotonation of the thiol group play an important role in the rate determining step of the acryl-thiol reaction. Assuming that the thiolate anion reacts much faster with the acryl group of ACAIN than the undissociated thiol group, eq. 4 can be applied (the derivation of eq. 4. is shown in the Supporting Information)

$$\frac{1}{k_{\text{obs}}} = \frac{1}{k_{\text{thiolate}}K_a[S]_0}[\text{H}^+] + \frac{1}{k_{\text{thiolate}}[S]_0} \quad (4)$$

where k_{thiolate} and K_a are rate constants for thiolate-acryl reaction and the acid dissociation constant of the thiol group.

Plotting $1/k_{\text{obs}}$ as a function of $[\text{H}^+]$ yielded straight lines (see Supporting Information Fig.S12). From the slopes and intercepts of these lines, the values of K_a were obtained. The $\text{p}K_a$ values determined for cysteine and KAC using eq. 4. are 8.1 and 8.2, respectively. It is important to note that these values are very close to the literature values of $\text{p}K_a$ reported for the free cysteine that are in the range of 8.2-8.5 [26,27]. However, the $\text{p}K_a$ value determined for the BSA is ~ 7 , which is lower than those obtained for cysteine and KAC. This result, however, seems to be in line with the reported $\text{p}K_a$ values for BSA being in the range of 7-8.5 [28,29].

It is known that the acryl moiety is not only electrophilic towards sulfhydryl groups but can also react with amines. Therefore, acryl containing fluorophores such as Acrylodan can also modify lysine residues during the biolabeling of proteins [16,30]. Our model protein BSA contains **35** cysteines and **58** lysines [31,32]; however, all of the cysteine residues, with the exception of **Cys-34**, are involved in intramolecular disulfide bonds. The lysine/free cysteine ratio is 58 to 1 in BSA, therefore for practical applications it is important to determine the selectivity of ACAIN towards free cysteine units. Mass spectrometric methods such as ESI-MS and MS/MS provide an efficient way to study not only the kinetics but also the binding point of ACAIN during the labeling reaction therefore MS studies were also carried out to get deeper insight into the reaction of ACAIN with KAC and BSA. For these experiments unbuffered solutions were applied in order to avoid using of buffers incompatible with ESI and ion suppression. To investigate the selectivity of ACAIN with SH and NH_2 groups it was reacted with a simple tripeptide, KAC where an N-terminal lysine (K) is separated by an alanine (A) from a C-terminal cysteine (C) with amide functionality. KAC therefore contains 2 NH_2 (lysine) and 1 SH (cysteine) group. ACAIN (1.04 mM) was mixed with KAC (0.52 mM) in water/DMSO (50/50 V/V) at 25 °C. Samples were taken out at predetermined times, up to 6 hours reaction time, and were analyzed by ESI-MS/MS method.

Similar fragmentation pathway was found for the KAC and the labeled KAC, which shows that the connected ACAIN has no significant effect on the fragmentation pattern.

Figure 5. shows the ESI-MS/MS spectrum of the ACAIN-KAC adduct recorded after 6 hours reaction time.




Figure 5. The tandem mass spectrum of the ACAIN-KAC adduct. The fragment ions marked by asterisks are generated with the loss of NH₃ from the corresponding product ions.

Figure inset shows the structures of the ACAIN and the KAC and the possible generated fragment ions of the KAC molecule.

As can be seen in Fig. 5, the presence of the ACAIN-KAC adduct is supported by the accurate mass of the protonated adduct ion, [ACAIN-KAC+H]⁺ with *m/z* 634.256 (634.257 calculated) and the fragmentation pattern. The majority of the product ions come from the fragmentation of the tripeptide backbone. Based on the tandem mass spectrum the ACAIN binding point can be determined by the analysis of the product ions. The fragment ions generated in the higher mass region are produced by the loss of the lysine and the alanine. These amino acids appear in the mass spectrum as the a₁, a₁^{*}, a₂, b₁, b₂ and b₂^{*} type product ions. According to the ESI-MS spectrum, neither cysteine, nor any product ion containing cysteine with free thiol group can be detected, while all of the fragment ions which contain cysteine are labelled with ACAIN. Furthermore, no

lysine bound to ACAIN can be detected, while the **x**, **y** and **z** type fragment ions can only be found with the ACAIN moiety. It should be noted that these **x**, **y** and **z** type fragment ions appeared in the tandem mass spectrum of unlabelled KAC. The ESI-MS/MS spectrum of the unlabelled KAC tripeptide can be found in the supporting information as FigS13. Based on the MS/MS investigations, it can be concluded that the ACAIN reacts only with the cysteine moiety of the KAC tripeptide and no reaction with the amine groups under the applied reaction conditions could be observed.

The ESI MS investigation of the labeling of BSA with ACAIN using the same experimental conditions as in the case of KAC followed. Attachment of one ACAIN molecule to a BSA molecule increases the mass of BSA with 315 Da compared to that of the starting protein. According to Figure 6., the BSA-ACAIN reaction can be followed by ESI-MS method. At $t=0$ min the typical mass spectrum of the protonated starting protein at different charge states can be observed. The measured masses and the charges of the ions are marked over the peaks. After 40 min reaction time another series of peaks is clearly visible in the middle panel of Figure 6, which belongs to the BSA-ACAIN reaction product. The charge state of these ions is the same as that of the corresponding BSA ions. In order to gain the masses of the compounds deconvolution was used (FigS14 in the Supporting Information). As it can be seen the differences between the masses of labelled and unlabelled BSA is 315 Da, which is in very good agreement with the mass of the ACAIN molecule (315 Da). However, after 260 min reaction time a third series with a molecular weight of 630 Da higher appeared which belongs to the BSA-2ACAIN adducts. Although it was shown that ACAIN does not bind to the lysine moiety of KAC it reacts with one lysine of BSA. This behavior was previously shown in the reaction of Acrylodan with BSA [16] and with Interleukin-1 β [30]. Since ACAIN is a hydrophobic molecule just as Acrylodan, it tends to incorporate into the hydrophobic pockets of proteins. BSA has two such hydrophobic pockets: one at Cys-34 where the first ACAIN molecule binds and the other at Lys-240 where the lysine moiety is activated because of its nonpolar environment and the second ACAIN binding most likely happens. The assumption that ACAIN binds to the only activated lysine (Lys240) of BSA is further supported by the fact that no additional series appeared in the mass spectrum, however long the reaction was allowed to proceed. Further details including the kinetic curves for the reactions of ACAIN with KAC and BSA are presented in the Supporting Information as FigS15 and FigS16, respectively.




Figure 6. The mass spectra of the reaction mixture of the BSA and ACAIN at different reaction times. The peaks are marked with the corresponding masses and the charge state of the ions. Experimental conditions: $[BSA]_0=0.52$ mM, $[ACAIN]_0=1.04$ mM in water/DMSO (50/50 V/V) at 25 °C.

DFT calculations

For practical applications and for the design of more effective fluorophores it is important to know why and how the ACAIN molecule has negligible fluorescence in the least polar and observable emission in the most polar solvents. Therefore we selected hexane and DMSO as model solvents and calculated the ground state (S_0) and excited state (S_1 and T_1) geometries with the electronic spectra of ACAIN and its thiol adduct. The DMSO system was expanded with an explicit solvent molecule to account for the H-bonding between the ACAIN and the DMSO.

Depending on the polarity of the solvent molecules, intramolecular or solvent-solute interactions can dominate which give rise to different molecular structures in different solvents. The calculated equilibrium structures of ACAIN in n-hexane at different electronic states are shown in FigS17 in the Supporting Information. We found that in the triplet state a hydrogen atom (or proton) transfer from the $-NH$ group ($T_1(N)$) to the acrylic oxygen is possible ($T_1(O)$). (In the ground and S_1 states we found no local minima with O-localized proton.) All the structures have an intramolecular H-bond between the amino group and the acrylic oxygen. Also, in the S_1 and $T_1(O)$ states the acryl group becomes almost coplanar with the naphthalene ring.

These geometry optimizations revealed a possibility for the lack of emission because of intersystem crossing (ISC) which is most likely to happen between S_1 and $T_1(O)$ as they are similar in geometry and very close in energy (5.4 kJ/mol) making the overlap of their vibrational energy levels possible. The energy diagram in Fig. 7 shows the electronic energy of the ground, S_1 and T_1 electronic states in the S_1 optimized geometry along with the energy minimum of the $T_1(N)$ and $T_1(O)$ states. The dashed and dotted lines show approximations of the potential energy surfaces. It can be seen, however, that the energies of S_1 and $T_1(O)$ are close to each other and the conical intersection point for the relaxation into the $T_1(N)$ state lies even closer to S_1 .




Figure 7. Electronic energies of the S_1 structure of ACAIN (in hexane) in its S_0 , S_1 and T_1 ($T_1(S_1)$) electronic states. The other two energies correspond to the two triplet minima. The lines are approximations of the S_1 (\cdots) and T_1 ($---$) potential energy surfaces.

For the thiol adduct although the energy difference between the S_1 and $T_1(O)$ is also relatively small (12.0 kJ/mol) ISC cannot happen because the geometry of S_1 is much closer to the other triplet minimum ($T_1(N)$) with an energy separation of 78.7 kJ/mol. Furthermore, the $T_1(O)$ state is much higher in energy than the other two states in Fig. S19.

The situation in DMSO is different as there is an H-bond between the amino group and a DMSO molecule, while the acryl unit is π -stacked to the naphthalene ring (Fig. S18). The structures of S_0 , S_1 and $T_1(N)$ are very similar to each other, while in the $T_1(O)$ state the hydrogen favors the acrylic oxygen over the DMSO and upon protonation the acryl unit twists out from the π -stacking. This is similar to what was observed in the case of the thiol adduct in hexane because the energy spacing of the S_1 and $T_1(O)$ states is also small (8.6 kJ/mol). Moreover, they are still far away on the potential energy surface due to their geometries. The energy difference between the similar structures of S_1 and $T_1(N)$ is 56.2 kJ/mol (Fig. S20) which is about ten times higher than the one calculated in hexane. Therefore, the non-radiative relaxation to the T_1 state in DMSO is less probable than in hexane.

The calculated electronic spectra are in good agreement with the experimental results except for the emission of ACAIN (Fig. S21), which is due to the near degenerate triplet and singlet states. It is important to note that the main emission band of the thiol adduct consists of only one transition, while the one in the calculated spectra of ACAIN has two, one at higher wavelength with very low oscillator strength and one at lower wavelength with oscillator strength similar to that of the thiol adduct. The spacing of these two transitions is larger in hexane. This result implies that the excited ACAIN molecules relax to a dark S_1 state in both hexane and DMSO.

The difference in the emission of the ACAIN and its thiol adduct lies in the fact that ACAIN and thio-ACAIN has different frontier molecular orbitals. In the amino-isocyanonaphthalene moiety the electronic transition responsible for the fluorescence is a charge transfer between the donor amino and the acceptor isocyano groups [33]. However in ACAIN the HOMO-LUMO transition involves an intramolecular charge transfer from the main ring to the acryl unit as it can be seen in Fig. S22-S23. In the thiol adduct the acrylic double bond gets saturated eliminating the possibility for this low energy π - π^* transition, hence the molecule becomes fluorescent.

In order to support our model we performed calculations on molecules with modified acryl units. First the ACAIN molecule was simplified by removing the chloromethyl group (replaced by a hydrogen atom) to reduce the calculation times. This truncating has no effect on the electronic transitions investigated here therefore the chloromethyl group plays no role in the quenching of the emission observed in ACAIN. However, when an acrylic hydrogen atom is substituted with an electron-donating methyl group, it is expected to destabilize the molecular orbitals containing the acryl π -system thereby increasing the HOMO-LUMO gap. The calculations indeed show a higher energy S_1 state with increased singlet-triplet gap which means a less probable internal conversion. In addition, when an electron withdrawing chlorine atom is used as the substituent instead of the methyl group calculations show the opposite effect (Table S5).

After these results we performed calculations on MACAIN and found good correlation between the experimental quantum yield and the calculated singlet-triplet energy difference (Table S6).

Conclusions

1-(2-acryloyloxy-3-chloro-prop-1-yl)-amino-5-isocyanonaphthalene (ACAIN), 1-(2-methacryloyloxy-3-chloro-prop-1-yl)-amino-5-isocyanonaphthalene (MACAIN) and (2-chloroacryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (CACAIN) were prepared

by introducing acryl, methacryl and chloroacryl moieties using epichlorohydrin as spacer on our previously developed solvatochromic fluorophore ICAN. All three compounds exhibited real solvatochromic behavior ($\Delta\lambda_{em}\sim 100$ nm) which could be quantitatively described by the Catalán and Kamlet-Taft models as well. Interestingly, the quantum yields were found to be rather low, especially in nonpolar solvents, decreasing in the order of $QY_{MACAIN} > QY_{ACAIN} > QY_{CACAIN}$. ACAIN reacts rapidly and quantitatively with simple thiols in a thiol-ene type click reaction. Using propanethiol, thiophenol, and cysteine as thiols complete disappearance of the acryl double bond was observed in 5 minutes at room temperature in DMSO without any catalyst using the thiols in a molar excess. The hydrothiolation reaction was accompanied by a significant increase in fluorescence intensity depending on the solvent used. The brightening of the addition product is due to the saturation of the acrylic group and can be applied in the fluorometric monitoring of the hydrothiolation reactions. In the case of MACAIN no such reaction could be observed, while CACAIN proved to react 100 times faster in DMSO than ACAIN. Unfortunately, the improved reactivity owing to the chloroacryl moiety drastically reduces the hydrolytic stability of CACAIN, whereas ACAIN is stable under the same conditions. The biolabeling properties of ACAIN were investigated in detail by both fluorometry and ESI mass spectrometry using cysteine, lysyl-alanyl-cysteine (KAC) and BSA as a model protein. The selectivity of the labeling was tested using KAC as a simple tripeptide containing a lysine and a cysteine moiety separated by an alanine unit. Investigating the reaction by ESI-MS/MS method no fragment ions could be found where ACAIN labelled the lysine NH_2 group, only the hydrothiolation reaction with the cysteine SH group was observed. However, when labeling BSA two series of labelled products could be identified in the ESI-MS spectrum belonging to the molecules containing one and two ACAIN molecules attached covalently. The second attachment most probably occurs on the only activated lysine unit of BSA in a hydrophobic pocket as was described earlier for Acrylodan, a well-known biolabeling dye. The optical properties of ACAIN and its SH adducts were described by high level quantum chemical calculations. The optimized structures revealed the formation of an intramolecular hydrogen bond between the NH and C=O moieties. This intramolecular H-bond gives rise to a dark nonfluorescent state of ACAIN due to a close energy triplet state where electron transition can happen between the acrylic double bond and the aromatic core. When the double bond is saturated by the SH addition the internal conversion pathway leading to nonradiative decay ceases to exist resulting in the increase of the fluorescence quantum yields of the thiolated products. The

brightening is more pronounced (15-19 fold) in nonpolar solvents where the majority of the starting ACAIN molecules is locked in the internal H-bonding position and only moderate (1.5-2 fold) in polar solvents where the solvent can break up the H-bond. Based on our DFT calculations the structure of the spacer has little effect on the electronic transitions as long as the internal H-bond can form. The situation significantly changes when the modification is carried out on the acrylic part. When the acryl group is replaced by a methacryl group i.e. an electron-donating group is introduced on the double bond the calculations showed an increased singlet-triplet gap which means a less probable internal conversion. Indeed, after preparing and investigating the properties of MACAIN a good correlation between the experimental quantum yield and the calculated singlet-triplet energy difference was found. Also if an electron withdrawing chlorine atom is used instead of the methyl group calculations show the tightening of the singlet-triplet gap, which could be applied in the construction of molecular switches based on ACAIN.

Experimental

Materials

Acetone, dichloromethane (DCM), hexane, 2-propanol (iPrOH), toluene, triethyl amine (reagent grade, Molar Chemicals, Hungary) were purified by distillation. Acetonitrile (MeCN), tetrahydrofuran (THF), methanol (MeOH), dimethyl-sulfoxide (DMSO), pyridine (HPLC grade, VWR, Germany), chloroform, ethyl-acetate (EtOAc), cyclohexane, 1,4-dioxane (reagent grade, Reanal, Hungary), epichlorohydrin, acryloyl chloride, methacryloyl chloride, 2-chloroacrylic acid, 4-N,N-dimethylaminopyridine, N,N-dicyclohexyl carbodiimide (Sigma-Aldrich, Germany) were used without further purification. Buffer solution pH 7.01 (HI 7007, Hanna Instruments) was used as received. Buffer solution pH 8.0 and 9.0 (0.1 M) were prepared by mixing 1 M K_2HPO_4 and KH_2PO_4 solutions (pH 8.0) or 1 M K_2HPO_4 and 1 M HCl solutions (pH 9.0) at appropriate ratios diluted with water and their final pH was adjusted under a pH meter.

Synthesis

1-N-(3-chloro-2-hydroxyprop-1-yl)-amino-5-isocyanonaphthalene (CHAIN)

A 250 ml ACE pressure flask was charged with 1-amino-5-isocyanonaphthalene (1.00 g, 5.95 mmol) dissolved in ethanol (100 ml) and with epichlorohydrin (930 μ l, 11.9 mmol) and vigorously stirred with a magnetic stirrer at 90 °C for 3 days. After cooling down, the solvent was removed on a rotary evaporator. The crude product was purified on a column filled with normal-phase silica gel, using dichloromethane as eluent. Yield: 0.82 g, 53 % (light brown waxy solid).

^1H NMR (360 MHz, CDCl_3) δ = 7.90 (d, J = 8.6 Hz, 1H), 7.59 – 7.42 (m, 3H), 7.35 (t, J = 8.0 Hz, 1H), 6.71 (d, J = 7.4 Hz, 1H), 4.85 (s, 1H), 4.25 (s, 1H), 3.80 – 3.67 (m, 2H), 3.57 – 3.46 (m, 1H), 3.36 (dd, J = 12.7, 7.4 Hz, 1H), 2.80 (d, J = 4.7 Hz, 1H) ppm.

^{13}C NMR (95 MHz, CDCl_3) δ = 162.64 (C_{NC}), 140.52 ($\text{C}_{9,10}$), 126.70 (C_5), 126.46 (C_2), 122.44 (C_7), 121.66 (C_1), 121.43 (C_6), 119.94 (C_3), 110.85 (C_8), 104.68 (C_4), 70.98 (C_{CHOH}), 51.74 (C_{CH_2}), 46.31 ($\text{C}_{\text{CH}_2\text{Cl}}$) ppm.

ESI-TOF MS (m/z): calculated m/z of $[\text{C}_{14}\text{H}_{13}\text{ClN}_2\text{O}+\text{Na}]^+$ adduct ion is 283.061; found; 283.060

1-N-(2-acryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (ACAIN)

Into a 250 ml round-bottom flask charged with 1-N-(3-chloro-2-hydroxyprop-1-yl)-amino-5-isocyanonaphthalene (1.00 g, 3.84 mmol) and triethyl amine (2.14 ml, 15.3 mmol) dissolved in tetrahydrofuran (150 ml) acryloyl chloride (623 μ l, 7.68 mmol) was added dropwise at 0 °C, then stirred with a magnetic stirrer for 12 hours. The solvent was removed on a rotary evaporator and the crude product was purified on a column filled with normal-phase silica gel, using dichloromethane as eluent. Yield: 0.86 g, 71 % (orange crystals).

^1H NMR (360 MHz, CDCl_3) δ = 7.84 (d, J = 8.6 Hz, 1H), 7.64 – 7.46 (m, 3H), 7.39 (t, J = 8.0 Hz, 1H), 6.80 (d, J = 7.5 Hz, 1H), 6.51 (d, J = 17.3 Hz, 1H), 6.19 (dd, J = 17.2, 10.5 Hz, 1H), 5.94 (d, J = 10.4 Hz, 1H), 5.53 – 5.43 (m, 1H), 4.84 (t, J = 5.2 Hz, 1H), 3.83 (d, J = 4.8 Hz, 2H), 3.75 – 3.60 (m, 2H) ppm.

^1H NMR (360 MHz, DMSO) δ = 8.32 (d, J = 8.7 Hz, 1H), 7.83 – 7.72 (m, 1H), 7.54 (t, J = 8.1 Hz, 1H), 7.45 (dd, J = 8.6, 7.5 Hz, 1H), 7.29 (d, J = 8.3 Hz, 1H), 6.85 (d, J = 7.7 Hz, 1H), 6.74 (t, J = 5.7 Hz, 1H), 6.36 (dd, J = 17.2, 1.6 Hz, 1H), 6.19 (dd, J = 17.2, 10.3 Hz, 1H), 5.97 (dd, J = 10.3, 1.6 Hz, 1H), 5.39 (td, J = 9.7, 6.0 Hz, 1H), 3.99 (ddd, J = 17.9, 12.0, 4.8 Hz, 2H), 3.67 – 3.49 (m, 2H) ppm.

^{13}C NMR (95 MHz, CDCl_3) δ = 162.93 (C_{NC}), 161.96 ($\text{C}_{\text{C=O}}$), 140.15 ($\text{C}_{9,10}$), 130.04 (C_5), 126.76 (C_1), 125.37 (C_3), 122.55 ($\text{C}_{\text{Acryl CH}_2}$), 121.62 ($\text{C}_{\text{Acryl CH}}$), 119.79 ($\text{C}_{6,7}$), 111.00 ($\text{C}_{2,8}$), 104.65 (C_4), 71.67 (C_{CHO}), 46.52 (C_{CH_2}), 45.14 ($\text{C}_{\text{CH}_2\text{Cl}}$) ppm.

ESI-TOF MS (m/z): calculated m/z of $[\text{C}_{17}\text{H}_{15}\text{ClN}_2\text{O}_2+\text{Na}]^+$ adduct ion is 337.071; found; 337.070

Thiol adducts of 1-N-(2-acryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (ACAIN)

A 25 ml round-bottom flask was charged with 1-N-(2-acryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (0.050 g, 0.16 mmol), the thiol (0.16 mmol) dissolved in 5 ml DMSO and was stirred with a magnetic stirrer for 10 minutes to 6 hours at room temperature, then extracted with water and dichloromethane. The organic phase was separated, dried on anhydrous magnesium sulfate, and the solvent was removed on a rotary evaporator. Yield: 46-95 %.

Adduct formed with 1-propanethiol:

^1H NMR (360 MHz, CDCl_3) δ = 7.89 (d, J = 8.6 Hz, 1H), 7.56 (dt, J = 16.0, 8.4 Hz, 3H), 7.38 (t, J = 8.0 Hz, 1H), 6.78 (d, J = 7.5 Hz, 1H), 5.45 (dq, J = 9.8, 4.9 Hz, 1H), 4.86 (s, 1H), 3.80 (d, J = 5.1 Hz, 2H), 3.72 – 3.58 (m, 2H), 2.86 – 2.75 (m, 2H), 2.73 – 2.70 (m, 1H), 2.48 (t, J = 7.3 Hz, 2H), 1.63 – 1.50 (m, 3H), 0.94 (t, J = 7.3 Hz, 3H) ppm.

ESI-TOF MS (m/z): calculated m/z of $[\text{C}_{20}\text{H}_{23}\text{ClN}_2\text{OS}+\text{Na}]^+$ adduct ion is 413.106; found; 413.106

Adduct formed with thiophenol:

^1H NMR (360 MHz, DMSO) δ = 8.33 (d, J = 8.7 Hz, 1H), 7.75 (d, J = 7.1 Hz, 1H), 7.53 (t, J = 8.1 Hz, 1H), 7.44 (dd, J = 8.5, 7.5 Hz, 1H), 7.37 – 7.16 (m, 4H), 7.11 (t, J = 7.2 Hz, 1H), 6.83 (d, J = 7.8 Hz, 1H), 6.70 (t, J = 5.8 Hz, 1H), 5.39 – 5.31 (m, 2H), 3.94 (qd, J = 11.9, 4.8 Hz, 2H), 3.62 – 3.46 (m, 2H), 3.12 (t, J = 7.0 Hz, 2H), 2.63 (t, J = 7.0 Hz, 2H) ppm.

ESI-TOF MS (m/z): calculated m/z of $[\text{C}_{23}\text{H}_{21}\text{ClN}_2\text{O}_2\text{S}+\text{K}]^+$ adduct ion is 463.064; found; 463.064

Adduct formed with cysteine:

^1H NMR (360 MHz, DMSO) δ = 13.95 (s, 1H), 8.23 (d, J = 8.6 Hz, 1H), 7.72 (d, J = 7.3 Hz, 1H), 7.54 (t, J = 8.1 Hz, 1H), 7.44 (t, J = 8.0 Hz, 1H), 7.27 (d, J = 8.3 Hz, 1H), 6.80 (d, J = 7.8 Hz, 1H), 5.28 (d, J = 4.6 Hz, 1H), 3.89 – 3.77 (m, 2H), 3.52 (ddd, J = 21.3, 14.4, 6.2 Hz, 2H), 3.41 (s, 2H), 2.96 (dd, J = 14.2, 3.5 Hz, 1H), 2.86 – 2.70 (m, 3H), 2.66 (s, 2H), 2.63 – 2.54 (m, 2H) ppm.

ESI-TOF MS (m/z): calculated m/z of $[C_{20}H_{22}ClN_3O_4S+Na]^+$ adduct ion is 458.091; found; 458.091

Adduct formed with 1-thio-2,3,4,6-tetra-O-acetyl- β -D-glucose:

1H NMR (400 MHz, $CDCl_3$) δ = 7.91 (dd, J = 8.5, 2.6 Hz, 1H), 7.62 – 7.50 (m, 3H), 7.47 – 7.37 (m, 1H), 6.78 (d, J = 7.5 Hz, 1H), 5.49 – 5.40 (m, 1H), 5.19 (tt, J = 9.6, 4.8 Hz, 1H), 5.10 – 5.01 (m, 1H), 5.01 – 4.91 (m, 1H), 4.48 (dd, J = 10.0, 6.4 Hz, 1H), 4.19 – 4.06 (m, 3H), 3.80 (d, J = 4.9 Hz, 2H), 3.71 – 3.57 (m, 3H), 3.02 (tt, J = 11.1, 7.1 Hz, 1H), 2.90 (td, J = 13.7, 7.1 Hz, 1H), 2.77 (dt, J = 7.3, 3.6 Hz, 2H), 2.07 – 1.98 (m, 12H) ppm.

ESI-TOF MS (m/z): calculated m/z of $[C_{31}H_{35}ClN_2O_{11}S+Na]^+$ adduct ion is 701.154; found; 701.154

1-N-(2-methacryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (MACAIN)

Into a 250 ml round-bottom flask charged with 1-N-(3-chloro-2-hydroxyprop-1-yl)-amino-5-isocyanonaphthalene (1.00 g, 3.84 mmol) and triethyl amine (2.14 ml, 15.3 mmol) dissolved in tetrahydrofuran (150 ml) methacryloyl chloride (740 μ l, 7.68 mmol) was added dropwise at 0 °C, then stirred with a magnetic stirrer for 12 hours. The solvent was removed on a rotary evaporator and the crude product was purified on a column filled with normal-phase silica gel, using dichloromethane as eluent. Yield: 0.28 g, 22 % (orange crystals).

1H NMR (360 MHz, $CDCl_3$) δ = 7.83 (d, J = 8.6 Hz, 1H), 7.62 – 7.48 (m, 3H), 7.39 (t, J = 7.9 Hz, 1H), 6.81 (d, J = 7.4 Hz, 1H), 6.21 (s, 1H), 5.67 (t, J = 1.4 Hz, 1H), 5.39-5.51 (s, 1H), 4.86 (s, 1H), 3.84 (d, J = 4.8 Hz, 2H), 3.60-3.74 (m, 2H), 1.98 (s, 3H). ppm.

^{13}C NMR (95 MHz, $CDCl_3$) δ = ^{13}C NMR (91 MHz, $CDCl_3$) δ 167.17 (C_{NC} , $C_{C=O}$), 143.23 ($C_{9,10}$), 135.69 (C_5), 129.20 (C_1), 128.92 (C_3), 127.05 ($C_{Methacryl\ CH_2}$), 124.69 (C_7), 123.84 ($C_{Methacryl\ C}$), 123.74 (C_6), 121.79 (C_8), 112.67 (C_2), 106.00 (C_4), 71.76 (C_{CHO}), 45.34 (C_{CH_2}), 43.87 (C_{CH_2Cl}), 18.25 ($C_{Methacryl\ CH_3}$).

ESI-TOF MS (m/z): calculated m/z of $[C_{18}H_{17}ClN_2O_2+Na]^+$ adduct ion is 351.087; found; 351.087

1-N-(2-chloroacryloyloxy-3-chloroprop-1-yl)-amino-5-isocyanonaphthalene (CACAIN)

Into a 50 ml round-bottom flask charged with 1-N-(3-chloro-2-hydroxyprop-1-yl)-amino-5-isocyanonaphthalene (500 mg, 1.92 mmol), 4-N,N-dimethylaminopyridine (26.0 mg, 0.210 mmol) and 2-chloroacrylic acid (225 mg, 2.11 mmol) dissolved in dichloromethane (15 ml) N,N-

dicyclohexyl carbodiimide (435 mg, 2.11 mmol) dissolved in dichloromethane (5 ml) was added dropwise at 0 °C, then stirred with a magnetic stirrer for 2 days. The solvent was removed on a rotary evaporator and the crude product was purified on a column filled with normal-phase silica gel, using dichloromethane as eluent. Yield: 0.33 g, 49 % (pale yellow viscous oil).

¹H NMR (360 MHz, CDCl₃) δ = 7.84 (d, *J* = 8.6 Hz, 1H), 7.65 – 7.46 (m, 3H), 7.41 (dd, *J* = 8.6, 7.4 Hz, 1H), 6.83 (d, *J* = 7.5 Hz, 1H), 6.59 (d, *J* = 10.3 Hz, 1H), 6.09 (d, *J* = 10.2 Hz, 1H), 5.52 – 5.40 (m, 1H), 4.77 (s, 1H), 3.86 (d, *J* = 4.8 Hz, 2H), 3.73 (t, *J* = 5.8 Hz, 2H) ppm.

¹³C NMR (91 MHz, CDCl₃) δ = 167.15 (C_{C=O}), 161.75 (C_{NC}), 142.91 (C_{9,10}), 130.80 (C₅), 129.19 (C₁), 128.88 (C_{Chloroacryl CH₂}), 127.29 (C_{Chloroacryl CH}), 126.61 (C₃), 124.80 (C₇), 123.92 (C₆), 121.73 (C₈), 113.01 (C₂), 106.15 (C₄), 73.56 (C_{CHO}), 44.87 (C_{CH₂}), 43.38 (C_{CH₂Cl}) ppm.

ESI-TOF MS (*m/z*): calculated *m/z* of [C₁₇H₁₄Cl₂N₂O₂-H]⁺ ion is 346.036; found; 346.036

Lysyl-alanyl-cysteine (KAC)

The KAC tripeptide with terminal amino and amide functionalities (see Figure 2) was synthesized by solid phase peptide synthesis using a microwave-assisted Liberty 1 Peptide Synthesizer (CEM, Matthews, NC). Fmoc-protected amino acid derivatives were introduced according to the Fmoc/*t*Bu technique and the TBTU/HOBt/DIEA strategy. Cleaving on the α-amino protecting group of amino acids and resin was performed by 30 Watts microwave power for 180 s at 80 °C using 20 V/V% piperidine and 0.1 M HOBt·H₂O in DMF. Four times excess of amino acids and 30 Watts microwave power for 300 s was used for coupling at 80 °C in the presence of 0.5 M HOBt and 0.5 M TBTU in DMF as activator and 2 M DIPEA in NMP as activator base. After building up the peptide sequences, Fmoc group was removed similarly while the side chain protecting groups were cleaved with the treatment of TFA/TIS/H₂O/ 2,2'-(ethylenedioxy)diethanethiol (94/2.5/2.5/1 V/V) mixture, simultaneously with the removal of the peptide from the resin, at room temperature for 2 h. The resin was filtered from the trifluoroacetic acid solution of the peptide which was precipitated in cold diethyl ether. The crude product was also washed using this solvent and dried after separation followed by dissolution in water and freezing for lyophilization.

The purity of the prepared products was checked by analytical rp-HPLC using a Jasco instrument equipped with a Jasco MD-2010 plus multiwavelength detector monitoring the absorbance at 222 nm.

Gradient elution was carried out using solvent A (0.1 V/V% TFA in acetonitrile) and solvent B (0.1 V/V% TFA in water) at a flow rate of 0.8 ml/min. From 1 min to 15 min 0 to 25% of A, from 15 min to 16 min 25% of A and from 16 min to 20 min 25 to 0% of A was applied and a Teknokroma Europa Peptide C18 chromatographic column (250 x 4.6 mm, 120 Å pore size, 5 µm particle size) was used.

Chemicals and solvents used for synthetic purpose were purchased from commercial sources in the highest available purity and used without further purification. Rink Amide AM resin, 2-(1-H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TBTU) and all N-fluorenylmethoxycarbonyl (Fmoc)-protected amino acids (Fmoc-Cys(Trt)-OH, Fmoc-Lys(Boc)-OH and Fmoc-Ala-OH) are Novabiochem (Switzerland) products. 2-methyl-2-butanol, N-hydroxybenzotriazole (HOBt), N-methyl-pyrrolidone (NMP), 2,2'-(ethylenedioxy)diethanethiol (DOTD) and triisopropylsilane (TIS) were purchased from Sigma-Aldrich Co., while N,N-diisopropyl-ethylamine (DIEA) and trifluoroacetic acid (TFA) were Merck Millipore Co. products. Peptide-synthesis grade N,N-dimethylformamide (DMF) and acetic anhydride (Ac₂O) were from VWR International and piperidine, dichloromethane (DCM), diethyl ether (Et₂O), acetic acid (AcOH) and acetonitrile (ACN) were from Molar Chemicals Ltd.

Methods

NMR

¹H and ¹³C-NMR spectra were recorded in CDCl₃ and DMSO-d₆ at 25 °C on a Bruker Avance DRX-400 and a Bruker AM 360 spectrometer at 400 MHz and 360 MHz, respectively with tetramethylsilane as the internal standard.

Electrospray Quadrupole Time-of-Flight MS/MS (ESI-Q-TOF)

The MS and MS/MS measurements were carried out by a MicroTOF-Q type Qq-TOF MS instrument (Bruker Daltonik, Bremen, Germany). The instrument was equipped with an electrospray ion source where the spray voltage was 4 kV. N₂ was utilized as drying gas. The temperature of the drying gas was 200 °C and the flow rate was 4.0 L/min. For the tandem MS measurements collision gas was nitrogen. The precursor ions for MS/MS were selected with an

isolation width of 4 m/z units. The mass spectra were recorded by means of a digitizer at a sampling rate of 2 GHz. The mass spectra were calibrated externally using the exact masses of cluster ions of sodium trifluoroacetate $[(\text{NaTFA})_n+\text{Na}]^+$ generated under electrospray condition. In the case of the BSA and BSA derivatives the collision RF of the quadrupole analyzer was set to 700. The spectra were evaluated with the DataAnalysis 3.4 software from Bruker. The samples were introduced by a syringe pump (Cole-Parmer Ins. Co., Vernon Hills, IL, USA) at a flow rate of 10 $\mu\text{L}/\text{min}$. For the mass spectrometric measurements the samples were dissolved in methanol in the concentration of 0.1 mM. In order to generate the protonated adduct ions in the case of the BSA and BSA derivatives samples low amount of formic acid was added ($c_{\text{acid}}=0.01 \text{ M}$).

UV-vis

The UV-vis spectra were recorded on an Agilent Cary 60 spectrophotometer (Agilent, Santa Clara, CA, USA) in a quartz cuvette of 1.00 cm optical length. 3.00 cm^3 solution was prepared from the sample.

Fluorescence measurements

Steady-state fluorescence measurements were carried out using a Jasco FP-8200 fluorescence spectrophotometer equipped with a Xe lamp light source. The excitation and emission spectra were recorded at 20 °C, using 2.5 nm excitation, 5.0 nm emission bandwidth, and 200 nm/min scanning speed. Fluorescence quantum yields were calculated by using quinine-sulfate in 0.1 mol/L sulfuric acid as the reference absolute quantum efficiency ($\Phi_n = 55\%$).

For UV-vis and fluorescence measurements the investigated compounds were dissolved in acetonitrile at a concentration of 1.19 mM and was diluted to $2.38 \times 10^{-5} \text{ M}$ and $4.76 \times 10^{-6} \text{ M}$ in the solvents in interest.

Density Functional Theory (DFT) calculations

All calculations were performed using the Gaussian09 C.01 software package [34]. We employed DFT and TDDFT for geometry optimizations at the ground and excited states respectively. The chosen functional was the M06 [35] with the TZVP basis set [36] because it is able to describe some amount of the dispersion forces in contrast to B3LYP [37] and CAM-B3LYP [38] which was

found to be necessary to properly optimize π -stacked structures [33]. The calculations were done using the IEF-PCM solvent model [39-42].

Acknowledgment

This work was financially supported by the grant K-116465 given by OTKA (National Found for Scientific Research Development, Hungary). This work was partially supported by the European Union and the European Social Fund through project Supercomputer, the National Virtual Lab, grant no.: TÁMOP-4.2.2.C-11/1/KONV-2012-0010. The authors also would like to express their thanks to Ms. Katalin Várnagy for providing the KAC tripeptide sample.

Supporting Information


Copies of NMR, MS and optical spectra, computational details and derivation of equations.

References

- [1] Nair DP, Podgórski M, Chatani S, Gong T, Xi W, Fenoli CR, Bowman CN. *Chem Mater* 2014;26:724–744
- [2] Kolb HC, Finn MG, Sharpless KB. *Angew Chem Int Ed* 2001;40:2004–2021.
- [3] Baskin JM, Bertozzi CR, *QSAR Comb Sci* 2007;26:1211–1219.
- [4] Nwe K, Brechbiel MW. *Cancer Biother Radio* 2009;24:289-302.
- [5] Wang SQ, Wu QH, Wang HY, Zheng XX, Shen SL, Zhang YR, Miao JY, Zhao BX, *Biosens Bioelectron* 2014;55:386–390.
- [6] Hoyle CE, Lowe AB, Bowman CN. *Chem Soc Rev* 2010;39:1355-1387.
- [7] Li GZ, Randev RK, Soeriyadi AH, Rees G, Boyer C, Tong Z, Davis TP, Becer CR, Haddleton DM. *Polym Chem* 2010;1:1196–1204.
- [8] Matsumoto T, Urano Y, Shoda T, Kojima H, Nagano T. *Org Lett* 2007;9:3375-3377.
- [9] Lin W, Yuan L, Cao Z, Feng Y, Long L. *Chem Eur J* 2009;15:5096-5103.
- [10] Sun YQ, Chen M, Liu J, Lv X, Li JF, Guo W. *Chem Commun* 2011;47:11029-11031.
- [11] Lallana E, Sousa-Herves A, Fernandez-Trillo F, Riguera R, Fernandez-Megia E. *Pharm Res* 2012;29:1–34.
- [12] Lowe AB. *Polym Chem* 2010;1:17-36.
- [13] Jones MW, Gibson MI, Mantovani G, Haddleton DM. *Polym Chem* 2011;2:572-574.
- [14] Loving GS, Sainlos M, Imperiali B. *Trends Biotechnol* 2009;28:73-83.
- [15] Sommer A, Gorges R, Kostner GM, Paltauf F, Hermetter A. *Biochemistry* 1991;30:11245–11249.
- [16] Mims MP, Sturgis CB, Sparrow JT, Morrisett JD. *Biochemistry* 1993;32:9215–9220.
- [17] Flora K, Brennan JD, Baker GA, Doody MA, Bright FV. *Biophys J* 1998;75:1084-1096.
- [18] Baudier J, Glasser N, Duportail G. *Biochemistry* 1986;25:6934–6941.
- [19] Oztug Durer ZA, Kudryashov DS, Sawaya MR, Altenbach C, Hubbell W, Reisler E. *Biophys J* 2012;103:930-939.
- [20] Simard JR, Getlik M, Grütter C, Pawar V, Wulfert S, Rabiller M, Rauh D. *J Am Chem Soc* 2009;131:13286–13296.
- [21] Bottari G, Leigh DA, Perez EM. *J Am Chem Soc* 2003;125:13360–13361.

- [22] Lewis PA, Inman CE, Maya F, Tour JM, Hutchison JE, Weiss PS. *J Am Chem Soc* 2005;127:17421–17426.
- [23] Saha S, Stoddart JF. *Chem Soc Rev* 2007;36:77-92.
- [24] Olofsson J, Önfelt B, Lincoln P. *J Phys Chem A* 2004;108:4391–4398.
- [25] Rácz D, Nagy M, Mándi A, Zsuga M, Kéki S. *J Photoch Photobio A* 2013;270:19–27.
- [26] Szajewski RP, Whitesides GM. *J Am Chem Soc* 1980;102:2011-2066.
- [27] Wilson JM, Wu D, Motiu-DeGroot R, Hupe DJ. *J Am Chem Soc* 1980;102:359-363.
- [28] Pedersen AO, Jacobsen J. *Eur J Biochem* 1980;106:291-295.
- [29] Noble DR, Williams DLH. *J Chem Soc Perkin Trans* 2001;2:13–17.
- [30] Yem AW, Epps DE, Mathews WR, Guido DM, Richard KA, Staite ND, Deibel MR Jr. *J Biol Chem* 1992;267:3122-3128.
- [31] Brown JR. *Fed Proc Fed Am Soc Exp Biol* 1975;34:591.
- [32] Hirayama K, Akashi S, Furuya M, Fukuhara K. *Biochem Biophys Res Commun* 1990;173:639-646.
- [33] Nagy M, Rácz D, Lázár L, Purgel M, Ditrói T, Zsuga M, Kéki S. *ChemPhysChem* 2014;15:3614–3625.
- [34] Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR. *Gaussian 09*, revision A.02. Wallingford CT: Gaussian, Inc.; 2009.
- [35] Zhao Y, Truhlar DG. *Theor Chem Acc* 2008;120:215-241.
- [36] Schaefer A, Huber C, Ahlrichs RJ. *Chem Phys* 1994;100:5829-5835.
- [37] Becke AD. *J Chem Phys* 1993;98:1372-1377.
- [38] Yanai T, Tew D, Handy N. *Chem Phys Lett* 2004;393:51-57.
- [39] Tomasi J, Mennucci B, Cammi R. *Chem Rev* 2005;105:2999-3093.
- [40] Scalmani G, Frisch MJ. *J Chem Phys* 2010;11:114-110.
- [41] Tomasi J, Mennucci B, Cancès E. *THEOCHEM* 1999;464:211.
- [42] Pascual-Ahuir JL, Silla E, Tunon I. *J Comput Chem* 1994;15:1127-1138.

Graphical Abstract



Highlights

- Acrylated isocyanonaphthalene based solvatochromic click reagent was prepared
- Reacts easily and quantitatively with a range of different thiols
- The emission intensity significantly increases during the hydrothiolation reaction
- The biolabeling abilities were demonstrated using biomolecules with free thiol group
- DFT calculations revealed an internal acryl-naphthalene electron transfer