

Cladocera remains and vegetation types to assess the state of oxbows

István Gyulai¹, Csilla Lakatos², János Tamás Kunderát², Zsuzsanna Balogh²,
Edina Simon², Béla Tóthmérész³

1 Department of Hydrobiology, University of Debrecen, Debrecen, Egyetem tér 1, H-4032 Hungary

2 Department of Ecology, University of Debrecen, Debrecen, Egyetem tér 1, H-4032 Hungary **3** MTA-DE Biodiversity and Ecosystem Services Research Group, Debrecen, Egyetem tér 1, H-4032 Hungary

Corresponding author: Edina Simon (edina.simon@gmail.com)

Academic editor: M. Pinna | Received 24 April 2018 | Accepted 29 September 2018 | Published 18 September 2018

<http://zoobank.org/98E1D00F-2827-4ADF-B64B-FAA1C80E414>

Citation: Gyulai I, Lakatos C, Kunderát JT, Balogh Z, Simon E, Tóthmérész B (2018) Cladocera remains and vegetation types to assess the state of oxbows. *Nature Conservation* 29: 27–39. <https://doi.org/10.3897/natureconservation.29.26221>

Abstract

We assessed the usefulness of Cladocera remains for establishing the ecological status of oxbows and also tested the association of Cladocera species with various vegetation types. Cladocera remains were collected from the surface sediment of four habitat types (tangled vegetation, open water, reeds and tunnels) and 15 physical and chemical parameters of surface water were studied. In the surface sediment samples, we identified 32 Cladocera taxa. There was a significant difference in the number of species amongst habitat types as per ANOVA. The benthic and plant associated Cladocera communities of reeds, tangled vegetation, open water and tunnels were clearly separated from each other by NMDS ordination. CCA showed that habitat types had characteristic Cladocera species: *Pleuroxus* species were frequent in the tangled vegetation habitat, while *Chydorus* species were frequent in the open water. Remarkably, in reeds, *Bosmina* species were frequent, although these species are usually common in open water. Specimens of the *Alona* genus were found everywhere. Our findings suggest that the remains of Cladocera species may be useful indicators to assess and monitor the structure of freshwater lakes.

Keywords

Water-Sediment chemistry, Macrophytes, Zooplankton Indicator species

Introduction

There are several organic and inorganic remains in sediments which reflect the history of oxbows. In lake sediments, some of the most common animal remains are those of Cladocera which derive from both water and sediment (Kurek et al. 2010). The taxonomic structure of Cladocera remaining in sediment cores indicates past changes in the environment, such as eutrophication (Visconti et al. 2008, Nevalainen and Luoto 2013), acidification (Jeziorski et al. 2008) and changes in the water level (Korponai et al. 2016). Cladocera communities play an important role in a lake's food web because of their intermediate position, which means that Cladocera species have significant effects on the ecology status and water quality of lakes (Jeppesen et al. 2011, Zhi et al. 2012). Due to this position in the food web, Cladocera are sensitive indicators of environmental changes (Kurek et al. 2010).

Earlier studies indicated that the boundary zone between macrophyte beds and open water is particularly important as a refuge for cladocerans (Lauridsen et al. 1996; Davidson et al. 2010). At the same time, Cladocera communities vary with macrophyte bed size and open water which is an important daytime refuge for potentially migrating pelagic cladocerans (Lauridsen et al. 1996).

In this study, we tested the effect of habitat types of oxbows on Cladocera communities in the Upper Tisza Region, in northeast Hungary. We also studied the correlation between the water chemical parameters of Cladocera species. We hypothesised that there were Cladocera species characteristic of typical oxbow habitats and that they are useful indicators for assessing and monitoring the structure and ecological state of lakes.

Methods

Study sites

Many oxbows were formed during the 19th century with the controlling of the River Tisza. To assure shipping and flood-control, more than 100 meanders were cut. As a result, many artificial oxbow lakes were formed along the River Tisza (Babka et al. 2011, Balogh et al. 2016, 2017, Kundrát et al. 2017). In the Upper-Tisza region, there are more than 40 oxbows. This region is mainly cultivated using traditional agricultural systems with meadows, orchards and some cereal cultivation. The characteristic land use of this countryside has changed considerably during the last 200 years (Varga et al. 2013).

The oxbows studied were in the Upper Tisza region, near the town of Vásárosnamény in Hungary. The following oxbows were studied: Keskeny Holt-Tisza (48°9'5.64"N, 22°20'9.30"E), Foltos-kerti Holt-Tisza (48°5'47.58"N, 22°23'47.64"E) and Patkó Holt-Tisza (48°6'27.66"N, 22°23'1.56"E). In each oxbow, four sampling points were chosen to represent typical habitat types: tangled vegetation, open water, reeds and

tunnels. In the tangled habitat, *Ceratophyllum demersum* (about 90%) and *Potamogeton natans* (about 10%) were the most abundant plant species. In the open water habitats, there were no aquatic plants. In the reed habitat, the main aquatic plant species was *Phragmites australis*. The tunnel habitats were at least 1.2 m deep and wide open surfaces with little vegetation towards the sides were typical (*Ceratophyllum demersum*).

Cladocera identification

For Cladocera identification, surface sediment subsamples (1 cm³) were treated with 100 ml 10% KOH (Normapur, VWR) solution and heated at 100 °C for about 1 hour. Hydrofluoric acid (HF) (38%, VWR) was used to remove the inorganic material following identification. Then we added safranin O (ALFA (AESAR)) to the sample to stain the remains. We prepared quantitative slides by pipetting 100 µl of each subsample on to a microscope slide and then examined it under a microscope (B-183, OPTIKA Microscopes, Italy) at magnifications of 100 and 400; about 200 Cladocera remains were counted from each sample. Usually two slides are sufficient for identifying at least 200 remains, which is the recommended number for counting (Kurek et al. 2010). The identification was based on Bledzki and Rybak (2016) and Szeroczyńska and Sarmaja-Korjonen (2007).

Water analyses

Surface water samples were collected in plastic bottles and parallel measurements were performed at the study sites. Water depth, transparency, temperature, conductivity (WTW cond. 340i) and pH (WTW pH 315i) were measured. Samples were stored at 4 °C until the laboratory process. In the laboratory, the content of suspended solids, chlorophyll-a, Chemical Oxygen Demand (COD) and the concentrations of carbon dioxide, ammonium-nitrogen, nitrite-nitrogen, nitrate-nitrogen and orthophosphate were measured. Laboratory analyses of water samples were based on APHA (2000) and Nollert and De Gelder (2011).

Sediment analyses

To determinate the organic matter content of surface sediment, the loss on ignition method was used. After drying at 105 °C, 0.2 g samples were cremated at 550 °C for 4 h in a muffle furnace (Nabertherm L5/C6, Germany). The loss on ignition was calculated with the following equation: $LOI_{550} = 100 \cdot (DW_{105} - DW_{550}) / WS$, where LOI_{550} was the percentage loss on ignition at 550 °C, DW_{105} was the dry weight of samples at 150 °C and DW_{550} was the weight of the sample at 550 °C (Heiri et al. 2001, Matthews 2014). To determine the content of calcium carbonate in the surface

sediment, the samples were burnt at 950 °C for 4 h. After cooling, we measured them with analytic scales. The calculation of the loss on ignition was conducted with the following equipment: $LOI_{950} = 100 \cdot (DW_{550} - DW_{950}) / WS$, where LOI_{950} is the percentage of loss on ignition at 950 °C and DW_{950} is the weight of the sample after heating at 950 °C (Heiri et al. 2001, Matthews 2014).

Statistical analyses

The benthic and plant associated Cladocera communities were studied, based on vegetation types, by non-metric multidimensional scaling (NMDS) ordination. CCA was used to display the correlation between water chemistry and the Cladocera community (Lepš and Šmilauer 2003). One-way ANOVA was used to test the effect of habitat types on Cladocera diversity and water chemistry. In the case of significant differences, Tukey's Multiple Comparison test was used (Abbott 2016).

Results

Cladocera diversity

In total, we counted 1324 Cladocera specimens in the samples; altogether, we identified 32 taxa (Table 1). There was a significant difference in the number of Cladocera species amongst the vegetation types by ANOVA ($F_{3,8} = 4.744$, $P = 0.034$) (Fig. 1). A significantly higher number of Cladocera species was found in the open water than in the reed vegetation type ($P < 0.05$). There was no significant difference in the number of Cladocera individuals amongst the vegetation types (oxbows: $F_{3,8} = 0.500$, $P = 0.693$ Fig. 2).

The benthic and plant associated Cladocera communities of reeds, tangled vegetation, open water and tunnels were clearly separated from each other by NMDS ordination. The communities of benthic Cladocera in tangled vegetation, open water and tunnels were similar to each other (Fig. 3). A similar result was found in the cases of plant associated Cladocera communities when using NMDS ordination (Fig. 4).

Water physico-chemistry and sediment chemistry differences amongst vegetation types

There were no significant differences in the water physico-chemistry parameters studied (depth: $F = 1.234$, $p = 0.359$; visibility: $F = 0.591$, $P = 0.638$; temperature: $F = 0.164$, $P = 0.918$; pH: $F = 2.433$, $P = 0.140$; conductivity: $F = 0.029$, $P = 0.993$; suspended solids: $F = 1.038$, $P = 0.427$; CO_2 : $F = 2.519$, $P = 0.132$; COD: $F = 0.004$, $P = 1.000$; NH_4^+ : $F = 1.406$, $P = 0.310$; NO_3^- : $F = 0.696$, $P = 0.580$; Chlorophyll-a: F




Figure 1. Number of Cladocera species by vegetation type.




Figure 2. Number of Cladocera individuals by vegetation type.




Figure 3. NMDS ordination of the benthic Cladocera remains of vegetation types (stress=0.1785).




Figure 4. NMDS ordination of the plant associated Cladocera remains of vegetation types (stress=0.1922).

= 2.823, $P = 0.107$) amongst vegetation types (Table 2). Similar to the water physico-chemical parameters, significant differences were not found amongst vegetation types based on the organic matter content ($F = 3.159$, $P = 0.086$), nor on the calcium carbonate content ($F = 0.134$, $P = 0.937$) of sediment (Table 2).

Correlation between sediment and water chemistry and Cladocera communities

Based on the CCA ordination, our results show positive correlations between the organic matter and the calcium carbonate content of sediment and *A. elongatus*, *A. rectangularis*, *L. leydigii* and *A. quadrangularis* species (Fig. 5). In the cases of water chem-

Table 2. Physical and chemical parameters of surface water and sediment (mean \pm SD) according to vegetation type.

Parameters	Vegetation type			
	tangled vegetation	open water	reeds	tunnel
Water				
depth, cm	59 \pm 20	80 \pm 20	135 \pm 35	75 \pm 20
visibility, cm	49 \pm 21	54 \pm 7	73 \pm 23	47 \pm 4
temperature, °C	12 \pm 1	11 \pm 1	11 \pm 1	11 \pm 1
pH	8.3 \pm 0.3	8.6 \pm 0.2	8.5 \pm 0.3	7.9 \pm 0.1
conductivity, $\mu\text{S cm}^{-1}$	334 \pm 78	346 \pm 73	343 \pm 73	370 \pm 920
suspended solid, mg l^{-1}	8 \pm 4	10 \pm 2	12 \pm 8	3 \pm 1
CO ₂ , mg l^{-1}	19 \pm 5	10 \pm 2	24 \pm 3	15 \pm 4
COD, mg l^{-1}	6 \pm 5	4 \pm 2	5 \pm 3	6 \pm 4
NH ₄ ⁺ , mg l^{-1}	3 \pm 1	1 \pm 1	2 \pm 1	1 \pm 1
NO ₃ ⁻ , mg l^{-1}	0.2 \pm 0.1	0.3 \pm 0.1	0.1 \pm 0.1	0.2 \pm 0.1
Chlorophyll-a, mg l^{-1}	6 \pm 2	9 \pm 3	15 \pm 7	4 \pm 1
Sediment				
organic matter, %	4.1 \pm 0.4	3.4 \pm 0.2	2.4 \pm 1.0	2.9 \pm 0.8
CaCO ₃ , %	0.7 \pm 0.2	0.8 \pm 0.3	0.7 \pm 0.5	0.6 \pm 0.2




Figure 5. CCA ordination of benthic Cladocera taxa and sediment parameters.




Figure 6. CCA ordination of Cladocera taxa and water chemical parameters.

istry, data positive correlations were found between conductivity and *K. latissima* species, between chemical oxygen demand and *A. nana*, *A. harpae* and *A. exigua* species and between temperature and *P. truncatus*, *O. tenuicaudis* and *A. emarginatus* (Fig. 6).

Discussion

Our study demonstrated the usefulness of Cladocera remains in the assessment of the ecological status of oxbows. Similarly to our study, Gulyás and Forró (1999) and Korhola and Rautio (2001) also demonstrated a correlation between habitat types and Cladocera species. The CCA results corroborated the habitat preferences reported by Gulyás and Forró (1999) and Korhola and Rautio (2001). At the tunnel of the Kes-

keny Holt-Tisza oxbow, we found the kind of Cladocera species which usually live in vegetation zones. In the tangled vegetation of the Keskeny Holt-Tisza oxbow, we found *Alonella exigua*, *Acroperus harpae* and *Alona guttata* which were typical tangled vegetation species; however, *Acroperus elongatus* and *Alona affinis* were mostly living in the biotecton of vegetation. In the tangled vegetation of the Foltos-kerti Holt-Tisza, *Alonella excisa* was a typical tangled vegetation species. We also found *Daphnia longispina* remains there; this species is characteristically an open water and/or generalist species. In tunnels, *Alona quadrangularis* was a benthic species as reported by literature. The Cladocera species we found in the tangled vegetation of the Patkó Holt-Tisza oxbow usually lived in tangled vegetation and in sediment. In the open water, we found the kind of species which usually live in sediment and in vegetation and not usually in open water. Probably these species are able to adapt quickly to the modified environmental conditions caused by human disturbance (i.e. the intense utilisation of the oxbow for recreational fishing).

There were significant differences amongst oxbows and the habitat types based on water chemistry parameters. Similar to earlier studies (Lukács et al. 2009, 2011), we found that aquatic plants influenced the water chemistry parameters. Lukács et al. [(2011) demonstrated that the amount of chlorophyll-a was very high in sweet grass beds communities, but a small amount of chlorophyll-a was found in chestnut and water lily beds. Our findings also demonstrated that there was a strong interaction between water chemistry parameters and reed habitats.

We found that temperature was in a positive correlation with the number of Cladocera individuals. Nevalainen and Luoto (2010) also reported that many Cladocera species are sensitive to seasonal temperature changes. Zawisza et al. (2016) and Wojewódka et al. (2016) reported that several studies described strong correlations between pH, conductivity and Cladocera taxa. Bjerring et al. (2009) found negative correlations between temperature and chlorophyll-a and several Cladocera taxa. We found a positive correlation between conductivity and Cladocera taxa, while a negative correlation was found between pH and Cladocera taxa.

Conclusions

Our results show that Cladocera taxa are usually associated with characteristic habitat types; however, human disturbance can change the habitat association of these species by changing the local environment conditions. Based on our results, Cladocera are useful indicators for assessing and monitoring the structure of freshwater lakes.

Acknowledgements

Csilla Lakatos and István Gyulai were supported by the TÁMOP 4.2.1/B-09/1/KONV-2010-0007 and TÁMOP- 274 4.2.2/B-10/1-2010-0024 projects of the European Union and the State of Hungary, co-financed by the European Social Fund in the

framework of TÁMOP 4.2.4.A/2-11-1-2012-0001 ‘National Excellence Program’. E. Simon was supported by the TÁMOP-4.2.2.A-11/1/KONV. Research was supported by OTKA K 116639, KH 126481 and KH 126477 grants.

References

- Abbott ML (2016) *Using Statistics in the Social and Health Sciences with SPSS and Excel*. John Wiley & Sons.
- American Public Health Association (APHA) (2000) *Standard methods for the examination of water and wastewater*. Washington (DC).
- Babka B, Futó I, Szabó S (2011) Clustering oxbow lakes in the Upper-Tisza Region on the basis of stable isotope measurements. *Journal of Hydrology (Amsterdam)* 410(1–2): 105–113. <https://doi.org/10.1016/j.jhydrol.2011.09.026>
- Balogh Zs, Harangi S, Gyulai I, Braun M, Hubay K, Tóthmérész B, Simon E (2017) Exploring river pollution based on sediment analysis in the Upper Tisza region (Hungary). *Environmental Science and Pollution Research International* 24(5): 4851–4859. <https://doi.org/10.1007/s11356-016-8225-5>
- Balogh Zs, Harangi S, Kundrat JT, Gyulai I, Tóthmérész B, Simon E (2016) Effects of anthropogenic activities on the elemental concentration in surface sediment of oxbows. *Water, Air, and Soil Pollution* 227(1): 13–21. <https://doi.org/10.1007/s11270-015-2714-x>
- Bjerring R, Becares E, Declerck S, Gross EM, Hansson LA, Kairesalo T, Nykanen M, Halkiewicz A, Kornijow R, Conde-Porcuna JM, Seferlis M, Nöges T, Moss B, Amsinck SL, Odgaard BV, Jeppesen E (2009) Subfossil Cladocera in relation to contemporary environmental variables in 54 Pan-European lakes. *Freshwater Biology* 54(11): 2401–2417. <https://doi.org/10.1111/j.1365-2427.2009.02252.x>
- Bledzki LA, Rybak JI (2016) *Freshwater Crustacean zooplankton of Europe*. Springer.
- Davidson TA, Sayer CD, Perrow M, Bramm M, Jeppesen E (2010) The simultaneous inference of zooplanktivorous fish and macrophyte density from sub-fossil cladoceran assemblages: A multivariate regression tree approach. *Freshwater Biology* 55(3): 546–564. <https://doi.org/10.1111/j.1365-2427.2008.02124.x>
- Gulyás P, Forró L (1999) *A guide for the identification of Cladocera occurring in Hungary*. KGI, Budapest. [In Hungarian]
- Heiri O, Lotter AF, Lemcke G (2001) Loss on ignition as a method for estimating organic and carbonate content in sediments: Reproducibility and comparability of results. *Journal of Paleolimnology* 25(1): 101–110. <https://doi.org/10.1023/A:1008119611481>
- Jeppesen E, Nöges P, Davidson TA, Haberman J, Nöges T, Blank K, Lauridsen TL, Søndergaard M, Sayer C, Laugaste R, Johansson LS, Bjerring R, Amsinck SL (2011) Zooplankton as indicators in lakes: A scientific-based plea for including zooplankton in the ecological quality assessment of lakes according to the European Water Framework Directive (WFD). *Hydrobiologia* 676(1): 279–297. <https://doi.org/10.1007/s10750-011-0831-0>
- Jeziorski A, Yan ND, Paterson AM, Desellas AM, Turner MA, Jeffries DS, Keller B, Weeber RC, Mcnicol DK, Palmer ME, McIver K, Arseneau K, Ginn BK, Cumming BF, Smol

- JP (2008) The widespread threat of calcium decline in fresh waters. *Science* 322(5906): 1374–1377. <https://doi.org/10.1126/science.1164949>
- Korhola A, Rautio M (2001) Cladocera and other Branchiopod Crustaceans. In: Tracking Environmental Change Using Lake Sediments. Smol JP, Birks HJB, Last WM (Eds) Kluwer Academic Publishers, Dordrecht, The Netherlands, 5–41. https://doi.org/10.1007/0-306-47671-1_2
- Korponai J, Gyulai I, Braun M, Kövér C, Papp I, Forró L (2016) Reconstruction of flood events in an oxbow lake (Marótzugi-Holt-Tisza, NE Hungary) by using subfossil cladoceran remains and sediments. *Advances in Oceanography and Limnology* 7: 131–141. <https://doi.org/10.4081/aiol.2016.6168>
- Kundrát JT, Balogh Zs, Harangi S, Tóthmérész B, Simon E (2017) Assessment of anthropogenic, seasonal and aquatic vegetation effects on the contamination level of oxbows. *Community Ecology* 18(3): 237–243. <https://doi.org/10.1556/168.2017.18.3.2>
- Kurek J, Korosi JB, Jeziorski A, Smol JP (2010) Establishing reliable minimum count sizes for cladoceran subfossils sampled from lake sediments. *Journal of Paleolimnology* 44(2): 603–612. <https://doi.org/10.1007/s10933-010-9440-6>
- Lauridsen TL, Pedersen LJ, Jeppesen E, Sønergaard M (1996) The importance of macrophyte bed size for cladoceran composition and horizontal migration in a shallow lake. *Journal of Plankton Research* 18(12): 2283–2294. <https://doi.org/10.1093/plankt/18.12.2283>
- Lukács BA, Dévai G, Tóthmérész B (2009) Aquatic macrophytes as bioindicators of water chemistry in nutrient rich backwaters along the Upper-Tisza river (in Hungary). *Phytocoenologia* 39: 287–293. <https://doi.org/10.1127/0340-269X/2009/0039-0287>
- Lukács BA, Dévai G, Tóthmérész B (2011) Small scale macrophyte-environment relationship in an oxbow-lake of the Upper-Tisza valley (Hungary). *Community Ecology* 12(2): 259–263. <https://doi.org/10.1556/ComEc.12.2011.2.15>
- Matthews JA (2014) *Encyclopedia of Environmental Change*. SAGE.
- Nevalainen L, Luoto TP (2013) Limnological deterioration forces community and phenotypic changes in cladocera: Tracking eutrophication of mallusjärvi, a lake in southern Finland. *Boreal Environment Research* 18: 209–222.
- Nevalainen L, Luoto TP (2010) Temperature sensitivity of gamogenesis in littoral cladocerans and its ecological implications. *Journal of Limnology* 69(1): 120–125. <https://doi.org/10.4081/jlimnol.2010.120>
- Nollet LM, De Gelder LSP (2011) *Handbook of water analysis*. CRC Press.
- Szeroczyńska K, Sarmaja-Korjonen K (2007) *Atlas of Subfossil Cladocera from Central and Northern Europe*. Friends of the Lower Vistula Society, Świecie, Poland.
- Varga K, Dévai G, Tóthmérész B (2013) Land use history of a floodplain area during the last 200 years in the Upper-Tisza region (Hungary). *Regional Environmental Change* 13(5): 1109–1118. <https://doi.org/10.1007/s10113-013-0424-8>
- Visconti A, Manca M, De Bernardi R (2008) Eutrophication-like response to climate warming: An analysis of Lago Maggiore (N. Italy) zooplankton in contrasting years. *Journal of Limnology* 67(2): 87–92. <https://doi.org/10.4081/jlimnol.2008.87>
- Wojewódka M, Zawisza E, Cohuo S, Macario-González L, Schwalb A, Zawiska I, Pérez L (2016) Ecology of Cladocera species from Central America based on subfossil assemblages. *Advances in Oceanography and Limnology* 7: 151–162. <https://doi.org/10.4081/aiol.2016.6266>

- Zawisza E, Zawiska I, Correa-Metrio A (2016) Cladocera Community Composition as a function of physicochemical and morphological parameters of Dystrophic Lakes in NE Poland. *Wetlands* 36(6): 1131–1142. <https://doi.org/10.1007/s13157-016-0832-x>
- Zhi W, Zhiyong Z, Junqian Z, Yingying Z, Haiqing L, Shaohua Y (2012) Large-scale utilization of water hyacinth for nutrient removal in Lake Dianchi in China: The effects on the water quality, macrozoobenthos and zooplankton. *Chemosphere* 89(10): 1255–1261. <https://doi.org/10.1016/j.chemosphere.2012.08.001>