PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link. http://hdl.handle.net/2066/20441

Please be advised that this information was generated on 2018-07-07 and may be subject to change.

1210 Bartels et al.

Interhemispheric Subdural Hematoma in Adults: Case Reports and a Review of the Literature

Electrocardiography showed atrial fibrillation.

CT revealed a left ISH extending over the falx toward the tentorium (*Fig.* 1). Two weeks after admission, this had completely resolved.

It was decided to conservatively man-

Ronald H.M.A. Bartels, M.D., Wim I.M. Verhagen, M.D., Ph.D., Mathé J.J. Prick, M.D., Ph.D., Johanna E. Dalman, M.D.

Department of Neurology, Canisius Wilhelmina Hospital, Nijmegen, The Netherlands

THE INTERHEMISPHERIC SUBDURAL hematoma is a relatively uncommon type of subdural hematoma, especially seen in patients with blood clotting disturbances. When its mass becomes sufficiently large, specific neurological abnormalities such as hemiparesis and signs of the falx syndrome are seen. Treatment can consist of conservative observation or craniotomy and is dictated by the clinical course. Conservative management is the treatment of choice for patients without disturbances of consciousness and for patients with stable clinical conditions. Surgical treatment is necessary in patients

age the patient. Immediately after admission, she received medication to counteract the anticoagulant agent. No changes were observed for 2 weeks. After this period, gradual improvement was noticed in her speech and in her right arm strength. Three days later, there was also improvement in her right leg. Six weeks after admission, she was able to walk without any assistance. She was discharged to her own home and regained the ability to perform almost all of her normal activities.

Patient 2


A 70-year-old woman was recovering from an intracerebral hemorrhage in the right internal capsule 112 days before admission. Ten hours before admission, she fell out of bed and hit her head. She had amnesia regarding this event, but it was uncertain whether she had lost consciousness. Afterwards, she complained of right-sided headaches. She vomited several times and noticed progressive gait disturbance, which deteriorated so that she was no longer able to walk, even with assistance. Her medications

with progressive deterioration. Three case reports are presented, as well as a review of 64 cases described in the literature. The salient aspects of this clinical entity are discussed. (Neurosurgery 36:1210–1214, 1995)

Key words: Blood coagulation, Computer-assisted tomography, Interhemispheric, Subdural hematoma

The interhemispheric subdural hematoma (ISH) in adults is a rare type of subdural hematoma. Since the first description by Aring and Evans in 1940 (1), only 64 cases have been described in detail (1–4, 6–8, 10–16, 18–23, 25–46, 48– 50). Zimmerman et al. (52) described the brain computed tomography (CT) findings, but not the clinical features, of 25 adult patients. Particularly because of its location, the ISH is considered to be a distinct entity. Recent experience with three patients prompted us to carefully review the literature. days, these complaints gradually progressed. Twelve hours before admission, she also noticed problems with talking and speech. These symptoms also worsened. She was using an oral anticoagulant agent and 0.125 mg of Lanoxin (Burroughs Wellcome, Research Triangle Park, NC) daily because of atrial fibrillation.

At admission, her consciousness was undisturbed. She had symptoms of nuchal rigidity and was suffering from dysphasia. Except for right central facial paresis, no other signs of cranial nerve involvement were present. Her right arm and leg were equally paretic (motor strength = 1/5, Medical Research Council scale). Stretch reflexes were exaggerated on both sides with a right extensor plantar reflex.



CASE REPORTS

Patient 1


Four days before admission, a 78year-old woman struck her head during a fall. One day later, she noticed difficulties with walking. Over the next 3

Laboratory examinations were within normal values except for an International Normalized Ratio of 3.9.

FIGURE 1. CT without contrast at admission showing an ISH on the left side.

Interhemispheric Subdural Hematoma 1211

were 100 mg of acetylsalicylic acid daily and 20 mg of isosorbide dinitrate thrice daily. At admission, she was fully alert. The cranial nerves were intact, except for a left central facial paresis and a slight deviation of the tongue to the left. She had left hemiparesis involving the leg (3/5) more than the arm (4/5), which was more pronounced than it was before the fall (arm and leg, 4/5). On both sides, brisk stretch reflexes and positive Hoffmann's, Trömner's, and extensor plantar reflexes were seen. There was no sensory loss. The results of laboratory investigations were within normal limits. CT was compatible with a dense interhemispheric collection of fresh blood on the right side. A minimal amount of blood was also seen in the subarachnoid space (right sylvian fissure). We decided to conservatively manage the patient. Three weeks after admission, the ISH had almost completely resolved; 2 weeks later, the results of the CT were unremarkable. In the meantime, the patient's muscle strength had gradually increased. After 3 months, she was able to walk with the assistance of two persons, and after 6 months, she was able to walk alone.



occurred most often in the 6th, 7th, and 8th decades (*Fig.* 3). The incidence in children might be much higher than in adults because of child abuse by shaking; Zimmerman et al. (51) found ISHs in 17 of 28 children.

Pathogenesis

Preceding head trauma was mentioned in 61 ISH cases (91.0%). In two cases, the ISH occurred spontaneously;

in two cases, it was caused by a ruptured intracranial aneurysm, and in the remaining two cases, no underlying mechanism was mentioned. The site of impact of the head trauma in 7 cases was frontal, in 18, occipital, in 3, vertex, and in 2, parietal, and it was not mentioned in the remaining cases.

In Table 1, accompanying factors or diseases are shown. One of the patients in whom the ISH spontaneously appeared had hemophilia B (40); the other had been using an oral anticoagulant agent for arteritis (23), whereas another patient had been using an anticoagulant agent for cardiological reasons (22). The generally accepted causative mechanism for the development of ISHs is tearing of the fixed bridging veins between the medial cerebral cortex and the superior sagittal sinus. Torsion forces are known to be important, which is in accordance with the high incidence of frontal and occipital sites of impact. Tight connection of the arachnoid trabeculae between the brain and the parasagittal dura prevents further extension of a fresh clot to the ipsilateral

Patient 3

A 75-year-old man had been complaining of headaches for 2 weeks and was admitted because of a decreasing level of consciousness that had started the day before. On that day he had fallen twice. He had been using an oral anticoagulant agent because of a myocardial infarction 6 weeks before. At admission, he was also suffering from nausea and vomiting and was stuporous. Cranial nerve functions were undisturbed. He had mild (4/5) hemiparesis on the right side. The arm stretch reflexes were symmetrical and normal, whereas the right patellar reflex was exaggerated. He had a flexor plantar response on the left side and an equivocal response on the right. The results of laboratory examinations were within normal limits except for an International Normalized Ratio of 5.0. CT was compatible with an acute ISH on the left side within a more chronic subdural hematoma, because "pseudoFIGURE 2. *A* and *B*, CT without contrast at admission disclosing a left-sided ISH with compartments of various densities and "pseudomembranes" compatible with a recent hemorrhage in a more chronic ISH.

membranes" were present (*Fig. 2*). The oral anticoagulant was stopped and antagonized immediately. His clinical condition very rapidly deteriorated. However, his relatives refused surgical treatment, and 22 hours after admission, he died. Autopsy was not permitted.

TABLE 1. Accompanying Diseases orFactors in 67 Patients withInterhemispheric Subdural Hematoma

Disease or Factor

NO. OF


DISCUSSION

Incidence

The exact incidence of ISHs is unknown. Only 67 adult patients, including our patients, have been described in detail (1–4, 6–8, 10–16, 18–23, 25–46, 48– 50). The male-to-female ratio in adult patients was 1.87:1. In both sexes, ISHs

Alcoholism	
Bleeding diathesis	
von Willebrand's disease	
Hemophilia B	
Thrombocytopenia	
Oral anticoagulants	
Hydrocephalus caused by	
aquaduct stenosis	
Hypertension	
Chronic renal failure	and the second sec
Total	32

1212 Bartels et al.



CT has now replaced cerebral angiography. Angiography is mandatory only in nontraumatic ISHs (35). CT was performed on 55 patients (82.1%). In accordance with the series of Zimmerman et al. (52), hyperdensity in the interhemispheric fissure was seen in most patients (87.3%). Several conditions may cause hyperdensity in the interhemispheric fissure, including a normal falx, dural calcification, a subarachnoid hemorrhage, a subdural hematoma, and, in children, diffuse cerebral edema (52). Hypodensity and isodensity were each described once (23, 36). Pseudomembranes in an ISH, such as those observed in Patient 3, were described once (23). In three patients, the CT findings were not reported in detail. The right and left interhemispheric subdural spaces were affected equally. Bilateral ISHs were rare. Only three cases have been described in detail (6, 28, 48), although Zimmerman et al. (52) described the CT scans of another three cases.

male female

In two cases, sex or age was not mentioned.

FIGURE 3. Distribution of age related to sex.

subdural space over the convexity (6, 11, 19, 21, 23, 48, 52). If a solid clot liquefies, it may migrate into the space over the convexity and behave like a chronic subdural hematoma, as has been demonstrated by Ogsbury et al. (34).

tected in only 5 patients (27.8%). The frequency of cranial fractures in patients with subdural hematomas reported in the literature varies from 38 to 77% (5, 9, 17).

For many years, cerebral angiography was the diagnostic tool of choice. Characteristic features on the anterior-posterior view are an avascular space between the falx and the medial surface of the brain and a separation of the right and left callosal marginal branches above the free edge of the falx, if both anterior cerebral arteries are opacified (3, 41). Cerebral arteriography was performed in 28 patients and suggested a space-occupying lesion in the interhemispheric subdural space.

Treatment and outcome

Surgical treatment, including craniotomies (30 of 46 patients) and burr holes (13 of 46 patients), was performed on 46 patients, and in 3 patients, the surgical treatment was not specified. The remaining 21 cases were conservatively managed (*Table 2*). Conservative management and surgical intervention have been equally advocated. If the neurological condition of the patient is stable or if any accompanying diseases contraindicate surgery, conservative therapy is generally preferred. For the whole group of patients, the mortality rate after conservative

Signs and symptoms

The symptoms of the patients with ISHs included disturbances of consciousness (38.8%), hemiparesis (55.2%), monoparesis of the contralateral leg (10.5%), epileptic manifestations (generalized, 11.9%, and focal seizures, 3.0%), gait ataxia (3.0%), dementia (6.0%), language disorders (4.5%), and oculomotor nerve dysfunction (3.0%). The characteristic falx syndrome, consisting of contralateral hemiparesis affecting the leg much more than the arm, was present in 62.2% of the patients with hemiparesis. None of the patients were symptom free.

Many patients (23 of 60) were admitted within 24 hours after the trauma or onset of symptoms; 53 patients were admitted within 1 week, and for 7 patients, the time interval was not reported.

TABLE 2. Treatment versus Outcome^a

Tracture	Recovery				
Treatment	Complete Incomplete		Dead	Total	
Surgery					
Craniotomy	16	2	9	27	
Burr holes	7	2	4	13	
Unspecified	2			2	
Subtotal	25	4	13	42	
Conservative	14	3	2	19	
Total	39	7	15	61	
(%)	63.9%	11.5%	24.6%	100%	

Neuroradiology

Plain cranial radiographs were taken in 18 patients; a cranial fracture was de-

^a Outcome not mentioned in six cases.

Interhemispheric Subdural Hematoma 1213

management was lower than that after surgical treatment (10.5% and 31.7%, respectively). The difference in outcome did not reach statistical significance $(\chi^2 = 5.8572; P = 0.1188)$. However, the clinical conditions of the patients who were surgically treated were worse than those of the group who were conservatively treated. Twenty-five patients who underwent surgery experienced disturbed consciousness of fairly long duration preoperatively, whereas only five of the conservatively managed patients had clouded consciousness (Table 3). Similar proportions of patients without long-term clouding of the consciousness were treated conservatively or surgically. The outcomes were similar. The level of consciousness immediately after the traumatic event did not have any influence on the outcome of surgical or conservative treatment (*Table* 4). Craniotomies and burr holes resulted in similar outcomes. From these data, it is not possible to conclude which treatment is best for ISHs. It is clear that more patients with disturbed consciousness undergo operations. More detailed information on the level of consciousness of these patients is necessary to make a good comparison.

TABLE 4. Correlation between Loss of Consciousness at the Event Responsible for the Development of Interhemispheric Subdural Hematoma (Initial Loss of Consciousness) and the Outcome of Therapy^a

	Outcome			T !	
	Complete Recovery	Incomplete Recovery	Dead	Total	
Loss of consciousness		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			
Surgery	11	2	2	15	
Conservative	6	2	1	9	
No loss of consciousness					
Surgery	11	2	8	21	
Conservative	7	1	1	9	
Total	35	7	12	54	

^aIn six cases the outcome was unknown, whereas in another seven cases loss of consciousness was unknown.

Some authors say that surgery is always warranted, irrespective of the clinical condition of the patient, whereas others only advocate surgery if the clinical condition deteriorates. On the basis of personal experience and our review of the literature, we advise conservative management for patients without consciousness disturbances and for patients in stable clinical conditions at admission. However, if there is any deterioration, especially in consciousness (Patient 3), surgery should be immediately performed.

The overall mortality was 15 of 61 patients, or 24.6% (in 6 patients, the outcome was not mentioned). When only acute and subacute ISHs (traumatic event within the past 10 days) were considered, the mortality increased to 15 of 56 patients or 26.8%. This is lower than the lowest mortality rate described for traumatic acute subdural hematoma, which ranged from 30 to 90% (24, 47). In the latter studies, the patients underwent aggressive surgical treatment. Acute and subacute ISHs seem to be more "benign" forms of acute subdural

REFERENCES

- Aring CD, Evans JP: Aberrant location of subdural hematoma. Arch Neurol Psychiatry 44: 1296–1306, 1940.
- Borzone M, Rivano C, Capuzzo T, Altomonte M: Interhemispheric subdural hematoma: Case report. Zentralbl Neurochir 47:354–361, 1986.
- 3. Campbell JA, Campbell RL: Angiographic diagnosis of traumatic head and neck lesions. JAMA 175:761–768, 1961.
- 4. Clein LJ, Bolton CF: Interhemispheric subdural haematoma: A case report. J Neurol Neurosurg Psychiatry 32:389–392, 1969.
- 5. Cooper PR, Ho V: Role of emergency skull x-ray films in the evaluation of the head-injured patient: A retrospective study. Neurosurgery 13: 136–140, 1983.
- 6. Cronin TG, Shippey DU: Bilateral interhemispheric subdural hematoma: A case report. AJNR Am J Neuroradiol 8:909–910, 1987.
- 7. Delfini R, Santoro A, Innocenzi G, Ciappetta P, Salvati M, Zamponi C: Interhemispheric subdural hematoma (ISH): Case report. J Neurosurg Sci 35:217–220, 1991. 8. Drábek P, Kuba J, Wolný E: Interhemisferální subdurální hematom. Cesk Neurol Neurochir 47:268–272, 1984. 9. Echlin FA, Sordillo SVR, Garvey TQ Jr: Acute, subacute, and chronic subdural hematoma. JAMA 161:1345-1350, 1956. 10. Esposito DP, Devkota J, El Gammal T, Sullivan HG: Interhemispheric subdural hematoma. South Med J 77:379-381, 1984. 11. Fearnside MR, Hall K, Sengupta RP: Interhemispheric subdural haematoma following head injury. Aust N Z J Surg 49:678-680, 1979. 12. Fein JM, Rovit RL: Interhemispheric subdural hematoma secondary to hemorrhage from a calloso-marginal artery aneurysm. Neuroradiology 1:183–186, 1970. 13. Friedman MB, Brant-Zawadzki M: Interhemispheric subdural hematoma from ruptured aneurysm. Comput Radiol 7:129-134, 1983.

hematoma.


Received, June 3, 1994.

Accepted, November 11, 1994. **Reprint requests:** W.I.M. Verhagen, M.D., Ph.D., Department of Neurology, Canisius Wilhelmina Hospital, Weg door Jonkerbos 100, 6532 SZ Nijmegen, The Netherlands.

TABLE 3. Outcome in Relation to Long-term Clouding of the Consciousness and Therapy^a

	Outcome			
	Complete Recovery	Incomplete Recovery	Dead	Total
Clouding				**************************************
Surgery	12	2	10	24
Conservative	1	2	2	5
No clouding				
Surgery	13	2	2	17
Conservative	13	7		14

14. Fruin AH, Juhl GL, Taylon C: Interhemispheric subdural hematoma: Case report. J Neurosurg



Bartels et al. 1214

port of three autopsy cases. No Shinkei Geka 9:349-355, 1981.

- 16. Furui T, Iwata K, Usui K: Interhemispheric subdural haematoma complicated by intracerebral haematoma: Case report. Acta Neurochir (Wien) 91:139–141, 1988.
- 17. Galbraith S, Smith J: Acute traumatic intracranial haematoma without skull fracture. Lancet 1:501-503, 1976.
- 18. Gannon WE: Interhemispheric subdural hematoma: Case report. J Neurosurg 18:829-830, 1961.
- 19. Gartman JJ Jr, Atstupenas EA, Vollmer DG, Powers SK: Traumatic laceration of pericallosal artery resulting in interhemispheric subdural hematoma: A case report. J Emerg Med 7:603-610, 1989.
- 20. Gentile SL, Rivano C, Rosa M, Viale GL: Sugli

- 34. Ogsbury JS, Schneck SA, Lehman RAW: Aspects of interhemispheric subdural haematoma, including the falx syndrome. J Neurol Neurosurg Psychiatry 41:72-75, 1978.
- 35. Pozzati E, Gaist G, Vinci A, Poppi M: Traumatic interhemispheric subdural hematomas. J Trauma 22:241–243, 1982.
- 36. Pujadas F, Ricart C, Oliveras C, Pou A: Hematoma subdural agudo interhemisférico. Rev Clin Esp 179:70–71, 1986 (letter).
- 37. Russell NA, del Carpio-O'Donovan R, Mallya KB, Benoit BG, Belanger G: Interhemispheric subdural hematoma. Can J Neurol Sci 14:172-174, 1987.
- 38. Samii M, Schindler E, Hey O: Ein subdurales Hämatom des Interhemisphärenspalts. Acta Neurochir (Wien) 30:319–326, 1974.
- 39. Satoh T, Yamamoto Y, Asari S, Sadamoto K:

CT: Part II—Recognition and differentiation of interhemispheric subarachnoid and subdural hemorrhage. AJNR Am J Neuroradiol 3:635-642, 1982.

COMMENTS

The authors have provided three case reports of patients with interhemispheric subdural hematomas, as well as a comprehensive review of the current literature. The uniqueness of interhemispheric subdural hematomas in the general context of acute subdural hematomas is emphasized. Interhemispheric

- ematomi sottodurali acuti interemisferici. Rass Arch Chir 7:165–170, 1969.
- 21. Glista GG, Reichman OH, Brumlik J, Fine M: Interhemispheric subdural hematoma. Surg Neurol 10:119–122, 1978.
- 22. Ho SU, Spehlmann R, Ho HT: CT scan in interhemispheric subdural hematoma: Clinical and pathological correlation. Neurology 27:1097-1098, 1977.
- 23. Houtteville JP, Toumi K, Theron J, Derlon JM, Benazza A, Hubert P: Interhemispheric subdural haematomas: Seven cases and review of the literature. Br J Neurosurg 2:357–367, 1988.
- 24. Howard MA III, Gross AS, Dacey RG Jr, Winn HR: Acute subdural hematomas: An age-dependent clinical entity. J Neurosurg 71:858--863, 1989.
- 25. Isfort A: Angiographische Befunde beim Hämatom und Empyem im Interhemisphärenspalt. Fortschr Geb Röntgenstr 107:127–130, 1967.
- 26. Izawa M, Takahashi K, Sentoh S: A case of interhemispheric subdural hematoma associated with thrombocytopenia. No Shinkei Geka 10:557-560, 1982.

- Traumatic interhemispheric subdural hematoma: Report of a case and analysis of 7 cases. No Shinkei Geka 10:667-672, 1982.
- 40. Shigemori M, Kawaba T, Shirahama M, Tokutomi T, Hara K: Acute interhemispheric subdural hematoma: Report of 2 cases. No Shinkei Geka 10:647–652, 1982.
- 41. Sibayan RQ, Gurdjian ES, Thomas LM: Interhemispheric chronic subdural hematoma: Report of a case. Neurology 20:1215–1218, 1970.
- 42. Takeda N, Kurihara E, Matsuoka H, Kose S, Tamaki N, Matsumoto S: Three cases of acute interhemispheric subdural hematoma. No Shinkei Geka 16:87–92, 1988.
- 43. Tietjen DP, Moore J Jr, Gouge SF: Hemodialysis-associated acute subdural hematoma: Interim management with peritoneal dialysis. Am J Nephrol 7:478–481, 1987.
- 44. Tijssen CC, Tavy DLJ, Hekster REM, Bots GTAM, Endtz LJ: Aphasia with a left frontal interhemispheric hematoma. Neurology 34:1261-1264, 1984.
- 45. Toffol GJ, Biller J, Adams HP Jr: Interhemispheric subdural hematoma. J Emerg Med
- subdural hematomas primarily are a disease of the elderly. They primarily occur as a result of low velocity impacts, and in two-thirds of the reported cases, their occurrence has been associated with various bleeding disturbances (i.e., anticoagulation). Although a mortality rate of 25% as quoted by the authors would seem to be generally low for acute subdural hematomas, those patients with interhemispheric subdural hematomas and altered levels of consciousness actually had a 42% mortality rate. The authors' cogent review of the incidence, pathogenesis, signs and symptoms, neuroradiological features, treatments, and outcomes from interhemispheric subdural hematomas in adults should be of interest to all who

- 27. Jacobsen HH: An interhemispherically situated haematoma: Case report. Acta Radiol 44:235-236, 1955.
- 28. Kasdon DL, Magruder MR, Stevens EA, Paullus WS: Bilateral interhemispheric subdural hematomas. Neurosurgery 5:57–59, 1979.
- 29. Koyama S, Nishimura T: A case of bilateral interhemispheric subdural hematoma. No Shinkei Geka 18:289–294, 1990.
- 30. Kusunoki T, Nogaki H, Masurama M: Traumatic (para) interhemispheric hematoma. Prog CT (Tokyo) 2:433–435, 1980.
- 31. Lusins JO, McChesney JD: Nonsurgical management of an isolated interhemispheric subdural hematoma sequentially followed by CT. Mt Sin J Med 56:68–70, 1989.
- 32. Nagasawa S, Ohtsuki H, Nagayasu S, Kikuchi H: Acute interhemispheric subdural hematoma: A case report with good result by small craniotomy during the chronic stage. No Shinkei Geka 18:643–646, 1990.
- 33. New PFJ, Scott WR, Schnur JA, Davis KR, Taveras JM: Computerized axial tomography with the EMI scanner. Neuroradiology 110:109–123,

4:103-107, 1986.


- 46. Vilalta J, Sahuquillo J, Rubio E: Hematoma subdural interhemisférico. Rev Clin Esp 180:410-411, 1987 (letter).
- 47. Wilberger JE Jr, Harris M, Diamond DL: Acute subdural hematoma: Morbidity, mortality, and operative timing. J Neurosurg 74:212–218, 1991.
- 48. Woimant F, Thurel C, Roux FX, Cremer F, Bertrand HJ, Haguenau M, Houdart R, Pépin B: Hématomes sous-duraux aigus interhémisphériques. Rev Neurol (Paris) 139:299-303, 1983.
- 49. Wollschlaeger PB, Wollschlaeger G: The interhemispheric subdural or falx hematoma. Am J Roentgenol 92:1252-1254, 1964.
- 50. Yamagami T, Handa H, Nagasawa S, Nagata H, Chung KJ: Acute interhemispheric subdural hematoma: Case report and review of literature. Arch Jpn Chir 56:69–75, 1986.
- 51. Zimmerman RA, Bilaniuk LT, Bruce D, Schut L, Uzzell B, Goldberg HI: Interhemispheric acute subdural hematoma: A computed tomographic manifestation of child abuse by shaking. Neuroradiology 16:39-40, 1978.
- 52. Zimmerman RD, Russell EJ, Yurberg E, Leeds

treat head trauma.

Jack E. Wilberger Pittsburgh, Pennsylvania

The authors have reported 3 cases of their own and have reviewed the reports to date on 64 other cases of interhemispheric subdural hematomas in adults. The article provides much useful information, including the high incidence of this condition in patients with bleeding disorders and in patients taking oral anticoagulants. The data show that many of these patients can be conservatively managed, although patients with clouding of consciousness or deteriorating neurological deficits require surgical treatment.

Charles H. Tator



NE: Falx and interhemispheric fissure on axial

Toronto, Ontario, Canada