Radio continuum emission in the northern Galactic plane: Sources and spectral indices from the THOR survey*

Y. Wang¹, S. Bihr¹, M. Rugel¹, H. Beuther¹, K. G. Johnston², J. Ott³, J. D. Soler¹, A. Brunthaler⁴, L. D. Anderson^{5,6,7}, J. S. Urquhart⁸, R. S. Klessen^{9,10}, H. Linz¹, N. M. McClure-Griffiths¹¹, S. C. O. Glover⁹, K. M. Menten⁴, F. Bigiel⁹, M. Hoare², and S. N. Longmore¹²

- ¹ Max-Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany e-mail: wanq@mpia.de
- School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, UK
- ³ National Radio Astronomy Observatory, PO Box O, 1003 Lopezville Road, Socorro, NM 87801, USA
- ⁴ Max-Planck-Institut f
 ür Radioastronomie, Auf dem H
 ügel 69, 53121 Bonn, Germany
- ⁵ Department of Physics and Astronomy, West Virginia University, Morgantown, WV 26506, USA
- Adjunct Astronomer at the Green Bank Observatory, P.O. Box 2, Green Bank WV 24944, USA
- Center for Gravitational Waves and Cosmology, West Virginia University, Chestnut Ridge Research Building, Morgantown, WV 26505, USA
- ⁸ School of Physical Sciences, University of Kent, Ingram Building, Canterbury, Kent CT2 7NH, UK
- ⁹ Universität Heidelberg, Zentrum für Astronomie, Institut für Theoretische Astrophysik, Albert-Ueberle-Str. 2, 69120 Heidelberg, Germany
- ¹⁰ Universität Heidelberg, Interdisziplinäres Zentrum fur Wissenschaftliches Rechnen, INF 205, 69120, Heidelberg, Germany
- 11 Research School of Astronomy and Astrophysics, The Australian National University, Canberra, ACT, Australia
- Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool Science Park, 146 Brownlow Hill, Liverpool L3 5RF, UK

Received xxxx; accepted xxxx

ABSTRACT

Context. Radio continuum surveys of the Galactic plane can find and characterize H II regions, supernova remnants (SNRs), planetary nebulae (PNe), and extragalactic sources. A number of surveys at high angular resolution ($\leq 25''$) at different wavelengths exist to study the interstellar medium (ISM), but no comparable high-resolution and high-sensitivity survey exists at long radio wavelengths around 21 cm.

Aims. Our goal is to investigate the 21 cm radio continuum emission in the northern Galactic plane at <25" resolution.

Methods. We observed a large fraction of the Galactic plane in the first quadrant of the Milky Way ($l = 14.0 - 67.4^{\circ}$ and $|b| \le 1.25^{\circ}$) with the Karl G. Jansky Very Large Array (VLA) in the C-configuration covering six continuum spectral windows. These data provide a detailed view on the compact as well as extended radio emission of our Galaxy and thousands of extragalactic background sources. Results. We used the BLOBCAT software and extracted 10916 sources. After removing spurious source detections caused by the sidelobes of the synthesised beam, we classified 10387 sources as reliable detections. We smoothed the images to a common resolution of 25" and extracted the peak flux density of each source in each spectral window (SPW) to determine the spectral indices α (assuming $I(\nu) \propto \nu^{\alpha}$). By cross-matching with catalogs of H II regions, SNRs, PNe, and pulsars, we found radio counterparts for 840 H II regions, 52 SNRs, 164 PNe, and 38 pulsars. We found 79 continuum sources that are associated with X-ray sources. We identified 699 ultrasteep spectral sources ($\alpha < -1.3$) that could be high-redshift galaxies. Around 9000 of the sources we extracted are not classified specifically, but based on their spatial and spectral distribution, a large fraction of them is likely to be extragalactic background sources. More than 7750 sources do not have counterparts in the SIMBAD database, and more than 3760 sources do not have counterparts in the NED database.

Conclusions. Studying the long wavelengths cm continuum emission and the associated spectral indices allows us to characaterize a large fraction of Galactic and extragalactic radio sources in the area of the northern inner Milky Way. This database will be extremely useful for future studies of a diverse set of astrophysical objects.

Key words. catalogs – surveys – radio continuum: general – techniques: interferometric

1. Introduction

A number of surveys at high angular resolution ($\leq 20''$) at different wavelengths exist to study the interstellar medium

(ISM), from infrared (e.g., UKIDSS¹, Lucas et al. 2008; *Spitzer*/GLIMPSE², Benjamin et al. 2003; Churchwell et al. 2009, *Spitzer*/MIPSGAL³, Carey et al. 2009), to (sub)mm (e.g.,

^{*} The full continuum catalog, full table of Table 4 and all the fits files of the continuum data are available at the project website http://www2.mpia-hd.mpg.de/thor/DATA/www/

UKIRT Infrared Deep Sky Survey

² Galactic Legacy Infrared Midplane Survey Extraordinaire

³ A 24 and 70 Micron Survey of the Inner Galactic Disk with MIPS

ATLASGAL⁴ and BGPS⁵, Schuller et al. 2009; Rosolowsky et al. 2010; Aguirre et al. 2011; Csengeri et al. 2014) and radio (e.g. MAGPIS⁶, CORNISH⁷, Helfand et al. 2006; Hoare et al. 2012) wavelengths. Previously, the best 21 cm H_I line survey was the HI Very Large Array Galactic Plane Survey (VGPS, Stil et al. 2006) which has a resolution of 60", significantly more coarse than the resolution of the aforementioned surveys. This was one of the motivations for initiating "The HI, OH, Recombination line survey of the Milky Way (THOR) 8" (Beuther et al. 2016). Using the Karl G. Jansky Very Large Array (VLA) in C-configuration, we achieve a spatial resolution of < 25". The WIDAR correlator at the VLA allows us to observe many spectral lines simultaneously, in particular several molecular OH transitions, a series of $Hn\alpha$ radio recombination lines (RRLs. n = 151 to 186), as well as eight spectral windows (SPWs) to cover the continuum emission between 1 and 2 GHz (Bihr et al. 2015; Beuther et al. 2016; Bihr et al. 2016; Rugel et al. 2018). We observed a large fraction of the Galactic plane in the first quadrant of the Milky Way $(l = 14.0 - 67.4^{\circ})$ and $|b| \le 1.25^{\circ}$ in several semesters (from 2012 to 2014). The continuum data from the first half of the survey $(l = 14.0 - 37.9^{\circ} \text{ and } l = 47.1 - 51.2^{\circ})$ have been published by Bihr et al. (2016). In this paper, we combine all the continuum data and present the results from the full survey.

The radio continuum emission from 1 to 2 GHz is dominated by thermal free-free emission from electrons and non-thermal synchrotron emission of the relativistic electrons in magnetic fields (e.g., Wilson et al. 2013). These can be distinguished by the spectral index α , assuming $I(\nu) \propto \nu^{\alpha}$, where $I(\nu)$ is the intensity at frequency ν . The thermal free-free emission shows an almost flat ($\alpha = -0.1$) spectrum if it is optically thin, or positive spectral index if it is optically thick with values varying between -0.1 and 2 (e.g., Keto 2003; Wilson et al. 2013). H II regions and planetary nebulae are often the sources for thermal free-free emission. In contrast to this, synchrotron emission shows a negative spectral index whose value depends, amongst other things, on the particle energy distribution. Towards extragalactic jets powered by an active galactic nucleus (AGN), one often finds the synchrotron emission with a spectral index $\alpha < -0.5$ (e.g., Hey 1971; Rybicki & Lightman 1979). Galactic SNRs often show synchrotron emission with a spectral index around -0.5 (e.g., Bhatnagar et al. 2011; Green 2014; Reynoso & Walsh 2015). Thus the spectral index can help us to characterize the nature of the continuum sources we detected in the survey. This allows us to determine whether they are Galactic or extragalactic, which is crucial for H I and OH absorption studies.

Compact galactic radio sources associated with X-ray emission are usually Pulsar Wind Nebulae (e.g., Brisken et al. 2005; Miller et al. 2005), or X-ray binaries (XRB or microquasars; Mirabel & Rodríguez 1998, 1999). By investigating the X-ray and radio flux ratios, the spectral indices and observations at other (optical/infrared) wavelengths of the Galactic sources in detail, we can also constrain the type of the sources, i.e., low mass XRB, PN, pulsar etc. (e.g., Seaquist 1993; Maccarone et al. 2012; Tetarenko et al. 2016). With the high-angular resolution (< 25") of our THOR continuum data, we can not only derive

the spectral indices of the sources, but also further study the variation in frequency and space.

The paper is structured as follows: In Sect. 2, we present the observations and data reduction. Sect. 3 presents the methods we used to extract sources and determine the spectral indices. Sect. 4 describes the continuum catalog, and the distribution of the continuum sources we extracted. The nature of continuum sources are discussed in Sect. 5. The conclusions and summary are presented in Sect 6. The appendix gives additional information of the continuum observations and tables.

2. Observations and data reductions

We observed a part of the first quadrant of the Galactic plane ($l=14.0-67.4^{\circ}$ and $|b| \leq 1.25^{\circ}$) with the *Karl G. Jansky* Very Large Array (VLA) in C configuration at L band from 1 to 2 GHz. The detailed observing strategy and data reduction is discussed and described in Bihr et al. (2016) and Beuther et al. (2016). With the WIDAR correlator, we cover the H _I 21 cm line, 4 OH lines, 19 H α recombination lines, as well as eight continuum bands, i.e., SPWs. Each continuum SPW has a band width of 128 MHz. Due to strong contamination from radio frequency interference (RFI), two SPWs around 1.2 and 1.6 GHz were not usable and discarded. The remaining six SPWs are centered at 1.06, 1.31, 1.44, 1.69, 1.82 and 1.95 GHz. For the fields at $l=23.1-24.3^{\circ}$ and 25.6 -26.8° , the SPW around 1.95 GHz is also severely affected by RFI and is therefore flagged (see Bihr et al. 2016).

The calibration and imaging were performed with the CASA⁹ software package. We employed the RFlag algorithm which was first introduced to AIPS by E. Greisen in 2011 to flag outliers in each visibility dataset before imaging. For imaging, we chose a pixel size of 2.5'' and robust = 0 as a weighting parameter, which results in a synthesized beam size varying from 9" to 25" depending on the SPW and the declination of different observing blocks and uv-coverage (see Table 1 for details). We also used the multiscale CLEAN in CASA in order to recover the large scale structure better, and chose the scale parameters as 1, 3 and 6 times the resolution element. The cleaning process was set to stop at a threshold of 5 mJy beam⁻¹ or 10⁵ iterations, whichever was reached first. The noise of our data is dominated by the artifacts resulting from residual sidelobes, and varies from ~ 0.3 to > 1 mJy beam⁻¹ depending on the frequency and sky position. The thermal noise is $\sim 0.1 \text{ mJy beam}^{-1}$. We will discuss the noise level of the images in Sect. 3.

THOR+VGPS data: The 1.4 GHz continuum data from the VGPS survey (Stil et al. 2006) combined VLA D-configuration with single-dish observations from Effelsberg, and have an angular resolution of 60". The $14.0^{\circ} < l < 17.5^{\circ}$ Galactic longitude range in the VGPS data is just comprised of single-dish observations. We smoothed the THOR continuum data at 1.4 GHz to a resolution of 25", and used the task "feather" in CASA to combine the THOR data with those from VGPS. While this combined dataset retains the high angular resolution of the THOR observations, it can recover the large scale structure. The combined image of the whole survey at 1.4 GHz in Fig. 1 shows that large scale H II regions and SNRs dominate the extended radio emission in the inner Galactic plane ($l < 55^{\circ}$), while compact sources are distributed across the whole survey area (more prominent in area $l > 55^{\circ}$). These compact sources are most likely extragalactic sources, and we will discuss them later.

⁴ APEX Telescope Large Area Survey of the Galaxy

⁵ Bolocam Galactic Plane Survey

⁶ Multi-Array Galactic Plane Imaging Survey

⁷ the Co-Ordinated Radio 'N' Infrared Survey for High-mass star formation

⁸ http://www.mpia.de/thor/Overview.html

⁹ http://casa.nrao.edu

Anderson et al. (2017) used this dataset and identified 76 new Galactic SNR candidates in the survey area. Although the spectral band of the VGPS continuum (~1 MHz) is different from our THOR continuum data (~128 MHz), Anderson et al. (2017) showed that the flux retrieved from the combined data is consistent with the literature by comparing the flux density of the known SNRs (Green 2014).

All the reduced continuum data including the THOR+VGPS dataset can be accessed from the THOR survey website located at http://www.mpia.de/thor.

3. Source extraction and spectral index determination

While for the 1.4 GHz the combined THOR+VGPS dataset exists, for the other bands only the THIR data are available. Therefore, all spectral index analysis is done on the THOR-only data. To achieve the best signal-to-noise ratio, we chose the two SPWs, spw-1820 and spw-1440, that are least effected by RFI and have the lowest noise, smoothed them into the same resolution, and then averaged them for source extraction. Since the noise of our data is dominated by the sidelobe noise, we followed the method described in Hales et al. (2012) and Bihr et al. (2016) and constructed noise maps using the averaged residual image from the clean process. The noise maps are shown in Fig. A.1. We furthermore calculated the cumulative fraction noise level map area at a specific 7σ noise level in mJy beam⁻¹ (Fig. 2). While the lowest 7σ noise level (≤ 1 mJy beam⁻¹, dominated by thermal noise) is achieved in ~ 20% of the survey area, more than 60% of the survey area has a 7σ noise level \leq 2 mJy beam⁻¹. Comparing to the first half of the survey (Bihr et al. 2016), the noise is better for the entire survey as in the large longitude Milky Way regions beyond Galactic longitudes of roughly 51° there are fewer strong sources (Fig. 1 and Fig. A).

We used the software BLOBCAT (Hales et al. 2012) to extract the sources from the averaged continuum images. Following the same criteria as in Bihr et al. (2016) for the first half of the survey, we set the detection threshold as 5σ and the flooding threshold to the standard value of 2.6σ (Hales et al. 2012), and extracted 10916 sources. We then inspected each source visually, and identified 530 sources as obvious observational sidelobe artifacts and removed them from the catalog. The remaining 10387 sources should be mostly real detections. It was difficult to determine whether some sources with a signal-to-noise ratio between 5 and 7σ were real. Therefore, we consider sources with a signal-to-noise ratio higher than 7σ with higher confidence. In total, out of the 10916 extracted sources, 7521 sources were detected with a signal-to-noise ratio higher than 7σ , 2866 have a signal-to-noise ratio between 5 and 7σ , 530 are observational sidelobe artifacts.

Following the same method described in Bihr et al. (2016), we performed a completeness test by extracting artificial compact sources from a $0.5^{\circ} \times 0.5^{\circ}$ region with a constant noise level. The result shows that we detected 94% of all sources with a peak intensity above 7σ (Fig. 3). Combining the noise map, we construct completeness maps for different peak intensities that are shown in Appendix B.

Since our observations cover a wide range in frequency from 1 to 2 GHz, we can determine the spectral indices for the sources we identified, assuming $I(v) \propto v^{\alpha}$. To do this, we first smooth all SPWs to a common resolution of 25", then extract the peak intensities. Since the integrated flux density suffers more from filtering effects for extended sources, we use only the peak intensity for the spectral index determination. We use the scipy

function "curve_fit" to fit the peak intensity between 1 to 2 GHz of each source we extracted and derive the spectral index α defined as $I(v) \propto v^{\alpha}$ together with the uncertainty of α . Since a single SPW image has a lower S/N than the averaged image we used for source extraction, we choose a lower threshold (3σ) for spectral index determination. We fit only the peak intensities that are higher than 3 times of the noise level of the respective SPW, i.e., a reliable intensity (see also Bihr et al. 2016). For 8228 sources we can extract a reliable peak intensity from at least two SPWs, and derive a spectral index. We list the number of SPWs that were used for spectral index determination in the continuum source catalog as "fit_spws". For instance, "fit_spws"=6 means the peak intensity for all 6 SPWs is reliable and used for spectral index determination, "fit spws"=4 means the peak intensity for only 4 SPWs is reliable and used for spectral index determination. Figure 4 shows the distribution of the uncertainties of the determined spectral index for all sources and for sources with a different number of SPWs. This clearly demonstrates that the more SPWs used to fit the spectral index, the smaller the uncertainty is. The mean uncertainty of the spectral indices are $\Delta \alpha = 0.18, 0.43, 0.70, 1.1, 2.2$ for sources detected in 6, 5, 4, 3 and 2 SPWs, respectively. We consider all spectral indices fitted with 4 or more SPWs (fit_spws \geq 4) reliable.

4. Catalog

Our continuum catalog contains 31 columns for each source as described in detail in Table 2. We summarize the number of extracted sources in Table 3. Due to the BLOBCAT algorithm, compact sources close to each other or superimposed on large extended emission are identified as one single source. Additionally, large, extended sources, such as SNRs, can be separated into several different sources. As the two examples shown in Fig. 5, two H II regions are identified into one single object in our catalog and the SNR W44 is resolved into 18 separate objects. Therefore, the exact numbers in Table 3 should be treated cautiously. We cross-matched the THOR continuum catalog with catalogs of different type of Galactic sources (H II regions, SNRs, etc., see Sect. 5) and list all the matched counterparts. For a source matched with multiple sources, we also list the number of counterparts that the continuum source is associated with in column "Ncounter" in the catalog (Table 2). We also categorise sources in groups based on counterparts they share, i.e., sources associated with the same H II regions or SNRs are in the same group with a "GroupID". The size of the groups are also listed as "GroupSize" (Table 2). In total, 126 sources have more than 1 counterparts, 153 sources are categorised into 49 groups with GroupSize varies from 2 to 19. We will discuss the association between different Galactic and extragalactic sources in detail in Sect. 5.

The distribution of the extracted sources along Galactic Longitude and latitude is shown in Fig. 6. The majority (79%, Table 3) of the extracted sources are unresolved, and a weak gradient for the those sources along Galactic longitude is shown in Fig. 6, where more weak unresolved sources are identified in the larger longitude regions. The resolved sources are evenly distributed along Galactic longitude, but larger sources ($n_{pix} > 500$, effective radius of $\sim 32^{\prime\prime}$) are more concentrated in the inner Galaxy, since these are mostly Galactic H II regions and SNRs. Another possible reason is that there are more mergers of smaller sources in the inner longitudes because of higher source density and longer path-lengths through the Galaxy. The distribution along Galactic latitude reveals that the resolved sources are concentrated close to the Galactic mid-plane, whereas the dis-




Fig. 1. THOR+VGPS 1.4 GHz continuum map of the whole survey. The yellow and black circles mark the WISE H π regions (Anderson et al. 2017) matched with our continuum sources (see also Sect. 5.3). The red circles mark the SNRs from Green (2014). The green circles mark the new SNR candidates identified in this combined dataset (Anderson et al. 2017). The synthesized beam size is 25". For regions $l < 17.5^{\circ}$, complementary D-configuration data do not exist. Therefore, in these regions we have only combined the THOR data with the Effelsberg data.

Table 1. Synthesized beams of SPWs.

SPW	Frequency range	Restoring beam size
	[MHz]	
spw-1060	989 – 1117	$24.4'' \times 15.1''$ to $15.1'' \times 14.7''$
spw-1310	1244 - 1372	$19.7'' \times 12.5''$ to $12.6'' \times 12.2''$
spw-1440	1372 - 1500	$18.1'' \times 11.1''$ to $12.0'' \times 11.6''$
spw-1690	1628 – 1756	$15.4'' \times 9.1''$ to $9.8'' \times 9.5''$
spw-1820	1756 – 1884	$14.5'' \times 8.9''$ to $9.2'' \times 9.1''$
spw-1950	1884 - 2012	$13.1'' \times 8.1''$ to $8.6'' \times 8.2''$
averaged image	spw-1440 & spw-1820	$18.1'' \times 11.1''$ to $12.0'' \times 11.6''$
THOR+VGPS	1420	$25'' \times 25''$

Notes. The averaged images are used for source extraction. The size of the restoring beam varies due to declination and uv-coverage differences among observing blocks.




Fig. 2. Cumulative fraction noise level diagram. The percentage of the map area as a function of the noise level at a S/N of 7σ in mJy beam⁻¹. More than 60% of the survey area has a noise level of $7\sigma \lesssim 2$ mJy beam⁻¹.




Fig. 3. Completeness test plot. Percentage of the added artificial sources detected as a function of the source peak intensity in units of the noise level σ .

tribution of the unresolved sources shows a dip at $|b| < 0.5^{\circ}$. The distribution of both resolved and unresolved sources drops at $|b| > 1.0^{\circ}$ due to higher noise at the edges of the survey area.

We did a Kolmogorov–Smirnov test (KS test) for the dip in the distribution of the unresolved sources at $|b| < 0.5^{\circ}$. We first re-binned the source counts into 100 bins and measured the mean value and the standard deviation of the distribution at $|b| > 0.5^{\circ}$ and $|b| < 1.0^{\circ}$. Then we generated a random artificial distribu-




Fig. 4. Histogram of the uncertainties of the determined spectral indices. The black line includes all sources for which we are able to determine a spectral index, the grey shaded area represents the sources for which we have an intensity measurement in all six SPWs and therefore the spectral index α is derived from fitting all six SPWs, the red line includes sources fitted with five SPWs, the blue line includes sources fitted with four SPWs, the green line includes sources fitted with three SPWs, and the purple line includes sources fitted with two SPWs.

tion with the same mean value and standard deviation. Finally we compute the KS statistic and P-value between the random artificial and the real distribution with the scipy function "ks_2samp". The mean KS statistic and p-value for 100 run is 0.34 ± 0.041 and 0.001 ± 0.0018 , respectively. So the dip at $|b|<0.5^{\circ}$ is statistically significant. A similar distribution along Galactic latitude is also found in the first half of our survey (Bihr et al. 2016) and in the MAGPIS survey (Helfand et al. 2006). Less unresolved sources are identified at $|b|<0.5^{\circ}$ and in the inner Galaxy which could be due to that the strong emissions from the extended Galactic sources such as H II regions and SNRs lower our detection completeness to weak sources towards the Galactic mdi-plane and inner Galaxy (see Appendix B).

5. Discussion

5.1. Comparison with other surveys

The comparison between the peak positions of the common sources in the THOR survey (first half, $l = 14.0 - 37.9^{\circ}$ and $l = 47.1 - 51.2^{\circ}$), the MAGPIS (Helfand et al. 2006), and the CORNISH survey (Hoare et al. 2012; Purcell et al. 2013) presented in Bihr et al. (2016) shows that the THOR survey has positional accuracy better than 2.5" (FWHM of the Gaussian fit

Table 2. Description of the continuum source catalog entries.

Col. Num.	Name	Unit	Description
1	Gal. ID		Name of the source the form G"Gal. longitude"±"Gal. latitude" ^a .
2	RA	deg	Right ascension in J2000 of the peak position.
3	Dec	deg	Declination in J2000 of the peak position.
4	S_p^b	Jy beam ⁻¹	Peak intensity of the aver. image used for source extraction (see Sect. 3).
5	S_int	Jy	Integrated flux density of the averaged image (see Sect. 3).
6	S/N		Signal-to-noise ratio in the averaged image.
7	BMAJ	arcsec	Major axis of the synthesized beam used for source extraction.
8	BMIN	arcsec	Minor axis of the synthesized beam used for source extraction.
9	BPA	deg	Position angle of the synthesized beam used for source extraction.
10	n_pix		Number of pixels flooded by BLOBCAT (see Sect. 3).
11	resolved_source		Resolved source label (see Sec 3). $1 = \text{Resolved}$, $0 = \text{Point}$.
12	$S_p(spw-1060)^c$	Jy beam ⁻¹	Peak intensity around 1.06 GHz used for spectral index (see Sect. 3).
13	$delta_S_p(spw-1060)^c$	Jy beam ⁻¹	Uncertainty of peak intensity around 1.06 GHz.
14	$S_p(spw-1310)^c$	Jy beam ⁻¹	Peak intensity around 1.31 GHz used for spectral index (see Sect. 3).
15	$delta_S_p(spw-1310)^c$	Jy beam ⁻¹	Uncertainty of peak intensity around 1.31 GHz.
16	$S_p(spw-1440)^c$	Jy beam ^{−1}	Peak intensity around 1.44 GHz used for spectral index (see Sect. 3).
17	delta_S_p(spw-1440) ^c	Jy beam ⁻¹	Uncertainty of peak intensity around 1.44 GHz.
18	$S_p(spw-1690)^c$	Jy beam ⁻¹	Peak intensity around 1.69 GHz used for spectral index (see Sect. 3).
19	$delta_S_p(spw-1690)^c$	Jy beam ⁻¹	Uncertainty of peak intensity around 1.69 GHz.
20	$S_p(spw-1820)^c$	Jy beam ⁻¹	Peak intensity around 1.82 GHz used for spectral index (see Sect. 3).
21	$delta_S_p(spw-1820)^c$	Jy beam ⁻¹	Uncertainty of peak intensity around 1.82 GHz.
22	$S_p(spw-1950)^c$	Jy beam ⁻¹	Peak intensity around 1.95 GHz used for spectral index (see Sect. 3).
23	$delta_S_p(spw-1950)^c$	Jy beam ^{−1}	Uncertainty of peak intensity around 1.95 GHz.
24	alpha	•	Spectral index of source we derived (see Sect. 3).
25	delta_alpha		Uncertainty of spectral index.
26	fit_spws		Number of SPWs used to fit the spectral index (see Sect. 3).
27	Note ^d		"HII", "SNR_green", "SNR_anderson", "PN", "PSR", "Xray", "jets" (see Sect. 5).
28	Counterparts		The counterparts of the H II region ^e , SNR^f , the planetary nebula ^g and the pulsars ^h .
29	Ncounter		The total number of counterparts that the continuum source is associated with.
30	GroupID		The Group of continuum sources associated with one or more same counterparts.
31	GroupSize		The number of the continuum sources in the same group.

Notes. (a) Indicating the peak position. (b) The synthesized beam is different for different fields and is given in rows 7-9. (c) The synthesized beam is smoothed to 25"×25". (d) We classified the continuum source into different categories. "HII": sources associated with H II regions from Anderson et al. (2014); "SNR_green": sources associated with SNRs from Green (2014); "SNR_anderson": sources associated with SNR candidates from Anderson et al. (2017); "PN:" sources classified as planetary nebula; "PSR": sources classified as pulsars; "Xray": sources associated with X-ray sources; "jets": sources classified as extragalactic jets candidates. (e) Anderson et al. (2014). (f) Green (2014); Anderson et al. (2017). (g) Parker et al. (2016). (h) Manchester et al. (2005).

Table 3. Numbers of the catalog.

Description	Numbers	Percentage
All	10387	100%
S/N> 7σ	7521	72%
Resolved sources	2210	21%
Unresolved sources	8177	79%
$n_{pix} > 500$	439	4%
fit_spws≥ 4	5857	56%

to the position offset distribution, see Fig. 15 in Bihr et al. 2016). Considering that the pixel size of our continuum images is only 2.5", the spatial accuracy of the THOR survey is consistent with previous observations.

To check for consistency in the flux density, we compared the THOR with MAGPIS. Within our survey area ($14.2^{\circ} < l < 67.4^{\circ}$, $b < |1.25|^{\circ}$), the MAGPIS catalog contains 2256 discrete sources. We cross-matched our catalog with the MAGPIS catalog with a matching radius of 5", and we found 1440 matches. We then selected sources with the following criteria: 1) S/N>

3 in spw-1440; 2) unresolved in the THOR catalog; 3) major axis smaller than 10"in the MAGPIS catalog. We selected 735 sources using the aforementioned criteria. We compared the 1.4 GHz continuum flux density from the spw-1440 in THOR and the flux densities in MAGPIS, as show in Fig. 7. The MAG-PIS/THOR flux ratio shows a tight distribution around 1, with a median value of ~ 0.97 . Most of the sources are within 5σ from the flux ratio equals to one. 67 sources show a deviation larger than 7σ from one, among which 56 sources have a ratio smaller than one. If we compare the integrated flux from MAGPIS to the peak flux from THOR for these 56 sources, the mean ratio is 1.08. This indicates that these sources could have been resolved by MAGPIS. For the remaining 11 sources the ratios are smaller than 1.18, therefore we can not rule out the possibility of variability for these sources. Among these 11 sources, 2 are matched with X-ray sources (ratios are 1.02 and 1.08, see also Sect.5.7). The remaining 9 sources do not match with any identified source and all have negative spectral index, which suggests that they can be extragalactic sources.

With a matching radius of 5", we found 1320 unresolved THOR sources matched with NRAO VLA Sky Survey (NVSS,





Fig. 5. Examples of source identification in the catalog. In both panels, the white contours represent the area of the source extracted by BLOB-CAT overlaid on the averaged image of spw-1440 and spw-1820, and the crosses mark the sources in the catalog. *Top:* Source G47.100+0.479 consists of two H II regions (marked with white circles, Anderson et al. 2014) close together. *Bottom:* The SNR W44 (e.g., Green 2014) is resolved into 19 sources in our catalog. The synthesized beam is shown in the bottom left corner of each panel. The beam in the right panel is scaled up to 10 times of its original size.

Condon et al. 1998) sources. We compared the 1.4 GHz flux densities from THOR and NVSS, and the THOR flux at spw-1440 is also tightly correlated with the one from NVSS but slightly higher. The median value of the NVSS/THOR flux ratio is about 0.93. By comparing the THOR 1.4 and 1.8 GHz average flux density of the first half of the survey with the NVSS flux density, Bihr et al. (2016) found similar results (see also Figure 17 in Bihr et al. 2016). This discrepancy is yet undetermined, but this is beyond the scope of this work. The NVSS images show that the NVSS catalog is severely contaminated with obvious false detections, which could be due to sidelobes from strong sources close to the Galactic plane or ghost artifacts (Grobler et al. 2014).

For the second half of the THOR survey, we applied the same data reduction method and source extraction algorithm as the first half (Bihr et al. 2016). Since there are overlapping areas between the first and second half of the survey, we further compare






Fig. 6. Histograms for the source distribution along Galactic Longitude (top) and latitude (bottom). In both panels, the black histogram shows the distribution of all the sources in the catalog, the other histograms show fractions of the catalog as marked in each panel.

the position and flux density of the common sources between the two halves. 94% of sources have position differences <2.5"(the pixel size of the maps), and 93% of the sources have flux differences <1 mJy beam $^{-1}$. Therefore, the second half of the survey is consistent with the first half and with the previous radio surveys.

To verify the reliability of our spectral index determination, we extrapolated the flux density of selected sources to 5 GHz and compared this with the CORNISH catalog. The CORNISH catalog contains 2493 sources within the THOR survey area. With a matching radius of 5", we found 1905 matches between THOR and CORNISH. We selected unresolved sources with a THOR spectral index between 0 and -0.2 (optically thin free-free emission has a spectral index -0.1) that are detected in all six spectral windows in THOR (fit_spws=6). 68 sources were selected this way. We estimated the flux density at 5 GHz of these 68 sources according to their spectral indices from the THOR catalog. The extrapolated flux densities are very close to the flux densities in the CORNISH catalog as shown in Fig. 8. Sources with a spectral index between -0.09 and -0.11 (red in Fig. 8) are closer to ideal optically thin, and the extrapolated flux for these sources also agree better with the ones from CORNISH as expected.

Furthermore, we also compared the THOR spectral indices with spectral indices Kalcheva et al. (2018) derived for the CORNISH UCH II regions. Kalcheva et al. (2018) used flux density




Fig. 7. Ratio of the MAGPIS and THOR flux density plotted against the THOR flux density. The points are colored according to their deviation from a flux ratio of unity in units of the uncertainty in the ratio, so the deviation= $(1 - \text{ratio})/\sigma_{\text{ratio}}$. The dashed line represents a one-to-one relation.




Fig. 8. Extrapolated flux densities at 5 GHz for selected sources according to their spectral indices from the THOR catalog plotted against the flux densities in the CORNISH catalog. Sources with a spectral index between 0 and -0.2 are shown in blue, and with a spectral index -0.09 and -0.11 are shown in red. The dashed line represents a one-to-one relation.

measurements of CORNISH 5 GHz and MAGPIS 20 cm to derive the spectral indices. 11 unresolved THOR sources are associated with the CORNISH UCH II regions and have a reliable spectral index measurement in Kalcheva et al. (2018). The comparison again reveals that the spectral indices from these two studies are tightly correlated.




Fig. 9. Histogram of the spectral index for all reliably detected sources (fit_spws \geq 4) with a reliable spectral index measurement (\sim 5857 sources).

5.2. Spectral index

As described in Sect. 3, we derived spectral indices for 8228 sources, of which we consider the values for 5857 sources as being reliable (fit spws≥ 4). Using the spectral index information we can distinguish the physical origin of the emission and classify the continuum sources. Figure 9 shows the distribution of the spectral indices for all detected sources with a reliable spectral index measurement. For all sources the spectral index distribution shows a strong peak around $\alpha \sim -1$, and a secondary peak around $\alpha \sim 0$. If we consider only the unresolved sources, the spectral index distribution peaks around $\alpha \sim -1$, and the secondary peak around $\alpha \sim 0$ diminishes. Most of the unresolved sources show a negative spectral index indicating that they are dominated by non-thermal synchrotron radiation. Considering the unresolved sources are also evenly distributed along Galactic longitude and latitude (Sect. 4), these unresolved sources with a negative spectral index are likely to be mostly extragalactic sources. The spectral distribution of the resolved sources shows two clear peaks around $\alpha \sim -1$ and $\alpha \sim 0$. Most of the resolved sources with a flat or positive spectral index are Galactic H II regions (see Sect. 5.3). The ones with negative spectral index are mostly SNRs (see Sect. 5.4) and radio galaxies (see Sect. 5.8), or overlapping unresolved sources that were classified as one single source. Considering that only the very large sources that have an area in pixels larger than 500 ($n_{pix} > 500$, effective radius $\gtrsim 32''$), the spectral index distribution peaks around $\alpha \sim 0$ and extends to $\alpha \sim -1$. Among these large sources, about 80% of them are associated with HII regions, and the rest of them are classified as SNRs and radio galaxies (see Sect. 5.3, 5.4, and 5.8).

5.3. Н п regions

Using data from the all-sky *Wide-Field infrared Survey Explorer* (*WISE*) satellite, Anderson et al. (2014) made the most complete catalog of H $\scriptstyle\rm II$ regions to date, with a total of more than 8000 sources¹⁰. Within the region of the THOR survey, the *WISE* H $\scriptstyle\rm II$ catalog contains ~2400 sources, and ~1500 of them show radio continuum emission. The size of the H $\scriptstyle\rm II$ regions varies from 10" to 20' from the mid-infrared (MIR) images, so multiple Galac-

http://astro.phys.wvu.edu/wise/

tic and extragalactic sources could easily be enclosed within one single large H π region. Bihr et al. (2016) shows that matching the THOR continuum sources with the WISE catalog sources with r < 150'' in an automated fashion would produce less than 10% false matches. We take the same radius threshold when doing the automated matching, then we visually inspected the results and removed the false detections. We further visually compared the remaining large WISE sources (r > 150'') with our continuum images, and identified the continuum sources that are associated with H π regions.

In addition to the *WISE* catalog, 239 UCH II regions are identified in the CORNISH survey (Kalcheva et al. 2018), and 205 lie within the THOR coverage. With a matching radius of the effective radius equal to each THOR continuum source, 202 UCH II regions have counter parts in the THOR survey. For sensitivity reason, we do not detect three CORNISH UCH II regions (G024.1839+00.1199, G026.1094-00.0937, and G030.0096-00.2734, see Kalcheva et al. 2018). All the continuum sources that are associated with UCH II regions are also associated with at least one H II region in the *WISE* catalog except G26.008+0.137 and G37.735-0.113, so we list the CORNISH UCH II region counterparts for only these two continuum sources.

In total, we matched 713 continuum sources with H $\scriptstyle\rm II$ regions. Among the matched H $\scriptstyle\rm II$ regions, 16 are in the radio quiet group in the *WISE* catalog, which means no radio continuum emission is detected in the MAGPIS (Helfand et al. 2006) and VGPS (Stil et al. 2006) surveys. H $\scriptstyle\rm II$ regions close to each other could be identified as one single source in our catalog (left-panel, Fig. 5), in total 231 continuum sources are matched with more than one H $\scriptstyle\rm II$ region. In particular, G43.171+0.007 is matched with 19 H $\scriptstyle\rm II$ regions in W49, G19.610-0.235 is matched with 7 H $\scriptstyle\rm II$ regions, G45.122+0.132 encompasses 6 H $\scriptstyle\rm II$ regions, G48.610+0.027 encompasses 9 H $\scriptstyle\rm II$ regions, G49.370-0.302 encompasses 8 H $\scriptstyle\rm II$ regions (W51), and G49.488-0.380 encompasses 11 H $\scriptstyle\rm II$ regions (W51). The radius of the matched H $\scriptstyle\rm II$ regions measured at MIR in the *WISE* catalog varies from 12" to 20' with a median value of \sim 60".

We show in Fig. 10 one of the extreme cases, G43.171+0.007, which is associated with 19 H II regions in W49. By comparing ATLASGAL and CORNISH surveys, Urquhart et al. (2013) identified 18 UCH II regions associated with this region, and that this region has the highest UCH II surface density in the 1st quadrant. The spectral index map of W49 (Fig. 10) reveals a positive or flat spectral index towards most of the area of the source, except the edge regions, where the spectral index fitting is not reliable due to low S/N. This indicates that the H II region is dominated by thermal free-free emission. The spectral indices towards some strong continuum peaks are even larger than 0.5, this could indicate optically thick free-free emission.

To derive the physical properties of the H π regions, we fitted the spectral energy distribution (SED) of the H π region sources with a simple homogeneous H π region model with optical depth to get the emission measure (EM) of the H π regions (Sánchez-Monge 2011). We selected continuum sources with a S/N larger than 7σ , fit_spws = 6 (see Sect. 5.2), and spectral index $\alpha > -0.3$. We further limited our fitting sample to sources with an effective radius smaller than 60", since tests done by Bihr et al. (2016) show that we are able to recover sources with size up to ~ 120 " reasonably well (80% flux recovery). Among 713 continuum sources that are matched with H π regions, 262 match our selection criteria and are fitted. We fit the peak flux of the sources assuming that the line of sight size of the sources to




Fig. 10. Spectral index map for the H II region W49. *Top-panel:* Spectral index fitting result of the peak of the continuum source G43.171+0.007 (marked with a cross in the *middle-panel*). *Middle-panel:* Averaged image of spw-1440 and spw-1820, the white contours represent the area of the source extracted by BLOBCAT, and the cross marks the peak of the source G43.171+0.007 in the catalog. The circles mark the H II regions from Anderson et al. (2014) matched with the continuum sources in the THOR catalog. *Bottom-panel:* The spectral index map produced by fitting the flux of the six SPWs pixel by pixel. The synthesized beams of the averaged image and the spectral index map are shown in the bottom-left corner of each panel.

be the same as the beam size (25'') to get the EM and the electron density (n_e) if the distance information is available in the *WISE* catalog (Anderson et al. 2014). For 168 sources we got a good fit to the model and list the fitting results in Table 4. The main uncertainties of the fitting results are from the uncertainties of the peak flux brought in by the flux calibrator ($\sim 5\%$ at these wavelengths), uncertainties from the fitting procedure (< 10%), and the uncertainties of the sizes of the sources on the line of sight. Two sources have T_e measurements from previous RRLs observations (Balser et al. 2011), and we assume a T_e of 8000 K for the remaining sources (the mean T_e estimated by Balser et al. 2011 is ~ 8000 K for H II regions in the THOR survey area).

For 151 sources we fitted an 10^4 < EM < 10^5 cm⁻⁶ pc. For 124 sources there is distance information in the *WISE* catalog so we estimated n_e , which is between 0.9×10^2 and 5.4×10^2 cm⁻³. Continuum source G49.477–0.328, which is associated with the H II region G049.490–00.381 in W51, has the highest value for EM and n_e from our fitting, which are 4.2×10^5 cm⁻⁶ pc and 3.1×10^4 cm⁻³, respectively.

Among the 168 sources where we can get a good H $\mbox{\sc i}$ region model fit, 39 are also associated with CORNISH UCH $\mbox{\sc i}$ regions. Assuming homogeneous optically thin and the line of sight size is the same as the angular size, Kalcheva et al. (2018) calculated EM and n_e for the CORNISH UCH $\mbox{\sc i}$ regions. Since CORNISH is at 5 GHz and the angular sizes of the 39 UCH $\mbox{\sc i}$ regions they derived are all smaller than 12", which is about half beam size of THOR, we can not compare the EM and n_e from the the two samples directly. However, if we assume all 39 UCH $\mbox{\sc i}$ regions have a size of 25", and convert the EM and n_e from the CORNISH observations accordingly, they are close to with the values we derived with THOR observations although with large scatter. We can fit the EM_{CORNISH} vs EM_{THOR} and $n_{eCORNISH}$ vs n_{eTHOR} plots with a linear function y = x * 0.8 + c. This is a very rough comparison, but shows our H $\mbox{\sc i}$ fifting procedure is correct.

5.4. Supernova remnants

Green (2014) provides the most complete catalog of Galactic SNRs with 294 sources, 67 of them lie, or partially lie, in our survey area. By visually comparing the catalog with our continuum images, we identify 92 continuum sources associated with 36 SNRs. As shown in Fig. 5, many of the SNRs are resolved into multiple sources by the source-finding algorithm. The rest of the SNRs from Green (2014) either have too weak radio continuum emission, which is below our sensitivity, or are too diffuse and are filtered out by the VLA C-configuration (see also Bihr et al. 2016). We marked all the matched continuum sources with "SNR_green" in the catalog and list the corresponding ID of the SNRs from Green (2014).

By studying the compact and extended THOR+VGPS 1.4 GHz continuum emission (in the region $l > 17.5^{\circ}$), Anderson et al. (2017) confirmed the radio emission for 52 SNRs from Green (2014). Anderson et al. (2017) found that six of the SNRs from the Green (2014) catalog (G20.4+0.1, G21.5 0.1, G23.6+0.3, G54.1+0.3, G59.8+1.2 and G065.8 0.5) are confused with H II regions and the radio emission appears to be thermal. Figure 11 shows the spectral index fitting result and the spectral index map of SNR W49B as an example, indicating that the region is dominated by negative spectral index with some variation, and further indicating non-thermal synchrotron emission.

By comparing the large scale diffuse radio continuum emission traced by the combined THOR+VGPS 1.4 GHz continuum data with the mid-infrared *Spitzer* GLIMPSE 8.0 μm (Benjamin et al. 2003; Churchwell et al. 2009) and MIPSGAL 24 μm (Carey et al. 2009) data, Anderson et al. (2017) identified 76 new SNR candidates. Since the radio continuum emission from most of the new SNR candidates is weak and diffuse, we could only detect 13 of them in our continuum catalog that uses only the THOR c-configuration data and hence do not trace the large-scale diffuse emission. Among the new SNR candidates, G17.80-0.02, G26.75+0.73 and G27.06+0.04 are matched with two continuum sources each in our catalog, and G51.21+0.11 is matched with 6 continuum sources. We marked all the matched continuum sources with "SNR_anderson" in the catalog and list the corresponding ID of the SNRs from Anderson et al. (2017).

The MAGPIS survey (Helfand et al. 2006) identified many SNR candidates, and 33 of them are covered by our THOR survey. Anderson et al. (2017) found that 17 MAG-PIS SNR candidates (G18.2536-0.3083, G19.4611+0.1444, G19.5800-0.2400, G19.5917+0.0250, G19.6100-0.1200, G19.6600-0.2200, G21.6417+0.0000, G22.7583-0.49171, G22.9917-0.3583, G23.5667-0.0333, G24.1803+0.2167, G25.2222+0.2917, G29.0667-0.6750, G30.8486+0.1333, G31.0583+0.4833, G31.6097+0.3347, and G31.8208-0.1222.) are spatially coincident with a known H II region from the WISE catalog. G18.2536-0.3083 is also reported as a known HII region by Bihr et al. (2016). Furthermore, G16.3583-0.1833 and G17.3361-0.1389 are spatially coincident with known H_{II} regions G016.360-00.211 and G017.336-00.146 from the WISE catalog. G29.0778+0.4542 is a known planetary nebula (PNG029.0+00.4, see also Anderson et al. 2017). For 10 MAG-PIS SNRs (Helfand et al. 2006) we find matched continuum sources in our catalog, however, these continuum sources are also matched with SNRs from Green (2014) or candidates from Anderson et al. (2017), we do not mark them separately in the catalog. For 3 MAGPIS SNRs we do not find matched continuum sources due to THOR sensitivity and the missing flux problem (see also Bihr et al. 2016).

5.5. Planetary nebulae

The Hong Kong/AAO/Strasbourg H α planetary nebula database (HASH, Parker et al. 2016) provides up-to date information for all known Galactic planetary nebulae (PNe). In the HASH database, there are 234 PNe with the PN status marked as T (True PN), L (Likely PN), and P (Possible PN) within our survey area. We detected 164 of them in our catalog, among which 75 are True PN, 60 are Likely PN and 29 are Possible PN. We mark them as "PN" in our catalog, and list the HASH ID (PNG). In particular PNG029.2+00.0 is associated with the continuum source G29.211–0.069, the extended emission is tracing the known H II region G029.165-00.035. Thus we mark the continuum source G29.211-0.069 as "HII;PN" and list both the PN and H_{II} IDs. Furthermore, 90 of the matched PNe do not have radio emission information at 20 cm in the database; our survey provides this important information. We list all the PNe detected in the THOR survey in Table D.1. For the remaining 74 sources the peak flux at spw-1440 from THOR observations are in agreement with the radio flux at 20 cm from the database in general, with a median flux difference of 31% relative to the ones in the HASH database. Since we are comparing flux density in the HASH database with the THOR peak flux, the large difference is expected.

In addition to the 164 detected PNe we include in our catalog, PNG 014.5+00.0, 017.4+00.3 and 018.7+00.0 are also detected in our observations. However, the continuum sources that these PNe are associated with are known H II regions (G014.598+00.019, G017.414+00.377 and G018.710+00.000). Since these three PNe are all "possible PN", we did not include them in our catalog.

We did not detect the remaining 67 PNe, most likely due to sensitivity reasons. Only 4 of the 67 PNe have 1.4 GHz radio flux information in the HASH catalog. The 1.4 GHz flux density of PNG029.8+00.5 in the HASH catalog is 33 mJy, however, we did not detect any radio counterpart in our THOR-only nor THOR+VGPS data. Bojičić et al. (2011) and Luo et al. (2005) also reported the radio detection for this PN as suspect. The 1.4 GHz flux of PNG028.7+00.7, 044.9+00.0 and 056.1-00.4 in the HASH catalog is 2.7, 3.5 and 6.6 mJy, respectively.

Table 4. Fitting results for H II regions

Gal.ID	α	Δα	Н п пате	$T_e{}^a$	d^b	EM	$n_e{}^b$
				[K]	[kpc]	$[\times 10^5 \text{ cm}^{-6} pc]$	$[\times 10^{2} \text{cm}^{-3}]$
G52.753+0.334	0.13	0.01	G052.750+00.334	8970	9.575	2.0	5.0
G53.187+0.209	-0.14	0.02	G053.188+00.209	_	9.96	0.7	2.9
G56.160+0.077	-0.21	0.04	G056.153+00.076	_	9.9	0.2	1.7
G56.420+0.423	-0.12	0.06	G056.412+00.423	_	9.93	0.2	1.6
G57.547-0.272	-0.05	0.01	G057.541-00.279	_	8.805	1.3	4.2
G58.773+0.646	-0.10	0.04	G058.769+00.648	_	4.405	0.2	2.5
G60.883-0.130	0.01	0.02	G060.881-00.135	7463	_	1.4	_

Notes. (a) T_e is assumed to be 8000K if it is not available from the WISE catalog. (b) We calculated n_e if distance is available from the WISE catalog. The full table is available at the project website http://www2.mpia-hd.mpg.de/thor/DATA/www/.




Fig. 11. Spectral index map for SNR W49B. *Top-panel:* Spectral index fitting result of the peak of the continuum source G43.257–0.161. *Middle-panel:* Averaged image of spw-1440 and spw-1820, the white contours represent the area of the source extracted by BLOBCAT, and the cross marks the peak of the source G43.257–0.161 in the catalog. *Bottom-panel:* The spectral index map produced by fitting the flux of the six SPWs pixel by pixel. The synthesized beams of the averaged image and the spectral index map are shown in the bottom-left corner of each panel.

5.6. Pulsars

The ATNF Pulsar Catalogue¹¹ (Manchester et al. 2005) provides the most complete catalog of all published rotation-powered pulsars. Within our survey region, the catalog contains 335 pulsars. We cross matched the pulsar catalog with the THOR continuum catalog taking into account the position uncertainty from the pulsar catalog. 38 THOR continuum sources are matched with pulsars, we mark them as "PSR" in the continuum catalog, and list the corresponding pulsar names. Except for G38.163-0.151, all matches have a separation smaller than 7.5". Considering the synthesized beam size of our observation is ~ 12" – 18", these matches are reliable. G38.163–0.151 is matched with J1901+0435 with a separation of 15", however, J1901+0435 has a relatively large position uncertainty of 10", which we consider it a good match. We list the matched continuum sources and pulsars in Table E.2.

Among all the pulsars detected in our survey, J1841-0500 is an extremely intermittent pulsar (Camilo et al. 2012), and has no 1.4 GHz radio flux information in the ATNF pulsar catalog. We detect a radio counterpart in our observation with a S/N \sim 6 (G27.322–0.033).

We did not detect the remaining 297 pulsars in our catalog, which is mainly due to low sensitivity in some sections of the maps. Among the non-detected pulsars, 247 have mean 1.4 GHz radio flux densities in the ATNF pulsar catalog, and their mean flux densities are all below 3 mJy, with a median value of 0.3 mJy. In comparison, the detected ones have a median mean flux density of 1.7 mJy from the ATNF catalog.

In general, the peak flux at spw-1440 from THOR observations are in agreement with the radio flux at 1.4 GHz in the ATNF catalog, with a median flux difference of 30% relative to the ones in the ATNF catalog.

5.7. X-ray sources

While diffuse radio and X-ray emission can trace H II regions (free-free emission and stellar wind shocks, Silich 2005) and SNRs (e.g., Decourchelle 2005), the compact emission is usually tracing Pulsar Wind Nebulae (e.g., Brisken et al. 2005; Miller et al. 2005), X-ray binaries (XRB or microquasars; Mirabel & Rodríguez 1998, 1999), or active galactic nuclei (AGN, e.g., Bridle & Perley 1984; Harris & Krawczynski 2006).

We cross-matched our continuum catalog with three X-ray source catalogs, 1SXPS SWIFT X-ray telescope point source catalog (Evans et al. 2014), XMM-Newton Serendipitous Source Catalog (3XMM-DR7 Version, Watson et al. 2009; Rosen et al.

¹¹ http://www.atnf.csiro.au/research/pulsar/psrcat

2016), and Chandra Source Catalog (CSC, v1.1, Evans et al. 2010). Within our survey area, 1SXPS has ~ 2800 sources, CSC has ~ 3900 sources, and 3XMM has ~ 11500 sources. With a matching radius of 10", we found that 79 of the remaining ~9300 THOR sources are associated with X-ray sources (see Table F.3). Among the 79 X-ray sources we detected, 43 do not have a radio counterpart within a radius of 15" on the SIMBAD Astronomical Database¹² or the NASA/IPAC Extragalactic Database (NED)¹³. 65 of the X-ray sources are unresolved, the remaining resolved sources are also relatively compact with an effective radius < 29". In particular, continuum source G45.366-0.219 is associated with the microquasar GRS1915+105 (Mirabel & Rodríguez 1999). Further H absorption studies towards these sources would allow us to identify which ones are galactic. By investigating the X-ray and radio flux ratios, the spectral indices and observations at other (optical/infrared) wavelengths of the Galactic sources in detail, we can further constrain the type of the sources, i.e., low mass XRB, PN, pulsar etc. (e.g., Seaquist 1993; Maccarone et al. 2012; Tetarenko et al. 2016).

5.8. Extragalactic radio sources

About 9300 sources in our catalog are not classified. Taking a matching radius of 15", which is the average size of the synthesized beam of the averaged images used for source extraction, more than 7750 sources do not match to anything in SIM-BAD. For the matched ones, about 84% of them are matched to only one or more radio sources. With the same matching radius (15"), \sim 3350 of the sources are matched to one or more infrared sources, and \sim 2140 of the sources are matched to one or more radio sources, and more than 3760 sources do not have any counterparts within a radius of 15". This means that for \sim 9000 sources detected in our survey, besides they have radio and/or infrared emission, we do not know exactly what they are.

As we mentioned in Sect. 5.2, besides the H II regions, SNRs, and PNe, the majority of the sources we detected are actually extragalactic. Many of the those sources are also resolved and show bipolar radio lobe structures. We checked all the resolved sources with a negative or flat spectral index in our catalog by eye, and identified ~300 sources that show clear bipolar jet structure, and mark them as "jet" in the catalog. For these "jet" sources, we can construct the spectral index maps, which could be used to estimate the source expansion velocity when combined with magnetic field strength information. Since the identification of these "jet" sources is done morphologically, two partially overlapped sources could also be categorised into "jet" sources. Since we do not have zero-spacing information, we filtered out large scale structures in our observations. Bihr et al. (2016) show that the THOR observations are able to recover sources with size up to ~120" reasonably well (≥80% flux recovery), spectral index maps for sources larger than that should be interpreted with caution. Figure 12 shows the spectral index maps of three "jet" sources. As the figure shows, the spectral index varies within the source, indicating that the optical depth of the synchrotron emission varies or possible thermal emission exists in different parts of the radio lobes.

In Sect. 5.2 we mentioned that the spatial distribution and the spectral index distribution of the unresolved sources indicate the majority of them are extragalactic sources. To further confirm this, we compare the normalized spatial distribution and spectral index distribution of the non-classified sources to the Galactic sources we identified. Figure 13 compares the normalized distribution along Galactic longitude and latitude of those non-classified sources with the Galactic objects (H II regions, PNe and pulsars). The normalized distribution along Galactic longitude (top-panel, Fig 13) shows that the Galactic objects are more concentrated in the lower longitude region, while more non-classified sources are detected in the higher longitude region. The distribution along Galactic latitude (top-panel, Fig 13) shows that the HII regions are concentrated very close to the Galactic mid-plane. Although not as much as the HII regions, the PNe and pulsars are also concentrated close to the mid-plane. In contrast, more non-classified sources are found in regions $|b| > 0.5^{\circ}$. This distribution difference confirms that the nonclassified sources are of mostly extragalactic origin. Furthermore, the spectral index distribution of the non-classified sources also shows clear differences from the Galactic sources and peaks around $\alpha \sim -1$ (Fig. 14), but there are still many of these nonclassified sources showing a flat SED ($\alpha \gtrsim 0$), which could be Galactic compact H II regions that are not listed in the WISE H II catalog. We classify the ~ 5300 sources with a negative spectral index $\alpha < -0.3$ as extragalactic source candidates, among which ~ 3970 sources have a reliable spectral index (fit_spws≥ 4).

Extragalactic radio sources often consist of a core and one or more radio lobes, which show different spectral profiles. The core dominates the emission at shorter wavelengths and shows a close to flat or positive spectral index, while the jet dominates at longer wavelengths and has shows negative spectral index (e.g., Hey 1971, chap. 9). This transition in the spectral profile happens around ~10 to 20 cm. Assuming the emission of the core and jet at 1 to 2 GHz can both be described with one single power law, then the total flux I(v) is proportional to $(c_1v^{\alpha_1}+c_2v^{\alpha_2})$, where α_1 and α_2 are the spectral indices of the jet and core, respectively, c_1 and c_2 are constants. We use the Markov chain Monte Carlo method (MCMC) to fit the extragalactic source candidates which have a S/N ratio larger than 50 with the two spectral indices using the *emcee* package (Foreman-Mackey et al. 2013). Then we select the ones that meet the following criteria as good fit: first, the reduced χ^2 of the two-spectral-indices fitting is smaller than the one with single spectral index fitting; second, the spectra of the two components cross with each other within our frequency coverage (1.05 to 1.95 GHz, such as Fig. 15). In total, we can derive a good fit with two spectral indices for 135 continuum sources.

We performed an F-test (e.g., PHILLIPS 1982; Lomax & Hahs-Vaughn 2013) to compare the two components fitting to the one power low fitting, and calculated the probability values (p-value) for all 135 sources. The p-values are all above 0.28, so the two components fitting is not significantly better than the simple one power law fitting. However, the two components fitting results still provide alternative insight for those sources. As shown in Fig. 15, the peak fluxes of G50.695–0.792 can be fitted with two components with a spectral indices of -1.47 and 0.03, respectively, which could be tracing the radio jet and core respectively. The averaged image in Fig. 15 shows G50.695–0.792 is resolved with an elongated structure, which might be tracing the jet.

Radio sources with Ultra Steep Spectra (USS, $\alpha \leq -1.3$) are efficient tracers of high redshift radio galaxies, since radio sources can be detected uniformly over all redshift ranges, and do not suffer dust extinction at high redshifts (Chambers et al. 1996; Hughes et al. 1997; Ivison et al. 1998; De Breuck et al. 2000, e.g.,). De Breuck et al. (2000) found a trend that steeper spectral index sources have higher redshifts, and 50% of the 4C

¹² http://simbad.u-strasbg.fr/simbad/

¹³ http://ned.ipac.caltech.edu/




Fig. 12. Spectral index maps for three "jet" sources. *Top-panels:* Spectral indices fitting results of the peak positions. *Middle-panels:* Averaged images of spw-1440 and spw-1820, the white contours represent the area of the sources extracted by BLOBCAT, and the crosses mark the peaks of the sources in the catalog. *Bottom-panels:* The spectral index map produced by fitting the flux of the six SPWs pixel by pixel. The synthesized beams of the averaged image and the spectral index map are shown in the bottom-left corner of each panel.

USS sample (Chambers et al. 1996) are z > 2 sources. This $z - \alpha$ correlation is considered to be a combined contribution of a K-correction and an increasing spectral curvature with z of a radio spectrum (Krolik & Chen 1991; Carilli et al. 1999; van Breugel et al. 1999).

In our continuum catalog, we have ~2170 sources that have an spectral index steeper than −1.3, which is our base sample of USS sources. De Breuck et al. (2000) further point out that 85% USS with X-ray detections are in galaxy clusters, so we chose the ones without X-ray emission. Furthermore, we select sources with a reliable spectral index determination (fit_spws≥ 4). This selection results 1362 sources that have a good spectral index determination and have a spectral index steeper than −1.3. 663 of these sources have a spectral index steeper than −1.6 and could be Galactic pulsars (De Breuck et al. 2000). The remaining 699 USS sources could be high-redshift radio galaxies; further spectroscopic observations are needed to confirm this.

6. Conclusions

We observed a large portion of the first Galactic quadrant ($l=14.0-67.4^{\circ}$, $|b| \leq 1.25^{\circ}$) using the VLA in C-configuration and achieved a spatial resolution of $\sim 10-25''$ at 1 to 2 GHz with the THOR Galactic plane survey. In this paper, we present the catalog of the continuum sources from the whole survey. We summarise numbers of different types of sources in the catalog in Table 5, and the main results below.

1. The catalog contains 10387 sources we extracted with the BLOBCAT software after removing the obvious observational sidelobe artifacts. About 72% (7521 sources) of the


Table 5. Summary of different types of sources in the catalog

Source Types	Number of continuum sources
H II regions	713
SNRs	92(+21)
PNe	164
Pulsars	38 (+663)
X-ray sources	79
extragalactic jets	299
USS	699

Notes. USS stands for sources with Ultra Steep Spectra ($\alpha \le -1.3$). The numbers of the candidates are in brackets.

extracted sources are detected at a significance higher than 7σ , and $\sim 79\%$ are unresolved. The catalog is complete to at least 94% above the 7σ detection limit. The noise of our data is dominated by the sidelobe noise and spatially varying, although more than 60% of the observed area has a noise level of $7\sigma \lesssim 2$ mJy beam⁻¹. We extracted the peak intensity of the six usable SPWs between 1 to 2 GHz, and we were able to determine a reliable spectral index (spectral index fitted with at least 4 SPWs) for 5657 sources.

2. We cross-matched the THOR catalog with the WISE H $\scriptstyle\rm II$ region catalog and found 713 continuum sources are associated with H $\scriptstyle\rm II$ regions. Among the matched H $\scriptstyle\rm II$ regions, 16 are in the radio quiet group in the WISE catalog which means they did not previously have radio continuum detected. 231 continuum sources are associated with more than one H $\scriptstyle\rm II$ region. The spectral index distribution shows a single peak around $\alpha=0$, indicating thermal free-free emission. For 168






Fig. 13. Normalized distribution of different types of objects along Galactic Longitude (*top-panel*) and Latitude (*bottom-panel*) for the non-classified sources, H $\scriptstyle\rm II$ regions, PNe, and pulsars. For H $\scriptstyle\rm II$ regions we plot only the ones associated with the THOR continuum sources. For PNe and pulsars, we plot all that lie within our survey area.

sources we can fit the SED with a simple homogeneous H $\scriptstyle\rm II$ region model and derive the emission measure (EM) and the electron density (n_e) where the distance information is available in the WISE catalog.

- 3. Although the diffuse emission from many of the large scale SNRs is filtered out by our interferometric observations, we identify 92 continuum sources associated with 39 SNRs from the SNR catalog by Green (2014). 13 of the new SNR candidates from Anderson et al. (2017) are detected in our continuum catalog.
- 4. By cross-matching the THOR catalog with the HASH database, we detect 164 PNe in our continuum catalog. As 90 of them do not have radio emission information at 20 cm in the database, our survey provides this important information in Table D.1. The spectral index distribution is similar to the one of the H $\scriptstyle\rm II$ regions and shows a single peak around $\alpha=0$, indicating thermal free-free emission.
- 5. We cross-matched the THOR catalog with the ATNF Pulsar Catalog and found 38 counterparts. One extremely intermittent pulsar J1841-0500 is also detected in our catalog. 663 sources with a spectral index $\alpha < -1.6$ could be Galactic pulsar candidates.
- 6. We cross-matched the THOR catalog with X-ray source catalogs 1SXPS, 3XMM- DR7 and CSC, and found 79 overlaps. 12 of the them have a spectral index steeper than −1.3




Fig. 14. Normalized spectral index distribution of the non-classified sources, H $\scriptstyle\rm II$ regions, and PNe.

and could be galaxy clusters. 43 of them do not have previous known radio counterparts within a radius of 15" on SIMBAD or NED.

- 7. About 300 sources show clear structure of bipolar jets, we mark them as "jet" in the catalog and construct spectral index maps. We identified 699 Ultra Steep Spectra (USS) sources and they could be high redshifted radio galaxies $(-1.3 > \alpha > -1.6)$. Further spectroscopic observations are needed to confirm this.
- 8. About 9000 sources in our catalog are not classified specifically. They are likely to be extragalactic background sources. More than 7750 sources do not have counterparts in the SIMBAD Astronomical Database, and more than 3760 sources do not have counterparts in the NED.

With the THOR continuum catalog, we provide a rich dataset to the community. All the fits images and catalogs can be downloaded from the project website

footnotehttp://www.mpia.de/thor/Overview.html. With the follow up observations of particular sources, such as absorption study of the extragalactic sources, or combining with other existing Galactic plane surveys, we can study the the structure of the Milky Way, and the ISM in different phases.

Acknowledgements. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. Y.W., H.B., S.B., and J.D.S. acknowledges support from the European Research Council under the Horizon 2020 Framework Program via the ERC Consolidator Grant CSF-648505, and RSK via the ERC AdvancedGrant 339177 (STARLIGHT). H.B. and M.R. acknowledge support from the Deutsche Forschungsgemeinschaft in the Collaborative Research Center (SFB 881) "The Milky Way 74 System" (subproject B1, B2, B8), and from the Priority Program SPP 1573 "Physics of the Interstellar Medium" (grant numbers KL 1358/18.1, KL 1358/19.2). F.B. acknowledges funding from the European Union's Horizon 2020 research and innovation programme (grant agreement No 726384). The authors thank J. Mottram and M. Fouesneau for the productive and fruitful discussions. This research made use of Astropy and affiliated packages, a community-developed core Python package for Astronomy (The Astropy Collaboration et al. 2018), Python package SciPy¹⁴, APLpy, an open-source plotting package for Python (Robitaille & Bressert 2012), and software TOPCAT (Taylor 2005).

References

Aguirre, J. E., Ginsburg, A. G., Dunham, M. K., et al. 2011, ApJS, 192, 4

¹⁴ https://www.scipy.org/




Fig. 15. One example source G50.695–0.792 fitted with two indices. The top two panels show each SPW separately. The bottom left large image represents the averaged image of spw-1440 and spw-1820, which we used for the source extraction (see Sect. 3). The white contours show the extent of the source determined by the BLOBCAT algorithm. The cross in each panel marks the peak position, which we used to determine the spectral index. The bottom right panel presents the peak intensity for each SPW and corresponding one single spectral index fitting (black solid line) and two spectral indices fitting (green dashed lines) from two components (red and blue dashed lines).

Anderson, L. D., Bania, T. M., Balser, D. S., et al. 2014, ApJS, 212, 1 Anderson, L. D., Wang, Y., Bihr, S., et al. 2017, A&A, 605, A58 Balser, D. S., Rood, R. T., Bania, T. M., & Anderson, L. D. 2011, ApJ, 738, 27 Benjamin, R. A., Churchwell, E., Babler, B. L., et al. 2003, PASP, 115, 953 Beuther, H., Bihr, S., Rugel, M., et al. 2016, A&A, 595, A32 Bhatnagar, S., Rau, U., Green, D. A., & Rupen, M. P. 2011, ApJ, 739, L20 Bihr, S., Beuther, H., Ott, J., et al. 2015, A&A, 580 Bihr, S., Johnston, K. G., Beuther, H., et al. 2016, A&A, 588, A97 Bojičić, I. S., Parker, Q. A., Filipović, M. D., & Frew, D. J. 2011, MNRAS, 412, 223 Bridle, A. H. & Perley, R. A. 1984, ARA&A, 22, 319 Brisken, W. F., Romani, R. W., & Ng, C. Y. 2005, in X-Ray and Radio Connections, ed. L. O. Sjouwerman & K. K. Dyer, #5.04 Camilo, F., Ransom, S. M., Chatterjee, S., Johnston, S., & Demorest, P. 2012, ApJ, 746, 63 Carey, S. J., Noriega-Crespo, A., Mizuno, D. R., et al. 2009, PASP, 121, 76 Carilli, C. L., Röttgering, H. J. A., Miley, G. K., Pentericci, L. H., & Harris, D. E. 1999, in The Most Distant Radio Galaxies, ed. H. J. A. Röttgering, P. N. Best, & M. D. Lehnert Chambers, K. C., Miley, G. K., van Breugel, W. J. M., & Huang, J.-S. 1996, ApJS, 106, 215 Churchwell, E., Babler, B. L., Meade, M. R., et al. 2009, PASP, 121, 213 Condon, J. J. 1992, ARA&A, 30, 575 Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, AJ, 115, 1693

Csengeri, T., Urquhart, J. S., Schuller, F., et al. 2014, A&A, 565, A75 De Breuck, C., van Breugel, W., Röttgering, H. J. A., & Miley, G. 2000, A&AS, 143, 303 Decourchelle, A. 2005, in X-Ray and Radio Connections, ed. L. O. Sjouwerman & K. K. Dyer, #4.02 Evans, I. N., Primini, F. A., Glotfelty, K. J., et al. 2010, ApJS, 189, 37 Evans, P. A., Osborne, J. P., Beardmore, A. P., et al. 2014, ApJS, 210, 8 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, Green, D. A. 2014, Bulletin of the Astronomical Society of India, 42, 47 Grobler, T. L., Nunhokee, C. D., Smirnov, O. M., van Zyl, A. J., & de Bruyn, A. G. 2014, MNRAS, 439, 4030 Hales, C. A., Murphy, T., Curran, J. R., et al. 2012, MNRAS, 425, 979 Harris, D. E. & Krawczynski, H. 2006, ARA&A, 44, 463 Helfand, D. J., Becker, R. H., White, R. L., Fallon, A., & Tuttle, S. 2006, AJ, 131, 2525 Hey, J. S. 1971, The radio universe Hoare, M. G., Purcell, C. R., Churchwell, E. B., et al. 2012, PASP, 124, 939 Hughes, D. H., Dunlop, J. S., & Rawlings, S. 1997, MNRAS, 289, 766 Ivison, R. J., Dunlop, J. S., Hughes, D. H., et al. 1998, ApJ, 494, 211 Kalcheva, I. E., Hoare, M. G., Urquhart, J. S., et al. 2018, ArXiv e-prints [arXiv:1803.09334]

Keto, E. 2003, ApJ, 599, 1196

Krolik, J. H. & Chen, W. 1991, AJ, 102, 1659

- Lomax, R. & Hahs-Vaughn, D. 2013, Statistical Concepts: A Second Course (Taylor & Francis)
- Lucas, P. W., Hoare, M. G., Longmore, A., et al. 2008, MNRAS, 391, 136
- Luo, S. G., Condon, J. J., & Yin, Q. F. 2005, ApJS, 159, 282
- Maccarone, T. J., Torres, M. A. P., Britt, C. T., et al. 2012, MNRAS, 426, 3057
- Manchester, R. N., Hobbs, G. B., Teoh, A., & Hobbs, M. 2005, AJ, 129, 1993
- Miller, N. A., Neff, S. G., & Mushotzky, R. F. 2005, in X-Ray and Radio Connections, ed. L. O. Sjouwerman & K. K. Dyer, #5.05
- Mirabel, I. F. & Rodríguez, L. F. 1998, Nature, 392, 673
- Mirabel, I. F. & Rodríguez, L. F. 1999, ARA&A, 37, 409
- Parker, Q. A., Bojičić, I. S., & Frew, D. J. 2016, in Journal of Physics Conference Series, Vol. 728, Journal of Physics Conference Series, 032008
- PHILLIPS, P. C. B. 1982, Biometrika, 69, 261
- Purcell, C. R., Hoare, M. G., Cotton, W. D., et al. 2013, ApJS, 205, 1
- Reynoso, E. M. & Walsh, A. J. 2015, MNRAS, 451, 3044
- Robitaille, T. & Bressert, E. 2012, APLpy: Astronomical Plotting Library in Python, Astrophysics Source Code Library
- Rosen, S. R., Webb, N. A., Watson, M. G., et al. 2016, A&A, 590, A1
- Rosolowsky, E., Dunham, M. K., Ginsburg, A., et al. 2010, ApJS, 188, 123
- Rugel, M. R., Beuther, H., Bihr, S., et al. 2018, ArXiv e-prints [arXiv:1803.04794]
- Rybicki, G. B. & Lightman, A. P. 1979, Radiative processes in astrophysics Sánchez-Monge, Á. 2011, Ph.D. Thesis
- Schuller, F., Menten, K. M., Contreras, Y., et al. 2009, A&A, 504, 415
- Seaquist, E. R. 1993, Reports on Progress in Physics, 56, 1145
- Silich, S. 2005, in X-Ray and Radio Čonnections, ed. L. O. Sjouwerman & K. K. Dyer, #3.03
- Stil, J. M., Taylor, A. R., Dickey, J. M., et al. 2006, AJ, 132, 1158
- Taylor, M. B. 2005, in Astronomical Society of the Pacific Conference Series, Vol. 347, Astronomical Data Analysis Software and Systems XIV, ed. P. Shopbell, M. Britton, & R. Ebert, 29
- Tetarenko, B. E., Bahramian, A., Arnason, R. M., et al. 2016, ApJ, 825, 10
- The Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, ArXiv e-prints [arXiv:1801.02634]
- Urquhart, J. S., Thompson, M. A., Moore, T. J. T., et al. 2013, MNRAS, 435, 400
- van Breugel, W., De Breuck, C., Stanford, S. A., et al. 1999, in Astronomical Society of the Pacific Conference Series, Vol. 193, The Hy-Redshift Universe: Galaxy Formation and Evolution at High Redshift, ed. A. J. Bunker & W. J. M. van Breugel, 44
- Watson, M. G., Schröder, A. C., Fyfe, D., et al. 2009, A&A, 493, 339
- Wilson, T. L., Rohlfs, K., & Hüttemeister, S. 2013, Tools of Radio Astronomy

Appendix A: Noise maps

Appendix B: Completeness maps

Appendix C: Compare with CORNISH

Sources with a positive spectral index between 1 to 2 GHz are mostly dominated by optically thick free-free emission at these frequencies, however, the emission should be optically thin at higher frequencies and flattening the spectral index there (e.g., Sánchez-Monge 2011). If we estimate the flux density at 5 GHz for these sources, we could easily overestimate the flux. Since we do not have flux density measurements between 2 GHz and 5 GHz, we are not sure whether such overestimation would be due to the uncertainty of spectral index or optically depth variation

Sources with a spectral index $\lesssim -0.5$ are mostly dominated by synchrotron emission. Depending on the nature of the source, the spectral index could get flatter between 2 GHz and 5 GHz as a possible contribution from free-free emission could increase (e.g., Condon 1992), since free-free emission has a flat or positive spectral index. If we estimate the flux density at 5 GHz for these sources, we would underestimate the flux.

Sources with a flat spectral index (\sim -0.1) in the THOR frequency range are dominated by optically thin free-free emission. The emission at 5 GHz would still be dominated by optically thin free-free emission, since dust continuum will not be relevant until \sim 100 GHz and synchrotron emission contribution is even smaller than at the lower frequencies. Thus, we can estimate the flux density at 5GHz from the THOR flux density measurements properly and reliably.

Among the 1905 THOR sources that are matched with a CORNISH source within a radius of 5", 1087 sources are unresolved and are detected in all six spectral windows in THOR (fit_spws=6). We extrapolated flux densities at 5 GHz for all these 1087 sources according to their spectral indices from the THOR catalog and plotted against the flux densities in the CORNISH catalog, see Fig. C.1. Figure C.1 shows that comparing to the flux densities in the CORNISH catalog, as expected we overestimate the flux density for many sources with a positive spectral index, and similarly we underestimate the flux density for many sources with a negative spectral index. This is expected for such a general extrapolation for the reasons explained above.

Appendix D: PNe in our continuum catalog

Appendix E: Detected pulsars in our continuum catalog

Appendix F: X-ray sources detected in our continuum catalog




Fig. A.1. Noise map of the THOR survey using the average of spw-1440 and spw-1820




Fig. B.1. Completeness map in percentage for sources with a peak intensity of 1 mJy beam⁻¹.




Fig. B.2. Completeness map in percentage for sources with a peak intensity of 2 mJy beam⁻¹.





Fig. B.3. Completeness map in percentage for sources with a peak intensity of 5 mJy beam⁻¹.



 $\textbf{Fig. B.4.} \ Completeness \ map \ in \ percentage \ for \ sources \ with \ a \ peak \ intensity \ of \ 10 \ mJy \ beam^{-1}.$




Fig. C.1. Extrapolated flux densities at 5 GHz for all 1087 compact sources that are detected in all six spectral windows in THOR (fit_spws=6) according to their spectral indices from the THOR catalog plotted against the flux densities in the CORNISH catalog. The green ones are sources with a spectral index smaller than 0, the blue ones are the ones with a spectral index larger than 0, the red ones are sources with a spectral index between -0.09 and -0.11 (optically thin free-free emission). The dashed line represents a one-to-one relation.

Table D.1. Detected planetary nebulae in THOR continuum catalog.

	[J2000]	Dec. [J2000]	${ m S_p} \ [m mJy\ beam^{-1}]$	\mathbf{Z}	ά	$\nabla \alpha$	fit_spws	PNG	Name
G14.585+0.462	18:15:21.10	-16:02:56.3	22.84	31	0.12	0.19	9	014.5+00.4	MPA J1815–1602
G14.658+1.012	18:13:29.12	-15:43:20.1	2.34	7	-0.84	0.51	4	014.6 + 01.0	PHR J1813-1543
G14.750-0.250	18:18:17.00	-16:14:31.2	23.97	28	-0.37	0.32	9	$014.7 - 00.2^*$	
G14.779-0.333	18:18:38.70	-16:15:20.9	15.27	15	2.25	0.72	4	$014.7 - 00.3^*$	
G14.896+0.484	18:15:53.16	-15:45:54.9	8.01	19	-0.26	0.71	4	$014.8+00.4^*$	
G15.200-0.086	18:18:34.22	-15:46:06.0	11.17	6	-0.49	0.64	7	$015.1-00.0^*$	
G15.518+1.034	18:15:06.47	-14:57:20.7	9.65	25	-0.46	0.20	9	015.5 + 01.0	PHR J1815-1457
G15.541+0.336	18:17:41.91	-15:16:04.0	13.95	30	0.55	0.29	9	015.5+00.3*	
G15.559+0.981	18:15:22.87	-14:56:43.9	4.04	14	-0.91	0.35	S	015.5+00.9*	Pa 99
G15.585+0.400	18:17:33.10	-15:11:56.5	11.13	25	0.81	0.38	5	015.5+00.3*a	
G15.800-0.006	18:19:27.38	-15:12:06.8	30.99	63	0.49	0.13	9	$015.7 - 00.0^*$	MSX6C G015.7987-00.0060
G16.055+0.828	18:16:54.90	-14:34:54.6	9.92	24	-0.05	0.26	9	016.0 + 00.8	MSX6C G016.0545+00.8279
G16.228-0.368	18:21:36.82	-14:59:40.8	7.91	15	-0.33	0.34	9	016.2-00.3	GPSR5 16.228-0.369
G16.428+1.007	18:16:59.72	-14:10:07.4	17.03	30	-0.58	0.21	9	$016.4+01.0^*$	
G16.499+0.115	18:20:22.76	-14:31:39.4	14.00	29	-0.78	0.17	9	$016.4+00.1^*$	
G17.015-0.190	18:22:29.76	-14:12:59.2	11.00	25	0.21	0.27	9	$017.0-00.1^*$	
G17.221+0.129	18:21:43.92	-13:53:03.9	1.73	9	-0.96	0.48	4	017.2 + 00.1*	PHR J1821-1353
G17.367+0.523	18:20:35.06	-13:34:15.0	14.84	49	0.24	0.15	9	017.3 + 00.5	HRDS G017.364+0.519
G17.449+0.115	18:22:13.47	-13:41:24.1	7.37	23	-0.38	0.28	9	$017.4+00.1^*$	
G17.588+1.068	18:19:02.18	-13:07:03.7	7.34	25	-0.23	0.26	5	017.5+01.0	MPA J1819–1307
G17.616–1.169	18:27:13.36	-14:08:34.0	75.50	152	0.03	0.03	9	017.6 - 01.1	VSP 2–18
G17.725-0.243	18:24:03.39	-13:36:50.6	50.26	141	0.11	0.04	9	$017.7 - 00.2^*$	
G17.822+0.987	18:19:47.13	-12:57:00.8	12.55	37	0.11	0.19	9	017.8+00.9*	
G17.865+0.212	18:22:40.55	-13:16:39.1	7.87	20	0.82	0.44	S	017.8+00.2*	
G18.066+0.854	18:20:44.31	-12:47:52.7	26.78	26	0.11	90.0	9	018.0 + 00.8	IRAS 18179–1249
G18.241-0.555	18:26:11.05	-13:18:12.8	3.62	6	1.85	4.16	7	$018.2 - 00.5^*$	
.241–0.915	18:27:29.77	-13:28:18.2	39.56	134	-0.14	0.03	9	018.2-00.9	MPA J1827–1328
G18.578-0.748	18:27:31.93	-13:05:45.6	5.52	20	0.01	0.42	9	018.5-00.7*	
G19.003+0.128	18:25:10.02	-12:18:38.5	5.73	12	-0.60	0.57	9	$019.0+00.1^*$	
G19.119+0.818	18:22:53.58	-11:53:09.3	10.91	46	-0.41	0.15	9	$019.1 + 00.8^*$	MPA J1822-1153
G19.468-0.015	18:26:34.37	-11:58:00.4	15.95	56	-0.10	0.24	9 1	$019.4-00.0^*$	
G19.533+0.731	18:23:59.96	-11:33:38.9	3.56	16	-0.65	0.47	S	$019.5+00.7^*$	GLIPN1823-1133
G19.610+1.187	18:22:30.08	-11:16:44.2	22.79	29	1.27	0.10	9	019.6 + 01.1	MSX6C G019.6095+01.1873
G19.649+0.774	18:24:03.96	-11:26:16.5	3.98	16	-0.22	0.29	9	019.6+00.7*	MPA J1824–1126
G19.930-0.664	18:29:47.85	-11:51:33.1	7.33	15	0.82	0.36	9	$019.9-00.6^*$	
G19.945+0.913	18:24:07.92	-11:06:41.7	26.63	109	0.24	0.05	9	019.9 + 00.9	M3-53
G20.432+0.357	18:27:03.55	-10.56:25.5	86.6	37	0.16	0.20	9	020.4+00.3*	MSX6C G020.4320+00.3571
G20.468+0.679	18:25:58.08	-10:45:29.9	63.93	232	0.69	0.03	9	020.4+00.6	PM 1–231
G20.517+0.478	18:26:47.27	-10:48:30.4	17.63	99	0.05	0.0	9	$020.5+00.4^*$	
G20.978+0.925	18:26:03.07	-10:11:31.2	13.49	46	-0.30	0.14	9	020.9 + 00.9	IRAS 18232-1013
G20.983+0.851	18:26:19.48	-10:13:21.8	2.70	10	-0.60	0.42	9	020.9 + 00.8	PHR1826-1013
G20.999–1.125	18:33:28.96	-11:07:25.5	215.77	372	0.40	0.01	9	020.9–01.1	M 1–51
GZ1.162+0.476	18:28:01.20	-10:14:09.8	50.44	717	77.1	0.03	0	021.1+00.4	GLMP /81

Table D.1. continued.

[J2000] [8:26:26.62	[J2000] -09:53:14.6	S _p [mJy beam ⁻¹]	$\frac{S}{Z}$	α 0.21	$\Delta \alpha$	nt_spws	PNG ³	Name Name DHR 11876_0953
		7.14 4.89	8 10	1.91	1.40	n 0	$021.2+00.9$ $021.3-00.8^*$	FHR J1820-0953 IRAS 18303-1043
_	-10:12:13.2	1.49	S	I	1	1	$021.4+00.0^*$	2MASS J18295609-1012119
	-09:38:14.4	21.79	71	1.43	0.10	9	021.6+00.8	PM 1–235
8:33:22.44 -	-10:20:13.4	28.96	53	-0.08	0.11	9	021.6-00.7*	M 2 55
	-10.15.18.7 -10.05.50.0	31.58	38 0	-0.45	0.14	9 ا	021.7 - 00.9 $021.8 - 00.4$	M 3-28
	-10:07:37.5	12.35	29	0.11	0.24	9	021.9-00.8*	
8:28:28	-09:06:13.1	18.02	77	0.15	0.08	9	022.2 + 00.9	IRAS 18257–0908
		27.58	49	0.29	0.13	9	022.5 - 00.1*	
		24.75	85	-0.01	0.08	9	022.5+01.0	MaC 1–13
	-08:59:49.4	8.63	25	0.14	0.27	9	022.6+00.2*	
	-08:05:42.2	96.6	56	0.55	0.28	S	$023.4+00.7^*$	PHR J1831-0805
•		23.33	51	0.31	0.18	S	$023.5-00.5^*$	
		55.10	171	0.89	0.04	ις, Έ	023.8–00.7	GLMP 805
		30.04	45	0.21	0.14	S	023.9+01.2	MA 13
		5.54	13	-0.26	0.49	ω	024.1+01.1	PHR J1831–0715
•		9.81	13	0.0	0.51	9	024.3+00.2*	MSX6C G024.3854+00.2867
		14.20	35	-0.18	0.15	9 1	024.4+00.9	MPA J1832–0706
· 		19.74	24	1.19	$0.33_{0.22}$	S,	024.6+00.5	MPA J1834-0706
		10.09	22	0.03	0.27	9 (024.7-01.0*	
•		19.00	94 9	0.86	0.15	9 4	025.0-00.6*	
8:34:10:01 8:42:07 00	06:40:49.0	59.62	93	-0.10	0.09	o v	023.8+01.1	D. 1 14
	-06.46.34.3	8.03 74.68	2 8	0.00	0.13) v	026 1+00 5*	+1-1-2-1
·	-05:28:14.0	13.01	21	0.52	0.41	S	$026.7+00.1^*$	
		5.01	17	-0.27	0.42	5	026.8 - 01.0	MPA J1843-0556
8:40:49.21	-05:29:45.9	9.05	29	-0.20	0.32	2	026.8 - 00.1*	MPA J1840–0529
•		92.9	11	-1.37	0.64	5	$027.4 - 00.2^*$	
	• •	2.25	S	I	I	ı	$027.5 + 01.0^{*}$	PHR J1838-0417
	-04:51:58.5	3.46	12	-1.40	3.84	7	$027.6 - 00.3^*$	MSX6C G027.6594-00.3838
·		6.47	28	0.13	0.22	9	027.6-00.8	PHR J1844-0503
		119.82	205	0.29	0.03	9	027.7+00.7	M 2–45
		5.00	6	1	I	1	$027.8+00.5^*$	MPA J1840-0415
·		22.00	33	-0.15	0.14	9	028.9 + 00.2*	PHR J1843-0325
'		5.46	19	-0.35	0.35	9	029.0 + 00.9*	Pa 112
		21.98	21	-0.19	0.08	9	029.0+00.4	Abell 48
-		117.99	156	0.52	0.04	9	029.2+00.0	TDC 1
	-03:05:41.3	76.91	149	0.92	0.05	9	029.8–00.8	IRAS 18461–0309
-		49.22	55	0.89	0.13	9	$030.6 - 00.3^*$	IRAS 18458-0213
		20.52	33	0.33	0.16	9	031.3-00.7*	GPSR 031.374-0.752
		6.82	13	0.05	0.25	9 \	031.9–00.3*	WeSb 4
8:49:45.11	-00:53:08.6	16.10	42	0.54	0.19	9	(14, 7, 4, 10)	154 - 4FAC

Table D.1. continued.

	175000	12000 12000	$[m]y beam^{-1}]$						
G32.498+0.162	18:50:04.27	-00:18:43.6	5.610	12	-1.04	0.39	9	$032.4+00.1^*$	MGE 032.4982+00.1615
G32.548-0.473	18:52:25.45	-00:33:25.9	39.54	61	0.72	90.0	9	032.5-00.4	MPA J1852-0033
G32.550-0.295	18:51:47.44	-00:28:28.5	3.88	∞	I	I	I	032.5-00.3*	Te 7
G32.614+0.797	18:48:01.27	+00:04:49.0	4.09	10	1.42	1.06	Э	032.6+00.7*	PM 1-258
G32.859+0.281	18:50:18.27	+00:03:46.4	6.44	∞	0.51	0.54	κ	032.8 + 00.2*	MGE 032.8593+00.2806
G32.939-0.747	18:54:06.74	-00:20:03.7	5.43	18	-0.51	0.38	9	032.9-00.7*	CBSS 3
G33.147+1.044	18:48:06.89	+00:40:03.6	26.93	105	-0.36	90.0	9	$033.1 + 01.0^*$	
G33.353+0.404	18:50:45.90	+00:33:31.3	7.87	13	1.10	0.95	4	$033.3 + 00.4^*$	
G33.454-0.615	18:54:34.74	+00:11:03.8	21.32	9/	1.38	0.11	9	033.4-00.6	GLMP 844
G33.959+1.243	18:48:53.01	+01:28:51.1	6.20	6	I	1	1	033.9 + 01.2	IPHAS J184853.00+012852.2
G33.978-0.985	18:56:51.09	+00:28:53.5	3.80	13	-0.81	0.45	2	033.9-00.9*	PHR J1856+0028
G34.179-0.178	18:54:20.74	+01:01:43.6	11.76	27	0.13	0.31	9	034.1 - 00.1*	
G34.420-0.318	18:55:17.09	+01:10:43.6	5.97	∞	-0.49	1.23	7	034.4-00.3*	PM 1–265
G34.862-0.063	18:55:10.93	+01:41:21.1	45.37	9	0.40	0.12	9	034.8 - 00.0*	IRAS 18526+0137
G35.565-0.491	18:57:59.60	+02:07:07.7	71.20	115	-0.01	0.05	9	035.5-00.4	PHR J1857+0207
G35.770-1.245	19:01:03.04	+01:57:23.8	4.21	9	I	I	I	035.7-01.2	UWISH2 PN 3
G35.814-0.253	18:57:36.10	+02:26:57.8	1.69	5	I	I	ı	$035.8 - 00.2^*$	UWISH2 PN 2
G36.012-0.256	18:57:58.29	+02:37:25.2	11.57	49	0.92	0.19	9	$036.0 - 00.2^*$	
G36.539+0.201	18:57:18.44	+03:18:05.4	18.89	42	-0.17	0.12	9	036.5 + 00.2*	
G36.954-1.181	19:02:59.44	+03:02:19.3	8.91	31	-0.52	0.16	9	036.9-01.1	HaTr 11
G37.903-0.275	19:01:30.36	+04:17:50.6	19.29	25	0.57	0.21	9	037.9-00.2*	
G37.960+0.454	18:59:00.53	+04:40:52.2	10.58	31	0.91	0.26	9	037.9+00.4	IRAS 18564+0436
G40.261-0.276	19:05:51.00	+06:23:32.4	3.73	17	0.59	0.58	9	$040.2 - 00.2^*$	
G40.336-1.010	19:08:36.72	+06:07:15.9	11.37	49	-0.14	0.12	9	$040.3 - 01.0^*$	
G40.370-0.474	19:06:45.51	+06:23:53.9	13.47	46	-0.05	0.04	9	040.3-00.4	Abell 53
G40.554-0.091	19:05:43.83	+06:44:13.9	4.84	24	-0.13	0.29	S	$040.5 - 00.0^*$	IPHASX J190543.8+064413
G41.271-0.697	19:09:13.60	+07:05:44.0	3.73	18	-0.88	0.29	9	041.2–00.6*	HaTr 14
G41.354+0.539	19:04:57.38	+07:44:16.2	36.43	131	0.32	0.05	9	041.3+00.5*	
G42.242+1.180	19:04:18.01	+08:49:14.9	1.89	9 ;	1	1 0	1 \	$042.2+01.1^*$	IPHASX J190417.9+084916
G42.663-0.865	19:12:25.41	+08:15:07.3	16.75	76	-0.24	0.10	9 \	042.6-00.8	PM 1–288
G42.767+0.822	19:06:33.70	+09:07:21.1	9.82	24	-0.32	0.18	9 \	042.7+00.8	[GKF2010] MN102
G43.029+0.140	19:09:30.19	+09:02:26.6	66.52		1.13	0.00	9 '	043.0+00.1	GLMP 8/9
G43.294-0.646	19:12:49.29	+08:54:48.0	38.97	68	1.28	0.08	9	043.2-00.6	IRAS 19104+0849
G43.580+0.026	19:10:56.62	+09:28:37.0	9.82	23	0.93	0.29	S.	043.5 + 00.0*	1631–598D
G44.638+0.483	19:11:17.20	+10:37:35.0	14.63	54	0.45	0.14	9	044.6+00.4	IRAS 19089+1032
G44.735+0.260	19:12:16.21	+10:36:33.4	8.02	23	0.32	0.34	9	044.7 + 00.2	AGP 1
G44.949+0.898	19:10:22.18	+11:05:38.6	7.78	56	-0.01	0.20	9	044.9 + 00.8	IPHASX J191022.1+110538
G45.283-0.627	19:16:30.44	+10:40:56.8	20.12	72	0.80	0.10	9	045.2-00.6	1648–2717
G45.661+1.040	19:11:11.93	+11:47:27.1	4.11	13	-0.23	0.46	9	$045.6 + 01.0^*$	Pa 128
G46.975+0.270	19:16:29.09	+12:35:51.5	17.15	9	90.0	0.12	9	$046.9 + 00.2^*$	
G47.611+1.082	19:14:45.09	+13:32:18.4	1.37	6	-0.54	99.0	4	$047.6 + 01.0^*$	IPHASX J191445.1+133219
G47.636-1.232	19:23:11.47	+12:28:37.3	17.02	36	-0.29	0.11	9	$047.6 - 01.2^*$	Pa 132
G47.688-0.302	19:19:55.74	+12:57:37.7	3.73	28	1.27	0.33	4	$047.6 - 00.3^*$	PM 1–296

Table D.1. continued.

s PNG ¹ Name ¹	047.9+00.2*	048.1+01.1 K 3-29	048.7+00.9 IPHASX J191727.2+142735	050.0+01.0 IRAS 19171+1536	050.5+00.0 NVSS J192414+153909	050.8+00.0*	051.0-00.4* MGE 051.0214-00.4885	051.5+00.2 KLW 1	$051.6+00.9^*$	051.8+00.2 IPHAS J192553.53+165331.4	$052.1-00.3^*$	054.7+00.4* UWISH2 PN 4	055.5-00.5 M 1-71	056.4–00.9 K 3–42	056.4-00.3 IPHASX J193740.4+203547	057.9–00.7 Kn 7	058.1–00.8 IPHASX J194301.3+215424	058.6+00.9 PM 1-309	059.4-00.7 PM 1-313	059.7-00.8 IPHASX J194633.0+231659	059.8–00.5 2MASS J19453289+2328105	059.9+00.6* PM 1-311	060.2+00.8 Kn 11	060.5-00.3 K 3-45	060.9–00.5 IRAS 19461+2419	061.2-00.0* PM 1-314	062.4-00.2 M 2-48	062.7+00.0 IPHASX J194940.9+261521	062.7-00.7 IPHASX J195248.8+255359	$063.0+00.5^*$	063.5+00.0 IPHASX J195126.5+265839	063.8+00.1 K 3-48	064.7-00.5* Pa 139	065.8-00.8 IPHASX J200041.5+283023	1 00 0 100
fit_spws	9	9	9	9	9	9	2	9	9	9	9	4	9	9	9	2	4	2	9	2	9	I	9	5	4	9	9	5	9	9	2	9	9	\mathcal{E}	7
$\nabla \alpha$	0.17	0.12	0.08	0.14	0.02	0.36	0.48	0.01	0.10	0.02	0.03	0.77	0.01	0.11	0.17	0.27	0.56	0.47	0.05	0.35	0.05	I	0.10	0.50	0.40	0.35	0.00	0.38	0.07	0.32	0.47	0.04	0.18	0.71	0.12
α	-0.22	1.32	0.40	0.17	0.72	-0.43	0.07	-0.04	0.80	0.61	-0.65	-1.20	1.12	0.80	-0.15	-0.72	-0.68	-0.12	0.26	-1.05	0.69	I	-0.58	-0.61	09.0	-1.02	-0.24	-0.67	0.0	-0.38	-0.10	0.03	-0.18	0.76	7.50
S/N	46	29	9/	4	215	23	14	314	72	252	251	10	560	9	21	21	∞	22	110	22	86	2	57	21	27	20	104	21	105	22	20	116	37	S	7
Sp [m.Iv beam ⁻¹]	8.07	17.81	17.12	15.18	93.33	7.97	1.92	107.19	11.96	54.21	40.02	1.06	96.49	09.6	3.60	2.66	1.52	3.54	27.54	3.31	17.95	0.64	7.78	3.32	6.65	4.70	19.84	4.15	15.39	5.73	3.59	29.92	7.17	0.68	900
Dec.	+13:27:39.9	+14:03:49.0	+14:27:35.8	+15:41:36.7	+15:39:12.4	+15:57:28.7	+15:48:37.5	+16:33:04.8	+16:59:21.5	+16:53:31.9	+16:51:17.4	+19:29:04.4	+19:42:23.6	+20:19:03.3	+20:35:43.1	+21:45:21.4	+21:54:25.9	+23:09:47.2	+22:58:33.6	+23:16:59.3	+23:28:11.2	+24:13:25.5	+24:30:53.2	+24:11:04.6	+24:27:27.8	+24:53:23.0	+25:54:30.5	+26:15:20.2	+25:53:58.1	+26:49:26.7	+26:58:37.7	+27:18:31.8	+27:41:53.5	+28:30:26.4	120.17.22 8
R.A.	19:18:40.40	19:15:30.59	19:17:27.34	19:19:22.93	19:24:14.60	19:24:52.01	19:27:06.85	19:25:40.57	19:23:07.27	19:25:53.53	19:28:56.77	19:31:10.80	19:36:26.87	19:39:35.73	19:37:40.14	19:42:26.08	19:43:01.32	19:37:29.40	19:45:34.20	19:46:33.11	19:45:32.94	19:41:16.63	19:41:19.09	19:46:15.60	19:48:14.27	19:46:53.58	19:50:28.47	19:49:40.93	19:52:48.91	19:48:23.27	19:51:26.54	19:52:09.18	19:56:35.12	20:00:41.34	10.55.01 03
Gal.ID	G47.987+0.202	G48.162+1.163	G48.732+0.930	G50.040+1.096	G50.556+0.045	G50.895+0.057	G51.022-0.489	G51.510+0.167	G51.606+0.914	G51.834+0.284	G52.150-0.376	G54.712+0.420	G55.507-0.558	G56.402-0.903	G56.422-0.373	G57.980-0.768	G58.179-0.811	G58.641+0.919	G59.399-0.788	G59.778-0.829	G59.824-0.536	G59.991+0.686	G60.249+0.822	G60.524-0.318	G60.987-0.570	G61.206-0.088	G62.494-0.270	G62.702+0.060	G62.755-0.727	G63.045+0.598	G63.523+0.089	G63.889+0.123	G64.730-0.518	G65.893-0.866	C65 0111+0 500

1 These information are from the HASH planetary nebula database (Parker et al. 2016). PNe that have no 20 cm radio flux information in the HASH catalog are marked with *.

Table E.2. Detected pulsars in our continuum catalog

Gal.ID	R.A.	Dec.	S_p	S/N	α	Δα	fit_spws	PSR Name
Gai.iD	[J2000]	[J2000]	[mJy beam ⁻¹]	5/14	α	Δa	nt_spws	1 SIC I Vallic
G16.405+0.610	18:18:23.69	-14:22:38.1	5.60	14	-2.95	0.30	6	B1815-14
G10.403+0.010 G17.160+0.483	18:20:19.69	-14.22.38.1 -13:46:17.0	2.40	7	-2.93	0.50	- -	B1817–13
G17.100+0.483 G18.001-0.691	18:26:13.18	-13:34:47.9	3.55	16	-1.29	0.48	5	B1823–13
G19.767+0.946	18:23:40.30	-13.34.47.9 -11:15:10.5	3.48	17	-1.29 -1.55	0.48	6	B1820–11
G19.707+0.940 G19.810+0.741	18:24:29.54	-11:13:10.3 -11:18:40.0	1.01	6	-1.05	1.73	2	B1821–11
G21.286+0.798	18:27:05.49	-09:58:43.4	1.74	8	-2.72	4.08	$\overset{2}{2}$	B1824–10
G21.280+0.798 G23.385+0.063	18:33:40.15	-09.38.43.4 -08:27:32.1	4.89	8	-2.72 -1.35	1.10	3	B1830-08
G25.383+0.003 G25.172+0.762	18:33:40.13	-06:33:03.3	1.77	5	0.06	1.10	2	J1834–0633
G23.172+0.762 G27.073-0.941	18:44:05.09	-00:33:03.3 -05:38:32.7	1.77	9	-2.03	0.59	4	B1841–05
G27.322-0.033	18:44:03.09	-05:38:32.7 -05:00:21.6	6.27	6		0.39		J1841–05
G27.818+0.279	18:41:05.74	-03.00.21.6 -04:25:21.6	2.11	6	0.97	1.57		B1838-04
G27.818+0.279 G28.193–0.785	18:45:34.76	-04:23:21.6 -04:34:29.2	2.11	8	0.97	1.93	2	B1842-04
G28.347+0.174	18:43:34.76	-04:34:29.2 -03:59:59.6	6.83	12	-1.88	0.89	3	B1839-04
							3	
G28.876–0.939	18:47:22.78	-04:02:15.3	3.01	10	-1.90	0.60 0.77	3	B1844-04
G31.339+0.039	18:48:23.56	-01:23:58.9	10.64	14	-1.15			B1845-01
G32.763+0.091	18:50:48.27	-00:06:31.1	12.99	17	-0.66	0.35	6 2	J1850-0006
G36.007+0.057	18:56:50.89	+02:45:45.4	1.50	6	-2.84	5.76		J1856+0245
G37.213-0.637	19:01:31.72	+03:31:05.4	3.27	9	-3.21	1.17	2	B1859+03
G38.163-0.151	19:01:32.37	+04:35:08.1	5.45	17	-2.86	0.31	5	J1901+0435
G39.814+0.335	19:02:50.39	+06:16:33.2	1.52	7	-4.68	0.64	2	B1900+06
G40.569+1.056	19:01:39.01	+07:16:34.4	1.78	10	-1.00	0.82	4	B1859+07
G40.604–0.304	19:06:35.18	+06:41:03.9	2.27	10	_	_	-	B1904+06
G40.944+0.065	19:05:53.73	+07:09:19.0	1.25	6	_	-	_	B1903+07
G41.520–0.871	19:10:18.84	+07:14:09.5	1.07	5	-0.89	1.23	2	J1910+0714
G41.740–0.772	19:10:22.07	+07:28:35.8	1.19	8	-1.08	1.31	2	J1910+0728
G43.501–0.684	19:13:20.74	+09:04:42.2	2.61	7	_	_	_	J1913+0904
G44.557–1.019	19:16:32.43	+09:51:26.8	1.53	11	0.01	2.34	2	B1914+09
G44.707-0.650	19:15:29.98	+10:09:43.9	1.15	7	-2.09	3.31	2	B1913+10
G44.832+0.992	19:09:48.76	+11:02:02.5	1.75	9	0.51	1.12	3	B1907+10
G47.576+0.451	19:16:58.69	+13:12:49.4	1.84	18	-0.73	0.48	5	B1914+13
G48.260+0.624	19:17:39.80	+13:53:56.6	4.68	33	-1.74	0.22	6	B1915+13
G49.096+0.866	19:18:23.54	+14:45:04.0	2.10	13	-0.45	1.08	3	B1916+14
G51.859+0.063	19:26:45.32	+16:48:32.0	1.42	10	0.76	1.37	3	B1924+16
G55.575+0.639	19:32:07.99	+20:20:47.6	0.88	7	-3.67	3.30	2	B1929+20
G57.509-0.290	19:39:38.59	+21:34:58.7	11.05	21	-3.78	0.13	6	B1937+21
G57.903+0.308	19:38:14.12	+22:13:11.6	0.74	5	2.73	1.29	2	J1938+2213
G65.839+0.443	19:55:27.97	+29:08:43.4	1.13	10	-0.69	1.62	3	B1953+29
G65.924+0.772	19:54:22.47	+29:23:16.3	8.10	43	-3.45	0.13	6	B1952+29

Table F.3. X-ray sources detected in our continuum catalog. Sources do not have radio counterparts in SIMBAD or NED are marked with *.

Table F.3. continued.

Gal.ID	R. A.	Dec.	S	S/N	α	Δα	fit spws	X-ray ID
	[J2000]	[J2000]	$[mJy beam^{-1}]$				•	•
G40.437-0.797*	19:08:02.35	+06:18:32.9	1.56	6	1	ı	ı	3XMM J190802.5+061833
G42.895+0.573	19:07:41.80	+09:07:19.2	352.36	415	-0.51	0.01	9	1SXPS J190741.6+090715;3XMM J190742.0+090713
G43.407-0.021	19:10:47.32	+09:18:09.5	40.93	40	-1.04	0.07	9	3XMM J191047.7+091806
G45.366-0.219	19:15:11.52	+10:56:44.4	108.14	188	0.10	0.03	9	3XMM J191511.4+105645
G49.093-0.462*	19:23:13.84	+14:07:27.7	11.9	10	-0.28	0.42	S	3XMM J192313.4+140727
G49.113+0.935*	19:18:10.57	+14:47:55.8	1.22	9	I	I	I	3XMM J191810.7+144754
G53.769+0.161*	19:30:13.64	+18:32:01.4	1.32	S	1	I	I	1SXPS J193013.6+183203
$G54.116-1.014^*$	19:35:16.00	+18:16:11.7	0.74	2	I	I	I	1SXPS J193516.3+181613
G55.237+0.387*	19:32:22.82	+19:55:45.4	1.31	12	-1.93	0.82	4	1SXPS J193222.6+195543
G56.082+0.105	19:35:10.44	+20:31:54.7	319.89	1206	0.44	0.00	9	1SXPS J193510.2+203156
G57.157+0.796*	19:34:49.41	+21:48:24.9	1.53	11	-0.36	1.00	4	3XMM J193449.4+214824
$G57.161+0.746^*$	19:35:01.48	+21:47:11.1	1.08	7	-1.44	3.13	2	3XMM J193501.3+214710
G57.414+0.855*	19:35:08.35	+22:03:36.7	69.0	2	I	I	I	3XMM J193508.2+220336
$G59.031 + 0.850^{*}$	19:38:35.28	+23:28:08.3	2.07	13	-0.55	0.55	4	1SXPS J193835.5+232804
G59.774+0.047*	19:43:14.14	+23:43:05.1	96.0	9	-2.29	3.21	2	CXO J194314.5+234307
G59.908+0.199*	19:42:57.05	+23:54:34.4	4.01	14	1.00	89.0	4	CXO J194257.3+235435
G61.444+1.227	19:42:22.78	+25:45:13.4	3.53	7	-1.82	1.35	2	1SXPS J194222.7+254518
G63.271-1.158	19:55:38.63	+26:07:10.1	3.23	13	-1.02	0.52	5	1SXPS J195538.5+260708;3XMM J195538.4+260708
$G63.356-0.976^{*}$	19:55:08.74	+26:17:11.9	0.95	7	I	I	I	1SXPS J195508.3+261712
G63.658+1.188	19:47:29.19	+27:39:01.6	34.15	70	-1.16	0.05	9	3XMM J194729.1+273903
G63.750+1.071*	19:48:09.22	+27:40:14.9	1.58	∞	0.99	1.12	8	3XMM J194809.5+274023
G63.791-0.176*	19:53:04.71	+27:04:15.1	0.65	9	2.21	1.38	2	3XMM J195304.4+270413
G63.905+0.998*	19:48:47.56	+27:46:04.7	1.59	10	-1.02	0.84	3	CXO J194847.4+274605;3XMM J194847.4+274603
G64.075+0.077*	19:52:45.54	+27:26:39.9	1.19	∞	-3.31	1.15	3	3XMM J195245.4+272637
G65.665+1.042*	19:52:42.21	+29:18:16.8	1.74	∞	-1.11	0.73	3	CXO J195241.8+291820;3XMM J195241.9+291821
G65.678+1.205*	19:52:05.48	+29:23:58.1	3.82	6	-1.31	0.81	3	CXO J195206.1+292353
G65.731+1.063	19:52:46.28	+29:22:19.7	6.19	24	-1.64	0.23	9	CXO J195246.4+292213
G65.779-0.337*	19:58:21.98	+28:41:17.0	1.45	6	-7.52	2.33	2	1SXPS J195821.6+284118
G65.950-0.184	19:58:10.66	+28:54:50.2	14.12	85	-1.03	0.07	9	1SXPS J195810.6+285449
G66.766+0.458	19:57:37.30	+29:56:39.4	7.20	20	-0.13	0.14	9	1SXPS J195737.5+295640
G66.891+0.083*	19:59:23.87	+29:51:19.9	3.24	22	-0.47	0.35	5	1SXPS J195923.8+295115