

ON CONNECTED PREIMAGES OF SIMPLY-CONNECTED DOMAINS UNDER ENTIRE FUNCTIONS

LASSE REMPE-GILLEN AND DAVE SIXSMITH

ABSTRACT. Let f be a transcendental entire function, and let $U, V \subset \mathbb{C}$ be disjoint simply-connected domains. Must one of $f^{-1}(U)$ and $f^{-1}(V)$ be disconnected?

In 1970, Baker [Bak70] implicitly gave a positive answer to this question, in order to prove that a transcendental entire function cannot have two disjoint completely invariant domains. (A domain $U \subset \mathbb{C}$ is completely invariant under f if $f^{-1}(U) = U$.)

It was recently observed by Julien Duval that there is a flaw in Baker's argument (which has also been used in later generalisations and extensions of Baker's result). We show that the answer to the above question is negative; so this flaw cannot be repaired. Indeed, for the function $f(z) = e^z + z$, there is a collection of *infinitely many* pairwise disjoint simply-connected domains, each with connected preimage. We also answer a long-standing question of Eremenko by giving an example of a transcendental meromorphic function, with infinitely many poles, which has the same property.

Furthermore, we show that there exists a function f with the above properties such that additionally the set of singular values $S(f)$ is bounded; in other words, f belongs to the *Eremenko-Lyubich class*. On the other hand, if $S(f)$ is finite (or if certain additional hypotheses are imposed), many of the original results do hold.

For the convenience of the research community, we also include a description of the error in the proof of [Bak70], and a summary of other papers that are affected.

1. INTRODUCTION

Almost half a century ago, Baker [Bak70] proved that a transcendental entire function cannot have two disjoint completely invariant domains; in particular, the Fatou set of such a function has either one or infinitely many connected components. (Since we do not focus on dynamics in this paper, we refer to [Ber93] for background and definitions.) However, in 2016 Julien Duval observed that there is a flaw in Baker's proof. It follows that the question of whether a transcendental entire function can have two disjoint completely invariant domains remains open. The same flaw is also found in several subsequent proofs, which aimed to sharpen or generalise Baker's original result.

Baker's proof in [Bak70] is topological, rather than dynamical, and (if correct) would give a positive answer to the following question.

1.1. Question (Connected preimages of simply-connected domains).

If f is a transcendental entire function, and G_1 and G_2 are disjoint simply-connected domains, then is it true that at least one of $f^{-1}(G_1)$ and $f^{-1}(G_2)$ is disconnected?

The main aim of this article is to give a negative answer to Question 1.1.

2010 *Mathematics Subject Classification.* Primary 30D20; Secondary 30D05, 37F10, 30D30.

The first author was partially supported by a Philip Leverhulme Prize.

1.2. Theorem (Connected preimages).

Let $f(z) = e^z + z$. Then there is an infinite sequence $(U_j)_{j=1}^{\infty}$ of pairwise disjoint simply-connected domains such that $f^{-1}(U_j)$ is connected for all j .

Although Theorem 1.2 does not answer Baker's original question about completely invariant domains, it shows that his purely topological argument cannot be repaired. New ingredients, which involve the dynamics of the function under consideration, would therefore be required to resolve this problem.

While the function f in Theorem 1.2 is rather simple, the structure of the domains (U_j) is very complicated; they are constructed through a careful recursive procedure that is somewhat reminiscent of the famous "Lakes of Wada". We show that some similar complexity is necessary, by establishing that the answer to Question 1.1 is positive when certain additional hypotheses are imposed on the domains G_1 and G_2 . In the statement of the following theorem, $S(f)$ denotes the closure of the set of all critical and finite asymptotic values of f in \mathbb{C} . Also, \mathcal{S} is the *Speiser class*, consisting of those transcendental entire functions f for which $S(f)$ is finite. Note that part (h) of this theorem is an immediate consequence of [BE08a, Theorem 1], which does not depend on Baker's argument, and is included here only for completeness.

1.3. Theorem (Disconnected preimages).

Suppose that f is a transcendental entire function, and that G_1, G_2 are disjoint simply-connected domains such that $f^{-1}(G_1)$ is connected. If any of the following conditions hold, then $f^{-1}(G_2)$ is disconnected.

- (a) G_1 is bounded and its closure does not separate G_2 from infinity.
- (b) $G_1 \cap S(f)$ is compactly contained in G_1 .
- (c) $f \in \mathcal{S}$.
- (d) The domain $f^{-1}(G_1)$ contains two asymptotic curves tending to different transcendental singularities of f^{-1} .
- (e) There exists $\xi \in \partial G_1 \cap \partial G_2$ such that ξ is accessible from both G_1 and G_2 .
- (f) Infinity is accessible from G_1 .
- (g) $\overline{G_1} \cap \overline{G_2} = \emptyset$.
- (h) f has an omitted value.

Remark. In condition (d), an asymptotic curve is a curve to ∞ along which f converges to an asymptotic value $a \in \hat{\mathbb{C}}$. Two curves tend to the same singularity over a if they tend to infinity within the same connected component of $f^{-1}(\Delta)$, for every connected open neighbourhood Δ of a ; compare [BE95]. In particular, (d) holds whenever G_1 contains two different asymptotic values, or contains one asymptotic value a such that the preimage of any small disc around a has at least two connected components that are mapped by f with infinite degree.

Turning briefly to the dynamics of transcendental entire functions, the following is an easy consequence of Theorem 1.3, and is at least a partial result towards the one proved in [Bak70]. (We again refer to [Ber93] for definitions.)

1.4. Corollary (Completely invariant domains).

Suppose that f is a transcendental entire function, and that G_1, G_2 are distinct Fatou

components of f . If any of the conditions of Theorem 1.3 hold, then at most one of G_1 and G_2 is completely invariant.

Observe that a Siegel disc cannot be completely invariant, as the map is injective thereon. Moreover, infinity is accessible from any Baker domain by definition. Hence, if f has more than one completely invariant Fatou component, each such component must be an attracting or parabolic basin.

Other examples. We also exhibit three other functions having properties similar to those in Theorem 1.2. First we consider the case of meromorphic functions. There are straightforward examples of meromorphic functions with two simply-connected domains each with connected preimage. For example, we can take $f(z) = \tan z$, and let G_1, G_2 be the upper and lower half-plane respectively. In a question closely related to Question 1.1, Eremenko [Ere13] asked whether a non-constant meromorphic function can have *three* disjoint simply-connected regions each with connected preimage. We show that this is indeed possible, even for a meromorphic function with infinitely many poles.

1.5. Theorem (Meromorphic functions and connected preimages).

Let $f(z) = \tan z + z$. Then there is an infinite sequence $(U_j)_{j=1}^{\infty}$ of pairwise disjoint simply-connected domains such that $f^{-1}(U_j)$ is connected for all j .

1.6. Remark.

Note that another meromorphic example can be obtained directly from Theorem 1.2 by post-composing the function $z \mapsto e^z + z$ with a fractional linear transformation that takes a point of $\mathbb{C} \setminus (U_1 \cup U_2)$ to infinity.

We next ask whether condition (c) in Theorem 1.3 can be weakened to require only that $f \in \mathcal{B}$. Here \mathcal{B} is the *Eremenko-Lyubich class* consisting of those transcendental entire functions for which the set of singular values is bounded. We show that this is not the case. We also use this example to show that the domains with connected preimages can be bounded.

1.7. Theorem (Connected preimages in \mathcal{B}).

There is a transcendental entire function $f \in \mathcal{B}$ such that there is an infinite sequence $(U_j)_{j=1}^{\infty}$ of pairwise disjoint simply-connected domains such that $f^{-1}(U_j)$ is connected for all j . Moreover, these domains can be taken to be bounded.

Our final example addresses the question whether condition (d) in Theorem 1.3 can be weakened to require only that $f^{-1}(G_1)$ contains some asymptotic curve. Again, this is not the case.

1.8. Theorem (Examples with asymptotic values).

There exists a transcendental entire function f and pairwise disjoint simply-connected domains U and V , each with connected preimage and each containing exactly one asymptotic value of f .

The asymptotic values in our example are in fact *logarithmic asymptotic values* (see Section 6). In [EL92, Lemma 11], it is stated that any completely invariant domain of

an entire function must contain all logarithmic asymptotic values of f . However, the proof uses Baker's flawed argument, and would again imply that any simply-connected domain with connected preimage contains all such values. Hence, Theorem 1.8 shows that this proof also cannot be repaired. Compare the discussion in Section 9.

Topological results. Theorems 1.2, 1.5 and 1.7 all follow from our main construction, which is topological in nature. We begin with some preliminary definitions.

1.9. Definition (Branched coverings).

A function $f: X \rightarrow Y$ between oriented topological surfaces is a *branched covering* if every point $w \in Y$ has a simply-connected neighbourhood $D \subset Y$ with the following property. If $\tilde{D} \subset X$ is a connected component of $f^{-1}(D)$, then \tilde{D} is simply-connected and $f: \tilde{D} \rightarrow D$ maps like $z \mapsto z^d$ (up to orientation-preserving homeomorphisms), for some $d \geq 1$.

If f is a branched covering, then we denote by $CV(f)$ the set of critical values of f , and by $C(f)$ the set of critical points of f .

Our construction applies to a certain class of branched coverings, which we define as follows.

1.10. Definition (Vanilla functions).

Let $f: X \rightarrow Y$ be a branched covering between non-compact simply-connected surfaces X and Y . We say that f is *vanilla* if all the following conditions hold.

- (a) All critical points of f are simple; in other words f maps topologically like $z \mapsto z^2$ in a neighbourhood of each critical point.
- (b) $CV(f)$ is a discrete infinite subset of Y .
- (c) For every $v \in CV(f)$, $f^{-1}(v)$ contains exactly one element of $C(f)$.

We are now able to state our main topological result.

1.11. Theorem (Connected preimages for vanilla functions).

Let $f: X \rightarrow Y$ be vanilla. Then there exist disjoint simply-connected domains $U, V \subset Y$ such that $f^{-1}(U)$ and $f^{-1}(V)$ are connected.

Observe that by part (b) of Theorem 1.3, the domains U and V from Theorem 1.11 must both contain infinitely many critical values of f . It follows easily that the restriction $f: f^{-1}(V) \rightarrow V$ is itself again vanilla, and hence we can apply Theorem 1.11 again, obtaining two simply-connected subdomains of V , each with connected preimage. Continuing inductively, we obtain the following corollary.

1.12. Corollary (Infinitely many domains with connected preimages).

Let $f: X \rightarrow Y$ be vanilla. Then there is an infinite sequence $(U_j)_{j=1}^{\infty}$ of pairwise disjoint simply-connected subdomains of Y such that $f^{-1}(U_j)$ is connected for all j .

1.13. Remark (Common boundaries).

By a relatively straightforward modification to the construction in Theorem 1.11, we can ensure additionally that $\partial U = \partial V$; see Observation 4.5. With a more complicated modification, we can even ensure that the infinitely many domains of Corollary 1.12 have

a common boundary. We omit the detail in order to keep our presentation as simple as possible.

Structure. The structure of this paper is as follows. To help orient the reader, we begin by giving an outline of the proof of Theorem 1.11 in Section 2. In Section 3 we collect preliminary results relating to branched coverings and vanilla maps. Section 4 is dedicated to the proof of Theorem 1.11. In Section 5 we deduce Theorems 1.2, 1.5 and 1.7. The proof of Theorem 1.8 is carried out in Section 6, using a modification of our main construction. In Section 7 we discuss the error in Baker's original proof, and then prove Theorem 1.3 in Section 8. Finally, in Section 9 we give details of the papers and results that are affected by the flawed proofs mentioned earlier.

Notation and terminology. The (Euclidean) open ball of radius $r > 0$ around $a \in \mathbb{C}$ is denoted by

$$B(a, r) = \{z: |z - a| < r\}.$$

If $X, Y \subset \mathbb{C}$ are sets such that X lies in a bounded component of the complement of Y , then we say that Y *surrounds* X .

Suppose that X is a surface (or any topological space), and let $A \subset X$. We denote the closure of A by \bar{A} or $\text{cl}(A)$. If $B \subset A$ is connected, the connected component of A containing B is denoted by $\text{comp}_B(A)$. When $x \in A$, we write $\text{comp}_x(A) := \text{comp}_{\{x\}}(A)$. Finally we set $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$.

Acknowledgements. We thank Alex Eremenko for making us aware of the error in Baker's proof discovered by Duval, and for many interesting and useful discussions in the preparation of this article. We also thank Julien Duval for allowing us to include a description of his argument (see Figure 7), and for reading an early draft of our paper. Finally, we thank the referee for several helpful comments, suggestions, and corrections.

2. AN OVERVIEW OF THE CONSTRUCTION

Recall that our main result is Theorem 1.11. For a vanilla function f , this theorem asserts the existence of two disjoint simply-connected domains U and V , each with connected preimage. Theorems 1.2, 1.5 and 1.7 follow quickly once Theorem 1.11 is established.

Theorem 1.11 will be proved by an explicit, but rather complicated, construction of the domains U and V . For the benefit of the reader, we give a rough outline of our strategy in this section. We stress that this sketch is not intended to be precise. We note also that more general constructions are possible, but these have additional complexity, which we seek to avoid; the principal goal of this paper is to establish the three examples in Theorems 1.2, 1.5 and 1.7.

Recall from the introduction that a *vanilla* function has a set of critical values with particularly simple properties. It follows from these properties any two vanilla functions are in fact topologically equivalent, and hence have the same combinatorial structure. One way of expressing this combinatorial structure is as follows (see Proposition 3.3); there is an increasing sequence of Jordan domains $(D_n)_{n=1}^\infty$ such that each D_n contains exactly n critical values, and such that there is a preimage component \tilde{D}_n of D_n that contains all the critical preimages of these critical values.

The domains U and V will be obtained as increasing unions of simply-connected domains $U_k, V_k \subset D_{n_k}$, where n_k is a sequence tending to infinity and the D_n are the Jordan domains mentioned above. Here the domains U_k and V_k are disjoint and have the following straightforward relationship to the critical values; there are connected components $\tilde{U}_k, \tilde{V}_k \subset \tilde{D}_{n_k}$ of $f^{-1}(U_k)$ and $f^{-1}(V_k)$, respectively, such that

$$f(\mathbb{C}(f) \cap (\tilde{U}_k \cup \tilde{V}_k)) = \text{CV}(f) \cap D_{n_k}.$$

(That is, all critical points corresponding to critical values in D_{n_k} belong to either \tilde{U}_k or \tilde{V}_k .) We shall call (D_{n_k}, U_k, V_k) a *partial configuration*. Compare Proposition 4.2 and Figure 1.

Given a partial configuration (D_{n_k}, U_k, V_k) , the crux of the proof is to construct a subsequent partial configuration $(D_{n_{k+1}}, U_{k+1}, V_{k+1})$ with $n_{k+1} > n_k$, $V_{k+1} = V_k$, and, most importantly, such that \tilde{U}_{k+1} contains all the preimage components of U_k in \tilde{D}_{n_k} .

This construction is given in Proposition 4.3; compare Figures 2 and 3. We choose n_{k+1} so that $D_{n_{k+1}} \setminus D_{n_k}$ contains the same number of critical values as V_k . We then create U_{k+1} by adding a thin “snake” to U_k that “wraps around” all these critical values and also V_k , and then includes the critical values. It is shown that this domain U_{k+1} has the required properties. Proving that there is a preimage component of U_{k+1} that does indeed contain all the preimage components of U_k is the point at which we rely on the simple properties of the preimages of partial configurations.

At the next stage we reverse the roles of U_k and V_k , and then iterate the two steps of this process infinitely often. Finally we set $U = \bigcup U_k$ and $V = \bigcup V_k$. It is then shown to follow from the construction that $f^{-1}(U)$ and $f^{-1}(V)$ are indeed connected, and this completes the proof of Theorem 1.11.

3. PRELIMINARY RESULTS

We require two simple results concerning branched coverings. The first we use frequently, without comment.

3.1. Proposition (Preimages of simply-connected domains).

Let $f: X \rightarrow Y$ be a branched covering between non-compact simply-connected surfaces. Suppose that $U \subset Y$ is a simply-connected domain, and that \tilde{U} is a component of $f^{-1}(U)$ such that $\tilde{U} \cap \mathbb{C}(f)$ is finite. Then $f: \tilde{U} \rightarrow U$ is a proper map, and \tilde{U} is simply-connected.

If additionally U is bounded by a Jordan curve in Y that contains no critical values of f , then \tilde{U} is also bounded by a Jordan curve in Y .

Remark. Note that any non-compact simply-connected surface is homeomorphic to the plane. In particular, f is topologically equivalent either to an entire function $\mathbb{C} \rightarrow \mathbb{C}$ or a holomorphic map $\mathbb{D} \rightarrow \mathbb{C}$.

Proof. The fact that any pre-image component of a simply-connected domain is simply-connected follows from the fact that f is an open mapping. If $f: \tilde{U} \rightarrow U$ was not a proper map, and hence had infinite degree, it would have to contain infinitely many critical points, essentially by the Riemann-Hurwitz formula. See [BFR15, Proposition 2.8], where an analogous result is stated for entire functions; the proof is purely topological and applies equally in our setting.

Likewise, the final claim of the proposition is proved for entire functions in [BFR15, Proposition 2.9 (3)], and again the proof applies in our setting. \blacksquare

3.2. Proposition (Entire functions and branched coverings).

Suppose that f is an entire function, and that $U \subset \mathbb{C}$ is a simply-connected domain such that U contains no asymptotic values and $U \cap \text{CV}(f)$ is discrete. Then f is a branched covering from each component of $f^{-1}(U)$ to U .

Proof. This is clear from the definition. \blacksquare

We also need two structural results that are useful when studying vanilla functions.

3.3. Proposition (Increasing sequence of Jordan domains).

Let $f: X \rightarrow Y$ be vanilla, and let c be a critical point of f . Then there is an increasing sequence $(D_n)_{n=1}^\infty$ of Jordan domains in Y such that $f(c) \in D_1$, $\bigcup_{n \geq 1} D_n = Y$, and the following hold for all $n \geq 1$:

- (a) $\overline{D_n} \subset D_{n+1}$;
- (b) $\partial D_n \cap \text{CV}(f) = \emptyset$;
- (c) $\#(D_n \cap \text{CV}(f)) = n$; and
- (d) $f(\tilde{D}_n \cap C(f)) = D_n \cap \text{CV}(f)$, where $\tilde{D}_n := \text{comp}_c(f^{-1}(D_n))$.

Proof. Without loss of generality, we can assume that $X = Y = \mathbb{C}$ and that $c = f(c) = 0$.

Claim. Let $R > 0$. Then there is a bounded Jordan domain $U \supset B(0, R)$ such that $\partial U \cap \text{CV}(f) = \emptyset$ and such that $\tilde{U} := \text{comp}_0(f^{-1}(U))$ satisfies

$$f(\tilde{U} \cap C(f)) = U \cap \text{CV}(f).$$

Proof. Let V be a ball around 0 such that

$$V \supset \text{comp}_0(f^{-1}(B(0, R))) \cup (f^{-1}(B(0, R)) \cap C(f)).$$

Note that this is possible since $B(0, R) \cap \text{CV}(f)$ is finite, and each point in this set only has one preimage in $C(f)$. Let U^1 be a Jordan domain containing $\overline{f(V)}$, and $\tilde{U}^1 := \text{comp}_0(f^{-1}(U^1))$.

For each critical value $v \in U^1$ whose critical preimage is not in \tilde{U}^1 , choose an arc $\gamma_v \subset U^1 \setminus \overline{B(0, R)}$ connecting v to ∂U^1 , in such a way that γ_v contains no other critical values and arcs for different critical values are disjoint. Form the simply-connected domain $U^2 := U^1 \setminus \bigcup \gamma_v \supset \overline{B(0, R)}$ and set $\tilde{U}^2 := \text{comp}_0(f^{-1}(U^2))$. Note that, for each critical value v chosen as above, any preimage of v in \tilde{U}^1 is not a critical point, and so any preimage component of an arc γ_v in \tilde{U}^1 is an arc $\tilde{\gamma}_v$ connecting a simple preimage of v to $\partial \tilde{U}^1$. In particular, $\tilde{U}^2 = \tilde{U}^1 \setminus \bigcup \tilde{\gamma}_v$.

Then U^2 contains $\overline{B(0, R)}$, and the preimage component \tilde{U}^2 of $f^{-1}(U^2)$ contains all critical preimages of critical values in U^2 . Choosing a Jordan domain $U \subset U^2$ slightly smaller than U^2 , if necessary, we can ensure that $\partial U \cap \text{CV}(f) = \emptyset$. This completes the proof of the claim. \triangle

Now we use the claim to construct a subsequence (D_{n_j}) of the desired sequence (D_n) inductively, as follows. Set $n_1 := 1$, and let D_1 be any small disc around 0 not containing any other critical values in its closure. If n_j and D_{n_j} have been defined, let $R \geq j$

be sufficiently large that $\overline{D_{n_j}} \subset B(0, R)$. Then apply the claim to obtain a domain $U \supset \overline{D_{n_j}}$. Set $n_{j+1} := \#(U \cap \text{CV}(f))$, and $D_{n_{j+1}} := U$. Then the domains (D_{n_j}) satisfy all requirements in the statement of the proposition. It remains to define D_n for the remaining values of n .

We next construct the domains D_n for $n_j < n < n_{j+1}$ by removing thin slits containing critical values in $U_{n_{j+1}}$. More precisely, suppose that N_1, N_2 are such that $N_1 < N_2 - 1$, D_{N_1} and D_{N_2} have been defined, but D_n has not yet been defined for $N_1 < n < N_2$. Set $n := N_2 - 1$, and construct D_n from D_{N_2} as follows. For each critical value v in $D_{N_2} \setminus D_{N_1}$, let γ_v again be an arc connecting v to ∂D_{N_2} , not intersecting ∂D_{N_1} , and such that different arcs are pairwise disjoint.

For each such v , let $\tilde{\gamma}_v = \text{comp}_{c(v)}(f^{-1}(\gamma_v))$, where $c(v)$ is the unique critical point of f over v . Then $\tilde{\gamma}_v$ is a cross-cut of \tilde{D}_{N_2} , not intersecting \tilde{D}_{N_1} . At least one of these crosscuts, say $\tilde{\gamma}_{v_0}$, does not separate \tilde{D}_{N_1} from any other $\tilde{\gamma}_v$. Observe that any other component of $f^{-1}(\gamma_{v_0})$ in \tilde{D}_{N_2} is an arc connecting $\partial \tilde{D}_{N_2}$ to some simple preimage of v_0 , and hence does not disconnect \tilde{D}_{N_2} .

Set $U := D_{N_2} \setminus \gamma_{v_0}$ and $\tilde{U} := \text{comp}_{\tilde{D}_{N_1}}(f^{-1}(U))$. Then, by construction, \tilde{U} contains $n = N_2 - 1$ critical points. Slightly shrinking U if necessary, we obtain a disc D_n with $D_{N_1} \subset D_n \subset D_{N_2}$, satisfying the desired properties. Proceeding inductively, we define D_n for all $n \geq 1$. \blacksquare

3.4. Proposition (Connecting to the critical point).

Let $f: X \rightarrow Y$ be vanilla and let $(D_n)_{n=1}^\infty$ and $(\tilde{D}_n)_{n=1}^\infty$ be sequences as in the statement of Proposition 3.3. Fix $n \geq 1$ and $\zeta \in \partial \tilde{D}_n$.

Then there is a Jordan arc τ connecting $f(\zeta)$ to ∂D_{n+1} and lying in the annulus $D_{n+1} \setminus \overline{D_n}$ apart from its endpoints, such that $\text{comp}_\zeta(f^{-1}(\tau))$ contains the unique critical point c' of f in $\tilde{D}_{n+1} \setminus \tilde{D}_n$.

Proof. Note that $f: \tilde{D}_n \rightarrow D_n$ has degree $n + 1$, while $f: \tilde{D}_{n+1} \rightarrow D_{n+1}$ has degree $n + 2$, by the Riemann-Hurwitz formula. Hence \tilde{D}_{n+1} contains exactly one component U of $f^{-1}(D_n)$ different from \tilde{D}_n , and U is mapped conformally to D_n .

Choose an arc τ_0 connecting $f(\zeta)$ to ∂D_{n+1} lying in the annulus $D_{n+1} \setminus \overline{D_n}$ and passing through the critical value $f(c')$. Then $\text{comp}_{c'}(f^{-1}(\tau_0))$ contains two points of $f^{-1}(f(\zeta))$. One of these points must be the unique preimage of ζ on ∂U , while the other is some point $\zeta_0 \in \Xi := f^{-1}(f(\zeta)) \cap \partial D_n$.

Now consider what happens when we apply j Dehn twists to τ_0 near ∂D_n , for $0 \leq j \leq n - 1$, obtaining curves τ_j . Each τ_j has the same properties as τ_0 above; let ζ_j be the unique element of Ξ in $\text{comp}_{c'}(f^{-1}(\tau_j))$. Since $f: \partial \tilde{D}_n \rightarrow \partial D_n$ is a degree n covering map of circles, the values of ζ_j cycle through all the elements of Ξ . Hence, there is some j such that $\zeta_j = \zeta$, and we can take $\tau = \tau_j$. \blacksquare

As mentioned in the previous section, it follows that any two vanilla functions are topologically equivalent. Although we will not use this fact directly in the following, we give a proof for completeness.

3.5. Corollary (Topological uniqueness).

Suppose that $f^1: X^1 \rightarrow Y^1$ and $f^2: X^2 \rightarrow Y^2$ are vanilla. Then there are homeomorphisms $\varphi: X^1 \rightarrow X^2$ and $\psi: Y^1 \rightarrow Y^2$ such that $\psi \circ f^1 = f^2 \circ \varphi$.

Proof. Let $(D_n^j)_{n=1}^\infty$ and $(\tilde{D}_n^j)_{n=1}^\infty$ be as in Proposition 3.3, for $j = 1, 2$. We inductively specify homeomorphisms

$$\psi_n: \text{cl}(D_n^1) \rightarrow \text{cl}(D_n^2) \quad \text{and} \quad \varphi_n: \text{cl}(\tilde{D}_n^1) \rightarrow \text{cl}(\tilde{D}_n^2)$$

such that

$$(3.1) \quad \psi_n \circ f^1 = f^2 \circ \varphi_n,$$

$\psi_{n+1}|_{\text{cl}(D_n^1)} = \psi_n$, and $\varphi_{n+1}|_{\text{cl}(\tilde{D}_n^1)} = \varphi_n$. The claim follows by defining ψ and φ to be the common extension of the maps ψ_n and φ_n , respectively.

Let $\psi_1: \text{cl}(D_1^1) \rightarrow \text{cl}(D_1^2)$ be any homeomorphism that fixes 0, and let φ_1 be either of the two lifts of ψ_1 , such that $\psi_1 \circ f^1 = f^2 \circ \varphi_1$.

Now suppose that the homeomorphisms ψ_n and φ_n have already been defined. Fix any point $\zeta_n^1 \in \partial\tilde{D}_n^1$, and set $\zeta_n^2 := \varphi_n(\zeta_n^1)$. Applying Proposition 3.4 to f^1 (with $\zeta = \zeta_n^1$) and to f^2 (with $\zeta = \zeta_n^2$), we obtain Jordan arcs τ_n^1 and τ_n^2 . We extend ψ_n to a homeomorphism ψ_{n+1} from $\text{cl}(D_{n+1}^1)$ to $\text{cl}(D_{n+1}^2)$, in such a way that ψ_{n+1} maps τ_n^1 to τ_n^2 , taking the critical value of f^1 in τ_n^1 to the critical value of f^2 in τ_n^2 . It is now straightforward to verify that there is a lift φ_{n+1} , extending φ_n , satisfying 3.1. \blacksquare

4. PROOF OF THEOREM 1.11

As mentioned in Section 2, we consider certain triples of Jordan domains to facilitate the construction. These are defined as follows.

4.1. Definition (Partial configuration).

Let $f: X \rightarrow Y$ be vanilla. A triple (D, U, V) of Jordan domains in Y is called a *partial configuration* for f if all the following hold:

- (a) $\bar{U} \cap \bar{V} = \emptyset$;
- (b) $\bar{U} \cup \bar{V} \subset D$;
- (c) $D \cap \text{CV}(f) \subset U \cup V$;
- (d) there is a component \tilde{D} of $f^{-1}(D)$ such that $f(\tilde{D} \cap \text{C}(f)) = D \cap \text{CV}(f)$;
- (e) there are components \tilde{U} and \tilde{V} of $f^{-1}(U)$ and $f^{-1}(V)$, respectively, such that $\tilde{D} \cap \text{C}(f) \subset \tilde{U} \cup \tilde{V}$.

These partial configurations are easy to work with because their preimages have a very specific and extremely simple structure, as described in the following proposition. This structure is illustrated in Figure 1. In this proposition, and subsequently, if U is a domain, then we denote by m_U the number of critical values of f in U . (Of course $m_U = m_U(f)$ depends on the function f , but we suppress f from the notation since it will be fixed whenever this notation is used.)

4.2. Proposition (Structure of partial configurations).

Let $f: X \rightarrow Y$ be vanilla, and suppose that (D, U, V) is a partial configuration. Let γ be

a crosscut of D that separates \bar{U} from \bar{V} . Let z and w be its endpoints on ∂D , labeled such that U is on the left of γ and V is on the right of γ when γ is oriented from z to w .

Label the preimages $\tilde{z}_1, \dots, \tilde{z}_d$ of z on $\partial \tilde{D}$ and the preimages $\tilde{w}_1, \dots, \tilde{w}_d$ of w on $\partial \tilde{D}$ in positive orientation such that \tilde{w}_1 is between \tilde{z}_1 and \tilde{z}_2 ; here $d = m_U + m_V + 1$ is the degree of f on \tilde{D} . This determines the labeling up to the choice of \tilde{z}_1 .

Then the choice of \tilde{z}_1 can be made in such a way that the preimage components of $f^{-1}(\gamma) \cap \tilde{D}$ connect these preimages as follows.

- (a) \tilde{z}_1 is connected to \tilde{w}_{m_V+1} .
- (b) For $j = 1, \dots, m_V$, \tilde{w}_j is connected to \tilde{z}_{j+1} .
- (c) For $j = m_V + 2, \dots, d$, \tilde{z}_j is connected to \tilde{w}_j .
- (d) Each component of $\tilde{D} \setminus f^{-1}(\gamma)$ (which we shall call a face) contains either exactly one component of $f^{-1}(U)$ or exactly one component of $f^{-1}(V)$.
- (e) The faces containing \tilde{U} and \tilde{V} are adjacent to each other, and separated by the arc connecting \tilde{z}_1 and \tilde{w}_{m_V+1} . More precisely, the face containing \tilde{U} is bounded by this arc and those in (c), while that containing \tilde{V} is bounded by this arc and those in (b).
- (f) Each connected component of $f^{-1}(U)$ in \tilde{D} , apart from \tilde{U} , is contained in a face bounded solely by one of the arcs from (b). Similarly, each component of $f^{-1}(V)$ in \tilde{D} , apart from \tilde{V} , is contained in a face bounded by one of the arcs from (c).

Remark. We allow the case where $m_U = 0$ or $m_V = 0$, in which case the preimage domains \tilde{U} and \tilde{V} from Definition 4.1 may not be unique. In this case, the claim should be understood as follows: for all valid choices of \tilde{U} and \tilde{V} , there is a choice of \tilde{z}_1 satisfying the properties listed.

Proof. Recall that all critical values of f in D lie in U or V . Let D_L be the component of $D \setminus \gamma$ containing U . Then $A := D_L \setminus \bar{U}$ is an annulus containing no singular values. Every component \tilde{A} of $f^{-1}(A)$ is mapped as a finite-degree covering map, and hence is also an annulus. It follows that \tilde{A} is a face with exactly one component of $f^{-1}(U)$ removed. Clearly we can apply the same argument to V , replacing D_L by the other component D_R of $D \setminus \gamma$. Let us call preimage components of D_L *L-faces*, and other faces *R-faces*.

By the above, every face contains exactly one preimage component of U or of V , and is mapped with the same degree as this component. It follows that the set \tilde{U} from Definition 4.1 is contained in an L-face of degree $m_U + 1$, and there are m_V further L-faces, all of which are simple. Similarly, \tilde{V} is contained in an R-face of degree $m_V + 1$, and there are a further m_U R-faces, all simple.

Consider the dual graph G to this picture, where each face represents a vertex, and two faces are connected if they are adjacent; in other words if they have a common component of $f^{-1}(\gamma)$ in their boundary. By the remarks above, G is a connected graph with d edges and $m_U + m_V + 2 = d + 1$ vertices. So G is a tree. Moreover, G has one vertex of degree $m_U + 1$ corresponding to \tilde{U} , one of degree $m_V + 1$, corresponding to \tilde{V} , and a further $d - 1$ vertices, all of which are leaves (vertices of degree 1).

Since G is connected, the vertices representing \tilde{U} and \tilde{V} must be connected by a simple path in G , and since all other vertices are leaves, this path must in fact be an edge. This




FIGURE 1. The initial preimage (left) and image (right) in Proposition 4.2. Note that the domains U and V may look very complicated geometrically, but topologically we are speaking simply of a configuration of two discs within a larger disc, as shown here. Black discs on the left are preimage components of U , while grey discs are preimage components of V . For this example we have taken $m_U = 1$ and $m_V = 2$. We stress that, in general, D will have other preimage components; these are not illustrated.

edge corresponds to some component of $f^{-1}(\gamma)$. We choose \tilde{z}_1 to be the preimage of z contained in this component. It then follows that the description is indeed as above. ■

We will deduce Theorem 1.11 from the following proposition, which is the crux of our construction.

4.3. Proposition (Extending partial configurations).

Let $f: X \rightarrow Y$ be vanilla, and let $(D_n)_{n=1}^\infty$ be the sequence from Proposition 3.3. Suppose that $n \geq 1$ and that (D_n, U, V) is a partial configuration for f . Set $n' = n + m_V$.

Then there exists $U' \supseteq U$, such that $(D_{n'}, U', V)$ is a partial configuration, and such that \tilde{U}' contains all the connected components of $f^{-1}(U) \cap \tilde{D}_n$.

Proof. For simplicity of notation, set $m := m_V$. Observe that if $m = 0$, then \tilde{U} is connected by Proposition 4.2, and there is nothing to prove. So we may assume that $n' > n$. We must describe how U is extended to U' .

We apply Proposition 4.2 to the partial configuration (D_n, U, V) , and use the notation given there in the following paragraphs. We begin by constructing a curve α connecting the point $w \in \partial D_n$ to $\partial D_{n'}$ in such a way that there is a curve contained in $f^{-1}(\alpha)$ connecting the point \tilde{w}_{m+1} to $\partial \tilde{D}_{n'}$ and passing through all the critical points in $\tilde{D}_{n'} \setminus \tilde{D}_{n_k}$. This construction proceeds as follows. First let τ be the arc obtained from Proposition 3.4, with $\zeta = \zeta_1 := \tilde{w}_{m+1}$. Then set $\alpha_1 = \tau$.

Let ω_1 be the endpoint of τ on ∂D_{n+1} . Recall that all critical points of f are simple. It follows that the component of $f^{-1}(\tau)$ containing ζ_1 contains two points of $f^{-1}(\omega_1)$, say $\tilde{\omega}_1^1$ and $\tilde{\omega}_1^2$. One of the two arcs of $\partial \tilde{D}_{n+1} \setminus \{\tilde{\omega}_1^1, \tilde{\omega}_1^2\}$ does not contain any other points of $f^{-1}(\omega_1)$. We may assume that the points are labeled such that this arc, when oriented from $\tilde{\omega}_1^1$ to $\tilde{\omega}_1^2$, traverses $\partial \tilde{D}_{n+1}$ in positive orientation. We now set $\tilde{\omega}_1 := \tilde{\omega}_1^2$, and apply Proposition 3.4 again, with $\zeta = \zeta_2 := \tilde{\omega}_1$, obtaining a curve $\alpha_2 = \tau$.

Continuing inductively, we construct a curve $\alpha = \alpha_1 \cup \dots \cup \alpha_m$ connecting ∂D_n to $\partial D_{n'}$, and passing through the m critical values in $D_{n'} \setminus D_n$; see the right-hand side of Figure 2. This completes the construction of the curve α .

Let v_1, \dots, v_m be the critical values on α , ordered as they are encountered when traversing α from w to $\partial D_{n'}$. Let c_1, \dots, c_m be the corresponding critical points. Then, for each $j = 1, \dots, m$, there is a simple preimage \tilde{D}_n^j of D_n , attached to c_j by a simple preimage of the piece β_j of α connecting D_n to v_j , and with a preimage \tilde{w}'_j of w on its boundary; see the left-hand side of Figure 2. Note, in particular, that β_m is an arc in $D_{n'} \setminus D_n$ that connects w to all the critical values v_1, \dots, v_m .

Let Γ_0 be a Jordan curve in $D_{n'} \setminus (\overline{U} \cup \overline{V} \cup (\beta_m \setminus \{w\}))$ obtained as follows. We begin at w , proceed along an arc outside of D_n that connects w to z , running around β_m in negative orientation, and returning to w from z via the curve γ from Proposition 4.2. So Γ_0 surrounds V and $\beta_m \setminus \{w\}$, but not U .

Then the preimage of Γ_0 consists of m_U simple preimages of Γ_0 , and one loop that is mapped by f as a degree $2m + 1$ covering of circles. Each of the simple preimages contains the component of $f^{-1}(\gamma)$ connecting \tilde{z}_j and \tilde{w}_j for some $j \in \{m+2, \dots, n+1\}$, and is a loop surrounding the corresponding simple R-face. The remaining loop passes (in positive orientation) through $\tilde{w}_{m+1}, \tilde{z}_1, \tilde{w}_1, \tilde{z}_2, \tilde{w}_2, \dots, \tilde{w}_m, \tilde{z}_{m+1}$, and then through the components $\tilde{D}_n^1, \dots, \tilde{D}_n^m$.

Now consider an arc Γ , not intersecting $\overline{V} \cup \text{CV}(f)$ and not intersecting \overline{U} except in one endpoint, defined as follows. The arc starts at some point $\mu \in \partial U$, and runs around β_m and \overline{V}_k m times in negative orientation; in other words, in the same manner as Γ_0 . On its last loop it enters D_n at the point z , traversing along the arc γ and ending at w . Hence Γ is homotopic (in $D_{n'} \setminus \text{CV}(f)$) to a curve that connects μ to w within the left half of $D_n \setminus \gamma$, and then traverses Γ_0 m times.

Since Γ does not contain any critical values, every component of $f^{-1}(\Gamma)$ is an arc beginning on the boundary of some preimage component of U , and ending in a preimage of w . Let $j \in \{1, \dots, m\}$ and consider the component $\tilde{\Gamma}_j$ of Γ that starts on the boundary of the simple preimage of U contained in the L-face bounded by an arc connecting \tilde{w}_j




FIGURE 2. Application of Proposition 4.3 to the configuration from Figure 1. The larger disc $D_{n'}$ in the image (right) has two additional critical points, shown as a triangle and a square, and a curve α indicated as a dashed line. The preimage $\tilde{D}_{n'}$ (left) shows preimages of the critical values as triangles and squares, with the solid figure indicating the preimage that is actually the critical point. The preimage of the curve α is indicated as a dashed line. Note also two additional preimages of w , labeled \tilde{w}'_1 and \tilde{w}'_2 , each on the boundary of a new preimage of the original disc.

and \tilde{z}_{j+1} . Our discussion of the structure of $f^{-1}(\Gamma_0)$ shows that, for $j < m$, $\tilde{\Gamma}_j$ ends at the preimage \tilde{w}'_{j+1} of w on the boundary of the disc \tilde{D}_n^{j+1} , while $\tilde{\Gamma}_m$ ends at \tilde{w}_{m+1} . Furthermore, if $\tilde{\Gamma}$ is the component of $f^{-1}(\Gamma)$ ending at \tilde{w}'_1 , then $\tilde{\Gamma}$ begins on \tilde{U} .

Recall from the construction of α that $\tilde{w}'_1, \dots, \tilde{w}'_m$ and \tilde{w}_{m+1} , as well as the critical points c_1, \dots, c_m all belong to the same component $\tilde{\beta}$ of $f^{-1}(\beta_m)$. Take a thin channel W containing the Jordan arc $\Gamma \cup \beta_m$, in such a way that $U' := U \cup W$ is simply-connected; see the right-hand side of Figure 3.




FIGURE 3. The thin channel W has been added to the image (right) from Figure 2; the part containing Γ is solid, and the part containing β_m is dashed. The preimage of W has been added to the preimage on the left, shown solid or dashed appropriately

It then follows that the component of $f^{-1}(U')$ containing \tilde{w}_{m+1} contains the connected set

$$\tilde{\beta} \cup \tilde{\Gamma} \cup \bigcup_{j=1}^m \tilde{\Gamma}_j \cup (f^{-1}(U) \cap \tilde{D}_n);$$

see the left-hand side of Figure 3. This completes the proof. ■

4.4. Corollary (Sequence of partial configurations).

Let $f: X \rightarrow Y$ be vanilla, and let $(D_n)_{n=1}^{\infty}$ be the sequence from Proposition 3.3. Then there is a sequence $(D_{n_k}, U_k, V_k)_{k=1}^{\infty}$ of partial configurations such that (U_k) and (V_k) are increasing sequences. Furthermore, the sequences can be chosen such that every component of $f^{-1}(U_1)$ is contained in \tilde{U}_k for sufficiently large k , and similarly for the components of $f^{-1}(V_1)$.

Proof. We construct a sequence $(D_{n_k}, U_k, V_k)_{k=1}^\infty$ of partial configurations, with $n_k \rightarrow \infty$, such that when k is odd we have

$$(4.1) \quad U_k \subsetneq U_{k+1}, V_k = V_{k+1}, \text{ and } f^{-1}(U_k) \cap \tilde{D}_{n_k} \subset \tilde{U}_{k+1},$$

and when k is even we have

$$V_k \subsetneq V_{k+1}, U_k = U_{k+1}, \text{ and } f^{-1}(V_k) \cap \tilde{D}_{n_k} \subset \tilde{V}_{k+1}.$$

This sequence is defined recursively using Proposition 4.3. To anchor the recursion, we simply set $n_1 := 1$, let V_1 be a sufficiently small disc around the critical value of f in D_1 , and let U_1 be any Jordan domain whose closure is contained in D_1 and disjoint from \bar{V}_1 . It is easy to check that this is indeed a partial configuration.

Now suppose that (D_{n_k}, U_k, V_k) has been defined for $k \geq 1$. If k is odd, we apply Proposition 4.3 with $n = n_k$, $U = U_k$ and $V = V_k$. Setting $n_{k+1} := n'$, $U_{k+1} := U'$ and $V_{k+1} := V_k$, we obtain another partial configuration satisfying 4.1. If k is even, then we similarly apply Proposition 4.3, but interchanging the roles of U_k and V_k . This completes the construction.

Let W be a connected component of $f^{-1}(U_1)$. Then there is some N such that $W \subset \tilde{D}_n$ for $n \geq N$. Let K be odd such that $n_K \geq N$. Then

$$W \subset f^{-1}(U_1) \cap \tilde{D}_{n_K} \subset f^{-1}(U_k) \cap \tilde{D}_{n_k} \subset \tilde{U}_k$$

for $k > K$, as desired (and similarly for $f^{-1}(V_1)$). This completes the proof. \blacksquare

The proof of Theorem 1.11 is now quite straightforward.

Proof of Theorem 1.11. Set

$$U := \bigcup_{k \in \mathbb{N}} U_k \quad \text{and} \quad V := \bigcup_{k \in \mathbb{N}} V_k,$$

where U_k and V_k are as in Corollary 4.4. Then U and V are increasing unions of simply-connected domains and so are simply-connected. Moreover, $U \cap V = \emptyset$.

Let \tilde{U} be the component of $f^{-1}(U)$ containing \tilde{U}_k for all k . Then $f^{-1}(U_1) \subset \tilde{U}$ by assumption. Choose $x \in f^{-1}(U)$, and let γ be a curve in U connecting $f(x)$ and some point y of U_1 without passing through any critical values. Then there is a curve $\tilde{\gamma} \subset f^{-1}(U)$ connecting x to a point in $f^{-1}(y) \subset f^{-1}(U_1) \subset \tilde{U}$, and hence $x \in \tilde{U}$. So $f^{-1}(U) = \tilde{U}$ is connected. Likewise $f^{-1}(V)$ is connected. \blacksquare

We also observe that we can ensure that the domains U and V of Theorem 1.11 have a common boundary, as promised in Remark 1.13.

4.5. Observation (Common boundary).

The domains U and V in Theorem 1.11 can be chosen such that $\partial U = \partial V$.

Proof. Let d be a metric on X . In the setting of Proposition 4.3, we claim that the distance $d(U', v)$ can be chosen arbitrarily small for all $v \in \partial V$.

To prove this, we let $\varepsilon > 0$, and choose a simply-connected domain U^1 such that $U \subset U^1 \subset \bar{U}^1 \subset D_n \setminus \bar{V}$, and such that every point of ∂V has distance at most ε from U^1 .

Then (D_n, U^1, V) is also a partial configuration, and we can apply Proposition 4.3. The resulting domain U' has the required properties.

Now let (ε_k) be a sequence of positive real numbers tending to zero. In the k -th step of the inductive construction in Corollary 4.4, apply the above observation for $\varepsilon = \varepsilon_k$. For the resulting domains U and V , let $v \in \partial V$, and let $\delta > 0$. Then, for all sufficiently large k , $B(v, \delta)$ intersects V_k , and hence $\partial V_k \cap B(v, \delta) \neq \emptyset$. If k is chosen odd and sufficiently large that $\varepsilon_k < \delta$, then it follows that U_k contains a point of distance at most 2δ from v . As v and δ were arbitrary, we have shown $\partial V \subset \partial U$. The converse inclusion follows analogously. ■

Proof of Corollary 1.12. First we apply Theorem 1.11 to obtain two disjoint simply-connected domains U and V each with connected preimage. Since f has infinite degree on $f^{-1}(V)$, and f is vanilla, $V \cap CV(f)$ is infinite. Set $U_1 := U$ and then again apply Theorem 1.11 to the restriction $f: f^{-1}(V) \rightarrow V$. We obtain two simply-connected subdomains of V , each with connected preimage. The result follows by induction. ■

5. EXAMPLES

In this section, we show how to deduce Theorem 1.2, Theorem 1.5 and Theorem 1.7 from Theorem 1.11.

Proof of Theorem 1.2. Let $f(z) = e^z + z$. The critical points of f are $c_m = (2m + 1)\pi i$, with $m \in \mathbb{Z}$, and the corresponding critical values are $v_m = (2m + 1)\pi i - 1$. Furthermore, it is easy to see that f has no finite asymptotic values. By Proposition 3.2, f is a branched covering map from \mathbb{C} to \mathbb{C} , and by the above statements on critical values, f is vanilla. The result then follows by Corollary 1.12. ■

Proof of Theorem 1.5. Let $f(z) = \tan z + z$. Let U be the upper half-plane given by $U := \{z: \operatorname{Im} z > 0\}$, and we note that $f^{-1}(U) = U$. Observe also that f has no asymptotic values, and that U contains no poles of f .

The critical points of f in U , which are all simple, are

$$c_m = \left(m + \frac{1}{2}\right) \pi + i \operatorname{arsinh} 1, \quad \text{for } m \in \mathbb{Z}.$$

The critical values of f in U are

$$v_m = c_m + i\sqrt{2}, \quad \text{for } m \in \mathbb{Z},$$

and so each critical value has a unique critical preimage.

By Proposition 3.2, $f: U \rightarrow U$ is vanilla, and the result follows by Corollary 1.12. ■

Remark. The dynamics of the function f from Theorem 1.5 were studied in [BFJK17].

Proof of Theorem 1.7. Bergweiler [Ber02] introduced the transcendental entire function

$$f(z) = \frac{12\pi^2}{5\pi^2 - 48} \left(\frac{(\pi^2 - 8)z + 2\pi^2}{z(4z - \pi^2)} \cos \sqrt{z} + \frac{2}{z} \right).$$

It was shown in [Ber02] that f has a completely invariant Fatou component U , such that $0 \in \partial U$ and $(0, \infty) \subset U$. It was also shown that 0 is the only finite asymptotic value of f , and that f has infinitely many critical values, all of which lie in a real interval of the form $(0, c) \subset U$, and which accumulate only on the origin. All critical points of f are simple, real and positive, and clearly also lie in U ; see [Ber02, Figure 2].

By Proposition 3.2, to prove that $f: U \rightarrow U$ is vanilla we would need to show that each critical value of f only has one critical preimage. Although this is likely to be the case, it seems quite complicated to prove.

Instead, we use quasiconformal maps to find a function “close” to f with the properties we require. Let the critical points of f be $(c_k)_{k \in \mathbb{N}}$, and choose a strictly decreasing sequence of positive real numbers $(r_k)_{k \in \mathbb{N}}$ tending to zero. For each k consider the balls $B_k := B(f(c_k), r_k)$ and the component $\tilde{B}_k := \text{comp}_{c_k}(f^{-1}(B_k))$. If r_k is chosen sufficiently small, then

$$\text{diam}(\tilde{B}_k) < \frac{\text{dist}(c_k, \partial U \cup (\mathbb{C}(f) \setminus \{c_k\}))}{2}.$$

Hence \tilde{B}_k and B_k are in U , the \tilde{B}_k are pairwise disjoint, and $f: \tilde{B}_k \rightarrow B_k$ is proper.

Let $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ be a quasiconformal map such that $\varphi(\mathbb{D}) = \mathbb{D}$, such that $\varphi(z) = z$ for $z \in \partial\mathbb{D}$, and such that $\varphi(0) = 1/2$. We then define a quasiregular map $G: \mathbb{C} \rightarrow \mathbb{C}$ by

$$G(z) = \begin{cases} r_k \varphi\left(\frac{f(z) - f(c_k)}{r_k}\right) + f(c_k), & \text{if } z \in \tilde{B}_k \text{ for some } k \in \mathbb{N}, \\ f(z), & \text{otherwise.} \end{cases}$$

It is easy to see that G is indeed quasiregular, with the same quasiconstant as φ . Moreover, G has the same critical points as f , all of which are simple, and the critical values of G are $f(c_k) + r_k/2$, for $k \in \mathbb{N}$.

Since G is a quasiregular mapping of the plane, it follows by Stoilow factorisation that there is a transcendental entire function g and a quasiconformal map $\psi: \mathbb{C} \rightarrow \mathbb{C}$ such that $g = G \circ \psi$.

It can be seen that $\text{AV}(g) = \{0\}$. The critical points of g are simple, and given by $\psi^{-1}(c_k)$, for $k \in \mathbb{N}$. The critical values of g are equal to $f(c_k) + r_k/2$, for $k \in \mathbb{N}$, and all lie in U . Thus the (infinitely many) critical values of g accumulate only at 0, and g is in the Eremenko-Lyubich class. Moreover, each critical point of g has exactly one critical preimage.

We have shown that $g: \psi^{-1}(U) \rightarrow U$ is vanilla. By Corollary 1.12, it follows that there are disjoint simply-connected domains $U_1, U_2, \dots \subset U$ each with connected preimage in $\psi^{-1}(U)$. Since $g^{-1}(U) = \psi^{-1} \circ G^{-1}(U) = \psi^{-1}(U)$, this completes the proof.

It remains to show that each U_j can be assumed to be bounded. That this is possible is easily seen as follows. Let $T > 0$ be such that $\text{CV}(g) \subset (0, T]$, and let $U^1 \subset U$ be a bounded simply-connected domain containing $(0, T]$. It is easy to see that $g^{-1}(U^1)$ is also connected. Then the claim follows by applying Corollary 1.12 to the vanilla function $g: g^{-1}(U^1) \rightarrow U^1$. \blacksquare

6. PROOF OF THEOREM 1.8

We now turn to our construction of an entire function f having two simply-connected domains with connected preimages, each containing a logarithmic asymptotic value of f . (An asymptotic value a is *logarithmic* if there is a neighbourhood U of a and a connected component \tilde{U} of $f^{-1}(U)$ such that $f: \tilde{U} \rightarrow U \setminus \{a\}$ is a universal covering map.) Similarly as in Theorem 1.11, we could introduce a topological class of functions to which our methods apply. For definiteness and simplicity, let us instead study the

explicitly given entire function

$$(6.1) \quad f(z) := \frac{2}{e^{1/4}\sqrt{\pi}} \int_0^z \cosh w \exp(-w^2) dw = \frac{1}{2} \left(\operatorname{erf} \left(z + \frac{1}{2} \right) + \operatorname{erf} \left(z - \frac{1}{2} \right) \right).$$

Here erf is the *error function* $\operatorname{erf}(z) := \frac{2}{\sqrt{\pi}} \int_0^z \exp(-w^2) dw$; the equality in (6.1) is obtained by an explicit calculation. We shall use the following properties of f .

6.1. Lemma (Properties of f).

The function f is an odd entire function, real on the real axis with $f(\mathbb{R}) = (-1, 1)$, and satisfying

$$f(\{iy: y > 0\}) = f(\{iy: y < 0\}) = \{iy: y \in \mathbb{R}\}.$$

The critical points of f are all simple and given by $z_n = i \cdot \frac{2n+1}{2}\pi$, for $n \in \mathbb{Z}$. The critical values $\zeta_n := f(z_n)$ are all purely imaginary, and distinct critical points have distinct critical values. Moreover, $\operatorname{Im} \zeta_n$ is positive for even n , and negative for odd n .

In addition, there are exactly two asymptotic values, namely 1 and -1 , and with $f(x) \rightarrow \pm 1$ as $x \rightarrow \pm\infty$. These are logarithmic asymptotic values and there is exactly one tract over each. That is, let U be a sufficiently small Jordan domain containing $a \in \{-1, 1\}$, and let \tilde{U} be the connected component of $f^{-1}(U)$ containing an infinite piece of the real axis. Then $f: \tilde{U} \rightarrow U \setminus \{a\}$ is a universal covering, while all other components of $f^{-1}(U)$ are mapped conformally to U by f .

Proof. The function f is odd as it is the integral of an even function with $f(0) = 0$. Since the integrand is real and positive on the real axis, f is real and strictly increasing on \mathbb{R} . Since $\operatorname{erf}(x) \rightarrow \pm 1$ as $x \rightarrow \pm\infty$, the same is true for f . In particular, $f(\mathbb{R}) = (-1, 1)$, and -1 and 1 are asymptotic values of f .

The critical points of f are the zeros of \cosh , which are as stated in the lemma; they are simple since \cosh has only simple zeros. Furthermore, f is imaginary on the imaginary axis. Indeed, if $t \in \mathbb{R}$, then

$$(6.2) \quad \begin{aligned} f(it) &= \frac{2}{e^{1/4}\sqrt{\pi}} \int_0^{it} \cosh w \exp(-w^2) dw \\ &= \frac{2i}{e^{1/4}\sqrt{\pi}} \int_0^t \cos y \exp(y^2) dy =: \frac{2i}{e^{1/4}\sqrt{\pi}} \alpha(t). \end{aligned}$$

For $n \geq 0$, set $t_n := \frac{2n+1}{2}\pi$ and $\alpha_n := \alpha(t_n)$. Observe that $\alpha_0 > 0$. Since $\zeta_{-n} = -\zeta_{n-1}$ for all n , the following claim implies that all ζ_n are indeed pairwise distinct, and that $\operatorname{Im} \zeta_n$ is positive if and only if n is even.

Claim. The sequence $|\alpha_n|$ is strictly increasing and α_n is positive exactly for even n .

Proof. Note that $\alpha_{n+1} = \alpha_n + \beta_n$, where

$$\beta_n := \int_{\frac{2n+1}{2}\pi}^{\frac{2n+3}{2}\pi} \cos y \exp(y^2) dy.$$

Note that β_n is positive exactly when n is odd. We now estimate β_n from below in terms of α_n . Note that $|\alpha(t)| \leq \exp(t^2)$ for all t . A simple estimate shows that

$$|\beta_n| > \exp\left(\left(\frac{3n+2}{3}\right)^2 \pi^2\right) > 2 \exp\left(\left(\frac{2n+1}{2}\right)^2 \pi^2\right) \geq 2|\alpha(t_n)| = 2|\alpha_n|.$$

Hence $|\alpha_{n+1}| > |\alpha_n|$ for all n , and α_{n+1} has the same sign as β_n . This proves the claim. \triangle

Note that the order of f (see [Nev53, p. 219, ¶ 181]) is $\rho(f) = 2$. By the Denjoy-Carleman-Ahlfors theorem [Nev53, p. 313, ¶ 269], the number m of asymptotic values of f is finite. More precisely, $2m_{\text{direct}} + m_{\text{indirect}} \leq 4$, where m_{direct} and m_{indirect} are the numbers of *direct* and *indirect* singularities of f^{-1} , respectively, over finite asymptotic values. (Compare [Nev53, p. 289, ¶ 245], [BE95] or Section 8 for definitions.) As the set of critical values of f is discrete, every asymptotic value a of f is an isolated point of $S(f)$. Hence any singularity over a is logarithmic, and therefore direct. So $m = m_{\text{direct}} \leq 2$, and there are no asymptotic values except -1 and 1 , as claimed. \blacksquare

In summary, f has a structure similar to a vanilla function, apart from the two asymptotic values. Let $\text{CV}^+(f) = \{\zeta_n : n \geq 0\}$ denote the set of critical values corresponding to critical points with positive imaginary part, and let $\text{CV}^-(f) = \text{CV}(f) \setminus \text{CV}^+(f)$ denote the set of critical values corresponding to critical points with negative imaginary part.

Recall that we defined the notion of partial configurations only for vanilla functions; we shall next introduce a version of this notion specific to our function f .

6.2. Definition (Partial configuration).

A triple (D, U, V) of Jordan domains is called a *partial configuration* (for the function f from (6.1)) if all the following hold.

- (a) $[-1, 1] \subset D$.
- (b) $-1 \in U$ and $1 \in V$.
- (c) $\bar{U} \cap \bar{V} = \emptyset$ and $\bar{U} \cup \bar{V} \subset D$.
- (d) $D \cap \text{CV}(f) \subset U \cup V$ and $\partial D \cap S(f) = \emptyset$.
- (e) The component $\tilde{D} := \text{comp}_{\mathbb{R}}(f^{-1}(D))$ satisfies $f(\tilde{D} \cap \text{C}(f)) = D \cap \text{CV}(f)$.
- (f) Let \tilde{U} and \tilde{V} be the components of $f^{-1}(U)$ and $f^{-1}(V)$, respectively, that contain an infinite piece of the real axis. Then $\tilde{D} \cap \text{C}(f) \subset \tilde{U} \cup \tilde{V}$.

The mapping properties of f on a partial configuration are illustrated in Figure 4 and in the following analogue of Proposition 4.2.

6.3. Proposition (Structure of partial configurations).

Suppose that (D, U, V) is a partial configuration, and let γ be a crosscut of D that separates \bar{U} from \bar{V} . Then:

- (a) The boundary $\partial \tilde{D}$ consists of two injective curve tending to infinity in both directions: one in the upper half-plane (the upper boundary $\partial^+ \tilde{D}$) and one in the lower half-plane (the lower boundary $\partial^- \tilde{D}$).




FIGURE 4. An illustration of Proposition 6.3. The structure of the partial configuration (D, U, V) is the same as shown in the right-hand side of Figure 1, so we do not repeat it here. Shown is the preimage component \tilde{D} of D , where shades of grey indicate preimages of U and V , and white and black circles indicate preimages of w and z respectively. We stress again that this picture is topological rather than geometrically accurate. We also remind the reader that D has other preimage components under f ; these are not illustrated.

- (b) Each component of $\tilde{D} \setminus f^{-1}(\gamma)$ (which we shall call a face) contains either exactly one component of $f^{-1}(U)$ or exactly one component of $f^{-1}(V)$. A simple face is one that is mapped conformally by f .
- (c) The faces containing \tilde{U} and \tilde{V} are adjacent to each other, separated by a preimage $\tilde{\gamma}$ of $f^{-1}(\gamma)$ that connects the upper and lower boundaries of \tilde{D} .
- (d) Every component of $f^{-1}(U) \cap \tilde{D}$ apart from \tilde{U} is contained in a simple face adjacent to the one containing \tilde{V} . Similarly, each component of $f^{-1}(V) \cap \tilde{D}$ apart from \tilde{V} is contained in a simple face adjacent to the one containing \tilde{U} .

The components of $(f^{-1}(U) \setminus \tilde{U}) \cap \tilde{D}$ contained in a face whose boundary intersects the lower boundary of \tilde{D} will be labelled $(\tilde{U}_j^-)_{j=1}^\infty$, with \tilde{U}_1^- closest to $\tilde{\gamma}$ and

proceeding in positive orientation. We similarly define (\tilde{U}_j^+) , (\tilde{V}_j^-) , and (\tilde{V}_j^+) (see Figure 4).

Proof. Recall that all singular values of f in D lie in U or V . Let D_L be the component of $D \setminus \gamma$ containing U . Then $A := D_L \setminus \bar{U}$ is an annulus containing no singular values. Hence f is a covering map on every component of $f^{-1}(A)$. Thus each such component is either simply-connected and mapped as a universal covering, or doubly-connected and mapped as a finite covering of annuli. In particular, every component of $f^{-1}(A)$ is adjacent to exactly one component of $f^{-1}(U)$, and every component of $f^{-1}(D_L)$ contains exactly one component of $f^{-1}(U)$.

Recall that any component of $f^{-1}(U)$ different from \tilde{U} is mapped conformally by f , and hence contained in a simple face that is mapped conformally to D_L . The analogous statement holds for the preimages of D_R , so we see that there are exactly two non-simple faces, namely the face \tilde{D}_L containing \tilde{U} and the face \tilde{D}_R containing \tilde{V} . Since no simple face is adjacent to more than one other face, and \tilde{D} is connected, we see that \tilde{D}_L and \tilde{D}_R are adjacent, and that every simple face is adjacent to one of these two.

Let $\tilde{\gamma}$ be the component of $f^{-1}(\gamma)$ that separates \tilde{U} and \tilde{V} in \tilde{D} . Recall that $\mathbb{R} \subset \tilde{D}$, and that \tilde{U} (resp. \tilde{V}) contain an infinite piece of the negative (resp. positive) real axis. It follows that $\tilde{\gamma}$ must have one endpoint in the upper half-plane and one endpoint in the lower half-plane.

It remains to establish (a). Since ∂D contains no singular values and \tilde{D} is unbounded, every component of $\partial \tilde{D}$ is an injective curve tending to infinity in both directions, and f maps this component to ∂D as a universal covering. We must show that there are only two such components. To do so, we claim that there is exactly one component of $f^{-1}(A)$ adjacent to \tilde{U} (and similarly for \tilde{V}). In other words, $\partial \tilde{U}$ and $\partial \tilde{V}$ are connected.

Let U' be obtained from U by removing, for each critical value $c \in U$, an arc δ_c connecting c to ∂U , in such a way that these arcs are pairwise disjoint. Let \tilde{U}' be the connected component of $f^{-1}(U')$ containing an infinite piece of the real axis. By Lemma 6.1, $f: \tilde{U}' \rightarrow U'$ is a universal covering, and every other component of $f^{-1}(U')$ is mapped conformally. Since \tilde{U} contains only finitely many critical points, it follows that there are only finitely many preimage components of U' in \tilde{U} . The claim follows easily, as does the fact that $\partial \tilde{D}$ has only two connected components. \blacksquare

The central part of our construction is a method for extending partial configurations, in the same spirit as Proposition 4.3.

6.4. Proposition (Extending partial configurations).

Suppose that (D, U, V) is a partial configuration, and let P be a connected component of $f^{-1}(U) \cap \tilde{D}$. Then there is a partial configuration (D', U', V) such that $D \subset D'$, $U \subset U'$ and $P \subset \tilde{U}'$.

Moreover, for any $R > 0$, this configuration can be chosen such that $B(0, R) \subset D'$.

Proof. We first prove the proposition without the final statement. We may assume that $\tilde{P} \neq \tilde{U}$ (otherwise, there is nothing to prove). Let γ be as in Proposition 6.3 and recall that P is contained in a simple face adjacent to the face \tilde{D}_R containing \tilde{V} . The boundary of this simple face consists of a preimage of γ and a piece of $\partial \tilde{D}$; we may assume without

loss of generality that this piece belongs to the lower boundary of \tilde{D} . (Otherwise, since $f(\bar{z}) = \overline{f(z)}$, for $z \in \mathbb{C}$, we can replace D , U and V by their reflections in the real axis.)

Let $p \geq 1$ be such that the $P = \tilde{U}_p^-$. (In Figure 5, $p = 2$.) We now extend D to a Jordan domain $D' \supset \bar{D}$ such that:

- $\partial D' \cap \text{CV}(f) = \emptyset$;
- $\#((D' \setminus D) \cap \text{CV}^+(f)) = p$;
- $\#((D' \setminus D) \cap \text{CV}^-(f)) = 0$;
- $f(\tilde{D}' \cap \text{C}(f)) = D' \cap \text{CV}(f)$, where $\tilde{D}' = \text{comp}_{\mathbb{R}}(f^{-1}(D'))$.

This can be achieved by the same technique as in the proof of Proposition 3.3: First extend D to D_1 such that $\tilde{D}_1 := \text{comp}_{\mathbb{R}}(f^{-1}(D_1))$ contains at least p points of $\text{C}(f) \setminus \tilde{D}$ whose images have positive imaginary part. Then remove all critical values whose critical preimages are not in \tilde{D}_1 as in Proposition 3.3. We can also remove arcs connecting the points of $\text{CV}^-(f) \cap D_1$ to ∂D_1 without intersecting D . Observe that the critical preimage of any such arc lies in the lower half-plane, and hence does not separate $\tilde{D} \supset \mathbb{R}$ from any of the critical points in the upper half-plane. Finally, remove any excess critical values in $\text{CV}^+(f)$ exactly as in Proposition 3.3.

It now remains to extend U' to U . Let w be the endpoint of γ in the upper half-plane, and z its endpoint in the lower half-plane. As in the proof of Proposition 4.3, we construct an arc α joining w to $\partial D'$ so that all the critical values in $D' \setminus D$ lie on α , and such that one connected component $\tilde{\alpha}$ of $f^{-1}(\alpha)$ contains all points of $\text{C}(f) \cap \tilde{D}' \setminus \tilde{D}$. Then we let β be the sub-arc of α starting at w and ending at the last critical value on α .

Consider an arc Γ , not intersecting $\bar{V} \cup \text{CV}(f)$ and not intersecting \bar{U} except in one endpoint, defined as follows. The arc starts at some point $\mu \in \partial U$, and runs around β and \bar{V} exactly p times in negative orientation. On its last loop it enters D at the point z , traversing along the arc γ and ending at w .

Take a thin channel W containing the Jordan arc $\Gamma \cup \beta$, in such a way that $U' := U \cup W$ is simply-connected. By an argument similar to that in the proof of Proposition 4.3 it can be shown that \tilde{U}' contains both \tilde{U} and \tilde{U}_p^- ; see Figure 5. This completes the construction.

Now let us prove the final claim of the proposition. As in Proposition 3.3, we can find $D'' \supset D'$ such that $\tilde{D}'' := \text{comp}_{\mathbb{R}}(f^{-1}(D''))$ contains all critical preimages of points in $\text{CV}(f) \cap D''$, and such that furthermore $B(0, R) \subset D''$. We can find an arc $\alpha^+ \subset D'' \setminus D'$, having one endpoint on $\partial D'$ and another in $D'' \setminus D'$, and passing through all points of $(D'' \setminus D') \cap \text{CV}^+(f)$, with the following property: there is a connected component of $f^{-1}(\alpha^+)$ that intersects $\partial \tilde{D}'$ and contains all critical preimages of the critical values in α^+ . We can find a similar curve α^- (disjoint from α^+) for the critical values in $(D'' \setminus D') \cap \text{CV}^-(f)$. By extending U' appropriately to include these two arcs, we easily obtain $U'' \supset U'$ such that $\tilde{U}'' := \text{comp}_{\tilde{U}'}(f^{-1}(U''))$ contains $\text{C}(f) \cap (\tilde{D}'' \setminus \tilde{D}')$. We leave the details to the reader. \blacksquare

6.5. Remark (Symmetry).

Recall that f is odd. Hence, if (D, U, V) is a partial configuration for f , then so is




FIGURE 5. The extended preimage in Proposition 6.3, in the case where $p = 2$. The image configuration is the same as the right-hand side of Figure 3. The preimage of the thin channel W starts at \tilde{U} , passes through two preimages of D , meets a critical point (shown as a square), and ends on \tilde{U}_2^- . Note that other preimages of critical values, shown in Figure 3, are not shown here, for simplicity.

$(-D, -V, -U)$. It follows that Proposition 6.4 holds also with the roles of U and V exchanged.

Applying the preceding step inductively, we obtain the following fact, which easily implies Theorem 1.8.

6.6. Proposition (Sequence of configurations).

There is a sequence $(D_k, U_k, V_k)_{k=1}^\infty$ of partial configurations such that (D_k) , (U_k) and (V_k) are increasing sequences, and such that every component of $f^{-1}(U_1)$ is contained in \tilde{U}_k for sufficiently large k , and similarly for the components of $f^{-1}(V_1)$.

Proof. Let U_1 be a small disc containing -1 , let V_1 be a small disc containing 1 , and let D_1 be a simply-connected neighbourhood of $[-1, 1]$ containing $U_1 \cup V_1$ and such that $D_1 \cap \text{CV}(f) = \emptyset$. It is easy to see that (D_1, U_1, V_1) is a partial configuration.

Let $(k(\ell), j(\ell), \sigma(\ell))_{\ell=1}^{\infty}$ be an enumeration of the countable set $\mathbb{N} \times \mathbb{N} \times \{-, +\}$. We may assume that $k(\ell) \leq \ell$ for all ℓ . We now describe how we construct (D_k, U_k, V_k) inductively from $(D_{k-1}, U_{k-1}, V_{k-1})$, for $k \geq 2$.

First suppose that k is even, say $k = 2\ell$ with $\ell \geq 1$. Set $V_k := V_{k-1}$. Let P be the connected component of $f^{-1}(U_{k-1})$ that contains $(\tilde{U}_{k(\ell)})_{j(\ell)}^{\sigma(\ell)}$. (Recall the notation for the connected components of $f^{-1}(U_{k(\ell)}) \cap \tilde{D}_{k(\ell)}$ from Proposition 6.3.) We apply Proposition 6.4 to obtain a partial configuration (D_k, U_k, V_k) such that $B(0, k) \subset D_k$ and

$$(\tilde{U}_{k(\ell)})_{j(\ell)}^{\sigma(\ell)} \subset \tilde{U}_k.$$

Now suppose that k is odd, say $k = 2\ell + 1$. Using Remark 6.5, we proceed exactly as in the previous step, but with the roles of U and V exchanged. So we obtain a partial configuration (D_k, U_k, V_k) such that $B(0, k) \subset D_k$ and

$$(\tilde{V}_{k(\ell)})_{j(\ell)}^{\sigma(\ell)} \subset \tilde{V}_k.$$

This completes the inductive construction. Let P be any connected component of $f^{-1}(U_1)$. We must show that $P \subset \tilde{U}_k$ for sufficiently large k .

By construction, $P \cap \tilde{D}_{k_0} \neq \emptyset$ for some k_0 . Since $P \cap \partial \tilde{D}_{k_0} = \emptyset$, in fact $P \subset \tilde{D}_{k_0}$. If $P \subset \tilde{U}_{k_0}$, then we are done. Otherwise, $P \subset (\tilde{U}_{k_0})_j^{\sigma}$ for some $j \geq 1$ and $\sigma \in \{-, +\}$. Let ℓ be such that $k(\ell) = k_0$, $j(\ell) = j$ and $\sigma(\ell) = \sigma$; recall that $\ell \geq k_0$. So, by construction,

$$P \subset (\tilde{U}_{k(\ell)})_{j(\ell)}^{\sigma(\ell)} \subset \tilde{U}_{2\ell} \subset \tilde{U}_k$$

for $k \geq 2\ell$. The same argument applies to preimage components of $f^{-1}(V_1)$, and the proof is complete. \blacksquare

Proof of Theorem 1.8. Set

$$U := \bigcup_{k \in \mathbb{N}} U_k \quad \text{and} \quad V := \bigcup_{k \in \mathbb{N}} V_k.$$

The result follows in exactly the same way as in the proof of Theorem 1.11. \blacksquare

6.7. Remark (More asymptotic values).

For any $2 \leq d \leq \infty$, it should be possible to use a similar construction to obtain an entire function f having d logarithmic asymptotic values $(a_j)_{j=1}^d$, and pairwise disjoint simply-connected domains $(U_j)_{j=1}^d$ with $a_j \in U_j$, such that $f^{-1}(U_j)$ is connected for all j .

7. THE ERROR IN BAKER'S PROOF

In this section we briefly outline the proof in [Bak70], and highlight where the error occurs. As mentioned earlier, the proof in [Bak70] amounts to a positive answer to Question 1.1, so we suppose that f is a transcendental entire function, and that G_1 and G_2 are disjoint simply-connected domains each with connected preimage. Baker attempts to deduce a contradiction from this.

Baker uses a well-known result, known as the Gross star theorem [Nev53, p. 292, ¶ 247], which we also use later.




FIGURE 6. This figure displays the idea in the proof in [Bak70]. The left-hand side shows the domain of the function, and the right-hand side shows the image. The domains D, G_1 and G_2 are shaded. The images $f(\beta_1)$ and $f(\beta_2)$ are shown as dotted lines.

7.1. Theorem (Gross star theorem).

Suppose that f is a transcendental entire function and that φ is a holomorphic branch of the inverse of f defined in a neighbourhood of a point w . Then, for almost all $\vartheta \in [0, 2\pi)$, the branch φ can be continued analytically along the ray $\{w + t \cdot e^{i\vartheta} : t \geq 0\}$.

Baker begins by choosing a point $z_1 \in G_1$, with simple preimages p_1 and q_1 . Let φ_p and φ_q be the branches of f^{-1} taking z_1 to p_1 and q_1 , respectively. By Theorem 7.1, there is a line segment γ joining z_1 to a point $z_2 \in G_2$ such that both branches can be continued analytically along γ . Then $\gamma_p := \varphi_p(\gamma)$ is an arc joining p_1 to a point $p_2 \in f^{-1}(z_2)$, and likewise $\gamma_q := \varphi_q(\gamma)$ joins q_1 to a different preimage $q_2 \in f^{-1}(z_2)$. For each $k \in \{1, 2\}$, we know that $f^{-1}(G_k)$ is connected by assumption, and so we can let $\beta_k \subset f^{-1}(G_k)$ be an arc joining p_k and q_k .

The curves β_1 and β_2 may intersect the arcs γ_p and γ_q in some interior points. Baker notes that by taking suitable subcurves, γ'_p of γ_p , γ'_q of γ_q , β'_1 of β_1 , and β'_2 of β_2 , there is a bounded quadrilateral D with boundary $\beta'_1 \cup \gamma'_p \cup \beta'_2 \cup \gamma'_q$; see the left-hand side of Figure 6.




FIGURE 7. This figure, which is based on one first drawn by Duval, displays the error in the proof in [Bak70]. The left-hand side shows the domain of the function, and the right-hand side shows the image. The domains D, G_1 and G_2 are shaded. The images $f(\beta_1)$ and $f(\beta_2)$ are shown as dotted lines. The boundary of the bounded region $f(D)$ is shown in solid, and is made up of part of γ together with part of $f(\beta_2)$.

Since G_1 and G_2 are simply-connected, the curves $f(\beta_1)$ and $f(\beta_2)$ cannot surround z_2 and z_1 , respectively. Observe that

$$\partial f(D) \subset f(\partial D) \subset f(\beta_1) \cup \gamma \cup f(\beta_2).$$

Baker's claim is that an entire function cannot map a quadrilateral in this manner, since otherwise $f(D)$ must be unbounded, which is impossible for bounded D ; see Figure 6.

However, as pointed out by Duval (personal communication), it is topologically quite possible for $f(D)$ to be bounded, in the absence of additional assumptions. This can be seen in the right-hand side of Figure 7.

It is interesting to note a key feature of Figure 7: although G_2 is simply-connected, it still “loops around” z_1 sufficiently far as to intersect γ again. It is this “looping” which leads to the boundedness of $f(D)$. By examining Figure 7, it can be seen this would not occur if we chose the point $z_2 \in \gamma \cap G_2$ in the other component of $\gamma \cap G_2$. This explains how the intricate “looping” behaviour arises in the proof in Proposition 4.3.




FIGURE 8. This figure, which is complementary to Figure 7, illustrates a second configuration that leads to Baker’s original proof breaking down.

Indeed, consider the case of that proposition where $n = 1$ and $m_V = 1$. In this case, our construction leads exactly to the structure from Figure 7.

We remark that there is another way in which Baker’s argument can fail, where instead of one of the domains looping entirely around the other, both partly loop around the curve γ ; see Figure 8. The following proposition shows that Baker’s method of proof does apply whenever neither kind of looping occurs. We shall use it in the next section.

7.2. Proposition (Baker’s argument).

Suppose that f is analytic in a simply-connected domain U . Suppose that $\gamma_p, \gamma_q \subset U$ are disjoint arcs on both of which f is injective, and with $\gamma := f(\gamma_p) = f(\gamma_q)$. Let p_1 and p_2 be the endpoints of γ_p , and let q_1 and q_2 be the endpoints of γ_q . We may choose the labelling such that $z_1 := f(p_1) = f(q_1)$ and $z_2 := f(p_2) = f(q_2)$.

Let $\beta_1, \beta_2 \subset U$ be two arcs such that β_k joins p_k and q_k , and such that $C_1 := f(\beta_1)$ and $C_2 := f(\beta_2)$ are disjoint.

Then every point $\zeta \in \gamma \setminus (C_1 \cup C_2)$ is on the boundary of a bounded connected component of $\mathbb{C} \setminus (C_1 \cup C_2 \cup \gamma)$.

Proof. Let $\zeta \in \gamma \setminus (C_1 \cup C_2)$, and let ζ' be the preimage of ζ on γ_p . Since $\zeta' \notin (\beta_1 \cup \beta_2)$, there is a bounded connected component $D \subset U$ of $\mathbb{C} \setminus (\gamma_p \cup \gamma_q \cup \beta_1 \cup \beta_2)$ with $\zeta' \in \partial D$.

Then $\partial f(D) \subset f(\partial D) \subset C_1 \cup C_2 \cup \gamma$. Let Δ be a disc around ζ' chosen small enough that $f(\Delta)$ does not intersect $C_1 \cup C_2$, and let V be the connected component of $f(D \cap \Delta)$ with $\zeta \in \partial V$. Since $V \subset f(D)$, it follows that V is contained in a bounded component of $\mathbb{C} \setminus (C_1 \cup C_2 \cup \gamma)$, as claimed. \blacksquare

7.3. Remark (Winding behaviour).

In order to illustrate how the conclusion of the proposition reflects the “looping” behaviour mentioned above (and illustrated in Figures 7 and 8), we remark that it can be reformulated as follows. Suppose that $\zeta \in \gamma \setminus (C_1 \cup C_2)$. Then there is $k \in \{1, 2\}$ such that either C_k intersects γ on both sides of ζ , or $C_k \cup \gamma$ contains a Jordan curve surrounding ζ .

Indeed, the reformulation clearly implies the conclusion as stated. Conversely, suppose that ζ is on the boundary of such a bounded component V of $\mathbb{C} \setminus (C_1 \cup C_2 \cup \gamma)$. We may suppose that C_1 and C_2 intersect γ on different sides of ζ , and that neither curve surrounds ζ , as otherwise there is nothing to show. Let γ' be the connected component of $\gamma \setminus (C_1 \cup C_2)$ containing ζ ; then γ' is a cross-cut of the unbounded connected component W of $\mathbb{C} \setminus (C_1 \cup C_2)$.

There must be a second piece γ'' of γ such that $\gamma' \cup \gamma''$ separates V from ∞ in W . By assumption, there is $k \in \{1, 2\}$ such that both endpoints of γ'' belong to C_k . Hence $C_k \cup \gamma''$ separates ζ from infinity, and the claim follows by choosing a suitable non-intersecting sub-curve.

Note that the conclusion of Proposition 7.2 can be strengthened as follows when the curves β_k and γ_j bound a quadrilateral. (We do not require this fact in the remainder of the paper.)

7.4. Remark (The case of a quadrilateral).

Suppose that, under the hypotheses of Proposition 7.2, additionally the arcs $\gamma_p, \gamma_q, \beta_1$ and β_2 intersect only in their endpoints, and hence bound a quadrilateral Q . Then $f(Q) \cup \text{int}(\gamma)$ is a neighbourhood of $\text{int}(\gamma)$, and hence no point of γ can be connected to infinity without intersecting either C_1 or C_2 . It follows that, in this case, one of the curves C_1 and C_2 must surround the other.

8. PROOF OF THEOREM 1.3

We begin with the following fact, which will be used to deduce Theorem 1.3(a).

8.1. Proposition (Compact full sets with connected preimages).

Let f be a transcendental entire function, and let K be a compact connected set, containing more than one point, such that $f^{-1}(K)$ is connected.

Then $S(f) \cap W = \emptyset$, where W is the unbounded connected component of $\mathbb{C} \setminus K$.

To prove Proposition 8.1, we use the following fact: Near any singular value, there is an inverse branch φ and a simple closed curve along which φ can be continued in such a way as to obtain a different inverse branch at the same point. More precisely:

8.2. Lemma (Inverse branches near a singular value).

Let f be a transcendental entire function, and let $s \in \mathbb{C}$. Then the following are equivalent.

- (a) $s \in S(f)$;
- (b) for every neighbourhood D of s there is a polygonal arc $\beta: [0, 1] \rightarrow \mathbb{C}$, not passing through any critical points of f , such that $f(\beta) \subset D$, $f(\beta(0)) = f(\beta(1))$, and such that $f \circ \beta$ is injective on $[0, 1)$.

Proof. Clearly (b) implies (a). So suppose that s is a singular value, and that D is an open disc centred at s . If D contains a critical value c of f , then there is a point $\tilde{c} \in \mathbb{C}$ and $d \geq 2$ such that f maps like $z \rightarrow z^d$ in a neighbourhood of \tilde{c} . The claim follows easily.

So we may suppose that D contains no critical value. Let \tilde{D} be a connected component of $f^{-1}(D)$ that is not mapped one-to-one by f . Since D contains no critical values, it follows that \tilde{D} is unbounded and $f^{-1}(w) \cap \tilde{D}$ is infinite for all $w \in D$, with at most one exception. (This follows from a result of Heins [Hei57, Theorem 4']; see [BFR15, Proposition 2.8].) In particular, there is $w \in D$ having two different (simple) preimages $\zeta_1, \zeta_2 \in \tilde{D}$. Let $\beta' \subset \tilde{D}$ be a polygonal arc from ζ_1 to ζ_2 . Then $f(\beta')$ is a closed curve, but may be self-intersecting. However, since f is locally injective on β' , it follows that β' contains a sub-arc β with the required properties. \blacksquare

We also need to observe that increasing a set with connected preimage to a larger domain does not change this property.

8.3. Lemma (Increasing sets with connected preimage).

Suppose that $A \subset \mathbb{C}$ is a connected set containing more than one point, such that $f^{-1}(A)$ is connected. Then $f^{-1}(G)$ is connected for every domain G containing A .

Proof. Let U be a connected component of $f^{-1}(G)$. Again using [Hei57, Theorem 4'], every point in G , with at most one exception, has at least one preimage in U . In particular, U intersects, and hence contains $f^{-1}(A)$. Thus there can be at most one such component. \blacksquare

Proof of Proposition 8.1. Suppose, by way of contradiction, that $S(f) \cap W \neq \emptyset$. By the Riemann mapping theorem, applied to $W \cup \{\infty\}$, there is a Jordan domain $G \supset K$, bounded by an analytic curve α , such that $S(f) \not\subset \overline{G}$. By Lemma 8.3, $f^{-1}(G)$ is connected.

Let $B(w, \varepsilon) \subset \mathbb{C} \setminus \overline{G}$ be a ball intersecting $S(f)$. By Lemma 8.2, there is a polygonal arc β_1 , joining points $p_1, q_1 \in \mathbb{C}$ with $z_1 := f(p_1) = f(q_1)$, such that f is injective on $\beta_1 \setminus \{q_1\}$, and such that $C_1 := f(\beta_1) \subset B(w, \varepsilon)$ is a Jordan curve through z_1 .

Using the Gross star theorem 7.1, we construct an arc γ that joins z_1 to a point $z_2 \in G$, along which the inverse branches φ_p and φ_q of f that map z_1 to p_1 and q_1 , respectively, can both be analytically continued. We can choose γ such that $\gamma \cap C_1 = \{z_1\}$ and such that $\gamma \cap \alpha$ consists of a single point, ζ .

We now have two preimages of z_2 , corresponding to the analytic continuation of φ_p and of φ_q , which we label p_2 and q_2 . These are joined to p_1 and q_1 , respectively, by disjoint preimage components γ_p and γ_q of γ . Also join p_2 and q_2 by an arc $\beta_2 \subset f^{-1}(G)$.

By construction, the piece of γ connecting z_1 to ζ is contained in the unbounded connected component of $\mathbb{C} \setminus C_1 \cup \alpha$. As α surrounds C_2 by definition, no point on this arc is on the boundary of a bounded connected component of $\mathbb{C} \setminus (C_1 \cup C_2 \cup \gamma)$. This contradicts Proposition 7.2. ■

We next show how to deduce Theorem 1.3(b) from Proposition 8.1, using the following simple fact.

8.4. Proposition (Shrinking a simply-connected domain).

Suppose that f is a transcendental entire function, and that D is a simply-connected domain such that $S(f) \cap D$ is compact, and let \tilde{D} be a connected component of $f^{-1}(D)$.

If $U \subset D$ is simply-connected with $S(f) \cap U = S(f) \cap D$, then $f^{-1}(U) \cap \tilde{D}$ is connected.

Proof. This is shown in the second paragraph of [BFR15, Proof of Proposition 2.9]. ■

We thus have the following strengthening of Theorem 1.3(b).

8.5. Corollary (Domains with compact intersection with $S(f)$).

Suppose that f is a transcendental entire function, and that G is a simply-connected domain such that $G \cap S(f)$ is compact. Then $f^{-1}(G)$ is connected if and only if $S(f) \subset G$.

Proof of Corollary 8.5. The “if” direction is immediate from Proposition 8.4, taking $D = \mathbb{C}$ and $U = G$. (See also [BFR15, Proposition 2.9 (2)].)

For the “only if” direction, suppose that $f^{-1}(G)$ is connected. Let $D \subset G$ be a bounded Jordan domain with $S(f) \cap G \subset D$ and such that $K := \overline{D} \subset G$. Then $f^{-1}(D)$ is connected by Proposition 8.4, as is $f^{-1}(K) = \overline{f^{-1}(D)}$. It follows from Proposition 8.1 that $S(f) \subset K \subset G$, as required. ■

We next record a strengthening of Theorem 1.3 (f) that concerns the accessibility of *direct asymptotic values* on ∂G_1 from G_1 . Here $a \in \hat{\mathbb{C}}$ is a direct asymptotic value of f if there is an open connected neighbourhood Δ of a in $\hat{\mathbb{C}}$ and a connected component $\tilde{\Delta}$ of $f^{-1}(\Delta)$ such that $a \notin f(\tilde{\Delta})$. Observe that every Picard exceptional value (i.e., a value a for which $f^{-1}(a)$ is finite) is direct; in particular, ∞ is always a direct asymptotic value of f .

8.6. Proposition (Accessible asymptotic values on boundaries).

Let f be a transcendental entire function and let $\xi \in \hat{\mathbb{C}}$ be a direct asymptotic value of f . Suppose that G_1, G_2 are disjoint simply-connected domains such that $f^{-1}(G_1)$ is connected. If ξ is an accessible boundary point of G_1 , then $f^{-1}(G_2)$ is disconnected.

Proof. Let Δ and $\tilde{\Delta}$ be as in the definition of a direct asymptotic value. If Δ is chosen sufficiently small, then every value of $\Delta \setminus \{a\}$ has infinitely many preimages in Δ (this follows once more from [Hei57, Theorem 4]). By assumption, $G_1 \cap \Delta$ has a connected component Δ_1 from which a is accessible. Choose a point $z_1 \in \Delta_1$ which is not a critical value. Then z_1 has infinitely many simple preimages in $\tilde{\Delta}$, each corresponding to a different branch of f^{-1} . Using Theorem 7.1, we construct a polygonal arc $\gamma \subset \Delta_1$ from z_1 to ξ , possibly with infinitely many pieces, such that these inverse branches can all be continued along $\gamma \setminus \{\xi\}$.

Let $p \in \tilde{\Delta} \cap f^{-1}(z_1)$, and let $\gamma_p \subset \tilde{\Delta}$ be the component of $f^{-1}(\gamma)$ containing p . Since ξ has no preimages in U , the arc γ_p connects p to ∞ . Pick a second preimage $q \neq p$ of z_1 in $\tilde{\Delta}$, and define γ_q analogously. We also join p and q by an arc τ in $f^{-1}(G_1)$ that does not intersect $\gamma_p \cup \gamma_q$. Note that $\gamma_p \cup \tau \cup \gamma_q \subset f^{-1}(G_1)$ separates the plane into two complementary components; we claim that each such component D intersects $f^{-1}(G_2)$. This implies that $f^{-1}(G_2)$ is disconnected, as required.

Let B denote the union of $f(\tau)$ and its bounded complementary components. Then B is a compact subset of G_1 . Let $w \in \gamma$ be a point so close to ξ that B does not intersect the arc of γ connecting w to ξ . Then w has a preimage on γ_p ; say \tilde{w} . Let $W \subset G_1$ be a small neighbourhood of w disjoint from B , and let \tilde{W} be the component of the preimage of W containing \tilde{w} . We may choose $\zeta \in D \cap \tilde{W}$ such that ζ is not a critical point and $f(\zeta) \notin \gamma$. By choice of w , and since G_1 is simply-connected, the set $B \cup \gamma$ does not separate $f(\zeta)$ from G_2 .

Using the Gross star theorem 7.1, we find a polygonal curve $\Gamma \subset \mathbb{C} \setminus (B \cup \gamma)$ from $f(\zeta)$ to a point of G_2 in such a way that the inverse branch of f that sends $f(\zeta)$ to ζ can be continued along Γ . Then the preimage of Γ is a curve in D that ends at a point of $f^{-1}(G_2)$. This completes the proof. \blacksquare

Proof of Theorem 1.3. Let f be a transcendental entire function, and let G_1, G_2 be disjoint simply-connected domains such that $f^{-1}(G_1)$ is connected. In turn, we shall conclude from each assumption in Theorem 1.3 that $f^{-1}(G_2)$ is disconnected.

We begin with (a), so suppose that G_1 is bounded and $\overline{G_1}$ does not separate G_2 from infinity. Set $K := \overline{G_1}$ and let W be the unbounded connected component of $\mathbb{C} \setminus K$. Then, by assumption, $f^{-1}(K) = \overline{f^{-1}(G_1)}$ is connected and $G_2 \subset W$. By Proposition 8.1, $G_2 \cap S(f) = \emptyset$. Hence f is univalent on every component of $f^{-1}(G_2)$, and in particular $f^{-1}(G_2)$ is disconnected.

Next, we turn to (b). If $G_1 \cap S(f)$ is compactly contained in G_1 and $f^{-1}(G_1)$ is connected, then by Corollary 8.5, $S(f) \subset G_1$. It follows as above that $f^{-1}(G_2)$ is disconnected. Clearly (c) is a direct consequence of (b).

Now suppose that (d) holds. We use a technique similar to the proof of part (iii) of the theorem of [BE08b]. By assumption, $f^{-1}(G_1)$ contains a Jordan curve Γ , unbounded in both directions, such that f tends to an asymptotic value $a_1 \in \hat{\mathbb{C}}$ in one direction, and to a (possibly different) asymptotic value $a_2 \in \hat{\mathbb{C}}$ in the other. Since both ends of Γ represent different transcendental singularities, there is $\varepsilon > 0$ such that Γ has unbounded intersection with two different components of $f^{-1}(\{z \in \mathbb{C} : \text{dist}^\#(z, \{a_1, a_2\}) < \varepsilon\})$. (Here $\text{dist}^\#$ denotes spherical distance.)

The Jordan curve Γ divides the plane into two components. As in the proof of Proposition 8.6, we claim that each such component D must meet $f^{-1}(G_2)$, which is therefore disconnected. Otherwise, choose a point $w \in G_2$, and consider the function

$$g(z) = \frac{1}{f(z) - w}.$$

Then g is bounded and holomorphic in D and has finite limits as z tends to infinity along Γ . By a theorem of Lindelöf [Lin15, p. 9, ¶ 5], we have $a_1 = a_2$ and $f(z) \rightarrow a_1$ as

$z \rightarrow \infty$ in D . (See also ¶¶ 36, 39 and 61 in [Nev53, Chapter III].) This is a contradiction, as $D \cap f^{-1}(\{z \in \mathbb{C} : \text{dist}^\#(z, \{a_1, a_2\}) = \epsilon\})$ has an unbounded connected component.

We now turn to (e). By way of contradiction, suppose that $f^{-1}(G_2)$ is connected and ζ is accessible from both G_1 and G_2 . We can assume, by Proposition 8.6, that ζ is not a Picard exceptional point. Let ζ_p and ζ_q be two distinct preimages of ζ , and let Δ be a disc around ζ . If Δ is sufficiently small, then $\partial\Delta$ meets both G_1 and G_2 , and for each $j \in \{p, q\}$ the component Δ_j of $f^{-1}(\Delta)$ that contains ζ_j has all the following properties;

- Δ_j is bounded and does not meet the other preimage component;
- $\overline{\Delta_j}$ is mapped to $\overline{\Delta}$ as a degree d branched covering, where d is the local degree of f at ζ_j ;
- in particular, Δ_j contains no critical points of f except possibly ζ_j , and also $\Delta_j \cap f^{-1}(\zeta) = \{\zeta_j\}$.

For $j \in \{1, 2\}$, let γ_j be an arc connecting ζ to a point of $G_j \cap \partial\Delta$ with the property that $\text{int}(\gamma_j) \subset \Delta \cap G_j$. Set $\gamma = \gamma_1 \cup \gamma_2$. Let z_1 and z_2 be the two endpoints of γ , with $z_j \in G_j \cap \partial\Delta$.

For each $j \in \{p, q\}$, let $\tilde{\gamma}_j$ be the component of $f^{-1}(\gamma)$ containing ζ_j . Then $\tilde{\gamma}_j$ is either an arc (corresponding to the case where ζ_j is not a critical point), or a tree with just one vertex of order greater than one (at ζ_j). In either case, there is an arc $\gamma_j \subset \tilde{\gamma}_j$ such that $f: \gamma_j \rightarrow \gamma$ is a homeomorphism. We let p_k ($k \in \{1, 2\}$) be the point on γ_p whose image is z_k . We similarly let q_k be the point on γ_q whose image is z_k .

For $k \in \{1, 2\}$, we can join p_k and q_k by an arc $\beta_k \subset f^{-1}(G_k)$. Set $C_k := f(\beta_k)$ and consider the set $A_k := C_k \cup \gamma_k \subset G_k \cup \{\zeta\}$. Since G_k is simply-connected, every bounded connected component of $\mathbb{C} \setminus A_k$ is contained in G_k , and does not contain ζ on its boundary. Furthermore, $A_1 \cap A_2 = \{\zeta\}$ is connected, so by Janiszewski's theorem there are no other bounded connected complementary components of $A_1 \cup A_2 = C_1 \cup C_2 \cup \gamma$. This is impossible by Proposition 7.2. We have obtained the desired contradiction.

Observe that (f) is an immediate consequence of Proposition 8.6.

Now assume that (g) holds; i.e. that $\overline{G_1} \cap \overline{G_2} = \emptyset$, and suppose by way of contradiction that $f^{-1}(G_2)$ is connected. It follows by [Why42, Theorem VI.3.1] that there is a Jordan curve Γ , which may pass through ∞ , that separates the plane into two components, one containing $\overline{G_1}$ and the other containing $\overline{G_2}$.

For $j \in \{1, 2\}$, let V_j be the connected component of $\mathbb{C} \setminus \Gamma$ containing G_j . Then $f^{-1}(V_j)$ is connected by Lemma 8.3. If Γ is bounded, then one of the V_j , say V_1 , is bounded. Clearly $\overline{V_1} = V_1 \cup \Gamma$ does not separate V_2 from infinity, contradicting (a).

On the other hand, if Γ is unbounded, then both V_j are simply-connected and every point of $\partial V_1 = \partial V_2 = \Gamma$ is accessible from both V_1 and V_2 . This contradicts (e).

Finally, as noted earlier, (h) is an immediate consequence of [BE08a, Theorem 1]. ■

9. APPENDIX

In this final part of the paper we list, for the convenience of the research community, various papers whose results are impacted by the problem found. Note that this list is not necessarily complete; some of the papers in this list are very heavily cited, as they include other influential results that are not in doubt, and so it is difficult to be sure that every issue has been identified.

Firstly, we have identified six papers that have a common flaw in a proof, and so certain results in these papers must be considered *open*. These are as follows:

- (i) The result of [Bak70] (this paper contains only one result).
- (ii) The proof of [Bak75, Theorem 2]. Note that the other results of [Bak75] are not affected by the problem.
- (iii) The proof of [EL92, Lemma 11]. No other results of [EL92] are affected by the problem.
- (iv) The proof of [Dom98, Theorem K]. No other results of [Dom98] are affected by the problem.
- (v) The proof of [Lin08, Lemma 2.2], and hence [Lin08, Theorem 2.1], which depends on this lemma.
- (vi) The proofs of [CW03, Theorem 3 and Theorem 4].

Most papers that use these theorems are dynamical, and so our constructions give no additional information on whether their results are correct. We have identified just one explicitly stated result that is now seen to be false. This is [BE08a, Theorem 2], which states that if f is an entire function of finite order, and $a \in \mathbb{C}$ is either a critical value or a locally omitted value, then any simply-connected region that does not contain a has disconnected preimage. Theorem 1.2 can be seen to be a counter-example to this assertion. No other papers use [BE08a, Theorem 2].

We mention also that the preprint [DS18] attempts to give a “complementary proof” of Baker’s original result regarding completely invariant Fatou components. Unfortunately, the proof does not use the dynamics of the function in an essential way. Therefore it also runs afoul of our construction, and would effectively contradict Theorem 1.2.

We end with three lists of results that use the potentially flawed theorems. The first list contains those results that use the potentially flawed theorems, but can be seen nonetheless to hold.

- (i) The proof of [BD00, Theorem H] uses [EL92, Lemma 11]. This result applies only to the class \mathcal{S} , and so in fact the result can be recovered by using Theorem 1.3(c) instead.
- (ii) The proof of [BDH01, Theorem B] uses [Bak70]. This result can also be recovered by using Theorem 1.3(c) instead.
- (iii) The proof of [RS11, Theorem 1.2] uses the result of [Bak70]. However, this dependence can easily be shown not to be essential (since a slight adaptation of the proof gives the result for f^p in the case that there is a completely invariant p -cycle of Fatou components), and so [RS11, Theorem 1.2] remains valid.
- (iv) The proof of [Os13, Lemma 3.3] uses the result of [Bak70], and this lemma is then used to prove [Os13, Theorem 3.1]. However, it is easy to see how to weaken the statements of [Os13, Lemma 3.3 and Theorem 3.1] in such a way that [Bak70] is no longer needed, but the proof of [Os13, Theorem 1.1], which is one of the main results of [Os13], still holds.
- (v) The proof of [Den15, Proposition 2.6] requires [EL92, Lemma 11]. Since the functions in [Den15] are in the class \mathcal{S} , this can be recovered by using Theorem 1.3(c) instead.

The second list contains those that need to be considered as still open.

- (i) Bhattacharyya [Bha83], Hinkkanen [Hin94] and Fang [Fan97] each used [Bak70] to prove the analogous result for analytic self-maps of the punctured plane. These results are not cited by any other paper.
- (ii) The result of [Bak70] is used to prove [CW03, Lemma 3.1], which in turn is used to prove [CW03, Theorem 1]. This paper also uses [EL92, Lemma 11] to prove [CW03, Lemma 4.5] and thence [CW03, Theorem 2]. These results are also quoted in [Lin08]. Note that [CW03, Theorem 1] was generalised to a wider class of functions in [NZC06, Lemma 3.3], and this result does not require any of the open theorems.

Finally we note two implications for survey papers:

- (i) The result of [Bak70] is stated as [Ber93, Theorem 17]; however, there are no significant implications elsewhere in this paper.
- (ii) The result [EL92, Lemma 11] is stated and proved as [EL89, Theorem 4.7]; however, there are no implications elsewhere in this paper.

REFERENCES

- [Bak70] I. N. Baker, *Completely invariant domains of entire functions*, Mathematical Essays Dedicated to A. J. Macintyre, Ohio Univ. Press, Athens, Ohio, 1970, pp. 33–35.
- [Bak75] ———, *The domains of normality of an entire function*, Ann. Acad. Sci. Fenn. Ser. A I Math. **1** (1975), no. 2, 277–283.
- [BD00] I. N. Baker and P. Domínguez, *Some connectedness properties of Julia sets*, Complex Variables Theory Appl. **41** (2000), no. 4, 371–389.
- [BDH01] I. N. Baker, P. Domínguez, and M. E. Herring, *Dynamics of functions meromorphic outside a small set*, Ergodic Theory Dynam. Systems **21** (2001), no. 3, 647–672.
- [BE95] W. Bergweiler and A. Eremenko, *On the singularities of the inverse to a meromorphic function of finite order*, Rev. Mat. Iberoamericana **11** (1995), no. 2, 355–373.
- [BE08a] W. Bergweiler and A. E. Eremenko, *Direct singularities and completely invariant domains of entire functions*, Illinois J. Math. **52** (2008), no. 1, 243–259.
- [BE08b] Walter Bergweiler and Alexandre Eremenko, *Meromorphic functions with two completely invariant domains*, Transcendental dynamics and complex analysis, London Math. Soc. Lecture Note Ser., vol. 348, Cambridge Univ. Press, Cambridge, 2008, pp. 74–89.
- [Ber93] W. Bergweiler, *Iteration of meromorphic functions*, Bull. Amer. Math. Soc. (N.S.) **29** (1993), no. 2, 151–188.
- [Ber02] ———, *A question of Eremenko and Lyubich concerning completely invariant domains and indirect singularities*, Proc. Amer. Math. Soc. **130** (2002), no. 11, 3231–3236.
- [BFJK17] Krzysztof Barański, Núria Fagella, Xavier Jarque, and Bogusława Karpińska, *Accesses to infinity from Fatou components*, Trans. Amer. Math. Soc. **369** (2017), no. 3, 1835–1867.
- [BFR15] W. Bergweiler, N. Fagella, and L. Rempe-Gillen, *Hyperbolic entire functions with bounded Fatou components*, Comment. Math. Helv. **90** (2015), no. 4, 799–829.
- [Bha83] P. Bhattacharyya, *On completely invariant domains and domains of normality*, J. Math. Phys. Sci. **17** (1983), no. 4, 427–434.
- [CW03] C-L. Cao and Y-F. Wang, *On completely invariant Fatou components*, Ark. Mat. **41** (2003), no. 2, 253–265.
- [Den15] A. Deniz, *Convergence of rays with rational argument in hyperbolic components—an illustration in transcendental dynamics*, Nonlinearity **28** (2015), no. 11, 3845–3871.
- [Dom98] P. Domínguez, *Dynamics of transcendental meromorphic functions*, Ann. Acad. Sci. Fenn. Math. **23** (1998), no. 1, 225–250.
- [DS18] P. Domínguez and G. Sienna, *A complementary proof of Baker’s theorem of completely invariant components for transcendental entire functions*, Preprint, arXiv:1803.04598v1 (2018).

- [EL89] A. E. Eremenko and M. Yu. Lyubich, *The dynamics of analytic transformations*, Algebra i Analiz **1** (1989), no. 3, 1–70.
- [EL92] ———, *Dynamical properties of some classes of entire functions*, Ann. Inst. Fourier (Grenoble) **42** (1992), no. 4, 989–1020.
- [Ere13] A. E. Eremenko, *Singularities of inverse functions*, Presented at the ICMS conference “The role of complex analysis in complex dynamics”, 2013.
- [Fan97] L. Fang, *Completely invariant domains of holomorphic self-maps on \mathbb{C}^** , J. Beijing Inst. Tech. **6** (1997), no. 3, 187–191.
- [Hei57] Maurice Heins, *Asymptotic spots of entire and meromorphic functions.*, Ann. Math. (2) **66** (1957), 430–439 (English).
- [Hin94] A. Hinkkanen, *Completely invariant components in the punctured plane*, New Zealand J. Math. **23** (1994), no. 1, 65–69.
- [Lin08] Q. Ling, *Some dynamical properties of transcendental meromorphic functions*, J. Math. Anal. Appl. **340** (2008), no. 2, 954–958.
- [Lin15] E. Lindelöf, *Sur un principe général de l’analyse et ses applications à la théorie de la représentation conforme*, Acta Soc. Fennicae **46**, no. 4 (1915), 1–35.
- [Nev53] Rolf Nevanlinna, *Eindeutige analytische Funktionen*, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete, Bd XLVI, Springer-Verlag, Berlin, 1953, 2te Aufl.
- [NZC06] T. W. Ng, J-H. Zheng, and Y Y. Choi, *Residual Julia sets of meromorphic functions*, Math. Proc. Cambridge Philos. Soc. **141** (2006), no. 1, 113–126.
- [Os13] J. W. Osborne, *Spiders’ webs and locally connected Julia sets of transcendental entire functions*, Ergodic Theory Dynam. Systems **33** (2013), no. 4, 1146–1161.
- [RS11] P. J. Rippon and G. M. Stallard, *Boundaries of escaping Fatou components*, Proc. Amer. Math. Soc. **139** (2011), no. 8, 2807–2820.
- [Why42] G. T. Whyburn, *Analytic Topology*, American Mathematical Society Colloquium Publications, v. 28, American Mathematical Society, New York, 1942.

DEPT. OF MATHEMATICAL SCIENCES, UNIVERSITY OF LIVERPOOL, LIVERPOOL L69 7ZL, UK,
ORCID: 0000-0001-8032-8580

E-mail address: l.rempe@liverpool.ac.uk

DEPT. OF MATHEMATICAL SCIENCES, UNIVERSITY OF LIVERPOOL, LIVERPOOL L69 7ZL, UK,
ORCID: 0000-0002-3543-6969

E-mail address: djs@liverpool.ac.uk