

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/21434>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

Clinical Records

Unilateral hearing loss due to a rhabdomyoma in a six-year-old child

J. P. P. M. VAN LEEUWEN*, M. PRUSZCZYNSKI†, H. A. M. MARRES*, J. A. GROTENHUIS‡, C. W. R. J. CREMERS*

Abstract

A case report of a six-year-old child is presented, who had had a unilateral sensorineural hearing loss for several years. Because of impairment in the ABR as well as in the caloric testing a MRI and CT scan were performed. A 17 mm tumour in the cerebellopontine angle (CPA) was detected, which after suboccipital surgery proved to be a rhabdomyoma. This tumour has not been described before in the CPA. Unilateral sensorineural hearing loss should, at all ages, be an indication for further (radiodiagnostic) investigations.

Key words: Rhabdomyoma; Hearing loss, unilateral; Cerebello-pontine angle; Child

Introduction

Unilateral sensorineural hearing loss and complete unilateral deafness seems to occur in one out of 1000 children of school age (Everberg, 1960). The cause most often remains unknown. Rarely a genetic origin can be traced (Marres, 1994). In adolescents and adults a small asymmetric sensorineural hearing loss requires an investigation of the cerebello-pontine angle to exclude a space-occupying lesion. A case report is presented of a six-year-old boy with a unilateral sensorineural hearing loss. After histological studies it was found to be a rhabdomyoma, which previously has been reported only once in an intracranial location in the trigeminal nerve (Zwick *et al.*, 1989). Nowadays, we have access to advanced imaging techniques of the inner ear and the cerebello-pontine angle, and modern radiology is non-expensive and comfortable for patients these additional investigations should also be considered in childhood.

Case report

In 1990, a six-year-old child was referred to the ORL department of the University Hospital, Nijmegen because of a unilateral hearing loss, found at a regular screening at school. There was no medical history. In a scheduled examination after the first year of life a hearing loss had been noticed, but no further action had been taken. The speech and motor development were normal.

At the ENT-examination a normal eardrum was seen, with air-containing middle ear. The tuning fork test according to Rinne and Weber were normal, and the Barany noise box test proved that both ears could hear. At pure tone audiometry a sensorineural hearing loss was found with a Fletcher Index (mean loss at 0.5, 1.0 and 2.0 kHz) of 50 dB in the left ear. Speech audiometry gave a word discrimination of 100 per cent with the same 50 dB shift. The hearing was normal in the right ear. Contral-

ateral stapedial reflexes could be elicited on both sides. The ABR on the left side revealed an interwave delay from J1-J3 and J1-J5. On the right side a much smaller interwave delay J1-J5 was found. Labyrinth testing with calorics indicated a reduced function of the left labyrinth.

MRI scanning had just become available in a nearby hospital and a tumour of the eighth nerve was found (Figure 1). It was not possible to use Gadolinium at that time. Later, a CT scan with the use of contrast showed a tumour of 17 mm in the left cerebello-pontine angle (Figure 2). Angiography showed no abnormal vascular structures.

The patient was referred to the neurosurgical department, and a suboccipital approach was used for the exploration of the cerebello-pontine angle. During the operation the tumour showed encapsulation with a smooth surface and firm consistency. There was an ingrowth in a dilated internal acoustic meatus and adhesions were found with the pons. It was not possible to remove the tumour totally. Post-operatively there was a fast recovery, but a total facial paralysis House grade VI (House and Brackman, 1985) remained. The left ear was completely deaf. There were no vertigo problems afterwards. MRI-scanning three years after the operation showed no growth of the remaining part of the tumour. Four years after the operation, a cross face nerve transplant was made using the left sural nerve to reinnervate the left facial nerve.

Pathology

During the operation a frozen section examination was performed and because of the presence of matured skeletal muscle a diagnosis of teratoma was suggested. In the permanent paraffin sections the tumour showed large well-developed matured skeletal muscle fibres with easily identifiable haematoxylin and eosin stain cross striations, accentuated by PTAH-Mallory stain (Figure 3). The fibres




FIG. 1

MRI without Gadolinium of the rhabdomyoma in the left cerebello-pontine angle.

were arranged in irregular fascicles and separated by various amounts of fibrocollagenous tissue. No cystic structures, vacuolated cells or spider cells were found. There was no mitotic activity. Muscle fibres showed diffuse cytoplasmic immunoreactivity for desmin, muscle-specific actin HHF-35 and myoglobin. No glial tissue was detected and there was no reaction for S100 protein and GFAP. Some tumour cells stained also for vimentin. The diagnosis rhabdomyoma was made.

Discussion

Hearing loss and even deafness of one ear in childhood is a regular finding in daily practice. Usually, the hearing loss is due to middle ear problems and will be solved in time, by using medication or middle ear ventilation tubes. When a sensorineural hearing loss is found there is a serious indication for further investigations. Tumours as in this young child are rare but can exist for several years giving symptoms which should not be overlooked.

Congenital intracranial tumours are rare and seldomly found in the cerebello-pontine angle (Werb *et al.*, 1992). A rhabdomyoma is defined as a benign and usually circumscribed tumour consisting of mature muscle cells (Kleihues *et al.*, 1993). Other benign mesenchymal tumours with intracranial locations such as leiomyoma of fibrous xanthoma have also been reported (Burger and Scheithauer, 1994).

The location of rhabdomyoma can be cardiac or extracardiac. Cardiac rhabdomyomas are rare congenital tumours resulting from early dysembryogenetic disorder of organogenesis. They are generally benign lesions considered as hamartomas which may be the first manifestation of Bourneville's tuberous sclerosis (Mehta, 1993). The extracardiac rhabdomyomas may assume a number of histological patterns. Tumours composed of uniform large polygonal eosinophilic cells with occasionally cross stria-

tion are classified as the adult-type. Those consisting of small mesenchymal cells and spindled skeletal muscle cells are classified as the foetal-type. Both tend to occur in the head and neck region (Knowles and Jacobiec, 1975; Fu and Perzin, 1976; Kapadia *et al.*, 1993; Kleihues *et al.*, 1993). A third form, the genital-type with cells bearing resemblance to both the adult and foetal type is most often seen as a polypoid mass in the vagina and vulva (Agamaolis *et al.*, 1986; Enzinger and Weiss, 1988). Foetal extracardiac rhabdomyomas occur principally in infants and young children and have a predilection for the postauricular region (Batsakis and Manning, 1986) but are even rarer than the adult-type rhabdomyomas (Enzinger and Weiss, 1988). They show close resemblance to rhabdomyosarcomas. Of particular interest is their association with multiple basal cell carcinomas and anomalies of the rib and the iridocorneal angle of the eye (Enzinger and Weiss, 1988). In the adult form the tumour can be found multifocal in the head and neck region (Shemen *et al.*, 1992; Kapakia *et al.* 1993). In the neck the adult type rhabdomyoma seems to arise from the branchial musculature of the third and fourth branchial arches. Therefore, the tumour is found most frequently in the region of the larynx, pharynx, floor of the mouth and base of the tongue (Solomon and Tolete-Velcek, 1975; Enzinger and Weiss, 1988). Both adult and foetal type rhabdomyomas are more common in males (Agamaolis *et al.*, 1986; Kapadia *et al.*, 1993). Recently a juvenile-type of rhabdomyoma has been described which appears as an intermediate form between the foetal- and adult-type (Nakheh *et al.*, 1991; Crotty *et al.*, 1993).

Microscopically intracranial rhabdomyomas must be distinguished from rhabdomyosarcomas, medullomyoblas-




FIG. 2

CT scan with contrast of the rhabdomyoma in the left cerebello-pontine angle.



FIG. 3

Photomicrograph of the tumour illustrating well developed skeletal muscle fibres with easily identified cross striation accentuated by PTAH-Malory stain. The picture is consistent with rhabdomyoma ($\times 250$).

tomas, malignant triton tumours and rare intracranial rhabdoid tumours (Jakate *et al.*, 1988; Kodet *et al.*, 1991; Burger and Scheithauer, 1994; Parham *et al.*, 1994). Rhabdomyomas may be encountered as a component of the malignant germ cell tumours of the pineal gland and suprasellar region, particularly in immature teratomas. Skeletal muscle differentiation can also rarely be seen in glycosarcomas (Goldman, 1969), gliomas and meningiomas (Ferracini *et al.*, 1982; Burger and Scheithauer, 1994). In this case the tumour classified as the adult-type rhabdomyoma showed other distinctive features and no mitotic activity. Immature neuroblasts, immature mesenchymal cells, vacuolated or globular rhabdomyoblasts and glial cells were not found. Our immuno-histochemical findings were consistent with the rhabdomyomatous phenotype (Helliwell *et al.*, 1988; Kapadia *et al.*, 1993).

The four year follow-up time free of recurrent disease confirmed the benign character of the lesion. Overall incidence of rhabdomyomas is much lower than rhabdomyosarcomas both extracranially and intracranially (Enzinger and Weiss, 1988; Zwick *et al.*, 1989; Burger and Scheithauer, 1994). Rhabdomyosarcoma is a highly malignant tumour and patients stay alive no longer than two years despite aggressive chemotherapy and radiotherapy (Taratuto *et al.*, 1985).

It is a question of debate whether the presented case should be considered as a choristoma or a true neoplasm. We were able to find in the literature only one other case of a primary intracranial rhabdomyoma which involved the

trigeminal nerve. The authors regarded the lesion as choristoma and postulated that the tumour arose from embryologically disrupted and dislocated masticatory myogenic tissue following innervation of the mandibular motor division of the trigeminal nerve (Zwick *et al.*, 1989). The foci of skeletal muscles can be found accidentally in normal meninges, but their relationship to rhabdomyomas and rhabdomyosarcomas is not clear (Hoffman and Rorke, 1971; Fix *et al.*, 1989; Burger and Scheithauer, 1994). Rhabdomyosarcomas have been reported to arise in both the leptomeninges and the brain parenchyma (Korinthenberg *et al.*, 1984; Taratuto *et al.*, 1985; Ferracini *et al.*, 1992;). Focal skeletal muscle ectopia in leptomeninges has been reported predominantly in children with other developmental anomalies of the central nervous system and sometimes chromosomal abnormalities which suggests faulty mesenchymal differentiation secondary to a genetic error in the tissue regulation. In several cases skeletal muscle has been found in leptomeninges of the pontine region or ponto-medullary junction (Johnson and Ludwin, 1984; Fix *et al.*, 1989). Interestingly, the tumour reported by us was also located in the pontine angle area.

Conclusion

Unilateral sensorineural hearing loss may be an alarming symptom in childhood. Further investigations using CT scanning and MRI should be considered to find the cause of the phenomenon. In childhood rare congenital tumours such as a rhabdomyoma are sometimes found in these cases and early exploration increases the chance of successful radical surgery. The presented child is alive and well four years after operation, although only an incomplete resection could be performed. An additional surgical procedure for the total facial paralysis was performed.

Acknowledgement

The authors wish to thank Dr P. C. Burgers (Department of Pathology, Duke University Medical Centre, Durham USA) for his opinion on the presented case.

References

- Agamanolis, D. P., Dasu, S., Krill, C. E. (1986) Tumors of skeletal muscle. *Human Pathology* **17**: 778-795.
- Batsakis, J. G., Manning, J. T. (1986) Soft tissue tumors: unusual forms. *Otolaryngologic Clinics of North America* **19**: 659-683.
- Burger, P. C., Scheithauer, B. W. (1994) Tumors of the central nervous system. In *Atlas of Tumor Pathology*. Armed Forces Institute of Pathology, Washington, pp 302-308.
- Crotty, P. L., Nakhleh, R. E., Dehner, L. P. (1993) Juvenile rhabdomyoma: an intermediate form of skeletal muscle tumor in children. *Archives of Pathology and Laboratory Medicine* **117**: 43-47.
- Enzinger, F. M., Weiss, S. W. (1988) Rhabdomyoma. *Soft Tissue Tumors*. Mosby, St. Louis, pp 433-447.
- Everberg, G. (1960) Unilateral anacusis. Clinical, radiological and genetic investigations. *Acta Oto-Laryngologica (Stockholm)* (suppl) **158**: 366-374.
- Ferracini, R., Poggi, S., Frank, G. (1992) Meningeal sarcoma with rhabdomyoblastic differentiation: case report. *Neurosurgery* **30**: 782-785.
- Fix, S. E., Nelson, J., Schochet, S. S. (1989) Focal leptomeningeal rhabdomyomatosis of the posterior fossa. *Archives of Pathology and Laboratory Medicine* **113**: 872-873.
- Fu, Y., Perzin, K. H. (1976) Nonepithelial tumors of the nasal cavity, paranasal sinuses and nasopharynx. *Cancer* **37**: 364-376.
- Goldman, R. L. (1969) Gliomyosarcoma of the cerebrum.

- Report of a unique case. *American Journal of Clinical Pathology* **52**: 741-745.
- Helliwell, T. R., Sissons, M. C. J., Stoney, P. J., Ashworth, M. T. (1988) Immunocytochemistry and electron microscopy of head and neck rhabdomyoma. *Journal of Clinical Pathology* **41**: 1058-1063.
- Hoffman, S. F., Rorke, L. B. (1971) On finding striated muscle in the brain. *Journal of Neurology and Psychiatry* **34**: 761-764.
- House, J. W., Brackmann, D. E. (1985) Facial nerve grading system. *Otolaryngology-Head and Neck Surgery* **93**: 146-147.
- Jakate, S. M., Marsden, H. B., Ingram, L. (1988) Primary rhabdoid tumor of the brain. *Virchows Archives* **412**: 393-397.
- Johnson, E. S., Ludwin, S. K. (1984) Rhabdoneuroglial heterotopias of the pontine leptomeninges in trisomy 13. *Archives of Pathology and Laboratory Medicine* **108**: 906-908.
- Kapadia, S. B., Meis, J. M., Frisman, D. M., Ellis, G. L., Heffner, D. K., Hyams, V. J. (1993) Adult rhabdomyoma in the head and neck: a clinicopathologic and immunophenotypic study. *Human Pathology* **24**: 608-617.
- Kleihues, P., Burger, P. C., Scheithauer, B. E. (1993) Histological typing of tumors of the central nervous system. *WHO International Classification of Tumors*, Springer-Verlag, Berlin/Heidelberg, 28-29.
- Knowles, D. M. H., Jacobiec, F. (1975) Rhabdomyoma of the orbit. *American Journal of Ophthalmology* **80**: 1011-1018.
- Kodet, R., Newton, W. A., Hamoudi, A. B., Asmar, L. (1991) Rhabdomyosarcomas with intermediate-filament inclusions and features of rhabdoid tumors. *American Journal of Surgical Pathology* **15**: 257-267.
- Korinthenberg, R., Edel, G., Palm, D., Muller, K. M., Brandt, M., Muller, R. P. (1984) Primary rhabdomyosarcoma of the leptomeninges: clinical, neuroradiological and pathological aspects. *Clinical Neurology and Neurosurgery* **86**: 301-305.
- Marres, H. A. M. (1994) Genetic deafness. Special reference to branchial arch syndromes. Thesis KU Nijmegen, the Netherlands.
- Mehta, A. V. (1993) Rhabdomyoma and ventricular pre-excitation syndrome. A report of two cases and review of the literature. *American Journal of Diseases in Children* **147**: 669-671.
- Nakheh, R. E., Swanson, P. E., Dehner, L. P. (1991) Juvenile (embryonal and alveolar) rhabdomyosarcoma of the head and neck in adults: a clinical, pathological and histochemical study of 12 cases. *Cancer* **67**: 1019-1024.
- Parham, D. M., Weeks, D. A., Beckwith, J. B. (1994) The clinicopathological spectrum of putative extrarenal rhabdoid tumors. *American Journal of Surgical Pathology* **18**: 1010-1029.
- Shemen, L., Spiro, R., Tuazon, R. (1992) Multifocal and adult rhabdomyomas of the head and neck. *Head and Neck* **14**: 395-400.
- Solomon, M. P., Tolete-Velcek, F. (1975) Lingual rhabdomyoma (adult variant) in a child. *Journal of Pediatric Surgery* **14**: 91-94.
- Tatatuto, A. L., Molina, H. A., Diez, B., Zuccaro, G., Monges, J. (1985) Primary rhabdomyosarcoma of brain and cerebellum. Report of four cases in infants: an immunohistochemical study. *Acta Neuropathologica (Berlin)* **66**: 98-104.
- Werb, P., Scurry, J., Ostor, A., Fortune, D., Attwood, H. (1992) Survey of congenital tumors in perinatal necropsies. *Pathology* **24**: 247-253.
- Zwick, D. L., Livingston, K., Clapp, L., Kosnik, E., Yates, A. (1989) Intracranial trigeminal nerve rhabdomyoma/choristoma in a child: a case report and discussion of possible histogenesis. *Human Pathology* **20**: 390-393.

Address for correspondence:

J. P. P. M. van Leeuwen,
Dept of Otorhinolaryngology,
University Hospital Nijmegen,
PO Box 9101,
6500 HB Nijmegen,
The Netherlands.

Fax (80) 540251