

**UNIVERSITY OF  
LEADING  
THE WAY  
WESTMINSTER** 

**WestminsterResearch**

<http://www.westminster.ac.uk/research/westminsterresearch>

***In vitro* assessment of the synergistic effects of antibiotics and wound dressings on biofilms from diabetic foot pathogens**

**George Gyamfi-Brobby  
Pamela Greenwell  
Patrick Kimmitt**

Faculty of Science and Technology, University of Westminster

This is a copy of the poster presented at the Society for General Microbiology Annual Conference, International Convention Centre, Birmingham, UK, 30 March – 2 April 2015.

Copyright © 2015 The Authors.

---

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

---

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch:

(<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail [repository@westminster.ac.uk](mailto:repository@westminster.ac.uk)



## Background

The impact of biofilm in the effective control of wound microbiome is an ongoing dilemma which has seen the use of different treatment strategies. The effects of wound dressings and antibiotics on both planktonic bacteria and biofilms have been separately evaluated in previous studies.<sup>1,2,3,4</sup> Some of the methods used include 96-well microtitre plate assay, 6-well plate assay, isothermal calorimetry, microscopy and the constant depth film fermentation.<sup>1,2,5,7,9</sup>

However, the quest for a more sensitive and reproducible method to mimic the biofilm phenotype is ongoing. One such method uses poloxamer gel to grow biofilm phenotype due to its proven ability to promote sessile growth and maintain the biofilm architecture to mimic clinically conditions.<sup>8</sup>

## Aim of Study

The aim of this study was to assess the combined effects of some selected wound dressings (silver-impregnated: Acticoat (ACT) and Silvercel (SIL); and honey-impregnated: Medihoney™ Apinate (MA)) and antibiotics (ceftazidime and levofloxacin) on *Klebsiella pneumoniae* and *Proteus mirabilis* biofilms using a standard agar assay. The ability of poloxamer gel cultures to support sessile growth and maintain biofilm architecture was also assessed in comparison with the standard agar method.

## Methods

The two multidrug resistant diabetic foot isolates were initially grown overnight and diluted to final broth suspensions of 10<sup>8</sup> colony forming unit (CFU)/mL.<sup>5</sup> 1 mL volumes of *K. pneumoniae* and *P. mirabilis* suspensions containing ceftazidime (CAZ) and levofloxacin (LEV) at final concentrations of 256, 512, 1024 and 5120 µg/mL were inoculated on Mueller Hinton agar (MHA) and 30% (w/v) Kolliphor® P 407 (poloxamer) gel plates and allowed to dry.

Wound dressings cut into circular shapes (2cm-diameter) were aseptically placed on the agar and gel plates and incubated at 35 – 37° C for 24 hours. ZOIs produced by the 3 antibacterial dressings after 24 hours were measured and compared with a control dressing (Atrauman (ATR) – with no antibacterial activity).

The ZOIs measurements are presented as means (±SEM) and statistically analyzed using GraphPad Prism software.

## Results

ZOIs associated with ACT, SIL and MA dressings augmented with CAZ and LEV were compared with no antibiotic that of ATR controls. All three dressings showed significant ( $p < 0.05$ ) biofilm-inhibiting activity against both bacteria at antibiotic concentrations of 1024 and 5120µg/mL with ZOI between 17.5 and 35mm on MHA (Figure 2. (A) and (B)). Similarly, significant ZOIs on Kolliphor® P 407 gels were between 22 and 30mm at the same concentrations (Figure 2. (C) and (D)).




Figure 1. *P. mirabilis* biofilm phenotypes with ZOI on (A). Kolliphor gel and (B). MHA




Figure 2. Effect of wound dressings in combination with antibiotics on (C). *K. pneumoniae* and (D). *P. mirabilis* biofilms. (\*) - shows significant inhibition of biofilms on Kolliphor gel ( $p < 0.05$ )

## Discussion

This study has demonstrated that clinical strains of *K. pneumoniae* and *P. mirabilis* biofilms are highly resistant to current antimicrobial agents on the market. They are however sensitive to higher concentrations of antibiotics which are not clinically applicable.

In this study, ACT produced greater ZOIs against both *K. pneumoniae* and *P. mirabilis* biofilms than SIL and MA. This suggests that the continuous treatment of infected diabetic foot with ACT can improve healing.

It was also observed that, all 3 dressings were less effective on Kolliphor-grown biofilms than on MHA. This is because standard agar only promotes the growth of colony forming units in a semi-sessile state as previously described.<sup>10</sup>

As already established and confirmed in this study, biofilms are difficult to treat.<sup>5,6</sup> Their presence in wounds prolong healing, increase cost of treatment and affect the quality of life of affected individuals. Therefore, the effective management of chronic wounds must encompass a holistic approach that includes antimicrobial chemotherapy, surgical debridement, physiotherapy and periodic reviews.

## References

- Chaw, K. C., Manimaran, M., Tay, F. E. H. (2005). Role of silver ions in destabilization of intermolecular adhesion forces measured by atomic force microscopy in *Staphylococcus epidermidis* biofilms. *Antimicrob Agents Chemother.* 49(12): 4853-59.
- Willix, D. J, Molan, P. C, Harfoot, C. G.(1992). A comparison of the sensitivity of wound-infecting species of bacteria to the antibacterial activity of manuka honey and other honey. *J. Appl. Bacteriol.* 73:388–394.
- Wright, B.J., Lam, K., Burrell, R. (1998) 'Wound management in an era of increasing bacterial antibiotic resistance: A role for topical silver treatment', *American Journal of Infection Control*, 26(6), pp. 572-577.
- Seckam, A., Cooper, R. (2013). Understanding how honey impacts on wounds: an update on recent research findings. *Wounds International*, 4(1), 20-24
- Percival, S. L., Bowler, P. G., & Dolman, J. (2007). Antimicrobial activity of silver-containing dressings on wound microorganisms using an in vitro biofilm model. *International wound journal*, 4(2), 186-191.
- Lopez, D., Vlamakis, H., Kolter, R. (2010). Biofilms. *Cold Spring Harb Perspect Biol* 2:a000398
- Hill, K. E., Malic, S., McKee, R., Rennison, T., Harding, K. G., Williams, D. W., & Thomas, D. W. (2010). An in vitro model of chronic wound biofilms to test wound dressings and assess antimicrobial susceptibilities. *Journal of Antimicrobial Chemotherapy*, dkq105.
- Gilbert, P., Jones, M. V., Allison, D. G., Heys, S., Maira, T., & Wood, P. (1998). The use of poloxamer hydrogels for the assessment of biofilm susceptibility towards biocide treatments. *Journal of applied microbiology*, 85(6), 985-990.
- Gaisford, S., Beezer, A. E., Bishop, A. H., Walker, M., & Parsons, D. (2009). An in vitro method for the quantitative determination of the antimicrobial efficacy of silver-containing wound dressings. *International journal of pharmaceuticals*, 366(1), 111-116.
- Freeman, D. J., Falkiner, F. R., & Keane, C. T. (1989). New method for detecting slime production by coagulase negative staphylococci. *Journal of clinical pathology*, 42(8), 872-874.