

1 **Characterization of virus-like particles associated with the human faecal and caecal**  
2 **microbiota**

3

4 Lesley Hoyles<sup>a</sup>, Anne L. McCartney<sup>b</sup>, Horst Neve<sup>c</sup>, Glenn R. Gibson<sup>b</sup>, Jeremy D. Sanderson<sup>d</sup>,  
5 Knut J. Heller<sup>c</sup>, Douwe van Sinderen<sup>a,e\*</sup>

6

7 <sup>a</sup>School of Microbiology, University College Cork, Western Road, Cork, Ireland

8 <sup>b</sup>Food Microbial Sciences Unit, Department of Food and Nutritional Sciences, The University  
9 of Reading, Whiteknights, Reading RG6 6UR, UK

10 <sup>c</sup>Institute of Microbiology and Biotechnology, Max Rubner-Institut, Federal Research  
11 Institute of Nutrition and Food, Kiel, Germany

12 <sup>d</sup>Department of Gastroenterology, Guy's and St Thomas' NHS Foundation Trust and King's  
13 College London, London, UK

14 <sup>e</sup>Alimentary Pharmabiotic Centre, Biosciences Institute, National University of Ireland,  
15 Western Road, Cork, Ireland

16

17 Lesley Hoyles [lesley.hoyles@reading.ac.uk](mailto:lesley.hoyles@reading.ac.uk)

18 Anne L. McCartney [a.l.mccartney@reading.ac.uk](mailto:a.l.mccartney@reading.ac.uk)

19 Horst Neve [horst.neve@mri.bund.de](mailto:horst.neve@mri.bund.de)

20 Glenn R. Gibson [g.r.gibson@reading.ac.uk](mailto:g.r.gibson@reading.ac.uk)

21 Jeremy D. Sanderson [jeremy.sanderson@kcl.ac.uk](mailto:jeremy.sanderson@kcl.ac.uk)

22 Knut J. Heller [knut.heller@t-online.de](mailto:knut.heller@t-online.de)

23 Douwe van Sinderen [d.vansinderen@ucc.ie](mailto:d.vansinderen@ucc.ie) \*Correspondence and reprints

24

24 **ABSTRACT**

25           This work represents an investigation into the presence, abundance and diversity of  
26 virus-like particles (VLPs) associated with human faecal and caecal samples. Various  
27 methodologies for the recovery of VLPs from faeces were tested and optimized, including  
28 successful down-stream processing of such samples for the purpose of an in-depth electron  
29 microscopic analysis, pulsed-field gel electrophoresis and efficient DNA recovery. The  
30 applicability of the developed VLP characterization method beyond the use of faecal samples  
31 was then verified using samples obtained from human caecal fluid.

32

33 **Keywords:** bacteriophages; microbial ecology; gastrointestinal tract; transmission electron  
34 microscopy; caecum; faeces.

35

36

37

## 37 1. INTRODUCTION

38 Viruses are the most numerous biological entities within the biosphere of our planet  
39 (being present at an estimated number of  $\sim 10^{31}$ ), with bacteriophages representing the most  
40 abundant group of environmental viruses [1, 2]. Bacteriophages are ubiquitous viruses that  
41 infect bacterial cells and disrupt their metabolism. Multiple bacteriophage types can infect a  
42 specific microbial isolate, with most bacteriophages infecting only certain species or even  
43 strains of bacteria [3].

44 Although they are abundant and potentially important to microbial populations  
45 indigenous to different ecological niches within the human gastrointestinal tract and to host  
46 health, little attention has been paid to bacterial virus-like particle (VLP) assemblages and  
47 their interactions with the gastrointestinal microbiota and/or human host until recently.  
48 Various publications have highlighted the potential importance of bacteriophages in  
49 inflammation states, including Crohn's disease [4, 5, 6], and as therapeutic agents [7, 8].  
50 Breitbart *et al.* [9] conducted the first metagenomic study on dsDNA-containing VLPs  
51 associated with the human faecal virome using a fresh sample from a healthy adult male. On  
52 the basis of previous estimates of gut microbial diversity, Breitbart *et al.* [9] predicted that  
53 there are two to five times more viral genotypes ( $\sim 1,200$  viral genotypes) present in the  
54 human gastrointestinal microbiota than the number of bacterial species, with the vast majority  
55 of these VLPs representing bacteriophages and prophages. RNA viruses present in human  
56 faeces have been found to be mostly associated with plant viruses, such as Pepper Mild  
57 Mottle Virus, with RNA bacteriophages making a minimal contribution to the diversity of the  
58 intestinal virome [10]. Using frozen samples from four adult female monozygotic twins and  
59 their mothers, Reyes *et al.* [11] demonstrated that each individual harbours a unique virome  
60 (ssDNA and dsDNA) regardless of their genetic relatedness to another individual, and that  
61 intrapersonal virome diversity is very low, with >95 % of virotypes retained over a one-year

62 period (between 71 and 2,773 viral genotypes identified among the samples). More recently,  
63 Minot *et al.* [12] confirmed inter-individual variation in the virome and, similar to  
64 modulation of the faecal bacteriome by foodstuffs, demonstrated that diet influences host  
65 faecal virome structure. Minot *et al.* [13] also confirmed the relative stability of an  
66 individual's virome, monitoring 24 faecal samples from one individual over a 2.5-year  
67 period.

68 The murine virome has been used to demonstrate enrichment of bacteriophage-  
69 encoded, antibiotic-resistance genes (related and unrelated to the drug used) after antibiotic  
70 therapy, with the adaptive capacity of the virome (specifically its bacteriophage component)  
71 suggested to protect gut bacteria, thereby preserving the microbiota's robustness during  
72 antibiotic stress [14].

73 It is clear from the aforementioned studies that the virome–bacteriome community in  
74 the gut is governed by complex and dynamic interactions in health, and that its balance may  
75 be disturbed when under stress, e.g. during antibiotic intervention [14]. Although  
76 metagenomic studies have greatly improved our understanding of the virome associated with  
77 the human and murine gut microbiomes, it has been notable how little bacteriophage  
78 material, and consequently DNA, has been isolated from samples when this information was  
79 included in a publication. Reyes *et al.* [11] reported the isolation of ~500 ng DNA from 2 to 5  
80 g of frozen faeces, whereas Thurber *et al.* [15] stated that between 500 and 3000 ng of DNA  
81 could be isolated from 500 g of human faeces, though the publication they cite [9] provides  
82 no information regarding the amount of DNA isolated from the 500 g faecal sample  
83 examined in the original study. In addition, to the best of our knowledge, no attempts have  
84 been made to enumerate bacteriophages in faecal filtrates prior to CsCl purification. The aims  
85 of this study were to develop reliable and effective methods for the recovery and

86 characterization of VLPs in human samples, and to apply these methods to human faecal and  
87 caecal samples to demonstrate the methods' efficiency.

88

## 88 2. MATERIALS AND METHODS

### 89 2.1 Processing of and isolation of VLPs from faecal and caecal samples

90 Faecal samples were obtained from six healthy adult (2 male, 4 female) donors of  
91 between 23 and 52 years of age, all of whom were members of the Department of Food and  
92 Nutritional Sciences, University of Reading. None of the donors had taken antibiotics,  
93 prebiotics and/or probiotics in the 3 months prior to sampling, and none had any history of  
94 gastrointestinal disorder. All donors gave informed oral consent for their faecal samples to be  
95 used for microbiological analyses. Ethical approval for the collection of caecal effluent from  
96 patients was obtained from St Thomas' Hospital Research Ethics Committee (06/Q0702/74)  
97 covering Guy's and St Thomas' Hospitals and transferred by agreement to London Bridge  
98 Hospital. Where available, clinical information for the colonoscopy patients is given in the  
99 text. Samples were collected at Reading (faeces) or transported from St Thomas' Hospital  
100 (caecum), and maintained under anaerobic conditions (faeces, MACS1000 anaerobic  
101 workstation, Don Whitley Scientific, gas composition 80 % N<sub>2</sub>, 10 % H<sub>2</sub>, 10 % CO<sub>2</sub>; caecum,  
102 on ice in a gas jar with an anaerobic gas-generating pack; Oxoid Ltd) for a maximum of 2 h  
103 before processing. Caecal samples were collected during routine colonoscopy following  
104 preparation of the bowel with sodium picosulphate and a reduced fibre diet for 3 days. Liquid  
105 residue in the caecum was aspirated via the colonoscope suction channel into a standard 30  
106 ml trap specimen container and immediately transferred to a gas jar.

107 Faecal homogenates (20 %, w/v, from 25 g of faecal material) were prepared in sterile  
108 TBT [0.1 µm filtered prior to autoclaving; 100 mM Tris/HCl, pH 8.0; 100 mM NaCl; 10 mM  
109 MgCl<sub>2</sub>.6H<sub>2</sub>O] or sterile 0.5 % 'Lab-Lemco'/6 % NaCl (LL [16]). Caecal homogenates (20 %,   
110 v/v, from 10 to 30 ml of caecal effluent) were prepared in LL. Faecal and caecal samples  
111 were placed into a filter stomacher bag and homogenized in a Stomacher 400 Lab System  
112 (Seward) for 120 s at low speed. The bag was removed from the stomacher and massaged

113 manually to further disperse any large particles remaining in the sample; the sample was then  
114 stomached for a further 120 s. The homogenate was kept on ice for 2 h to allow desorption of  
115 VLPs from solid material, then centrifuged at 11,180 *g* for 30 min at 10°C. The supernatant  
116 was transferred to a fresh tube and centrifuged again at 11,180 *g* for 30 min at 10°C.  
117 Supernatant [herein referred to as faecal filtrate (FF) for both faecal and caecal samples] was  
118 passed through sterile 0.45 µm cellulose acetate filters (Millipore) and the FF was then  
119 collected in a sterile container. An aliquot (10 µl) of each of the FFs was examined by  
120 epifluorescence microscopy (EFM) after viral particles had been stained with SYBR Gold  
121 (see below) to confirm that the samples were free of bacteria.

122 FF prepared in LL was used for enumeration of VLPs in faeces via EFM (see below)  
123 and 1 ml aliquots of FF prepared in TBT were for examination by transmission electron  
124 microscopy (TEM; see below).

125 Aliquots of FF and LL (two sets of three aliquots of 100 µl each: one for aerobic  
126 cultivation, one for anaerobic cultivation) were spread onto Columbia blood agar containing  
127 5 % laked horse blood (Oxoid), and incubated aerobically and anaerobically. Sterility of the  
128 filtrates was confirmed by the absence of microbial growth on plates following incubation for  
129 2 (aerobic) and 5 (anaerobic) days. Sterile brain–heart infusion broth was inoculated with 100  
130 µl of FF and LL, and incubated aerobically for 2 days: aliquots (100 µl) were spread in  
131 triplicate on plates to confirm sterility of the broth cultures. In addition, 10-ml aliquots of FF  
132 were stored at 4°C for 6 months after collection, and were found to be free of bacteria when  
133 checked (by plating) at monthly intervals.

134 Poly(ethylene glycol) (PEG; BioUltra, 8000; Sigma) was added to 2 × 20 ml aliquots  
135 of the LL-prepared 0.45 µm-filtered FFs (final concentration of PEG, 10 %, w/v). The  
136 samples were shaken gently to dissolve the PEG, then left at 4°C for 16 h ([16] used 16 h as  
137 they observed an improved recovery of RNA bacteriophage compared to shorter incubations;

138 [17] used 10 h, as they determined this to be longer than the minimum time required to obtain  
139 a constant and stable white layer of viral precipitate). PEG-precipitated VLPs were then  
140 harvested by centrifugation at 4,500 *g* for 30 min in a swing-out-bucket rotor at 4°C. The  
141 supernatant was removed and the pellet resuspended in 1 ml TBT for pulsed-field gel  
142 electrophoresis (PFGE) or 5 ml TBT for purification of VLPs on a CsCl gradient [15] with  
143 ultracentrifugation performed at 100,000 *g* for 2 h at 10°C in a fixed-angle Type 50 Ti rotor  
144 (Beckman Coulter). VLPs recovered by CsCl (1.35 and 1.5 g/cm<sup>3</sup> fractions) were dialysed  
145 (12,000 Da cut-off) twice against 400 ml sterile TBT, passed through a sterile 0.45 µm  
146 cellulose acetate filter and stored at 4°C until DNA was extracted. An aliquot (5 µl) of each  
147 of the CsCl-purified samples was viewed using EFM after viral particles had been stained  
148 with SYBR Gold (see below) to confirm that the samples were free of bacteria.

149

## 150 **2.2 Epifluorescence microscopy (EFM)**

151 A combination of the methods of Thurber and Patel [15, 18] was used to prepare FFs  
152 (faecal only) for EFM. VLPs present in FF were not fixed in paraformaldehyde prior to  
153 enumeration, as Wen *et al.* [19] showed that aldehyde fixation leads to a rapid loss in viral  
154 abundance. The SYBR Gold concentration of 1–5× recommended by [15] did not work with  
155 FFs; therefore, the 400× concentration recommended by [18] for use with planktonic aquatic  
156 samples was used in this study.

157 The filtration system (see [15] for specifics of the set-up employed), including a  
158 glass-graduated column, for collecting VLPs on filters was cleaned using 5 ml of 0.1 µm-  
159 filtered, sterile H<sub>2</sub>O and 5 ml of 0.1 µm-filtered ethanol. A sterile pair of flat-tipped forceps  
160 was used to remove a 0.02-µm white Anodisc 25 membrane (Whatman) from its box, and the  
161 filter was fitted to the glass frit of the filtration system under a low vacuum [ $<10$  psi ( $\sim 62$   
162 kPa)]. Duplicate 2 ml aliquots of 0.1 µm filtered, sterile H<sub>2</sub>O and 10 µl of sterile LL in 2 ml



163 0.1 µm filtered, sterile H<sub>2</sub>O were used as negative controls (to assess whether there was any  
164 contamination in the water, the LL, the filter tower or the filter surface) and passed through  
165 filters under low vacuum before any of the FF aliquots were filtered in duplicate. For each  
166 sample, a 10 µl aliquot of 0.45 µm-filtered FF was added to 2 ml of the 0.1 µm filtered,  
167 sterile H<sub>2</sub>O and the suspension gently mixed. The diluted sample was then introduced into the  
168 filter system with the low vacuum maintained. The liquid was passed through the filter, and  
169 VLPs were collected on the filter. The clip and glass column were carefully removed from  
170 the filtration system and the filter was gently removed from the glass frit (still under  
171 vacuum). While holding the filter with a forceps, the back of the filter was gently blotted dry  
172 on a clean Kimwipe; the filter was then left to dry on a Kimwipe in a sterile Petri dish for 10  
173 min in a dark box. The filtration system was cleaned with 5 ml of 0.1 µm-filtered, sterile H<sub>2</sub>O  
174 and 5 ml of 0.1 µm-filtered ethanol, and the next sample processed.

175         Filters were stained with 100 µl droplets of 400× SYBR Gold and dried as described  
176 by [18], prior to being applied in pairs to sterile glass slides. The mountant (20 µl per  
177 coverslip) ProLong<sup>®</sup> Gold antifade reagent (Invitrogen) was added to 25 mm glass coverslips.  
178 Coverslips were picked up with sterile forceps, inverted and placed on the Anodisc filters on  
179 the microscope slides. Slides were then left at room temperature in the dark for 16 h to allow  
180 the antifade to cure. Slides prepared in this manner can be stored at room temperature for 1  
181 week or at 4°C for 2 months.

182         Slides were viewed at 1000× magnification under a Nikon Microphot-SA microscope  
183 fitted with a B-2A (blue excitation) filter and attached to a CoolSNAP-Pro MONOCHROME  
184 (Media Cybernetics Inc.) camera. Images were captured using Image-Pro PLUS version  
185 4.5.0.19 (Media Cybernetics Inc.), and VLPs in 25 fields of view were enumerated by eye.  
186 The number of VLPs per millilitre of FF was calculated using the following equation: mean  
187 number of VLPs in 25 fields of view × 100 × 25760.205; where 100 represents taking the

188 sample back to per millilitre FF and 25760.205 represents the number of fields of view on a  
189 filter. To give an estimate of the number of VLPs per gram of faeces, results were multiplied  
190 by a factor of 5. The detection limit of the method was  $1.03 \times 10^5$  VLPs/ml FF or  $5.15 \times 10^5$   
191 VLPs/g faeces.

192

### 193 **2.3 TEM of FFs (faecal and caecal)**

194 Aliquots (1 ml) of FF prepared in TBT (faeces) or LL (caecum) were used for TEM  
195 analysis within 1 week of collection: these filtrates had not been concentrated using PEG.  
196 Carbon films (~3×3 mm in size) were floated from mica-sheets into a drop of filtrate (100  
197 µl). After an adsorption time of 5–10 min, samples were transferred into a drop of 1 % (v/v)  
198 of EM-grade glutaraldehyde (20 min) and subsequently into a drop of 2 % (w/v) uranyl  
199 acetate for negative staining (1–2 min). After two washes for a few seconds in drops of  
200 distilled water, samples were picked up with 400-mesh copper grids (Plano, Wetzlar, D).  
201 Electron micrographs were taken in a Tecnai 10 transmission electron microscope (FEI  
202 Company, Eindhoven, the Netherlands) at an accelerating voltage of 80 kV. Digital  
203 micrographs were taken with a Megaview G2 CCD camera (Olympus SIS, Münster,  
204 Germany). For estimation of VLP titers on the EM grids, a freshly prepared *Lactococcus*  
205 *lactis* bacteriophage preparation with defined titer and unique morphology (i.e. prolate-  
206 headed bacteriophage P001 [20]) was added in concentrations of  $10^7$  and  $10^8$  plaque-forming  
207 units (pfu) per ml to the faecal sample of donor 2 (with highest bacteriophage titer according  
208 to TEM analysis).

209

### 210 **2.4 PFGE**

211 Aliquots (40 µl) of PEG-precipitated samples resuspended in 1 ml TBT were used for  
212 PFGE. These were heated at 60 or 75°C as appropriate. An aliquot (25 µl) of each heated

213 (viscous) suspension and loading dye (5  $\mu$ l) were combined and loaded into wells, which  
214 were sealed with molten agarose. Run conditions for PFGE were as described by Fuhrman *et*  
215 *al.* [21]: a 1 % agarose gel (SeaKem LE agarose) was made in 0.5 $\times$  TBE [250 ml of 5 $\times$  TBE  
216 buffer (27 g Sigma 7–9, 13.75 g boric acid, 5 ml 0.5 M EDTA, 500 ml of H<sub>2</sub>O) added to 2.25  
217 l H<sub>2</sub>O] and a CHEF DR II apparatus (Bio-Rad) was run for 18 h at 6 V and 14°C in 0.5 $\times$   
218 TBE, with a 1–10 s switch time. Gels were stained with ethidium bromide (5  $\mu$ g/ml) for 20  
219 min and destained in distilled H<sub>2</sub>O for 10 min, or with 1 $\times$  SYBR Gold (Molecular Probes)  
220 made in 0.5 $\times$  TBE. Bands of DNA were visualized under UV light (ethidium bromide) or  
221 using a Dark Reader DR89X Transilluminator (Integrated Scientific Solutions Inc.) (SYBR  
222 Gold).

223

## 224 **2.5 Isolation of DNA from CsCl-purified VLPs**

225 DNA was extracted from 500  $\mu$ l portions of CsCl-purified VLP samples according to  
226 [15]. Prior to extracting DNA from purified VLPs, samples were treated with 20  $\mu$ l of 1  
227 mg/mL DNase I (from bovine pancreas, 552 Kunitz/mg protein; made in water passed  
228 through a 0.1  $\mu$ m filter prior to autoclaving) for 1 h at 37°C, and were then heated at 80°C for  
229 10 min to inactivate the DNase prior to DNA extraction. DNA was air-dried and  
230 resuspended in 50  $\mu$ l of TE buffer. Sterility of samples (i.e. absence of bacterial DNA) was  
231 confirmed by negative PCR from samples (10  $\mu$ l) with universal primers [22]. DNA from an  
232 in-house strain of *Bifidobacterium longum* (4-FAA1; [23]) was used as a positive control;  
233 sterile water was used as the negative control.

234

### 234 3. RESULTS AND DISCUSSION

#### 235 3.1 Examination of VLPs in FFs (faeces) by using EFM

236 Previous studies examining the faecal virome have used 0.22 µm filters to prepare  
237 samples [9, 11, 12, 13]. Preliminary work performed with 0.45 µm-filtered faecal samples  
238 and 0.45 µm-filtered samples originating from *in vitro* fermentation systems inoculated with  
239 faeces stained with 4',6-diamidino-2-phenylindole (DAPI) and viewed under an  
240 epifluorescence microscope suggested that the abundance of dsDNA VLPs in intestinal  
241 samples was high, and that these particles can be enumerated (unpublished observations).  
242 Contamination by bacteria was not observed in any of these samples, and confirmed by an  
243 absence of cultivable bacteria and no bacteria in TEM analyses, so we decided to continue  
244 using larger-pore filters for processing samples. Klieve & Swain [24] had previously used  
245 0.45 µm-filtered samples to characterize VLPs present in rumen contents, and reported no  
246 problems with contamination by bacteria.

247 To examine the VLPs present in FFs (faeces) by EFM, the method described by [9]  
248 and [15] was used initially. However, it was found that we were unable to visualize many, if  
249 any, VLPs present in the samples using 1–5× SYBR Gold as the DNA/RNA stain. Increasing  
250 the concentration of SYBR Gold to 400×, as used by [18] for planktonic aquatic samples,  
251 allowed us to enumerate and detect VLPs present in the FFs prepared from faecal samples of  
252 six donors (Fig. 1 and Fig. 2). The samples were extracted in LL, though the method worked  
253 as well with samples extracted in TBT (data not shown). To determine the relative abundance  
254 of the predominant VLP morphologies in FFs, samples were not concentrated by CsCl  
255 centrifugation (Fig. 1). All FFs were found to be free of bacteria by EFM (Fig. 2), with the  
256 number of VLPs present in samples ranging from  $2.4 \times 10^8$  to  $1.12 \times 10^9$  VLPs/ml FF (mean  
257  $5.58 \times 10^8$  VLPs/ml FF) (equivalent to  $\sim 1.2 \times 10^9$  to  $5.58 \times 10^9$  VLPs/g faeces, mean  $2.94 \times 10^9$ ).  
258 Lepage *et al.* [4] enumerated VLPs in gut mucosal samples from 14 healthy individuals and

259 19 Crohn's disease patients and found on average  $1.2 \times 10^9$  VLPs/biopsy (range  $4.4 \times 10^7$ –  
260  $1.7 \times 10^{10}$ ), in agreement with the mean value we present here for faecal VLPs. Of note,  
261 Crohn's disease patients harboured significantly ( $P = 0.024$ ) more VLPs than healthy  
262 individuals ( $2.9 \times 10^9$  vs  $1.2 \times 10^8$  VLPs/biopsy) in the study of Lepage *et al.* [4].

263 It is generally accepted that there are around 10 bacteriophages for every microbial  
264 cell in environmental samples investigated to date [25]. Extrapolating this figure to the gut  
265 microbiota, from fluorescence *in situ* hybridization studies it is estimated that the faecal  
266 microbiota harbours  $\sim 10^{11}$  bacteria/g faeces in healthy adults [26]. Consequently, one would  
267 expect the presence of at least  $10^{12}$  VLPs/g faeces. Enumerating VLPs in faeces via FFs (or  
268 any liquid medium) by EFM is highly subjective as a dot of very intense fluorescence may in  
269 fact represent a cluster of VLPs (Fig. 2), a phenomenon frequently encountered during this  
270 study. Patel *et al.* [18] stated that, to accurately enumerate VLPs by EFM, micro-adjustments  
271 using the fine focus of the microscope have to be made to ensure that all viruses in a  
272 particular grid-reticle box are counted. We agree with this statement, and furthermore add  
273 that many VLPs have very likely been lost during the preparation and filtering of samples,  
274 either by association with debris in the initial centrifugations or by being caught in the filters  
275 because of clogging or because the VLPs are too big to pass through the pores (e.g. members  
276 of the order *Megavirales* [27]). Even after centrifugation, the supernatants from several of the  
277 samples, while appearing relatively translucent, were highly viscous (perhaps due to host  
278 mucins) and clogged the  $0.45 \mu\text{m}$  filters with less than 2 ml of sample being filtered.  
279 Consequently, the values we provide for the numbers of VLPs in FFs and faecal samples are  
280 a conservative estimate. We believe the true number of VLPs present in faeces to be higher,  
281 possibly between  $10^{10}$  and  $10^{12}$  VLPs/g faeces. Lepage *et al.* [4] determined there to be  $10^{10}$   
282 VLPs/ $\text{mm}^3$  tissue in their study of mucosal VLPs.

283

### 284 3.2 Examination of VLP diversity in FFs by the use of TEM

285 VLPs were readily detected in the TBT-extracted faecal samples from all donors by  
286 TEM. Bacterial cells were never observed in any samples examined by TEM (limit of  
287 detection  $10^6$  per ml). It was striking how visibly different/distinct the VLP assemblages were  
288 between the donors, with no two donors sharing the same VLPs, at least on the basis of  
289 morphological appearance (Fig. 3 and Fig. 4; Supplementary Fig. 1). The vast majority of  
290 VLPs present in the samples appear to represent bacteriophages. Donor 1's VLP assemblage  
291 was predominated by small and large isometric-headed *Siphoviridae* with various tail length  
292 sizes (approx. 120 nm, 350 nm, 650 nm, or, in one extreme case, 1220 nm), with some small  
293 and large isometric-headed *Myoviridae* also present (Fig. 3a). Notably, donor 2's VLP  
294 assemblage was more diverse, predominated by numerous different morphotypes of  
295 *Myoviridae* with *Siphoviridae* also present (Fig. 3b). Interestingly, two detached *Myoviridae*  
296 tails of extreme length (480 nm) and thickness of the sheaths (40 nm) indicated the presence  
297 of giant *Myoviridae* phages (Fig. 3b). Sime-Ngando *et al.* [28] have reported isolation of  
298 bacteriophages with tails of 400 nm in length (heads 50–130 nm) from hypersaline lake  
299 samples, though images of these large bacteriophages are unavailable for direct comparison  
300 of the tails' structures with those found in this study. Donor 2's faecal sample contained  
301 distinct small isometric-headed *Myoviridae* phages with uncommon radial fibers (approx 150  
302 nm in length) attached to the capsids (~70 nm diameter) (Fig. 4). These fibers are clearly  
303 extending the 110-nm tails of these phages, suggesting a primary role in bacteriophage  
304 adsorption. This unique bacteriophage morphotype was also documented in low numbers in  
305 Donor 4's faecal sample (not shown), and in caecal samples L16 (not shown) and L18  
306 (Supplementary Fig. 2d). Fig. 4 also illustrates the apparent clustering of *Myoviridae* phages  
307 in samples. VLPs of various morphotypes derived from Donors 3, 4, 5, and 6 are shown in  
308 Supplementary Figure 1a–d. "Zeppelin"-like VLPs of constant thickness (56 nm) but

309 different lengths (370–630 nm) were detected in Donor 5 (Supplementary Fig. 1c) and Donor  
310 6's samples (Supplementary Fig. 1d). Donor 6's sample was predominated by these  
311 “zeppelin”-like VLPs, with only a few bacteriophages detected. Sime-Ngando *et al.* [28]  
312 reported the presence of rods of up to 22–24 × 1000 nm: similar to the ‘zeppelins’ identified  
313 in the present study, these rods had no visible internal or external structures. The largest  
314 *Myoviridae* phages found in this study were documented for the sample of Donor 5  
315 (Supplementary Fig. 1c), with the phages having prolate heads of 150 × 115 nm and tails of  
316 400 nm in length. The sample from Donor 6 did not reveal the same level of diversity as seen  
317 for the other faecal samples and the reason for this is not known (Supplementary Fig. 1d).  
318 Enumeration of VLPs in the sample of Donor 2 (highest number of VLPs as detected by  
319 TEM) demonstrated there to be approx.  $5 \times 10^7$  VLPs/ml FF.

320 For the majority of the caecal samples, a lower extent of viral biodiversity was  
321 documented. Sample L10 (no clinical data) was unique, containing exclusively small  
322 isometric-headed *Podoviridae* (50 nm diameter) and ~20-nm long appendages (Fig. 5b).  
323 Samples L08 (healthy) (Fig. 5a), and L02 and L03 (both IBD) (Suppl. Fig. 5b, c) appeared to  
324 exclusively contain *Myoviridae* phages. The greatest extent of morphological variation and  
325 the highest bacteriophage numbers were seen in caecal samples L16 (no clinical data) and  
326 L17 (no clinical data) (Fig. 5c, d) and – to a lesser extent – in samples L01 (diverticulosis)  
327 and L18 (no clinical data) (Supplementary Fig. 2a, d). A new type of a giant *Myoviridae*  
328 bacteriophage was exclusively found in sample L16 (Fig. 5c) with isometric heads of 125 nm  
329 in diameter, with tails of 340 nm in length and unique “curled” tail fibers of ~70-nm in  
330 length.

331 Lepage *et al.* [4] demonstrated that gut mucosal samples were predominated by  
332 morphotypes consistent with *Siphoviridae*, *Myoviridae* and *Podoviridae*, and that each

333 individual appeared to be colonized by one dominant bacteriophage family. Our TEM results  
334 for faecal and caecal VLP assemblages are consistent with these results.

335

### 336 **3.3 PFGE of VLPs present in FFs**

337         It was clear from the EFM and TEM analyses that, based on the number of VLPs  
338 present in our samples, it should be possible to isolate large amounts of VLP-derived DNA  
339 from faeces. Attempts to recover abundant VLP DNA from samples using the centrifugation  
340 method of Thurber *et al.* [15], and which was used by [9] and [11] to isolate VLP DNA from  
341 faecal samples, proved disappointing in terms of the recovered DNA yield. However,  
342 increasing the centrifugation speed to 100,000 *g*, routinely used in the laboratory to purify  
343 lactococcal bacteriophages for preparation of high-quality DNA and used by Kulikov *et al.*  
344 [29] to recover bacteriophages from horse faeces, markedly improved recovery of faecal  
345 VLPs.

346         To concentrate VLPs from 20-ml quantities of FF so that they could be applied to  
347 CsCl gradients in 5 ml aliquots, it was decided to use PEG precipitation. This method of  
348 recovery has previously been used with, for example, marine samples and faecal samples to  
349 improve detection of F-specific coliphages in faecal material [16, 17], and allows large  
350 starting volumes of sample to be used for recovering VLPs from human faeces. The method  
351 can be scaled easily so that the VLPs from larger volumes of FF are precipitated for  
352 collection by centrifugation: we used 20 ml of FF from each donor here to demonstrate the  
353 efficacy of the method, but the entire FF derived for each donor's 25 g of starting material  
354 could have been used for PEG precipitation of VLPs.

355         PEG-precipitated samples resuspended in 1 ml TBT were examined using PFGE.  
356 Aliquots (40  $\mu$ l) of the resuspended pellets were heated at 60°C [9] or 75°C ([24] used this  
357 temperature to inactivate nucleases) before PFGE. Profiles containing one or more bands



358 were observed for samples from all donors (Fig. 6). Following treatment at 60°C, Donor 1's  
359 sample produced a faint band at 105 kb, with this band being more prominent following  
360 sample treatment at 75°C; at the higher temperature, bands were also observed at 135 kb and  
361 just below 48.5 kb, with the most prominent band at 79 kb. Donor 2's 60°C-treated sample  
362 produced a profile with a faint smear and a thin band at 84 kb; with the 75°C-treated sample,  
363 the smear was more pronounced, with a thick band observed between 75 and 84 kb. Similar  
364 to Donor 2, the smear in Donor 4's sample became more pronounced following treatment at  
365 the higher temperature. For Donor 3's 60°C-treated sample, two strong bands (at 66 and 97  
366 kb) were observed; in the 75°C-treated samples, bands were observed at 57 kb, 75 kb and 97  
367 kb. The profile of Donor 5 was most affected by changing the temperature at which samples  
368 were treated prior to loading on the gel: the 60°C-treated sample had three bands visible  
369 (most prominent at 100 kb, fainter bands at 48.5 kb and 66 kb), whereas the 75°C-treated  
370 sample had these three bands with at least another 8 bands visible between 48.5 and 97.0 kb.  
371 Donor 6's sample had a faint band just below 48.5 kb after heating at 60°C, whereas the  
372 sample heated at the higher temperature produced two prominent bands that were smaller  
373 than 48.5 kb. The results from the gel were reproducible (triplicate gels run; data not shown).  
374 It is unsurprising that heating at 75°C produced more complex banding patterns, as we would  
375 expect the higher temperature to disassemble (some of) the capsids of VLPs resistant to  
376 heating at 60°C, thereby releasing packaged DNA.

377       Using this method, it was possible to visualize VLP DNA when stained with SYBR  
378 Gold (Fig. 6) or ethidium bromide (Supplementary Fig. 3). It was clear that increasing the  
379 temperature at which the samples were heated prior to loading onto the gel revealed greater  
380 diversity in the samples, and that each individual harboured a unique VLP assemblage (Fig.  
381 4). This is in agreement with the findings of the metagenomic studies of [11] and [12]. The  
382 sample from Donor 6 did not reveal the same level of diversity as seen for the other donors,

383 in agreement with the results shown by TEM (Supplementary Fig. 1d). This donor's sample  
384 was predominated by “zeppelin”-like VLPs that may not have released their nucleid acids by  
385 the conditions employed here.

386 Heating PEG-precipitated samples allowed us to generate PFGE images that  
387 demonstrated that each individual harbours a unique VLP assemblage, and encouraged us to  
388 continue with studies to improve recovery of VLP DNA from human faeces. It also showed  
389 that our method of recovery of VLPs present in faeces was superior to that used previously  
390 by [9] to generate a viral assemblage fingerprint. In that study, the authors stated that the  
391 limited amount of DNA recovered using tangential flow filtration with a 500 g faecal sample  
392 from a 33-year-old made it necessary to enhance the sample bands on the gel relative to the  
393 ladder. In contrast, no enhancement of the bands detected by SYBR Gold (Fig. 6) or ethidium  
394 bromide (Supplementary Fig. 3) was required using the methodology described herein,  
395 although SYBR Gold was, as expected, far more sensitive than ethidium bromide. PEG  
396 precipitation of VLPs present in 20 ml of FF, as used in this study, equates to the extraction  
397 of VLPs from ~4 g faeces (sample losses are seen during filtration, and vary from donor to  
398 donor due to the differences in viscosity of faecal supernatant obtained after removal of most  
399 bacteria and debris from samples after centrifugation), with 50 µl of the resuspended PEG  
400 precipitate representing VLPs isolated from ~200 mg of faeces.

401 The current study used a crude method of extracting DNA for PFGE. Using the  
402 method of Rohozinski *et al.* [30], in which VLPs are embedded in agarose blocks, with CsCl-  
403 purified VLPs prior to heating may allow better ‘fingerprinting’ of VLPs in faecal samples.  
404 This could, for example, be used as an inexpensive means of determining the effect of  
405 freeze–thawing faecal samples prior to the recovery of VLPs from faecal samples. All of the  
406 metagenomic studies conducted to date on the human faecal virome have used samples that  
407 have been frozen prior to recovery of VLPs from samples.

408

409 **3.4 Isolation of high-quality DNA from FFs**

410       Reyes *et al.* [11] and Thurber *et al.* [15] (based on the work of [9]) have reported  
411 recovery of ~500 ng (from 2–5 g of frozen faeces) and 500–3000 ng (from 500 g of fresh  
412 faeces), respectively. The method of Reyes *et al.* [11] is closest to that presented herein for  
413 the recovery of VLPs from faeces [i.e. they hand-filtered samples, whereas Breitbart *et al.* [9]  
414 used tangential flow filtration to process their sample]. As stated above, 20 ml of FF equates  
415 to ~4 g faeces. We processed 500  $\mu$ l portions of 3.5–4 ml CsCl-purified VLPs for DNA  
416 extraction, and resuspended the DNA in 50  $\mu$ l of TE (Fig. 7). If we had processed the entire  
417 PEG-precipitated, CsCl-purified VLP sample for each donor, we would have recovered  
418 between ~1000 and 1800 ng from ~4 g of faeces (theoretically ~6000–11000 ng from 25 g of  
419 faeces). This increase in recovery of DNA is thought to be mainly due to the use of 0.45  $\mu$ m  
420 filters rather than 0.22  $\mu$ m filters to process samples, supported by our observation that  
421 passing FFs through 0.22  $\mu$ m filters led to VLP counts by EFM that were approximately half  
422 of those of the 0.45  $\mu$ m-filtered samples. Increasing filter size and using PEG precipitation to  
423 increase the volume of sample that can be processed would allow greater sampling of the  
424 faecal virome in future metagenomics studies.

425       In addition to demonstrating the utility of the method with faecal samples, we have  
426 successfully isolated VLP DNA from human caecal effluent (Fig. 7). For each of the samples,  
427 20 ml of 0.45  $\mu$ m-filtered caecal filtrate prepared in LL was PEG-precipitated, with the VLPs  
428 then purified on CsCl gradients. These 20-ml aliquots equated to ~4 ml caecal effluent from  
429 each of the three samples examined. We started with 30, 30 and 10 ml, respectively, of  
430 effluent from caecal samples L07 (healthy), L08 (no clinical data) and L10 (no clinical data).  
431 Therefore, if the filtrate from entire homogenates of these samples had been processed, we  
432 would have expected to isolate between ~600 and 8000 ng VLP DNA from caecal effluent.

433 Consequently, it should be possible to conduct metagenomics studies of the VLP  
434 assemblages associated with the human caecum using the methods described herein.

435

### 436 **3.5 Conclusions**

437 We have presented a series of methods for enumerating and characterizing VLPs  
438 present in human faecal and caecal samples. On the basis of enumeration of VLPs using  
439 EFM, there are  $\sim 10^8$  VLPs/ml FF, with faeces thought contain up to  $10^{12}$  VLPs per gram of  
440 sample. TEM analysis of faecal VLPs indicates that an individual is colonized by one  
441 dominant bacteriophage family, with *Myoviridae* and *Siphoviridae* representing the main  
442 families of bacteriophage detected in faeces. PFGE analysis of PEG-concentrated samples  
443 has demonstrated that each individual harbours a unique VLP population, and that the  
444 temperature at which samples are treated greatly affects diversity observed. Using 0.45  $\mu\text{m}$   
445 filters to prepare samples, it is possible to isolate twice as much DNA as reported previously  
446 from  $\sim 4$  g of faeces. The inclusion of PEG precipitation in the methodology means that VLPs  
447 from large volumes of FF can be concentrated, allowing recovery of microgram quantities of  
448 VLP DNA from faecal samples. In addition, PEG precipitation and CsCl purification can be  
449 used to recover VLPs from human caecal samples, with nanogram quantities of VLP DNA  
450 being recovered from the processed samples.

451

452

453

453 **4. ACKNOWLEDGEMENTS**

454 Dr Bernard M. Mackey (University of Reading) is thanked for the use of  
455 epifluorescence microscopy facilities, as is Angela Back (MRI Kiel) for her expert technical  
456 support in electron microscopy.

457

458

458 **5. REFERENCES**

459 [1] Breitbart M, Rohwer F. Here a virus, there a virus, everywhere the same virus? Trends

460 Microbiol 2005;13:278-84.

461 [2] Edwards RA, Rohwer F. Viral metagenomics. Nat Rev Microbiol 2005;3:504-10.

462 [3] Rohwer F. Global phage diversity. Cell 2003;113:141.

463 [4] Lepage P, Colombet J, Marteau P, Sime-Ngando T, Doré J, LeClerc M. Dysbiosis in

464 inflammatory bowel disease: a role for bacteriophages? Gut 2008;57:424-25.

465 [5] Vishwakarma V, Periaswamy B, Bhusan Pati N, Slack E, Hardt WD, Suar M. A novel

466 phage element of *Salmonella* enterica serovar Enteritidis P125109 contributes to accelerated

467 type III secretion system 2-dependent early inflammation kinetics in a mouse colitis model.

468 Infect Immun 2012;80:3236-246.

469 [6] Wagner J, Maksimovic J, Farries G, Sim WH, Bishop RF, Cameron DJ, et al.

470 Bacteriophages in gut samples from pediatric Crohn's disease patients: metagenomic analysis

471 using 454 pyrosequencing. Inflamm Bowel Dis 2013;19:1598-608.

472 [7] Watanabe R, Matsumoto T, Sano G, Ishii Y, Tateda K, Sumiyama Y., et al. Efficacy of

473 bacteriophage therapy against gut-derived sepsis caused by *Pseudomonas aeruginosa* in

474 mice. Antimicrob Agents Chemother 2006;51:446-52.

475 [8] Meader E, Mayer MJ, Steverding D, Carding SR, Narbad A. Evaluation of bacteriophage

476 therapy to control *Clostridium difficile* and toxin production in an *in vitro* human colon

477 model system. Anaerobe 2013;22:25-30.

478 [9] Breitbart M, Hewson I, Felts B, Mahaffy JM, Nulton J, Salamon P, et al. Metagenomic

479 analyses of an uncultured viral community from human feces. J Bacteriol 2003;185:6220-

480 223.

481 [10] Zhang T, Breitbart M, Lee WH, Run JQ, Wei CL, Soh SW, et al. RNA viral community

482 in human feces: prevalence of plant pathogenic viruses. PLoS Biol 2006;4:e3.

- 483 [11] Reyes A, Haynes M, Hanson N, Angly FE, Heath AC, Rohwer F, et al. Viruses in the  
484 faecal microbiota of monozygotic twins and their mothers. *Nature* 2010;466:334-38.
- 485 [12] Minot S, Sinha R, Chen J, Li H, Keilbaugh SA, Wu GD, et al. The human gut virome:  
486 inter-individual variation and dynamic response to diet. *Genome Res* 2011;21:1616-625.
- 487 [13] Minot S, Bryson A, Chehoud C, Wu GD, Lewis JD, Bushman FD Rapid evolution of the  
488 human gut virome. *Proc Natl Acad Sci U S A* 2013;110:12450-5.
- 489 [14] Modi SR, Lee HH, Spina CS, Collins JJ. Antibiotic treatment expands the resistance  
490 reservoir and ecological network of the phage metagenome. *Nature* 2013;499:219-22.
- 491 [15] Thurber RV, Haynes M, Breitbart M, Wigley L, Rohwer F. Laboratory procedures to  
492 generate viral metagenomes. *Nat Protoc* 2009;4:470-83.
- 493 [16] Jones TH, Johns MW. Improved detection of F-specific RNA coliphages in fecal  
494 material by extraction and polyethylene glycol precipitation. *Appl Environ Microbiol*  
495 2009;75:6142-6.
- 496 [17] Colombet J, Robin A, Lavie L, Bettarel Y, Cauchie HM, Sime-Ngando T. Virioplankton  
497 'pegylation': use of PEG (polyethylene glycol) to concentrate and purify viruses in pelagic  
498 ecosystems. *J Microbiol Methods* 2007;71:212-9.
- 499 [18] Patel A, Noble RT, Steele JA, Schwalbach MS, Hewson I, Fuhrman JA. Virus and  
500 prokaryote enumeration from planktonic aquatic environments by epifluorescence  
501 microscopy with SYBR Green I. *Nat Protoc* 2007;2:269-76.
- 502 [19] Wen K, Ortmann AC, Suttle CA. Accurate estimation of viral abundance by  
503 epifluorescence microscopy. *Appl Environ Microbiol* 2004;70:3862-7.
- 504 [20] Mills S, Ross PR, Neve H, Coffey A. Bacteriophage and anti-phage mechanisms in  
505 lactic acid bacteria, in Salminen S, von Wright A, Lahtinen S, Ouwehand AC (Eds). *Lactic*  
506 *Acid Bacteria: Microbial and Functional Aspects*, 4th edition, Taylor & Francis, London,  
507 2011, pp 165-86.

- 508 [21] Fuhrman, JA, Griffith JF, Schwalbach, MS. Prokaryotic and viral diversity patterns in  
509 marine plankton. *Ecol Res* 2002;17:183-94.
- 510 [22] Hoyles L, Collins MD, Falsen E, Nikolaitchouk N, McCartney AL. Transfer of members  
511 of the genus *Falcivibrio* to the genus *Mobiluncus*, and amended description of the genus  
512 *Mobiluncus*. *Syst Appl Microbiol* 2004;27:72-83.
- 513 [23] Hoyles, L. *In vitro* examination of the effect of orlistat on the ability of the faecal  
514 microbiota to utilize dietary lipids. 2009, PhD thesis, University of Reading, UK.
- 515 [24] Klieve AV, Swain RA. Estimation of ruminal bacteriophage numbers by pulsed-field gel  
516 electrophoresis and laser densitometry. *Appl Environ Microbiol* 1993;59:2299-303.
- 517 [25] Brüssow H, Hendrix RW. Phage genomics: small is beautiful. *Cell* 2002;108:13-6.
- 518 [26] Hoyles L, McCartney AL. What do we mean when we refer to *Bacteroidetes*  
519 populations in the human gastrointestinal microbiota? *FEMS Microbiol Lett* 2009;299:175-  
520 83.
- 521 [27] Colson P, Fancello L, Gimenez G, Armougom F, Desnues C, Fournous G, et al.  
522 Evidence of the megavirome in humans. *J Clin Virol* 2013;57:191-200.
- 523 [28] Sime-Ngando T, Lucas S, Robin A, Tucker KP, Colombet J, Bettarel Y, et al. Diversity  
524 of virus-host systems in hypersaline Lake Retba, Senegal. *Environ Microbiol* 2011;13:1956-  
525 1972.
- 526 [29] Kulikov EE, Isaeva AS, Rotkina AS, Manykin AA, Letarov AV. Diversity and dynamics  
527 of bacteriophages in horse feces. *Mikrobiologiya* 2007;76:271-78.
- 528 [30] Rohozinski J, Girton LE, Van Etten JL. *Chlorella* viruses contain linear nonpermuted  
529 double-stranded DNA genomes with covalently closed hairpin ends. *Virology* 1989;168:363-  
530 69.
- 531



531 **Fig. 1.** Number of VLPs detected in FFs by using EFM. Values are shown as mean + SD ( $n =$   
532 2 per donor). Passing samples through a 0.22  $\mu\text{m}$  filter reduced the number of VLPs present  
533 in samples by approximately half (data not shown). White bars, per ml FF (faeces); grey bars,  
534 per g faeces.

535

536

536 **Fig. 2.** Image showing faecal VLPs as they appear under an epifluorescence microscope  
537 ( $\times 1000$  magnification). Bacteriophages appear as ‘pinpricks’ of light when stained with  
538 SYBR Gold. The brightest ‘pinpricks’ (black arrows) represent clusters of up to 30 VLPs.  
539 The black dot (shown in the white circle) on the images is due to a scratch on the microscope  
540 lens. (a) Contamination on filter, which gives an idea of the size and appearance of a  
541 bacterium in comparison with VLPs. (b) Negative control, 0.1  $\mu\text{m}$ -filtered, autoclaved  $\text{H}_2\text{O}$ .  
542 (c) Negative control, sterile 0.5 % ‘Lab-Lemco’/6 % NaCl. (d) Donor 1, 0.22  $\mu\text{m}$ -filtered  
543 sample; (e) Donor 1 0.45  $\mu\text{m}$ -filtered sample. (f) Donor 2, 0.22  $\mu\text{m}$ -filtered sample; (g)  
544 Donor 2, 0.45  $\mu\text{m}$ -filtered sample. Scale bar, 10  $\mu\text{m}$ .

545

545 **Fig. 3.** Transmission electron micrographs revealing the diversity of VLPs found in faeces.  
546 (a) Donor 1 (female, 41 years) and (b) Donor 2 (female, 36 years) after extraction of VLPs in  
547 TBT buffer.  
548

548 **Fig. 4.** Transmission electron micrographs of VLPs found in faeces of Donor 2 (female, 36  
549 years) after extraction of VLPs in TBT buffer showing *Myoviridae* phages with radial  
550 whiskers attached to the capsids (top) and clusters of *Myoviridae* phages adsorbing to  
551 membrane vesicle material (bottom).  
552

552 **Fig. 5.** Transmission electron micrographs revealing the diversity of VLPs found in caecal  
553 effluents. Sample (a) L08, (b) L10, (c) L16 and (d) L17 after extraction of VLPs in LL. No  
554 clinical data were available for these individuals.

555

555 **Fig. 6.** Use of PFGE to examine VLP populations in PEG-precipitated FFs (faeces), and  
556 demonstration that increasing the temperature at which samples are heated prior to loading  
557 onto the gel can affect the diversity uncovered. The gel was stained with SYBR Gold and  
558 visualized as described in Methods. This is a crude (but inexpensive) method of examining  
559 VLP populations in human faeces, and demonstrates that each individual harbours a unique  
560 VLP profile. Ladder, lambda ladder (#340; New England Biolabs).

561

561 **Fig. 7.** Isolation of high-quality DNA from CsCl-purified samples of human gastrointestinal  
562 VLPs. VLP preparations were CsCl-purified and DNA extracted as described in Methods.  
563 Aliquots (10  $\mu$ l) of sample were run on a 0.8 % agarose gel at 90 V for 30 min. Lane 1,  
564 molecular size ladder; lane 2,  $\phi$ KPLN1 positive control (a bacteriophage isolated from the  
565 human caecum that infects *Klebsiella pneumoniae* subsp. *pneumoniae* K2 strains; L. Hoyles,  
566 unpublished data), 63 ng DNA/ $\mu$ l; lane 3, VLP extract from faeces of Donor 4, 3.1 ng  
567 DNA/ $\mu$ l; lane 4, VLP extract from faeces of Donor 3, 4.5 ng DNA/ $\mu$ l; lane 5, VLP extract  
568 from caecal sample L07 (healthy), 0.2 ng DNA/ $\mu$ l; lane 6, VLP extract from caecal sample  
569 L08 (no clinical data), 2.7 ng DNA/ $\mu$ l; lane 7, VLP extract from caecal sample L10 (no  
570 clinical data), 3.8 ng DNA/ $\mu$ l. The two black arrows highlight the presence of RNA in the  
571 sample from Donor 3, confirmed by treating the CsCl-purified sample with 20  $\mu$ l of 1 mg/mL  
572 RNase A (bovine pancreas,  $\geq$ 70 Kunitz/mg protein, prepared in 0.1  $\mu$ m-filtered, sterile H<sub>2</sub>O)  
573 with the DNase in a second DNA extraction (not shown). VLPs and DNA were extracted  
574 from human caecum samples (1:4 LL, v/v) as described in Methods as part of a study  
575 examining the microbiota associated with the human caecum (L. Hoyles, unpublished data).  
576 With the exception of the sample in lane 5, DNA visible to the naked eye was extracted from  
577 all samples of gastrointestinal origin.

578