

**UNIVERSITY OF
LEADING
THE WAY
WESTMINSTER** 

WestminsterResearch

<http://www.westminster.ac.uk/research/westminsterresearch>

Identification of *Schistosoma mansoni* microRNAs.

Mariana C Simões^{1,6}, Jonathan Lee², Appolinaire Djikeng^{3,8}, Gustavo C Cerqueira^{4,5}, Adhemar Zerlotini^{1,6}, Rosiane A da Silva-Pereira⁶, Andrew R Dalby^{2*}, Philip LoVerde⁷, Najib M El-Sayed⁴, Guilherme Oliveira⁶

1 Graduate Program in Bioinformatics, Universidade Federal de Minas Gerais

2 Department of Statistics, Peter Medawar Building, South Parks Road, Oxford, UK.

3 J Craig Venter Institute (JCVI)

4 Department of Cell Biology and Molecular Genetics and Center for Bioinformatics and Computational Biology, University of Maryland

5 Institute for Genome Sciences, University of Maryland

6 CEBio, Instituto Nacional de Ciência e Tecnologia em Doenças Tropicais, Laboratory of Cellular and Molecular Parasitology, Centro de Pesquisas René Rachou, Fundação

7 University of Texas Health Science Center

8 Biosciences eastern and central Africa - International Livestock Research Institute (BecA-ILRI) Hub

*Andrew Dalby now works within the School of Life Sciences, University of Westminster

This is an electronic copy of an article originally published in *BMC Genomics* and is made available under the Creative Commons Attribution License. A copy of the license can be downloaded here:

<http://www.biomedcentral.com/html/info/about/license.pdf>

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Users are permitted to download and/or print one copy for non-commercial private study or research. Further distribution and any use of material from within this archive for profit-making enterprises or for commercial gain is strictly forbidden.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail repository@westminster.ac.uk

RESEARCH ARTICLE

Open Access

Identification of *Schistosoma mansoni* microRNAs

Mariana C Simões^{1,6}, Jonathan Lee², Appolinaire Djikeng^{3,8}, Gustavo C Cerqueira^{4,5}, Adhemar Zerlotini^{1,6}, Rosiane A da Silva-Pereira⁶, Andrew R Dalby², Philip LoVerde⁷, Najib M El-Sayed⁴, Guilherme Oliveira^{6*}

Abstract

Background: MicroRNAs (miRNAs) constitute a class of single-stranded RNAs which play a crucial role in regulating development and controlling gene expression by targeting mRNAs and triggering either translation repression or messenger RNA (mRNA) degradation. miRNAs are widespread in eukaryotes and to date over 14,000 miRNAs have been identified by computational and experimental approaches. Several miRNAs are highly conserved across species. In *Schistosoma*, the full set of miRNAs and their expression patterns during development remain poorly understood. Here we report on the development and implementation of a homology-based detection strategy to search for miRNA genes in *Schistosoma mansoni*. In addition, we report results on the experimental detection of miRNAs by means of cDNA cloning and sequencing of size-fractionated RNA samples.

Results: Homology search using the high-throughput pipeline was performed with all known miRNAs in miRBase. A total of 6,211 mature miRNAs were used as reference sequences and 110 unique *S. mansoni* sequences were returned by BLASTn analysis. The existing mature miRNAs that produced these hits are reported, as well as the locations of the homologous sequences in the *S. mansoni* genome. All BLAST hits aligned with at least 95% of the miRNA sequence, resulting in alignment lengths of 19-24 nt. Following several filtering steps, 15 potential miRNA candidates were identified using this approach. By sequencing small RNA cDNA libraries from adult worm pairs, we identified 211 novel miRNA candidates in the *S. mansoni* genome. Northern blot analysis was used to detect the expression of the 30 most frequent sequenced miRNAs and to compare the expression level of these miRNAs between the lung stage schistosomula and adult worm stages. Expression of 11 novel miRNAs was confirmed by northern blot analysis and some presented a stage-regulated expression pattern. Three miRNAs previously identified from *S. japonicum* were also present in *S. mansoni*.

Conclusion: Evidence for the presence of miRNAs in *S. mansoni* is presented. The number of miRNAs detected by homology-based computational methods in *S. mansoni* is limited due to the lack of close relatives in the miRNA repository. In spite of this, the computational approach described here can likely be applied to the identification of pre-miRNA hairpins in other organisms. Construction and analysis of a small RNA library led to the experimental identification of 14 novel miRNAs from *S. mansoni* through a combination of molecular cloning, DNA sequencing and expression studies. Our results significantly expand the set of known miRNAs in multicellular parasites and provide a basis for understanding the structural and functional evolution of miRNAs in these metazoan parasites.

Background

Small non-coding RNAs are increasingly providing insights into important aspects of the biology of many organisms [1,2]. They include small interfering RNAs (siRNAs) and microRNAs (miRNAs), which are hallmarks of two important processes involved in RNA

silencing [3,4]. RNA silencing is a general process in which small RNA molecules derived from precursor dsRNA molecules trigger sequence-specific repression of gene expression [4-6].

miRNAs comprise a family of non-coding RNAs with approximately 21-25 nucleotides that down-regulate gene expression at the post-transcriptional level. miRNAs are generated from endogenous hairpin structures in the nucleus and play an important role in controlling diverse cellular functions in eukaryotes, including cell differentiation, development, apoptosis, and genome

* Correspondence: oliveira@cqrr.fiocruz.br

⁶CEBio, Instituto Nacional de Ciência e Tecnologia em Doenças Tropicais, Laboratory of Cellular and Molecular Parasitology, Centro de Pesquisas René Rachou, Fundação Oswaldo Cruz, Av. Augusto de Lima 1715, Belo Horizonte, 30190-002, Brazil

Full list of author information is available at the end of the article

integrity [7-9]. *In vivo* experiments indicate a crucial role in cell proliferation and cell death processes for some miRNAs, including *lin-4* and *let-7* in *C. elegans*; *bantam* and *mir-14* in *Drosophila*; and *mir-23* in humans [10].

The current understanding of miRNA biogenesis involves a series of coordinated processes. Briefly, primary transcripts of miRNAs are processed in the nucleus by Drosha, an RNase III-like enzyme into pre-miRNA, which are first exported into the cytoplasm by exportin-5 and then processed into miRNAs by Dicer, another type III RNase [11-13].

The primary method of identifying miRNA genes has been to isolate, reverse transcribe, clone, and sequence small RNA molecules [14-16]. In animals, discovery of miRNA genes, by using molecular cloning based methods has been supplemented by systematic computational approaches that identify evolutionarily conserved miRNA genes. Bioinformatics tools search for patterns of sequence and secondary structure conservation that are characteristic of metazoan miRNA hairpin precursors [17-19]. However, considerable filtering must be performed to elucidate likely miRNA candidates. The 5' end of miRNAs is reported to have a perfect base alignment of at least 7 consecutive nucleotides, which enables their identification [14]. The most sensitive of these methods indicate that miRNAs constitute nearly 1% of all predicted genes in nematodes, flies, and mammals [19-21]. However, computationally predicted miRNAs must be experimentally confirmed.

Although the first miRNA was identified in 1993, it was not until 2001 that the breadth of the miRNA gene class was recognized with cloning and sequencing of more than one hundred miRNAs from worms, humans, mice, and other species [22,23]. However, no large-scale identification of miRNAs has been carried out in *Schistosoma mansoni*.

Schistosoma mansoni is a human parasite that is responsible for the neglected tropical disease schistosomiasis. The parasite infects approximately 90 million people worldwide, causing morbidity and eventually death in Central and South America and Africa [24]. Although schistosomicidal drugs and other control measures exist, the development of new control strategies is necessary. In recent years, increasing attention has emerged over siRNAs as therapeutic agents [25]. The emergence of gene ablation technologies based on the RNAi phenomenon has opened up new experimental opportunities. Recently, several reports on the use of RNAi for the studies of schistosomes were published [26,27]. In this context, we attempted to identify potential miRNAs in *S. mansoni*. We use complementary experimental and computational approaches. We developed a homology-filtering approach used in a high-

throughput pipeline in which all known miRNA genes were used as reference miRNAs. Fifteen potential miRNA candidates were discovered in *S. mansoni* using this analysis. The pipeline automated some of the manual steps, in particular a rule-based filtering approach for extracting the candidate pre-miRNA sequence, and it can also be applied to other genomes. By sequencing small-RNA cDNA libraries, we provide experimental evidence for 211 potential miRNAs candidates. The identification of new miRNA in the *S. mansoni* genome presents relevant information that is likely to be important for parasite development and sexual maturation.

Results and Discussion

Experimental identification of miRNAs

Cloning of short RNAs from S. mansoni adult worm pairs

An adult worm cDNA library of small RNAs was constructed using an established method based on a sequential ligation of oligonucleotide adapters to a size-fractionated sample of small RNAs [28]. Concatenated DNA fragments (each fragment from one putative miRNA) were cloned into a plasmid vector to generate a library. A total of 582 recombinant clones randomly selected from the library were sequenced. Twelve hundred sequences were analyzed and show to contain ~2-3 small RNA sequence in the same vector. Size distribution of the non-redundant miRNA set ranged from 17 to 25 nt, although the majority contained 20-24 nt, 21 nt being the most abundant. To identify the putative origin of the cloned sequences, a FASTA search was performed against GenBank <http://www.ncbi.nlm.nih.gov> and the *S. mansoni* genome (version 4.0) [29]. Sequences that had significant homology to breakdown products of abundant non-coding (nc) RNAs such as rRNA and tRNA were eliminated. A total of 584 ncRNAs were grouped into 211 clusters and were identified as possible miRNA candidates (see additional file 1, Table S1: clustering of 584 sequenced miRNAs). One hundred and sixty-one miRNAs were represented in the library by only one read and 50 were represented by clusters with up to 32 sequences. Since miRNAs are believed to occur at a frequency of approximately 0.5-1.5% of the total genes in the genome, the 13,200 genes predicted for *S. mansoni* should have generated between 66 and 198 miRNAs [30,31]. Thus, the number of miRNAs experimentally observed is in the expected range.

We further screened the candidate sequences against a database of known miRNAs, miRBase (<http://microrna.sanger.ac.uk>; release 13.0) to compare our candidate *S. mansoni* miRNAs to miRNAs from different species. Some miRNAs showed a high degree of conservation. Forty-two sequences had at least one match with mature miRNAs from different metazoan miRNA families, such as miR-832, miR-71, miR-297, and let-7 (Table 1). For

Table 1 Small *S. mansoni* RNAs with matches to known microRNAs present in the miRBase


<i>S. mansoni</i> miRNAs	Number of Hits	Best Hit Name	miRNA family
sma-miR-7	4	gma-miR171b	miR-171
sma-miR-9	2	ptr-miR-1303	miR-1303
sma-miR-16	1	gga-miR-1465	miR-1465
sma-miR-17	10	dre-miR-739	miR-197
sma-miR-21	1	ebv-miR-BART2	miR-BART2
sma-miR-24	10	ptr-miR-451	miR-451
sma-miR-31	10	hsa-miR-513b	miR-513
sma-miR-32	1	ath-miR832-5p	miR-832
sma-miR-33	3	tca-miR-71	miR-71
sma-miR-36	10	tca-miR-87	miR-87
sma-miR-37	1	cbr-miR-240	miR-240
sma-miR-40	4	ptr-miR-432	miR-432
sma-miR-47	1	sme-let-7c	let-7
sma-miR-77	1	ptc-miR472b	miR472
sma-miR-78	1	ath-miR832-5p	miR-832
sma-miR-86	1	hsa-miR-1268	miR-1268
sma-miR-96	10	ptr-let-7b	let-7
sma-miR-103	1	kshv-miR-K12-3	miR-K12
sma-miR-105	8	dya-miR-289	miR-289
sma-miR-122	1	gga-miR-1465	miR-1465
sma-miR-129	8	dya-miR-289	miR-289
sma-miR-141	2	osa-miR166i	miR-166
sma-miR-149	1	gga-miR-1810	miR-1810
sma-miR-150	1	mmu-miR-297b-3p	miR-297
sma-miR-156	3	tni-miR-101b	miR-101
sma-miR-170	1	cel-miR-78	miR-78
sma-miR-182	10	dre-miR-739	miR-739
sma-miR-187	1	mmu-miR-297b-3p	miR-297
sma-miR-205	1	ppt-miR896	miR-896
sma-miR-209	1	mmu-miR-297b-3p	miR-297
sma-miR-210	1	gga-miR-1810	miR-1810
sma-miR-213	1	ath-miR832-5p	miR-832
sma-miR-216	1	kshv-miR-K12-3	miR-K12
sma-miR-226	1	mmu-miR-297b-3p	miR-297
sma-miR-234	1	gga-miR-1810	miR-1810
sma-miR-241	1	odi-miR-1500	miR-1500
sma-miR-247	4	ptr-miR-432	miR-432
sma-miR-262	1	sme-let-7c	let-7
sma-miR-276	1	mmu-miR-297b-3p	miR-297
sma-miR-278	1	gga-miR-1810	miR-1810
sma-miR-283	2	ptr-miR-1299	miR-1299
sma-miR-284	1	mmu-miR-297b-3p	miR-297

example, sma-miR-36 perfectly matched miRNA family miR-87 from different species demonstrating miRNA conservation among more than 10 species. Previous studies in *C. elegans* showed that this miRNA family is expressed throughout development [20]. The yield of probable miRNA candidates was much lower for this

analysis with *S. mansoni* than analyses of species that contain closer relatives in miRBase. The closest relative to *S. mansoni* in miRBase is *Schmidtea mediterranea*. These two organisms belong to the same phylum, a relatively broad classification. The sequences that did not match any of the known miRNAs (170 sequences) were considered to be putative members of novel families of schistosome miRNAs.

Expression analysis of miRNAs in *S. mansoni*

The expression of miRNAs is tightly regulated in both time and space. Stage-specific or regulated miRNA expression suggests a role in development [20,21]. While high throughput techniques, such as microarray and next generation sequencing are being used, northern blot still remains the consensus method for validating miRNAs [32]. The frequency of reads of a specific miRNA in a non-normalized library can also be correlated with the expression level of that miRNA [33,34]. Based on that, the most abundant candidate miRNAs were further examined by northern blot to test for expression. Northern blots of total RNA from a mixture of male and female *S. mansoni* adult worms and the intramammalian larval stage, schistosomula, were hybridized with biotin-labeled probes. Previous studies analyzed the expression pattern of the Dicer gene during different life stages of *S. mansoni*. A threefold higher expression level was detected in seven day old schistosomula in comparison to the adult worm pairs [35]. It is possible that higher Dicer gene expression at this time was selected for the control of retrotransposon activation that may be more prone to occur during this period of active larval cell division and growth [36]. We detected the expression for 11 of 30 miRNAs in at least one of the 2 analyzed stages (Figure 1). We also analyzed in *S. mansoni* the expression of the five novel miRNAs recently identified in *S. japonicum* (sja-let-7, sja-miR-71, sja-bantam, sja-miR-125 and sja-miR-new1) [37]. Three (sja-miR-71 - non-specific, sja-bantam - schistosomula specific and sja-miR-125 - adult worm specific) of the five probes had a hybridization signal that was characteristic of miRNAs, demonstrating evolutionary conservation. Although the expression of sja-miR-71 and sja-bantam dropped quickly in *S. japonicum* lung-stage schistosomulum, we observed a strong hybridization signal for both miRNAs in *S. mansoni* (Figure 1) [37]. The other 2 candidates detected in *S. japonicum* (new-1 and let-7) may be expressed in other life cycle stages or in undetectable amounts in *S. mansoni* in the life cycle stages tested. We observed 2 miRNAs (mir-2 and mir-71) expressed in both life cycle stages tested, 7 in adult worms only (mir-4, mir-6, mir-9, mir-32, mir-125, mir-3, mir-5) and 5 in schistosomula only (mir-20, mir-18, mir-22, mir-26, Bantam). These results suggest a role for these miRNAs over the life cycle stages of



S. mansoni possibly mediating important processes in the parasite growth and development.

Sequence analysis indicated miR-1 as the most abundant miRNA (32 reads). Although sequencing-based miRNAs expression profiling is a tool for measuring the relative abundance of miRNAs, the expression of the miR-1 was not detected by northern blot. In contrast, miR-32 is represented only by 2 clones in our sequences, which indicates a 12-fold lower expression level compared to that of miR-3 (24 reads). However, our small RNA blot analysis indicated that miR-32 was more abundantly expressed than miR-3 in the adult worm stage (Figure 1). The discrepancies between the cloning frequency and small RNA blot results could not be attributed to variations in RNA content because the same RNA samples were used for both experiments. One possible

explanation could be bias in cloning efficiencies, or differential turnover rates of these miRNAs [38].

The best method to differentiate miRNA from other endogenous small RNA is the ability of flanking sequences to adopt a pre-miRNA fold-back structure with the mature miRNA properly positioned within one of its strands enabling Dicer processing [17]. Eleven (36%) of the 30 potential miRNA detected by northern blot were mapped to ~500 different locations on the genome. To assess which of the regions corresponded to the real location of the possible miRNA gene, their secondary structures were studied using the Vienna RNA-fold package <http://rna.tbi.univie.ac.at/cgi-bin/RNAfold.cgi>. Each image generated was visually inspected. A non-redundant set of 26 potential miRNA sequences were predicted to be capable of forming stem-loop structures characteristic of miRNA precursors, 11 of them were also confirmed by northern blot (Figure 2, see additional file 2, Figure S1: miRNA structures not confirmed by northern blot). Our results also show that multiple hairpin precursors for the same miRNA were observed in more than one location in the parasite genome (data not shown), pointing to the possibility that the same mature miRNA may be transcribed from more than one miRNA gene. Next, the miRNA genomic location was analyzed by BLAST against the *S. mansoni* genome. The selected miRNAs genes were observed to be located on intergenic regions, in agreement with published results [39-41].


Computational identification of miRNAs

Homology search

The high-throughput homology search pipeline was performed with all known miRNAs in miRBase (release 13.0). In total, 6,211 mature miRNAs were used as reference sequences. The e-value cutoff for this analysis was set at 0.01. A total of 180 hits were registered. We observed 110 unique *S. mansoni* sequences, and 15 sequences were represented multiple times. For the BLASTn results see additional file 3, Table S2: high-throughput pipeline homology search results. The existing mature miRNAs that produced these hits are reported, as well as the locations of the homologous sequences in the *S. mansoni* genome. All hits aligned with at least 95% of the miRNA sequence, resulting in alignment lengths of 19-24 nt. All of the 110 unique mature miRNA candidates returned by the BLASTn search were assigned an analysis identifier with prefix 'SMan'.

RNA folding

The extended sequences (with additional 50 nt flanking the mature sequence) were folded with RNASHAPES [42]. For the complete results for the extended sequence folding see additional file 4, Table S3: high-throughput pipeline extended sequence folding results.



Mean free energy (MFE) is a widely used criterion for filtering RNA folding results, and was observed to be an important filtering step in this analysis as well. The rationale for how MFE thresholds are derived, however, is not obvious when examining the literature. In fact, the guidelines proposed for uniform determination and annotation of miRNAs given by Ambros *et al.* do not mention MFE thresholds. Instead, the guidelines merely suggest that to be considered a miRNA, a candidate's lowest MFE fold should be a hairpin [17].

Determination of MFE threshold is dependent on tolerance to false positives, i.e. higher MFE thresholds result in inclusion of more candidates. A threshold of -20 kcal/mol has been generally used, but levels as high as -12 kcal/mol have also been explored [43]. We used a

middle value of -15 kcal/mol as this genome has not been previously explored.

From the 110 unique *S. mansoni* sequences returned from the BLASTn search, 66 displayed MFE values of -15 kcal/mol or less when folded with RNashapes. Forty three hairpins had MFE values greater than -15 kcal/mol.

All 66 of the extended sequences with MFE \leq -15 kcal/mol displayed hairpins in at least one portion of the sequence when folded by RNashapes. At ~122nt in length, the sequence is considerably longer than a typical miRNA hairpin, and as a result, only the ~70nt surrounding the mature sequence are of interest. This region was considered the candidate pre-miRNA sequence. In each of the 66 hairpins detected, the region surrounding the mature sequence was within a hairpin.

Of the 66 hairpins with MFE \leq -15 kcal/mol, 36 structures contained the mature sequence entirely within the stem. The other 30 sequences contained the mature sequence partially or completely in a loop.

Candidate pre-miRNA sequences were extracted from the 36 remaining hairpins and were refolded with RNashapes. After refolding, 15 candidate pre-miRNAs had MFE \leq -15 kcal/mol. These 15 sequences were considered to be likely pre-miRNA sequences. A summary of the results and the structures for the pre-miRNAs candidates are shown in Figures 3, 4, 5, 6 and 7.

The yield of probable miRNA candidates was much lower for this analysis with *S. mansoni* than analyses of species that contain closer relatives in miRBase. The findings suggest that it may have been difficult to find a large number of miRNAs in this analysis, due to the possibility of a large amount of sequence divergence between *S. mansoni* and its closest relative in miRBase is *S. mediterranea*. On the contrary, if one was interested in studying miRNAs in a mammal not found in miRBase, one would find 22 members from the same class and over 2,500 miRNA sequences in miRBase. However, for now, organisms such as *S. mansoni* must continue to have a mix of computational and experimental approaches, with an emphasis on experimental discovery.

Comparison of results to existing work

The high-throughput pipeline yielded fifteen probable pre-miRNA candidates. This number is comparable to the number of miRNAs found by Palokodeti *et al.*, who identified ten miRNA candidates in *S. mediterranea* by using all known human, *Drosophila* and *C. elegans* miRNAs as reference sequences [44]. However, the yields for the *S. mediterranea* analysis and for this analysis with *S. mansoni* are considerably smaller than other studies that have used homology methods with multiple genomes. Luo *et al.* identified 118 miRNAs in *Tribolium castaneum* (red flour beetle) using all available metazoan miRNAs as reference sequences [45]. Zhou *et al.* using a




Figure 3 Probable miRNA candidates SMan124, SMan30 and SMan119 identified by homology search. For each structure, the location in the *S. mansoni* genome (version 4.0), pre-miRNA sequence, miRNA family, sequence length and MFE are given. The start and end of the mature sequence are circled in the structure. The mature sequence is also bolded and underlined in the pre-miRNA sequence.






Figure 4 Probable miRNA candidates SMan65, SMan36 and SMan125 identified by homology search. For each structure, the location in the *S. mansoni* genome (version 4.0), pre-miRNA sequence, miRNA family, sequence length and MFE are given. The start and end of the mature sequence are circled in the structure. The mature sequence is also bolded and underlined in the pre-miRNA sequence.




Figure 5 Probable miRNA candidates SMan64, SMan48 and SMan60 identified by homology search. For each structure, the location in the *S. mansoni* genome (version 4.0), pre-miRNA sequence, miRNA family, sequence length and MFE are given. The start and end of the mature sequence are circled in the structure. The mature sequence is also bolded and underlined in the pre-miRNA sequence.






Figure 6 Probable miRNA candidates SMan133, SMan92 and SMan108 identified by homology search. For each structure, the location in the *S. mansoni* genome (version 4.0), pre-miRNA sequence, miRNA family, sequence length and MFE are given. The start and end of the mature sequence are circled in the structure. The mature sequence is also bolded and underlined in the pre-miRNA sequence.




Figure 7 Probable miRNA candidates SMan82, SMan42 and SMan75 identified by homology search. For each structure, the location in the *S. mansoni* genome (version 4.0), pre-miRNA sequence, miRNA family, sequence length and MFE are given. The start and end of the mature sequence are circled in the structure. The mature sequence is also bolded and underlined in the pre-miRNA sequence.

homology-based computational approach, found 300 human miRNA homologs in the domestic dog using only human miRNAs as the reference miRNAs [46]. Furthermore, Baev et al. identified 639 chimpanzee miRNAs with a homology-based approach, also using only human pre-miRNA sequences as a reference set [47]. Recently, novel miRNAs were identified in *Schistosoma japonicum*, a close relative of *S. mansoni* [37,48-50]. Chatterjee and Chaudhuri detected 489 homologous miRNA sequences in the *Anopheles gambiae* genome, using only *Drosophila* miRNA as the query sequence [51]. *Drosophila* is a well represented group of organisms in miRBase, but it is also in the same order as *A. gambiae*. As a result, this close relationship produced a large number of hits. Vertebrates, especially mammals, are currently the most represented organisms in miRBase, with respect to number of species and number of miRNA sequences.

It is worthwhile to note the extent to which the yield decreases as the distance between species grows. As a result, these findings suggest that the yield of a homology-based analysis is very dependent on the available content of miRBase. Artzi et al. used their recently released homology search web-server, miRNAMiner, to increase the number of miRBase miRNAs for seven mammals by 50%, identifying 790 new miRNAs [52]. The strategy and filtering steps used by miRNAMiner are very similar to those used in this paper, but moderately more comprehensive.

Analysis of homology search hits by species

As shown in Figure 8, when the homology search for the high-throughput pipeline was performed against all known miRNAs in miRBase, seventeen species had at least three or more hits with e-values < 0.01. An additional eleven species had two hits, and another seven species had one hit. *Mus musculus* displayed the highest number of hits with 56, over four times the number of hits for the next most represented species, *Triticum aestivum*. Although *M. musculus* has a high number (488) of miRNA sequences in miRBase, it does not appear that the sheer number of sequences is solely responsible for the higher frequency of hits. Other organisms are also well represented in miRBase, but displayed relatively few hits. For example, 695 human miRNAs are recorded in miRBase, but only five hits were observed in this study. Including the eight hits from *Rattus norvegicus*, over one-third of all hits observed were from the order, Rodentia. The cause of this observation is unclear. These findings may be merely the result of incomplete coverage of the database, which is only capturing a small number of the actual miRNAs present in mammals, or it may be possible that the miRNAs that have been identified in these Rodentia species are particularly well conserved across species.

Both metazoan and non-metazoan miRNAs were used as reference miRNAs in this analysis. It was assumed that metazoan miRNAs would be more likely to yield hits in *S. mansoni*, and that the non-metazoan sequences would




Figure 8 Number of microRNAs hits per species with three or more hits with e-value < 0.01. A homology search for the high-throughput pipeline was performed. Seventeen species had at least three hits with e-values < 0.01. An additional eighteen species had one or two hits.

provide somewhat of a negative control for the method, i. e. few hits should be observed with non-metazoan sequences. Interestingly, four of the top eight represented species are plants: *T. aestivum* (13 hits), *Physcomitrella patens* (11 hits), *Arabidopsis thaliana* (7 hits) and *Oryza sativa* (6 hits). The number of hits with e-values < 0.01 for each major taxon listed in miRBase (subphylum, phylum or kingdom) is shown in Figure 9. With 37 hits, plants (Viridiplantae) represent 20% of the total number of hits with e-values < 0.01. The most represented taxon is the subphylum, Vertebrata, with 122 hits or 68% of the total hits. This finding is not surprising as Vertebrata is also the taxon with the most number of miRNAs in miRBase (5157), more than three times the number of miRNAs in the next largest taxa, Viridiplantae (1638) and Arthropoda (1194).

The percentages of miRNAs from each major taxon in miRBase that returned a hit with e-value < 0.01 are shown in Figure 10. The kingdom Protistae (6.1%) and phylum Platyhelminthes (4.8%) display the highest percentages of hits. However, both of these taxa contain only one organism in miRBase, with each organism containing less than 65 miRNAs. Of the three taxa that have the most representatives in miRBase, Vertebrata and Viridiplantae display similar percentages (1.6-1.7%), and are both higher than Arthropoda (0.8%). It is interesting that a higher frequency of hits is observed for Viridiplantae than for Arthropoda, considering *S.*


mansoni and Arthropoda would be more closely related as both are metazoans. This is a further indication that the number of miRNAs is likely to be much higher than that currently represented within the database.


Observed miRNA families

As shown in Figure 11, 36 different miRNA families were observed in the homology search. Of these, 22 families were observed multiple times, either from different species or within the same species. The miRNA family observed most frequently was miR-19, with 22 hits. Also shown in Figure 11 is the number of probable miRNA candidates that were observed in each family. Five of the six families with the most hits displayed at least one probable miRNA candidate. Ten of the thirteen families that displayed probable miRNA candidates rank in the top sixteen families with respect to number of hits. These results suggest that miRNA families that are highly conserved, appearing in the most number of species, may be most likely to yield probable miRNA candidates.

Conclusions


The discovery of small regulatory RNA molecules, miRNAs, is undoubtedly one of the most important recent findings in biological research. This study demonstrates for the first time the presence of miRNAs in *S. mansoni* identified by complementary experimental and computational approaches. By cloning and sequencing of 1200 sequences from a small RNA library, 211 potential





miRNA candidates were identified, of which 26 were predicted to form stem-loop structures characteristic of miRNA precursors. The expression of 14 of them was confirmed by northern blot analysis. The homology search by the high-throughput pipeline was performed with all known miRNAs in miRBase and fifteen novel

likely miRNAs were detected in the parasitic organism *S. mansoni*. The identification of miRNA in the *S. mansoni* genome presents relevant information that is likely to be important to study various aspects as parasite development, gene regulation, evolutionary processes and sexual maturation.



Methods

Parasites and nucleic acid extraction

Total RNA was extracted from adult worm pairs and lung-stage schistosomula of *S. mansoni* with use of Trizol[®] (Invitrogen). Cercariae were obtained from infected *Biomphalaria glabrata* snails and isolated parasite bodies were prepared as previously described [53]. Schistosomula were cultured for 7 days in complete RPMI medium supplemented with 10 mM Hepes, 2 mM glutamate, 5% fetal calf serum and antibiotics (100 U/ml penicillin and 100 µg/ml streptomycin) at 37°C in a 5% CO₂ atmosphere.

RNA isolation and miRNA cloning

A total of 5 aliquots with 200 µg of total RNA isolated from adult worms by guanidine thiocyanate phenol-chloroform extraction were pooled [54]. The short RNA fraction ranging from 17 to 26 nt was purified and cloned as described in Chappell *et al.* [28]. Briefly, the concentration was quantified using the NanoDrop Spectrophotometer (NanoDrop Technologies, USA). Total RNA (1 mg) was resolved by electrophoresis on 15% denaturing polyacrylamide gel (8 M urea, 1 × TBE buffer), and short RNAs (17 to 26 nucleotides in length) were excised and eluted in 3 M NaCl solution at 4°C for 16 h. The gel purified small RNAs were dephosphorylated using APex[™] Heat-Labile Alkaline Phosphatase (Epicentre) and ligated directly to a 5'-phosphorylated 3'-adapter oligonucleotide with a blocked 3'-hydroxyl terminus (5'-pUUUaacgcctcctctcx-3'; uppercase, RNA; lowercase, DNA; p, phosphate; x, inverted deoxythymidine) (Dharmacon Research, Boulder, CO) to prevent self-ligation. The ligation products were separated from the excess of 3'-adapter on a 15% denaturing polyacrylamide gel and were subsequently ligated to a non-phosphorylated 5'-adater oligonucleotide (5'-tactaatacactcactAAA-3'; uppercase, RNA; lowercase, DNA) (Dharmacon Research, Boulder, CO) using T4 RNA ligase (Invitrogen). The final products were again gel purified by size fractionation and submitted to reverse transcription reaction using the RT primer (5'-TTTTCTGCAGAAGGATGCGGTAAA-3'; bold, *Pst*I site). This was followed by high fidelity PCR amplification using the reverse (RT primer) and forward (5'-AAACCATGGTACTAATACGACTCACTAAA-3'; bold, *Nco*I site). The PCR products were digested with *Pst*I and *Nco*I and subsequently concatenated using T4 DNA ligase. The ends of the concatamers were filled in with Klenow/AT-tailing and ligated into a 2.1 TOPO TA vector (Invitrogen). Ligated plasmids were transformed into TOP10 cells (Invitrogen). The libraries were plated on Luria-Bertani (LB) ampicillin (100 µg/ml) plates and individual colonies were picked and put into 96-well plates containing LB ampicillin and grown overnight at

37°C with continuous shaking. The recombinant clones were selected, sequenced and the data was analyzed as described below.

Computational analysis of microRNA library sequences

Base calling and quality trimming of sequence chromatograms were conducted using PHRED [55]. After masking of vector and adapter sequences using EMBOSS-restrict <http://bioweb2.pasteur.fr/docs/EMBOSS/restrict.html>, small RNA sequences ranging 17-25 nt in length were aligned by ClustalW2 program and redundant sequences removed [56].

The unique sequences were used in BLAST searches against the *S. mansoni* genome and miRBase database (<http://microrna.sanger.ac.uk>; release 13.0) to identify sequences from other species that closely match candidate *S. mansoni* miRNAs and removal of contaminating mRNAs, tRNAs, rRNAs, and other small RNAs. To predict the secondary structure of the remaining small RNA, Perl scripts were implemented to align the sequences to the genome of the parasite *S. mansoni* (<http://www.schistodb.net> Genome version 4.0) aiming at retrieving all possible genomic locations. In brief, the script executes BLAST to perform sequence similarity analysis and the result is parsed to retrieve the genomic positions to which each miRNA aligns [57]. In the next step, the script builds a FASTA file containing two sets of approximately 500 entries for each miRNA: one set of genomic sequences plus 40, 50, 60 or 70 nucleotides upstream and downstream. The secondary structures were predicted for each sequence using RNAfold from the Vienna RNA package [58]. Each image was further visually inspected to confirm the presence of a typical stem-loop conformation of pre-miRNAs. Among all structures created for each sequence, the one containing the mature miRNA in one arm of the hairpin precursor and with lowest folding free energy was selected. The final images were created using the VARNA <http://varna.lri.fr/index.html> to insert subtitles and highlight the mature miRNA sequence.

miRNA expression analysis

For the northern blot analysis, total RNA from adult worm pairs and 7-day *in vitro* cultured schistosomula were used. Sixty micrograms of total RNA were separated on 15% denaturing polyacrylamide gels and electrotransferred to Hybond N+ membranes (GE Healthcare) in 1x Tris Borate EDTA using the Mini Trans-Blot Cell apparatus (Bio-Rad), according to the manufacturer's instructions. Membranes were UV cross-linked in the *UV Stratalinker*[®] (Stratagene) and pre-hybridized in DIG Easy Hyb solution (Roche) at 37°C for 30 min. DNA oligonucleotides complementary to the miRNA sequences

selected candidate pre-miRNA sequence is shown underlined in Figure 12C.

When selecting the candidate pre-miRNA, if additional paired bases were directly adjacent to the selected sequence, the selection was extended to include these bases. This step prevents known stems of the hairpin from being truncated. The two additional bases, 'UC' (shown italicized), on the left of the selected sequence in Figure 12D illustrates this rule.

If either end of the selected hairpin sequence terminates in paired bases, while the other end of the sequence terminates in unpaired bases, the paired end of the sequence was extended by the number of unpaired bases on the other end. Extending the sequence required extracting the additional bases from the *S. mansoni* database. The rationale for this rule was that unpaired bases on the miRNA end may not actually be unpaired, but instead the bases that they pair with were merely not present in the original extended sequence. After the additional bases were added, the sequence was refolded. If the newly added bases were unpaired, the original fold was used. In the example in Figure 12E, the six bases on the left of the selected sequence, 'GAUCGA', were unpaired. However, the other end of the selected sequence ended in paired bases. As a result, six additional bases were added and the sequence was refolded as shown in Figure 12F.

In cases where two or more hairpins were present in the extended sequence, two sequence selections are made, i.e. on either side of the mature sequence. The rules described above were then followed as shown in Figure 12G.

Unpaired bases at the ends of the hairpin stems that are not part of the mature miRNA sequence or the 3nt extension were removed.

The candidate pre-miRNA sequences were folded using RNASHAPES. Structures with MFE \leq -15 kcal/mol were considered probable pre-miRNA sequences.

Additional material

Additional file 1: Table S1. Clustering of 584 sequenced miRNAs.

584 sequenced miRNAs were grouped into 211 clusters. The putative miRNA ID, sequence, length, the number of sequences (Frequency) and genomic locations are shown. The results in Northern blot analysis are shown (+ positive signal, - negative signal, NT not tested). Known genomic locations are hyperlinked to <http://www.schistodb.net>. This Table is also available at <http://www.cebio.org/content/2009/04/08/schistosoma-mansoni-micrnas>.

Additional file 2: Figure S1. Predicted precursor structures of new *S. mansoni* miRNAs.

The miRNAs shown were undetected by northern blot in adult worm and schistosomula stages. The RNA secondary structure of the precursors was predicted using RNAfold from the Vienna RNA package <http://ma.tbi.univie.ac.at/cgi-bin/RNAfold.cgi>. The file is also available at <http://www.cebio.org/content/2009/04/08/schistosoma-mansoni-micrnas>.

Additional file 3: Table S2. High-throughput pipeline homology search results. Results from BLASTn homology search using all known mature miRNA sequences to search the *S. mansoni* miRNA database (e-value < 0.01). The file is available at <http://www.cebio.org/content/2009/04/08/schistosoma-mansoni-micrnas>.

Additional file 4: Table S3. High-throughput pipeline extended sequence folding results. Results from RNASHAPES folding of 110 extended miRNA sequences. The file is available at <http://www.cebio.org/content/2009/04/08/schistosoma-mansoni-micrnas>.

Acknowledgements

This work was funded by NIH-NIAID Grant 5D43TW007012-04, FAPEMIG (CBB-1181/08 and 5323-4.01/07), Capes, CEBio, CNPq - INCT (573839/2008-5) and Fiocruz (GO), NIH Training Grant D43TW006580 (PLV), NIH Grant U01-AI48828 (NMES). GO is a CNPq fellow (306879/2009-3). The parasites were kindly provided by Fred Lewis (Biomedical Research Institute through NIH-NIAID, MD) and Laboratory of Mollusks, Centro de Pesquisas René Rachou-Fiocruz, Belo Horizonte, Brazil.

Author details

¹Graduate Program in Bioinformatics, Universidade Federal de Minas Gerais, Av. Antonio Carlos 6627, Belo Horizonte, MG, Brazil. ²Department of Statistics, Peter Medawar Building, South Parks Road, Oxford, UK. ³J Craig Venter Institute (JCVI), 9704 Medical Center Drive, Rockville, MD 20850, USA. ⁴Department of Cell Biology and Molecular Genetics and Center for Bioinformatics and Computational Biology, University of Maryland College Park, MD 20742, USA. ⁵Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, MD, USA. ⁶CEBio, Instituto Nacional de Ciência e Tecnologia em Doenças Tropicais, Laboratory of Cellular and Molecular Parasitology, Centro de Pesquisas René Rachou, Fundação Oswaldo Cruz, Av. Augusto de Lima 1715, Belo Horizonte, 30190-002, Brazil. ⁷University of Texas Health Science Center, 7703 Floyd Curl Dr. Mail Code 7760, San Antonio, Texas 78229-3900, USA. ⁸Biosciences eastern and central Africa - International Livestock Research Institute (BeCA-ILRI) Hub, P.O. Box 30709 Nairobi, Kenya.

Authors' contributions

MCS performed all experiments and wrote the paper. AD directed the miRNA library construction and MCS, GCC and AZ carried out the prediction and computational analysis of miRNA libraries. JL and ARD performed the bioinformatics experiments. RASP contributed to the northern blot experiments. PLV, GO and NMES designed and directed the project. All authors read and approved the final manuscript.

Received: 31 July 2010 Accepted: 19 January 2011

Published: 19 January 2011

References

1. Kim VN, Nam JW: **Genomics of microRNA.** *Trends Genet* 2006, **22**:165-173.
2. Huttenhofer A, Vogel J: **Experimental approaches to identify non-coding RNAs.** *Nucleic Acids Res* 2006, **34**:635-646.
3. He L, Hannon GJ: **MicroRNAs: small RNAs with a big role in gene regulation.** *Nat Rev Genet* 2004, **5**:522-31.
4. Valencia-Sanchez MA, Liu J, Hannon GJ, Parker R: **Control of translation and mRNA degradation by miRNAs and siRNAs.** *Genes Dev* 2006, **20**:515-524.
5. Lagos-Quintana M, Rauhut R, Meyer J, Borkhardt A, Tuschl T: **New microRNAs from mouse and human.** *RNA* 2003, **9**:175-179.
6. Wiznowerowicz M, Szulc J, Trono D: **Tuning silence: conditional systems for RNA interference.** *Nature Methods* 2006, **3**:682-688.
7. Lau NC, Lai EC: **Diverse roles for RNA in gene regulation.** *Genome Biol* 2005, **6**:315.
8. Ambros V: **The functions of animal's microRNAs.** *Nature* 2004, **431**:350-355.
9. Miska A: **How microRNAs control cell division, differentiation and death.** *Curr. Opin. Genet. Dev* 2005, **15**:563-568.
10. Bushati N, Cohen SM: **microRNA functions.** *Annual Review of Cell and Developmental Biology* 2007, **23**(I):175-205.

11. Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Radmark O, Kim S, Kim VN: **The nuclear RNase III Drosha initiates microRNA processing.** *Nature* 2003, **425**:415-419.
12. Winter J, Jung S, Keller S, Gregory RI, Diederichs S: **Many roads to maturity: microRNA biogenesis pathways and their regulation.** *Nat Cell Biol* 2009, **11**:228-34.
13. Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U: **Nuclear export of microRNA precursors.** *Science* 2003, **303**:95-98.
14. Lewis BP, Shi I, Jones-Rhoades MW, Bartel DP, Burge CB: **Prediction of mammalian microRNA targets.** *Cell* 2003, **115**:787-798.
15. Lee RC, Ambros V: **An extensive class of small RNAs in *Caenorhabditis elegans*.** *Science* 2001, **294**:862-864.
16. Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, Tuschl T: **Identification of tissue-specific microRNAs from mouse.** *Curr Biol* 2002, **12**:735-739.
17. Ambros V, Bartel B, Bartel DP, Burge CB, Carrington JC, Chen X, Dreyfuss G, Eddy SR, Griffiths-Jones S, Marshall M, Matzke M, Ruvkun G, Tuschl T: **A uniform system for microRNA annotation.** *RNA* 2003, **9**:277-279.
18. Kloosterman WP, Steiner FA, Berezikov E, de Bruijn E, van de Belt J, Verheul M, Cuppen E, Plasterk RH: **Cloning and expression of new microRNAs from zebrafish.** *Nucleic Acids Res* 2006, **34**:2558-2569.
19. Lai EC, Tomancak P, Williams RW, Rubin GM: **Computational identification of *Drosophila* microRNA genes.** *Genome Biol* 2003, **4**:R42.
20. Lim LP, Lau NC, Weinstein EG, Abdelhakim A, Yekta S, Rhoades MW, Burge CB, Bartel DP: **The microRNAs of *Caenorhabditis elegans*.** *Genes Dev* 2003, **17**:991-1008.
21. Lim LP, Glasner ME, Yekta S, Burge CB, Bartel DP: **Vertebrate microRNAs genes.** *Science* 2003, **299**:1540.
22. Lee RC, Feinbaum RL, Ambros V: **The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*.** *Cell* 1993, **75**:843-854.
23. Hertel J, Lindemeyer M, Missal K, Fried C, Tanzer A, Flamm C, Hofacker IL, Stadler PF, Students of Bioinformatics Computer Labs 2004 and 2005: **The expansion of the metazoan microRNA repertoire.** *BMC genomics* 2006, **7**:25.
24. Savioli L, Stansfield S, Bundy DA, Mitchell A, Bhatia R, Engels D, Montresor A, Neira M, Shein AM: **Schistosomiasis and soil-transmitted helminth infections: forging control efforts.** *Trans R Soc Trop Med Hyg* 2002, **96**:577-579.
25. Bumcrot D, Manoharan M, Koteliensky V, Sah DW: **RNAi therapeutics: a potential new class of pharmaceutical drugs.** *Nat Chem Biol* 2006, **2**:711-719.
26. Cheng G, Fu Z, Lin J, Shi Y, Zhou Y, Jin Y, Cai Y: **In vitro and in vivo evaluation of small interference RNA-mediated gynaecophoral canal protein silencing in *Schistosoma japonicum*.** *J Gene Med* 2009, **11**:412-421.
27. Krautz-Peterson G, Simoes M, Faghiri Z, Ndegwa D, Oliveira G, Shoemaker CB, Skelly PJ: **Suppressing glucose transporter gene expression in schistosomes impairs parasite feeding and decreases survival in the mammalian host.** *PLoS Pathog* 2010, **6**:e1000932.
28. Chappell L, Baulcombe D, Molnár A: **Isolation and cloning of small RNAs from virus-infected plants.** *Current Protocols in Microbiology* 2006, Chapter 16:Unit 16H.2.
29. Pearson WR: **Flexible sequence similarity searching with the FASTA3 program package.** *Methods Mol. Biol* 2000, **132**:185-219.
30. Carthew RW: **Molecular biology. A new RNA dimension to genome control.** *Science* 2006, **313**:305-306.
31. Berriman M, Haas BJ, LoVerde PT, Wilson RA, Dillon GP, Cerqueira GC, Mashiyama ST, Al-Lazikani B, Andrade LF, Ashton PD, et al: **The genome of the blood fluke *Schistosoma mansoni*.** *Nature* 2009, **460**:352-358.
32. Chaudhuri K, Chatterjee R: **MicroRNA detection and target prediction: integration of computational and experimental approaches.** *DNA Cell Biol* 2007, **26**:321-337.
33. Landgraf P, Rusu M, Sheridan R, Sewer A, Iovino N, Aravin A, Pfeffer S, Rice A, Kamphorst AO, Landthaler M, et al: **Amammalian microRNA expression atlas based on small RNA library sequencing.** *Cell* 2007, **129**:1401-1414.
34. Fahlgren N, Howell MD, Kasschau KD, Chapman EJ, Sullivan CM, Cumbie JS, Givan SA, Law TF, Grant SR, Dangl JL, Carrington JC: **High-throughput sequencing of *Arabidopsis* microRNAs: evidence for frequent birth and death of miRNA genes.** *PLoS ONE* 2007, **2**:e219.
35. Krautz-Peterson G, Skelly PJ: ***Schistosoma mansoni*: the dicer gene and its expression.** *Exp Parasitol* 2008, **118**:122-128.
36. Krautz-Peterson G, Radwanska M, Ndegwa D, Shoemaker CB, Skelly PJ: **Optimizing gene suppression in schistosomes using RNA interference.** *Mol Biochem Parasitol* 2007, **153**:194-202.
37. Xue X, Sun J, Zhang Q, Wang Z, Huang Y, Pan W: **Identification and characterization of novel microRNAs from *Schistosoma japonicum*.** *PLoS One* 2008, **3**:e4034.
38. Ding X, Weiller J, Großhans H: **Regulating the regulators: mechanisms controlling the maturation of microRNAs.** *Trends in Biotech* 2009, **27**:27-36.
39. Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, Baillie DL, Fire A, Ruvkun G, Mello CC: **Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing.** *Cell* 2001, **106**:23-34.
40. Ro S, Song R, Park C, Zheng H, Sanders KM, Yan W: **Cloning and expression profiling of small RNAs expressed in the mouse ovary.** *RNA* 2007, **13**:2366-2380.
41. Bartel DP: **MicroRNAs: genomics, biogenesis, mechanism, and function.** *Cell* 2004, **116**:281-297.
42. Steffen B, Voß B, Rehmsmeier M, Reeder J, Giegerich R: **RNASHapes: an integrated RNA analysis package based on abstract shapes.** *Bioinformatics* 2006, **22**:500-503.
43. Hsu P, Huang H, Hsu S, Lin L, Tsou A, Tseng C, Stadler P, Washietl S, Hofacker : **miRNAMap: genomic maps of microRNA genes and their target genes in mammalian genomes.** *Nucleic Acids Research* 2006, **34**:135-139.
44. Palakodeti D, Smielewski M, Graveley B: **MicroRNAs from the planarian *Schmidtea mediterranea*: a model system for cell biology.** *RNA* 2006, **12**:1640-1649.
45. Luo Q, Zhou Q, Yu X, Lin H, Hu S, Yu J: **Genome-wide mapping of conserved microRNAs and their host transcripts in *Tribolium castaneum*.** *J. Genetics and Genomics* 2008, **35**:349-355.
46. Zhou D, Li S, Wen J, Gong X, Xu L, Luo Y: **Genome wide computational analyses of microRNAs and their targets from *Canis familiaris*.** *Comput. Biol. Chem* 2008, **32**:61-66.
47. Baev V, Daskalova E, Minkov : **Computational identification of novel microRNA homologs in the chimpanzee genome.** *Comput. Biol. Chem* 2009, **33**:62-70.
48. Hao L, Cai P, Jiang N, Wang H, Chen Q: **Identification and characterization of microRNAs and endogenous siRNAs in *Schistosoma japonicum*.** *BMC Genomics* 2010, **11**:55.
49. Wang Z, Xue X, Sun J, Luo R, Xu X, Jiang Y, Zhang Q, Pan W: **An "in-depth" description of the small non-coding RNA population of *Schistosoma japonicum* schistosomulum.** *PLoS Negl Trop Dis* 2010, **4**:e596.
50. Huang J, Hao P, Chen H, Hu W, Yan Q, Liu F, Han ZG: **Genome-wide identification of *Schistosoma japonicum* microRNAs using a deep-sequencing approach.** *PLoS One* 2009, **4**:e8206.
51. Chatterjee R, Chaudhuri K: **An approach for the identification of microRNA with an application to *Anopheles gambiae*.** *Acta Biochimica Polonica* 2006, **53**:303-309.
52. Artzi S, Kiezun A, Shomron N: **miRNAMiner: A tool for homologous microRNA gene search.** *BMC Bioinformatics* 2008, **9**:39.
53. Skelly PJ, Da'dara A, Harn DA: **Suppression of cathepsin B expression in *Schistosoma mansoni* by RNA interference.** *Int J Parasitol* 2003, **33**:363-369.
54. Chomczynski P, Sacchi N: **Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction.** *Anal Biochem* 1987, **162**:156-159.
55. Ewing B, Hillier L, Wendl MC, Green P: **Base-calling of automated sequencer traces using phred. I. Accuracy assessment.** *Genome Res* 1998, **8**:175-185.
56. Thompson JD, Higgins DG, Gibson TJ: **CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice.** *Nucleic Acids Res* 1994, **22**:4673-4680.
57. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ, et al: **Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.** *Nucl Acids Res* 1997, **25**:3389-3402.
58. Hofacker IL: **Vienna RNA secondary structure server.** *Nuc. Acids Res* 2003, **31**:3429-3431.

doi:10.1186/1471-2164-12-47

Cite this article as: Simões et al.: Identification of *Schistosoma mansoni* microRNAs. *BMC Genomics* 2011 **12**:47.