

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Theodoratou, E; Campbell, H; Tenesa, A; Houlston, R; Webb, E; Lubbe, S; Broderick, P; Gallinger, S; Croitoru, EM; Jenkins, MA; Win, AK; Cleary, SP; Koessler, T; Pharoah, PD; Kry, S; Bzieau, S; Buecher, B; Ellis, NA; Peterlongo, P; Offit, K; Aaltonen, LA; Enholm, S; Lindblom, A; Zhou, XL; Tomlinson, IP; Moreno, V; Blanco, I; Capell, G; Barnetson, R; Porteous, ME; Dunlop, MG; Farrington, SM (2010) A large-scale meta-analysis to refine colorectal cancer risk estimates associated with MUTYH variants. *British journal of cancer*, 103 (12). pp. 1875-84. ISSN 0007-0920 DOI: <https://doi.org/10.1038/sj.bjc.6605966>

Downloaded from: <http://researchonline.lshtm.ac.uk/4649218/>

DOI: [10.1038/sj.bjc.6605966](https://doi.org/10.1038/sj.bjc.6605966)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

A large-scale meta-analysis to refine colorectal cancer risk estimates associated with *MUTYH* variants

E Theodoratou^{1,2}, H Campbell^{1,2}, A Tenesa¹, R Houlston⁴, E Webb^{4,18}, S Lubbe⁴, P Broderick⁴, S Gallinger⁵, EM Croitoru⁶, MA Jenkins⁷, AK Win⁷, SP Cleary⁶, T Koessler⁸, PD Pharoah⁸, S Küry⁹, S Bézieau⁹, B Buecher¹⁰, NA Ellis¹¹, P Peterlongo¹², K Offit¹³, LA Aaltonen¹⁴, S Enholm¹⁴, A Lindblom¹⁵, X-L Zhou¹⁵, IP Tomlinson¹⁶, V Moreno¹⁷, I Blanco¹⁷, G Capellà¹⁷, R Barnetson¹, ME Porteous^{1,3}, MG Dunlop¹ and SM Farrington^{*,1}

¹Colon Cancer Genetics Group and Academic Coloproctology, MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK; ²Public Health Sciences, University of Edinburgh, Teviot Place, Edinburgh, UK; ³Southeast of Scotland Clinical Genetic Services, University of Edinburgh, Edinburgh, UK; ⁴Section of Cancer Genetics, Institute of Cancer Research, Sutton, Surrey, UK; ⁵Ontario Familial Colorectal Cancer Registry, Toronto, Ontario, Canada; ⁶Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, Canada; ⁷Centre for Molecular, Environmental, Genetic and Analytic Epidemiology, The University of Melbourne, Victoria, Australia; ⁸Strangeways Research Laboratory, Department of Oncology and Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK; ⁹CHU de Nantes, pôle de Biologie, service de Génétique Médicale, 9 quai Moncoussu, Nantes, 44093 cedex 1, France; ¹⁰unité de Génétique Constitutionnelle, Institut Curie, service de Génétique Oncologique, 26 rue d'Ulm, Paris, 75248 cedex 05, France; ¹¹University of Chicago, 900 East 57th Street, Chicago, IL 60637, USA; ¹²IFOM, Fondazione Istituto FIRC di Oncologia Molecolare, and Unit of Genetic Susceptibility to Cancer, Department of Experimental Oncology and Molecular Medicine, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy; ¹³Clinical Cancer Genetics, Memorial Sloan-Kettering Cancer Centre, New York, USA; ¹⁴Department of Medical Genetics, Biomedicum Helsinki, University of Helsinki, PO Box 63 (Haartmaninkatu 8), FIN-00014, Finland; ¹⁵Department of Molecular Medicine and Surgery Karolinska Institute and Department of Clinical Genetics Karolinska University Hospital, Stockholm, Sweden; ¹⁶Molecular and Population Genetics Laboratory, London Research Institute, Cancer Research UK, London, UK; ¹⁷Catalan Institute of Oncology-ICO, IDIBELL and University of Barcelona, Av Gran Via 199, L'Hospitalet, Barcelona 08907, Spain

BACKGROUND: Defective DNA repair has a causal role in hereditary colorectal cancer (CRC). Defects in the base excision repair gene *MUTYH* are responsible for *MUTYH*-associated polyposis and CRC predisposition as an autosomal recessive trait. Numerous reports have suggested *MUTYH* mono-allelic variants to be low penetrance risk alleles. We report a large collaborative meta-analysis to assess and refine CRC risk estimates associated with bi-allelic and mono-allelic *MUTYH* variants and investigate age and sex influence on risk. **METHODS:** *MUTYH* genotype data were included from 20 565 cases and 15 524 controls. Three logistic regression models were tested: a crude model; adjusted for age and sex; adjusted for age, sex and study.

RESULTS: All three models produced very similar results. *MUTYH* bi-allelic carriers demonstrated a 28-fold increase in risk (95% confidence interval (CI): 6.95–115). Significant bi-allelic effects were also observed for G396D and Y179C/G396D compound heterozygotes and a marginal mono-allelic effect for variant Y179C (odds ratio (OR) = 1.34; 95% CI: 1.00–1.80). A pooled meta-analysis of all published and unpublished datasets submitted showed bi-allelic effects for *MUTYH*, G396D and Y179C (OR = 10.8, 95% CI: 5.02–23.2; OR = 6.47, 95% CI: 2.33–18.0; OR = 3.35, 95% CI: 1.14–9.89) and marginal mono-allelic effect for variants *MUTYH* (OR = 1.16, 95% CI: 1.00–1.34) and Y179C alone (OR = 1.34, 95% CI: 1.01–1.77).

CONCLUSIONS: Overall, this large study refines estimates of disease risk associated with mono-allelic and bi-allelic *MUTYH* carriers.

British Journal of Cancer (2010) **103**, 1875–1884. doi:10.1038/sj.bjc.6605966 www.bjcancer.com

Published online 9 November 2010

© 2010 Cancer Research UK

Keywords: colorectal cancer; base excision repair; *MUTYH*; carrier risk estimates; meta-analysis

Oxidative damage to DNA occurs with cell proliferation and increases with age. Certain organs such as the gut are heavily exposed to oxidising agents, which impacts on carcinogenic potential. Dysfunction of base excision repair, the major pathway

for repairing oxidative damage, has been implicated as a risk factor for the development of multiple colorectal adenomas and colorectal cancer (CRC; Al-Tassan *et al*, 2002; Croitoru *et al*, 2004; Farrington *et al*, 2005). Bi-allelic mutations of the *MUTYH* gene seem to be responsible for a high proportion of the multiple adenoma phenotype families (termed *MUTYH*-associated polyposis (MAP)) unaccounted for by germline *APC* mutations (Al-Tassan *et al*, 2002; Sampson *et al*, 2003; Sieber *et al*, 2003; Gismondi *et al*, 2004; Venesio *et al*, 2004; Nielsen *et al*, 2009) and predispose to CRC *per se* (Enholm *et al*, 2003; Croitoru *et al*, 2004; Fleischmann *et al*, 2004; Kambara *et al*, 2004; Wang *et al*, 2004; Farrington *et al*, 2005; Peterlongo *et al*,

*Correspondence: Dr SM Farrington;

E-mail: Susan.Farrington@hgu.mrc.ac.uk

¹⁸Present address: Infectious Disease Epidemiology Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London, UK

Received 2 July 2010; revised 16 September 2010; accepted 30 September 2010; published online 9 November 2010

2005; Zhou *et al*, 2005; Moreno *et al*, 2006; Tenesa *et al*, 2006; Webb *et al*, 2006; Küry *et al*, 2007; Cleary *et al*, 2009; Lubbe *et al*, 2009). Although an increased CRC risk associated with bi-allelic *MUTYH* mutations is incontrovertible, the risk associated with one *MUTYH* mutant allele is controversial (Croitoru *et al*, 2004; Farrington *et al*, 2005; Jenkins *et al*, 2006; Tenesa *et al*, 2006; Webb *et al*, 2006; Cleary *et al*, 2009; Jones *et al*, 2009; Lubbe *et al*, 2009). A statistically significant or close to significant effect, for a *MUTYH* mono-allelic effect, has been reported in different studies with possible age specific effects present, but the rarity of the alleles associated with the small increased risk for CRC have made it difficult to replicate study findings. A recent risk analysis of MAP family members agreed with previous family based findings (Jenkins *et al*, 2006) that mono-allelic carriers are at a two-fold increase in risk of CRC (Jones *et al*, 2009) providing further evidence of a mono-allelic effect of the gene. However, family based studies can be subject to ascertainment bias and any mono-allelic effect could potentially be modified by other inherited factors, including alleles at other loci. Furthermore, environmental risk factors also show familial aggregation and hence, studies in which there has been selection of cases based on family history may be confounded. Bi-allelic carriers may develop CRC because of the predominant effect of *MUTYH*, whereas the environmental effect is greater in affected siblings with mono-allelic mutations but the risk is ascribed to the *MUTYH* allele. Thus further work is required to resolve the mono-allelic carrier risk question.

To clarify the role of *MUTYH* in disease risk, we initiated a multi-centre collaboration allowing large-scale meta-analysis of the individual *MUTYH* variants, with special interest in determining if there were age and sex-specific effects on CRC association with *MUTYH* variants (Farrington *et al*, 2006). In this study, we present the results of this collaborative meta-analysis.

SUBJECTS AND METHODS

Participating studies

Relevant case-control studies to be invited for inclusion in the meta-analysis of the effect of *MUTYH* on CRC risk were identified by a literature search in the ISI Web of Science (<http://wok.mimas.ac.uk>) and PUBMED bibliographic databases (<http://www.ncbi.nlm.nih.gov/pubmed/>), using the search terms 'MYH or *MUTYH* and CRC'. In the initial search 55 studies were identified and eight of these were considered for our study (Enholm *et al*, 2003; Croitoru *et al*, 2004; Fleischmann *et al*, 2004; Kambara *et al*, 2004; Wang *et al*, 2004; Farrington *et al*, 2005; Peterlongo *et al*, 2005; Zhou *et al*, 2005). The inclusion criteria were as follows: the patients had to be diagnosed with CRC and the studies had to have genotype data for both cases and controls. Ten additional studies were identified during the progress of the project – Webb *et al* (2006), Moreno *et al* (2006), Küry *et al* (2007), Cleary *et al* (2009), Lubbe *et al* (2009); and unpublished data from Koessler T and Pharoah PD; and Tomlinson – personal communication. Colebatch *et al* (2006); Balaguer *et al* (2007); Avezzi *et al* (2008) were used in the pooled meta-analysis of all available published and unpublished datasets.

The principal investigators (PIs) of the selected studies were contacted and were asked to participate by providing a minimum dataset including variables necessary for the analysis (Supplementary Box 1: Study questionnaire; Supplementary Table 1: Data extraction table). In cases, in whom PIs failed to respond to our invitation to participate, reminder letters were despatched. It was not possible to include data from the following studies in the logistic regression analyses because (i) data was only available for cases that were heterozygous or homozygous for a *MUTYH* mutation (Enholm *et al*, 2003); (ii) co-variate data were only available for cases, as controls were anonymous blood donors (Zhou *et al*, 2005; Tomlinson, unpublished data); (iii) failure to

communicate with us (Kambara *et al*, 2004 and Wang *et al*, 2004). The study by Fleischmann *et al* (2004) and Webb *et al* (2006) were not used because they had been superseded by a later study (Lubbe *et al*, 2009), which was included.

Statistical analysis

Data from all collaborating centres were checked for completeness, coded and merged to form a core database. *MUTYH* defects were considered pathogenic only if there was published evidence of their pathogenicity. Individuals reported to have two defects of *MUTYH* in the original report were classified as mutated/mutated (MM), those with one defect as wild type/mutant (WM) and those with no mutation as wild type/wild type (WW). Descriptive statistics were produced on all subject characteristics, risk factors and event data. All populations described in the case-control studies were tested for Hardy-Weinberg equilibrium in controls and the genotype distributions between all groups were compared by χ^2 -test.

Three logistic regression models were applied to address confounding co-variables (model I: crude, model II: including co-variables for age and sex, model III: including co-variables for age, sex and study) on the combined datasets investigating the effect of *MUTYH* defects (WM vs WW and MM vs WW) as well as of the individual mutations Y179C (c.536A > G/p.Tyr179Cys; AA = WW, GG = MM) and G396D (c.1187G > A/p.Gly396Asp; GG = WW, AA = MM; previously known as Y165C and G382D), to identify any variant specific associations. The three logistic regression models were applied after sex and age (over 55 years and under or equal 55 years) stratification as previously described (Farrington *et al*, 2005), to assess the effect of age and sex on the association of the variants with disease risk. In all the studies, interaction associations between the *MUTYH* variants and study code (for each individual study) were estimated and similarly between *MUTYH* variants and hormone replacement therapy (HRT) among female participants in three studies (the Scottish SOCCS studies and the studies – Croitoru *et al*, 2004; Cleary *et al*, 2009). Association between both genetic (i.e., one of the *MUTYH* mutations) and the study code or environmental factor (i.e., HRT) and disease was assessed and interaction was tested by fitting interactive and nested multiplicative models. To assess for any small study effects, we performed Funnel plot analysis and tested for significance using the Harbord test.

Finally, the relationship between the genotype and CRC was analysed by meta-analysis, combining the effect estimates of all published and unpublished datasets.

All statistic analyses were conducted using Intercooled STATA version 10.0 (Stata Corp, College Station, TX, USA). For the logistic regression analyses, it is necessary to add a whole number to any fields containing 0 (see model I^a in Table 2), which reduces the final OR value, however, by using the META command in the STATA meta-analysis programme, a lower value can be added (0.5 as indicated by model I^b in Table 2) thereby giving a more accurate assessment of risk. However, this is a grouped analysis and therefore cannot be adjusted for confounding co-variables, such as age/sex and study as in models II and III. To account for multiple testing we applied the Bonferroni correction method, and the *P*-value threshold for significance was estimated to be 0.003.

RESULTS

Table 1 details summary data from the studies included in our combined analysis (comprising a total of 20 565 cases and 15 524 controls). The two variant alleles are rare with G396D variant allele having a frequency of 0.007 in controls and the Y179C variant allele a frequency of 0.002. Tests for deviation from Hardy-Weinberg equilibrium in controls were *P* = 0.99 and *P* < 0.00005 for G396D and Y179C variants, respectively.

Table 1 Summary table^a

Study	Number	G396D						Y179C			Genotype			Age			Sex			Ethnicity		
		GG	GA	AA	AA	AG	GG	WW	WM	MM	n (%)	Mean (SD)	n (%)	Mean (SD)	n (%)	Males (%)	n (%)	Caucasian/White (%)				
All studies																						
Cases	20 565	19 966	324	33	20 260	154	11	19 783	417	76	20 555 (99.9)	59.5 (10.1)	20 555 (99.9)	11 668 (56.8)	5374 (26.1)	5030 (93.6)						
Controls	15 524	15 145	211	0	15 074	68	2	14 723	280	2	15 287 (98.5)	59.0 (10.9)	15 521 (99.9)	6845 (44.1)	5597 (36.1)	4796 (85.7)						
Croitoru et al (2004)																						
Cases	1238	1209	25	4	1223	13	2	1197	29	12	1236 (99.8)	60.1 (8.7)	1236 (99.8)	544 (44.0)	Missing	Missing						
Controls	1255	1238	17	0	1251	4	0	1234	21	0	1244 (99.1)	63.6 (8.6)	1252 (99.8)	702 (56.1)	Missing	Missing						
Peterlongo et al (2005)																						
Cases	585	567	3	1	569	3	0	552	4	2	584 (99.8)	62.1 (13.1)	584 (99.8)	319 (54.6)	585 (100)	251 (42.9)						
Controls	1158	1040	5	0	1039	2	0	923	7	0	1148 (99.1)	53.9 (11.5)	1156 (99.8)	240 (20.8)	1158 (100)	363 (31.3)						
SOCCS (includes data from Farrington et al (2005), Tenesa et al (2006) and unpublished SOCCS prospective samples)																						
Cases	3278	3086	55	8	3192	23	0	3038	71	12	3277 (100)	59.5 (11.3)	3278 (100)	1883 (57.4)	3278 (100)	3268 (99.7)						
Controls	3318	3239	43	0	3061	15	0	2993	57	0	3313 (99.8)	61.4 (10.9)	3318 (100)	1861 (56.1)	3318 (100)	3312 (99.8)						
Moreno et al (2006)																						
Cases	356	343	8	1	348	0	0	336	8	1	356 (100)	66.5 (11.7)	356 (100)	217 (61.0)	Missing	Missing						
Controls	297	285	7	0	291	0	0	283	7	0	297 (100)	65.3 (12.5)	297 (100)	158 (53.2)	Missing	Missing						
Koesler T, (unpublished data obtained in 2007)																						
Cases	2262	2215	31	2	2227	19	1	2198	37	9	2253 (99.6)	59.1 (8.1)	2262 (100)	1287 (57.1)	Missing	Missing						
Controls	2253	2217	31	0	2236	11	0	2204	42	0	2053 (91.1)	53.4 (7.6)	2253 (100)	949 (42.0)	Missing	Missing						
Küry et al (2007)																						
Cases	1025	1003	22	0	1021	4	0	999	25	1	1023 (99.8)	68.7 (9.9)	1025 (100)	632 (61.7)	1025 (100)	1025 (100)						
Controls	1121	1105	16	0	1117	4	0	1100	21	0	1121 (100)	61.9 (10.0)	1121 (100)	609 (54.3)	1121 (100)	1121 (100)						
SOCCS retrospective cases (unpublished data obtained in 2008)																						
Cases	486	403	6	2	446	2	3	391	6	4	486 (100)	54.7 (17.2)	479 (98.6)	240 (50.1)	486 (100)	486 (100)						
Controls	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0						
Cleary et al (2009)																						
Cases	2076	2023	33	5	2053	7	1	2029	38	9	2070 (99.7)	56.6 (10.8)	2076 (100)	1114 (53.7)	Missing	Missing						
Controls	1049	1032	17	0	1042	6	1	1024	24	1	1047 (99.8)	56.4 (11.3)	1049 (100)	444 (42.3)	Missing	Missing						
Lubbe et al (2009) ^b																						
Cases	9268	9117	141	10	9181	83	4	9043	198	27	9268 (100)	59.0 (8.5)	9268 (100)	5432 (58.6)	Missing	Missing ^c						
Controls	5064	4989	75	0	5037	26	1	4962	101	1	5064 (100)	59.2 (10.8)	5064 (100)	1882 (37.2)	Missing	Missing ^c						

Abbreviations: MM = mutated/mutated; WM = wild type/wild type; WW = wild type/wild type. ^aThis table presents the raw data sent to us by each group that was then included in the analyses. G396D and Y179C are looked at independently and any compound bi-allelic carriers are presented as heterozygotes for each variant; MUTYH genotype data also includes other pathogenic mutations; unpublished data are included in Peterlongo P. and Moreno V. SOCCS prospective and Küry S. ^bIncludes data from Webb et al (2006) and Fleischmann et al (2004). ^cAll the cases and controls were UK residents and of European ancestry (self-reported).

Table 2 Logistic regression analysis of the combined datasets; G396D analysis was conducted for individuals that were Y179C AA; Y179C analysis was conducted for individuals that were G396D GG; combined genotype analysis was conducted for individuals with data for both Y179C and G396D

Gene	Cases	Controls	Model I ^a			Model I ^b		
			OR	95% CI	P-value	OR	95% CI	P-value
<i>G396D^c</i>								
<i>Whole sample</i>								
GG	19 767	14 723	1.00					
GA	292	210	1.04	0.87, 1.24	0.70			
AA	31	0	23.09	3.15, 169.15	0.002	46.93	2.87, 766.89	0.007
<i>≤ 55 Years old</i>								
GG	6269	5270	1.00					
GA	88	77	0.96	0.71, 1.31	0.80			
AA	13	0	10.93	1.43, 83.57	0.02	22.70	1.35, 381.91	0.03
<i>> 55 Years old</i>								
GG	13 498	9453	1.00					
GA	204	133	1.07	0.86, 1.34	0.52			
AA	18	0	12.60	1.68, 94.37	0.01	25.91	1.56, 430.03	0.02
<i>Males</i>								
GG	11 229	6460	1.00					
GA	161	95	0.98	0.76, 1.26	0.85			
AA	15	0	8.62	1.14, 65.25	0.04	17.84	1.07, 298.11	0.04
<i>Females</i>								
GG	8528	8259	1.00					
GA	131	115	1.10	0.86, 1.42	0.45			
AA	16	0	15.49	2.05, 116.86	0.008	31.96	1.92, 532.72	0.02
<i>Y179C^d</i>								
<i>Whole sample</i>								
AA	19 767	14 723	1.00					
AG	122	68	1.34	1.00, 1.80	0.05			
GG	11	2	4.10	0.91, 18.48	0.07	NA		
<i>≤ 55 Years old</i>								
AA	6269	5270	1.00					
AG	35	25	1.18	0.70, 1.97	0.54			
GG	9	0	7.57	0.96, 59.74	0.06	15.97	0.93, 274.491	0.06
<i>> 55 Years old</i>								
AA	13 498	9453	1.00					
AG	87	43	1.42	0.98, 2.04	0.06			
GG	2	2	0.70	0.10, 4.97	0.72	NA		
<i>Males</i>								
AA	11 229	6460	1.00					
AG	68	23	1.70	1.06, 2.73	0.03			
GG	8	0	4.60	0.58, 36.80	0.15	9.78	0.56, 169.48	0.12
<i>Females</i>								
AA	8528	8259	1.00					
AG	54	45	1.16	0.78, 1.73	0.46			
GG	3	2	1.45	0.24, 8.70	0.68	NA		
<i>Genotype^e</i>								
<i>Whole sample</i>								
WWW	19 767	14 723	1.00					
WM	418	280	1.11	0.95, 1.29	0.17			
MM	76	2	28.30	6.95, 115.26	3.1 × 10 ⁻⁶	NA		
G396D AA	31	0	23.09	3.15, 169.15	0.002	46.93	2.87, 766.89	0.007
Y179C GG	11	2	4.10	0.91, 18.48	0.07	NA		
Compound heterozygous ^f	29	0	21.60	2.94, 158.58	0.003	43.95	2.69, 719.26	0.008
<i>≤ 55 Years old</i>								
WWW	6269	5270	1.00					
WM	124	104	1.00	0.77, 1.30	0.99			
MM	43	0	36.15	4.98, 262.57	0.0004	73.14	4.50, 1188.3	0.003
G396D AA	13	0	10.93	1.43, 83.57	0.02	22.70	1.35, 381.91	0.03
Y179C GG	9	0	7.57	0.96, 59.74	0.06	15.97	0.93, 274.49	0.06
Compound heterozygous	17	0	14.29	1.90, 107.42	0.01	29.42	1.77, 489.38	0.02
<i>> 55 Years old</i>								
WWW	13 498	9453	1.00					
WM	294	176	1.17	0.97, 1.41	0.10			
MM	33	2	11.56	2.77, 48.17	0.001	NA		
G396D AA	18	0	12.60	1.68, 94.37	0.01	25.91	1.56, 430.03	0.02
Y179C GG	2	2	0.70	0.10, 4.97	0.72	NA		
Compound heterozygous	12	0	8.40	1.09, 64.64	0.04	17.51	1.04, 295.75	0.05

Table 2 (Continued)

Gene	Cases	Controls	Model I ^a			Model I ^b		
			OR	95% CI	P-value	OR	95% CI	P-value
<i>Males</i>								
WW	11 229	6460	1.00					
WM	232	119	1.12	0.90, 1.40	0.31			
MM	36	0	20.71	2.84, 151.09	0.003	42.00	2.58, 684.38	0.009
G396D AA	15	0	8.62	1.14, 65.25	0.04	17.84	1.07, 298.11	0.04
Y179C GG	8	0	4.60	0.58, 36.80	0.15	9.78	0.56, 169.48	0.12
Compound heterozygous	12	0	6.90	0.90, 53.10	0.06	14.38	0.85, 242.96	0.06
<i>Females</i>								
WW	8528	8259	1.00					
WM	186	161	1.12	0.90, 1.38	0.30			
MM	40	2	19.37	4.68, 80.17	4.3 × 10 ⁻⁵	NA		
G396D AA	16	0	15.49	2.05, 116.86	0.008	31.96	1.92, 532.72	0.02
Y179C GG	3	2	1.45	0.24, 8.70	0.68	NA		
Compound heterozygous	17	0	16.46	2.19, 123.70	0.006	33.90	2.04, 563.74	0.01

Abbreviations: CI = confidence interval; MM = mutated/mutated; NA = not available; OR = odds ratio; WM = wild type/mutant; WW = wild type/wild type. ^aCrude analysis. ^bEstimated by adding one control with the variant genotype. ^cEstimated using the meta command of STATA and for mathematical reasons, cells with zero frequencies were assumed to be 0.5 (as defaulted by the meta command). ^dAnalysis conducted only for the AA Y179C, that is, WW. ^eAnalysis conducted only for the GG G396D, that is, WW. ^fIncluding subjects with data for both Y179C and G396D. ^gThis category includes 29 G396D GA and Y179C AG cases; 5 cases with either G396D GA or Y179C AG and any other pathogenic *MUTYH* mutation were excluded.

Bi-allelic effect of *MUTYH*

All three models of the logistic regression analysis gave consistent results and so the results of the crude analysis (model I) are described below and presented in Table 2; results of the other two models can be found in Supplementary Table 2. Bi-allelic carriers for the MM genotype of the combined *MUTYH* defects, G396D and Y179C/G396D compound heterozygotes were associated with a significant increase in CRC risk (odds ratio (OR) = 28.3, 95% confidence limits (CIs): 6.95–115; 23.1 (95% CI: 3.15–169) and 21.6 (95% CI: 2.94–159), respectively). These risks are conservative, concentrating on the significant logistic regression results – model I^b results presented in Table 2 are likely a better reflection of risk and tend to be two-fold higher. There was a greater CRC risk for the MM genotype for the earlier age individuals when compared with the older age group (OR = 36.2 (95% CI: 4.98–263) for ≤55 years compared with 11.6 (95% CI: 2.77–48.2) for >55 years). However, their CIs overlapped and the results were not statistically significantly different. ANOVA analysis of mean age of carriers demonstrated that there are significant age differences between cases and controls when considering MM genotype carriers and Y179C bi-allelic carriers but not for G396D carriers ($P < 0.0005$, $P < 0.0005$ and $P = 0.27$, respectively – Supplementary Table 3). Indeed there is a significant age difference between mean age of bi-allelic Y179C and G396D carriers (48.9 vs 56.7, respectively, $P = 0.003$ based on *t*-test – Supplementary Table 4).

Colorectal cancer risk associated with mono-allelic *MUTYH* mutations

The results of the combined analysis demonstrate that there are no significant mono-allelic effects for either G396D or for combined *MUTYH* variants (Table 2). However, the specific Y179C variant was shown to increase risk of disease in the heterozygous state (OR = 1.34; (95% CI: 1.00–1.80)) in the whole sample set and also when stratified by sex, male sex demonstrated a mono-allelic effect (OR = 1.70; (95% CI: 1.06–2.73)). However, after Bonferroni correction, these mono-allelic effects did not remain significant.

The role of study population and HRT in modulating CRC risk

We hypothesised that origin of the data, that is, study population might modify the association between the genotype and CRC risk. However, there was no evidence for an interaction between study code and *MUTYH* genotypes (Supplementary Table 5). Similarly, HRT intake, a known risk factor for CRC (Chan *et al*, 2006; Theodoratou *et al*, 2008), might be influenced by genotype and therefore modulate female risk. Both the Scottish and Canadian datasets had recorded data on HRT intake and these were used to test for an interaction between HRT and *MUTYH* genotype. Across both datasets there was no evidence of any interaction between HRT and *MUTYH* genotype (Supplementary Table 6).

Meta-analysis of published and unpublished datasets

The results of a meta-analysis of published and unpublished datasets submitted to us, estimating the effect of the *MUTYH* whole gene defects demonstrated a pooled fixed bi-allelic effect of 10.8 (95% CI: 5.02–23.2) for the MM and a pooled fixed mono-allelic effect of 1.16 (95% CI: 1.00–1.34) for WM genotype (Table 3; Figures 1 and 2). Analysis of the specific variants by pooled meta-analysis demonstrated bi-allelic effects for both G396D and Y179C (OR = 6.47 (95% CI: 2.33–18.0) and OR = 3.35 (95% CI: 1.14–9.89), respectively) and in agreement with the logistic regression analysis results, Y179C variant also demonstrated a very similar pooled fixed mono-allelic effect of 1.34 (95% CI: 1.01–1.77; Tables 4 and 5; Supplementary Figures 1–4).

Assessment of study publication bias

Funnel plots for both the mono- and bi-allelic effect were created to assess whether study size was significantly influencing the results. These plots appeared asymmetric, but the Harbord's test for small study effect demonstrated that this was not statistically significant (Supplementary Figures 5 and 6).

Table 3 Meta-analysis of studies^a

Study	Genotype (cases)			Genotype (controls)			WM vs WW OR (95% CI)	MM vs WW OR (95% CI)
	WW	WM	MM	WW	WM	MM		
Enholm <i>et al</i> (2003)	994	5	4	424	0	0	4.70 (0.26, 85.10)	3.84 (0.21, 71.51)
Kambara <i>et al</i> (2004)	90	2	0	52	—	0	1.16 (0.10, 13.06)	NA
Wang <i>et al</i> (2004)	432	10	2	309	4	0	1.79 (0.56, 5.75)	3.58 (0.17, 74.79)
Zhou <i>et al</i> (2005)	432	6	0	466	3	0	2.16 (0.54, 8.68)	NA
Peterlongo <i>et al</i> (2005)	549	4	2	911	7	0	0.95 (0.28, 3.25)	8.29 (0.40, 173.08)
Tomlinson I (2006) ^b	662	15	1	197	2	0	2.23 (0.51, 9.84)	0.89 (0.04, 22.04)
Moreno <i>et al</i> (2006)	323	9	0	278	6	0	1.29 (0.45, 3.67)	NA
Colebatch <i>et al</i> (2006)	859	11	2	473	5	0	1.21 (0.42, 3.51)	2.75 (0.13, 57.49)
Koessler T (2007) ^c	2198	37	9	2204	42	0	0.88 (0.57, 1.38)	19.05 (1.11, 327.53)
Küry <i>et al</i> (2007)	999	24	1	1100	21	0	1.26 (0.70, 2.27)	3.30 (0.13, 81.18)
Balaguer <i>et al</i> (2007)	1089	19	8	912	22	0	0.72 (0.39, 1.34)	14.24 (0.82, 247.02)
SOCCS (2008) ^d	3429	77	16	2993	57	0	1.18 (0.83, 1.67)	28.80 (1.73, 480.32)
Avezzu <i>et al</i> (2008)	435	2	2	246	1	0	1.13 (0.10, 12.54)	2.83 (0.14, 59.19)
Cleary <i>et al</i> (2009) ^e	3697	87	27	2758	43	1	1.51 (1.04, 2.18)	20.14 (2.74, 148.32)
Lubbe <i>et al</i> (2009) ^f	9043	198	27	4962	101	1	1.08 (0.84, 1.37)	14.82 (2.01, 109.06)
M-H pooled effect (fixed) P-value	25231	506	101	18285	315	2	1.16 (1.00, 1.34) 0.05	10.80 (5.02, 23.21) <0.0005
Heterogeneity								
P-value							0.82	0.84
<i>I</i> ² (95% CI)							0 (0, 54)	0 (0, 60)

Abbreviations: CI = confidence interval; MM = mutated/mutated; NA = not available; OR = odds ratio; WM = wild type/mutant; WW = wild type/wild type. ^aThis table presents the data as they were published. Two unpublished studies included. ^bUnpublished data obtained in 2006. ^cUnpublished data obtained in 2007. ^dIncludes data from Farrington *et al* (2005), Tenesa *et al* (2006) and unpublished data from the SOCCS study obtained in 2008. ^eIncludes data from Croitoru *et al* (2004). ^fIncludes data from Webb *et al* (2006) and Fleischmann *et al* (2004).




Figure 1 Meta-analysis of studies comparing *MUTYH* MM vs WW. SOCCS data include Farrington *et al* (2005), Tenesa *et al* (2006) and unpublished data from the SOCCS study obtained in 2008; Cleary data include Croitoru *et al* (2004); Lubbe data include Webb *et al* (2006) and Fleischmann *et al* (2004). Unpublished studies included are Tomlinson I (2006) and Koessler T (2007).

DISCUSSION

This large meta-analysis study refines the estimates of CRC risk associated with mutations in the *MUTYH* gene to date. Bi-allelic carriers of the combined *MUTYH* mutations (MM) are associated with a 28-fold (95% CI: 6.95–115) increase in CRC risk from the

logistic regression analysis. Bi-allelic carriers of the G396D variant and Y179C/G396D compound heterozygotes were also significantly associated with a similar increase in CRC risk (OR = 23.1 (95% CI: 3.15–169) and 21.6 (95% CI: 2.94–159), respectively). Although the risk estimate was slightly lower from the overall larger pooled meta-analysis of published and unpublished datasets (OR = 10.8




Figure 2 Meta-analysis of studies comparing *MUTYH* WM vs WW. SOCCS data include Farrington *et al* (2005), Tenesa *et al* (2006) and unpublished data from the SOCCS study obtained in 2008; Cleary data include Croitoru *et al* (2004); Lubbe data include Webb *et al* (2006) and Fleischmann *et al* (2004). Unpublished studies included are Tomlinson I (2006) and Koessler T (2007).

Table 4 Meta-analysis of studies for the G396D genotypes^a

Study	G396D (cases)			G396D (controls)			GA vs GG OR (95% CI)	AA vs GG OR (95% CI)
	GG	GA	AA	GG	GA	AA		
Enholm <i>et al</i> (2003)	994	4	1	424	0	0	3.84 (0.21, 71.51)	1.28 (0.05, 31.50)
Kambara <i>et al</i> (2004)	90	2	0	52	—	0	Not enough data available	NA
Wang <i>et al</i> (2004)	432	5	0	309	2	0	1.79 (0.35, 9.28)	NA
Zhou <i>et al</i> (2005)	432	1	0	466	1	0	1.08 (0.07, 17.30)	NA
Peterlongo <i>et al</i> (2005)	549	2	0	911	5	0	0.66 (0.13, 3.43)	NA
Tomlinson I (2006) ^b	662	9	1	197	2	0	1.34 (0.29, 6.25)	0.89 (0.04, 22.04)
Moreno <i>et al</i> (2006)	323	9	0	278	6	0	1.29 (0.45, 3.67)	NA
Colebatch <i>et al</i> (2006)	859	8	0	473	4	0	1.10 (0.33, 3.67)	NA
Koessler T (2007) ^c	2198	25	2	2204	31	0	0.81 (0.48, 1.37)	5.01 (0.24, 104.49)
Küry <i>et al</i> (2007)	999	21	0	1100	16	0	1.45 (0.75, 2.79)	NA
Balaguer <i>et al</i> (2007)	1089	15	1	912	20	0	0.63 (0.32, 1.23)	2.51 (0.10, 61.75)
SOCCS (2008) ^d	3429	56	8	2993	42	0	1.16 (0.78, 1.74)	14.84 (0.86, 257.19)
Avezzu <i>et al</i> (2008)	435	2	1	246	0	0	2.83 (0.14, 59.19)	1.70 (0.07, 41.84)
Cleary <i>et al</i> (2009) ^e	3697	63	11	2758	32	0	1.47 (0.96, 2.25)	17.16 (1.01, 291.31)
Lubbe <i>et al</i> (2009) ^f	9043	128	10	4962	75	0	0.94 (0.70, 1.25)	11.52 (0.68, 196.69)
<i>M-H pooled effect (fixed)</i>	25 231	350	35	18 285	236	0	1.07 (0.90, 1.26)	6.47 (2.33, 17.97)
<i>P-value</i>							0.44	<0.0005
<i>Heterogeneity</i>								
<i>P-value</i>							0.74	0.73
<i>I² (95% CI)</i>							0 (0, 55)	0 (0, 68)

Abbreviations: CI = confidence interval; MM = mutated/mutated; NA = not available; OR = odds ratio; WM = wild type/mutant; WW = wild type/wild type. ^aThis table presents the data as they were published. Two unpublished studies included. ^bUnpublished data obtained in 2006. ^cUnpublished data obtained in 2007. ^dIncludes data from Farrington *et al* (2005), Tenesa *et al* (2006) and unpublished data from the SOCCS study obtained in 2008. ^eIncludes data from Croitoru *et al* (2004). ^fIncludes data from Webb *et al* (2006) and Fleischmann *et al* (2004).

(95% CI: 5.02–23.2)), both G396D and Y179C variants demonstrated bi-allelic effects in this pooled analysis (OR = 6.47 (95% CI: 2.33–18.0) and OR = 3.35 (95% CI: 1.14–9.89), respectively). A marginal significant mono-allelic effect was demonstrated for the specific variant Y179C (OR = 1.34 (95% CI: 1.00–1.80)) and indeed a marginally significant result was also observed in the pooled meta-analysis for *MUTYH* WM (OR = 1.16 (95% CI: 1.00–1.34))

and Y179C variant alone, 1.34 (95% CI: 1.01–1.77). The increased bi-allelic risk of CRC varied when stratified for age and sex but none of the differences were significant, although when stratified by sex, males showed a marginal significant mono-allelic effect for Y179C (OR = 1.70; 95% CI: 1.06–2.73). The results from this large dataset indicate that the two variants may be acting mechanistically differently; G396D appears to be a true example of

Table 5 Meta-analysis of studies for the Y179C genotypes^a

Study	Y179C (cases)			Y179C (controls)			AG vs AA	GG vs AA
	AA	AG	GG	AA	AG	GG	OR (95% CI)	OR (95% CI)
Enholm <i>et al</i> (2003)	994	1	0	424	0	0	1.28 (0.05, 31.50)	NA
Kambara <i>et al</i> (2004)	90	0	0	52	—	0	Not enough data available	NA
Wang <i>et al</i> (2004)	432	5	1	309	2	0	1.79 (0.35, 9.28)	2.15 (0.09, 52.87)
Zhou <i>et al</i> (2005)	432	3	0	466	2	0	1.62 (0.35, 9.28)	NA
Peterlongo <i>et al</i> (2005)	549	2	0	911	2	0	1.66 (0.23, 11.81)	NA
Tomlinson I (2006) ^b	662	6	0	197	0	0	3.88 (0.22, 69.10)	NA
Moreno <i>et al</i> (2006)	323	0	0	278	0	0	NA	NA
Colebatch <i>et al</i> (2006)	859	3	0	473	1	0	1.65 (0.17, 15.93)	NA
Koessler T (2007) ^c	2198	12	1	2204	11	0	1.09 (0.48, 2.48)	3.01 (0.12, 73.88)
Küry <i>et al</i> (2007)	999	3	0	1100	4	0	0.83 (0.18, 3.70)	NA
Balaguer <i>et al</i> (2007)	1089	4	2	912	1	0	3.35 (0.37, 30.02)	4.19 (0.20, 87.34)
SOCCS (2008) ^d	3429	21	3	2993	15	0	1.22 (0.63, 2.38)	6.11 (0.32, 118.34)
Avezù <i>et al</i> (2008)	435	0	0	246	1	0	0.19 (0.01, 4.65)	NA
Cleary <i>et al</i> (2009) ^e	3697	15	5	2758	10	1	1.12 (0.50, 2.50)	3.73 (0.44, 31.95)
Lubbe <i>et al</i> (2009) ^f	9043	70	4	4962	26	1	1.48 (0.94, 2.32)	2.20 (0.25, 19.64)
M-H pooled effect (fixed)	25 231	145	16	18 285	75	2	1.34 (1.01, 1.77)	3.35 (1.14, 9.89)
P-value							0.04	0.03
Heterogeneity								
P-value							0.98	0.99
I ² (95% CI)							0 (0, 57)	0 (0, 75)

Abbreviations: CI = confidence interval; MM = mutated/mutated; NA = not available; OR = odds ratio; WM = wild type/mutant; WWW = wild type/wild type. ^aThis table presents the data as they were published. Two unpublished studies included. ^bUnpublished data obtained in 2006. ^cUnpublished data obtained in 2007. ^dIncludes data from Farrington *et al* (2005), Tenesa *et al* (2006) and unpublished data from the SOCCS study obtained in 2008. ^eIncludes data from Croitoru *et al* (2004). ^fIncludes data from Webb *et al* (2006) and Fleischmann *et al* (2004).

recessive Mendelian disease, whereas the results for Y179C are more complex and there is therefore some argument against combining the two variants. However, the results from the Y179C/G396D compound heterozygotes analysis demonstrates an increase in risk similar to G396D bi-allelic carriers, suggesting that the two variants are complementary and analysis of combined *MUTYH* mutations as historically performed, appears appropriate to assess risk for the whole gene. The rarity of the Y179C allele has made it difficult to truly assess its effect on disease risk, however the large numbers analysed in this report have resulted in the demonstration that both bi-allelic and mono-allelic Y179C variants are associated with disease risk.

The study population did not appear to modulate disease risk and although the study replicated the reported decrease in disease risk in *MUTYH* wild-type females associated with HRT intake (Chan *et al*, 2006; Theodoratou *et al*, 2008), we found no interaction with the *MUTYH* gene and its variants. Therefore, it is unlikely that HRT intake is an explanation for any sex variation in risk and other genetic factors may be involved in modifying CRC risk.

Evidence of a mono-allelic *MUTYH* effect on CRC has been reported in several case-control studies (Croitoru *et al*, 2004; Wang *et al*, 2004; Farrington *et al*, 2005; Zhou *et al*, 2005; Tenesa *et al*, 2006; Cleary *et al*, 2009) and family-based studies (Jenkins *et al*, 2006; Jones *et al*, 2009), but not in other studies (Kambara *et al*, 2004; Webb *et al*, 2006; Balaguer *et al*, 2007; Lubbe *et al*, 2009). Our large meta-analysis has demonstrated a marginal significant association for the specific variant Y179C, highlighting the possible increased phenotypic severity of this allele. This is in agreement with other studies and biochemical and model organism studies, which indicate that this variant shows an increased detrimental effect on protein function (Al-Tassan *et al*, 2002; Parker *et al*, 2005; Lubbe *et al*, 2009; Nielsen *et al*, 2009; D'Agostino *et al*, 2010). The pooled analysis of published studies and unpublished datasets submitted to us also indicated a marginally significant mono-allelic *MUTYH* effect, as well as a mono-allelic Y179C effect.

However, there are a number of caveats that need to be considered; if any of the studied datasets contain cases recruited because of the familial clustering of disease, there may be ascertainment bias, artificially inflating the number of *MUTYH* WM variant allele carriers; secondly the screening of the *MUTYH* gene has predominantly been performed on the two most common pathogenic variants Y179C and G396D – in some studies, the rest of the gene may be explored in cases with a heterozygous allele for these variants but not usually in the controls, hence there is an overall screening bias and bi-allelic carriers may well have been missed in both cases and controls.

The demonstration of a mono-allelic effect specifically for Y179C should be considered with further caution, as analysis of the control datasets for the Y179C allele demonstrated that it was not in Hardy-Weinberg equilibrium. This may be because of several factors, the rarity of the allele and the fact that both female control subjects with bi-allelic mutations carry Y179C variants. One of these control subjects was shown to have polyps on colonoscopy (Cleary *et al*, 2009) and may therefore be considered a case. The other is relatively young, less than 60 years old (Lubbe *et al*, 2009), so potentially may develop cancer over the next few years. However, in this large dataset, we have also shown that bi-allelic carriers of Y179C predisposes to an earlier onset of disease than G396D, consistent with previous reports (Lubbe *et al*, 2009; Nielsen *et al*, 2009) and highlights a severer disease phenotype of this variant.

In conclusion, inactivation of the *MUTYH* gene is a recessive risk factor for CRC, with possible modifying effects indicated by increased risk in cases with early age of onset, although not significantly different in the current dataset. An increased risk associated with mono-allelic *MUTYH* mutation is indicated, albeit small and not currently clinically relevant, and likely specific for the variant Y179C. Despite the size of this study it has not been possible to definitively establish whether there are significant age and sex effects of increasing disease risk for G396D and Y179C carriers. The evidence presented raises the possibility of a mono-allelic effect for Y179C, but the effect is low (OR 1.34; 95% CI:

1.00–1.80) and is sensitive to variations in population allele frequency because of the rarity of the variant (allele frequency 0.002), as well as potential issues of subgroup analysis and multiple testing (indeed the overall significance is lost after Bonferroni correction). Nonetheless, it does appear that this study is the first to demonstrate that the Y179C variant does impart an increased risk of CRC.

ACKNOWLEDGEMENTS

We thank all the study participants and the many colleagues worldwide who contributed to recruitment. We are particularly grateful to Ruth Wilson, Rosa Bisset, Nicola Cartwright and Gisela Johnstone, and all those who contributed to recruitment, data collection and data curation for the Scottish COGS and SOCCS studies. This work was supported by grants from Cancer Research UK (C348/A3758, C348/A8896), Scottish Government Chief Scientist Office (K/OPR/2/2/D333, CZB/4/94, CZB/4/449); Medical Research Council (G0000657-53203); Centre Grant from CORE as part of the Digestive Cancer Campaign (<http://www.corecharity.org.uk>). ET is funded by a Cancer Research UK Fellowship C31250/A10107. The work at the Institute of Cancer Research is supported by grants from Cancer Research UK (supported by Bobby Moore), Steven Lubbe is supported by a Cancer Research UK PhD Studentship. The work in Canada was made possible through collaboration and cooperative agreements with the Colon Cancer

Family Registry and PIs. The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating institutions or investigators in the Colon CFR, nor does mention of trade names, commercial products, or organisations imply endorsement by the US Government or the Colon CFR. The work in Cambridge (SEARCH) is funded through a programme grant from Cancer Research UK, TK is funded by the Foundation Dr Henri Dubois-Ferriere Dinu Lipatti. The work in France was supported by a regional Hospital Clinical Research Program, the Regional Council of Pays de la Loire, the Groupement des Entreprises Françaises dans la Lutte contre le Cancer, the Association Anne de Bretagne Génétique and the Ligue Régionale Contre le Cancer. The work in Stockholm was supported by The Swedish Cancer Society, The Swedish Research Council and The Stockholm Cancer Foundation. The work in Spain was supported by the Ministerio de Educación y Ciencia (SAF09-7319), Spanish Networks RTICCC (RD06/0020/1050, 1051), Instituto de Salud Carlos III grants (FIS PI03/0114 and FIS 05/1006 and CIBERESP).

Conflict of interest

The authors declare no conflict of interest.

Supplementary Information accompanies the paper on British Journal of Cancer website (<http://www.nature.com/bjcr>)

REFERENCES

- Al-Tassan N, Chmiel NH, Maynard J, Fleming N, Livingston AL, Williams GT, Hodges AK, Davies DR, David SS, Sampson JR, Cheadle JP (2002) Inherited variants of MYH associated with somatic G:C→T:A mutations in colorectal tumors. *Nat Genet* 30(2): 227–232
- Avezù A, Agostini M, Pucciarelli S, Lise M, Urso ED, Mammi I, Maretto I, Enzo MV, Pastrello C, Lise M, Nitti D, Viel A (2008) The role of MYH gene in genetic predisposition to colorectal cancer: another piece of the puzzle. *Cancer Lett* 268(2): 308–313
- Balaguer F, Castellví-Bel S, Castells A, Andreu M, Muñoz J, Gisbert JP, Llor X, Jover R, de Cid R, Gonzalo V, Bessa X, Xicola RM, Pons E, Alenda C, Payá A, Piqué JM (2007) Identification of MYH mutation carriers in colorectal cancer: a multicenter, case-control, population-based study. *Clin Gastroenterol Hepatol* 5: 379–387
- Chan JA, Meyerhardt JA, Chan AT, Giovannucci EL, Colditz GA, Fuchs CS (2006) Hormone replacement therapy and survival after colorectal cancer diagnosis. *J Clin Oncol* 24: 5680–5686
- Cleary SP, Cotterchio M, Jenkins MA, Kim H, Bristow R, Green R, Haile R, Hopper JL, LeMarchand L, Lindor N, Parfrey P, Potter J, Youngusband B, Gallinger S (2009) Germline *MutY* human homologue mutations and colorectal cancer: a multi-site case-control study. *Gastroenterology* 136(4): 1251–1260
- Colebatch A, Hitchins M, Williams R, Meagher A, Hawkins NJ, Ward RL (2006) The role of MYH and microsatellite instability in the development of sporadic colorectal cancer. *Br J Cancer* 95(9): 1239–1243
- Croituru ME, Cleary SP, Di Nicola N, Manno M, Selander T, Aronson M, Redston M, Cotterchio M, Knight J, Gryfe R, Gallinger S (2004) Association between biallelic and monoallelic germline MYH gene mutations and colorectal cancer risk. *J Natl Cancer Inst* 96(21): 1631–1634
- D'Agostino VG, Minoprio A, Torrieri P, Marinoni I, Bossa C, Petrucci TC, Albertini AM, Ranzani GN, Bignami M, Mazzei F (2010) Functional analysis of *MUTYH* mutated proteins associated with familial adenomatous polyposis. *DNA Repair* 9: 700–707
- Enholm S, Hienonen T, Suomalainen A, Lipton L, Tomlinson I, Kärjä V, Eskelinen M, Mecklin JP, Karhu A, Järvinen HJ, Aaltonen LA (2003) Proportion and phenotype of MYH associated colorectal neoplasia in a population-based series of Finnish colorectal cancer patients. *Am J Pathol* 163(3): 827–832
- Farrington SM, Tenesa A, Barnetson R, Wiltshire A, Prendergast J, Porteous M, Campbell H, Dunlop MG (2005) Germline susceptibility to colorectal cancer due to base-excision repair gene defects. *Am J Hum Genet* 77(1): 112–119
- Farrington SM, Tenesa A, Barnetson R, Wiltshire A, Prendergast J, Porteous M, Campbell H, Dunlop MG (2006) Colorectal cancer risk in monoallelic carriers of MYH variants. *Am J Hum Genet* 79(4): 771–772
- Fleischmann C, Peto J, Cheadle J, Shah B, Sampson J, Houlston RS (2004) Comprehensive analysis of the contribution of germline MYH variation to early-onset colorectal cancer. *Int J Cancer* 109(4): 554–558
- Gismondi V, Meta M, Bonelli L, Radice P, Sala P, Bertario L, Viel A, Fornasarig M, Arrigoni A, Gentile M, Ponz de Leon M, Anselmi L, Mareni C, Bruzzi P, Varesco L (2004) Prevalence of the Y165C, G382D and 1395delGGA germline mutations of the MYH gene in Italian patients with adenomatous polyposis coli and colorectal adenomas. *Int J Cancer* 109(5): 680–684
- Jenkins MA, Croitoru ME, Monga N, Cleary SP, Cotterchio M, Hopper JL, Gallinger S (2006) Risk of colorectal cancer in monoallelic and biallelic carriers of MYH mutations: a population-based case-family study. *Cancer Epidemiol Biomarkers Prevention* 15(2): 312–314
- Jones N, Vogt S, Nielsen M, Christian D, Wark PA, Eccles D, Edwards E, Evans DG, Maher ER, Vasen HF, Hes FJ, Aretz S, Sampson JR (2009) Increased colorectal cancer incidence in obligate carriers of heterozygous mutations in *MUTYH*. *Gastroenterology* 137(2): 489–494
- Kambara T, Whitehall VL, Spring KJ, Barker MA, Arnold S, Wynter CV, Matsubara N, Tanaka N, Young JP, Leggett BA, Jass JR (2004) Role of inherited defects of MYH in the development of sporadic colorectal cancer. *Genes Chromosomes Cancer* 40(1): 1–9
- Küry S, Buecher B, Robiou-du-Pont S, Scoul C, Colman H, Lelièvre B, Olschwang S, Le Houérou C, Le Neel T, Faroux R, Ollivry J, Lafraise B, Chupin LD, Bézieau S (2007) The thorough screening of the *MUTYH* gene in a large French cohort of sporadic colorectal cancers. *Genet Test* 11(4): 373–379
- Lubbe SJ, Di Bernardo MC, Chandler IP, Houlston RS (2009) Clinical implications of the colorectal cancer risk associated with *MUTYH* mutation. *J Clin Oncol* 27: 3975–3980
- Moreno V, Gemignani F, Landi S, Gioia-Patricola L, Chabrier A, Blanco I, González S, Guino E, Capella G, Canzian F (2006) Polymorphisms in genes of nucleotide and base excision repair: risk and prognosis of colorectal cancer. *Clin Cancer Res* 12: 2101–2108
- Nielsen M, Joerink-van de Beld MC, Jones N, Vogt S, Tops CM, Vasen HF, Sampson JR, Aretz S, Hes FJ (2009) Analysis of *MUTYH* genotypes and colorectal phenotypes in patients with *MUTYH*-associated polyposis. *Gastroenterology* 136(2): 471–476

- Parker AR, Sieber OM, Shi C, Hua L, Takao M, Tomlinson IP, Eshleman JR (2005) Cells with pathogenic biallelic mutations in the human *MUTYH* gene are defective in DNA damage binding and repair. *Carcinogenesis* **26**(11): 2010–2018
- Peterlongo P, Mitra N, Chuai S, Kirchoff T, Palmer C, Huang H, Nafa K, Offit K, Ellis NA (2005) Colorectal cancer risk in individuals with biallelic or monoallelic mutations of MYH. *Int J Cancer* **114**(3): 505–507
- Sampson JR, Dolwani S, Jones S, Eccles D, Ellis A, Evans DG, Frayling I, Jordan S, Maher ER, Mak T, Maynard J, Pigatto F, Shaw J, Cheadle JP (2003) Autosomal recessive colorectal adenomatous polyposis due to inherited mutations of MYH. *Lancet* **362**: 39–41
- Sieber OM, Lipton L, Crabtree M, Heinemann K, Fidalgo P, Phillips RK, Bisgaard ML, Orntoft TF, Aaltonen LA, Hodgson SV, Thomas HJ, Tomlinson IP (2003) Multiple colorectal adenomas, classic adenomatous polyposis, and germ-line mutations in MYH. *N Engl J Med* **348**(9): 791–799
- Tenesa A, Campbell H, Barnetson R, Porteous M, Dunlop M, Farrington SM (2006) Association of *MUTYH* and colorectal cancer. *Br J Cancer* **95**(2): 239–242
- Theodoratou E, Campbell H, Tenesa A, McNeill G, Cetnarskyj R, Barnetson RA, Porteous ME, Dunlop MG, Farrington SM (2008) Modification of the associations between lifestyle, dietary factors and colorectal cancer risk by APC variants. *Carcinogenesis* **29**(9): 1774–1780
- Venesio T, Molatore S, Cattaneo F, Arrighoni A, Risio M, Ranzani GN (2004) High frequency of MYH gene mutations in a subset of patients with familial adenomatous polyposis. *Gastroenterology* **126**(7): 1681–1685
- Wang L, Baudhuin LM, Boardman LA, Steenblock KJ, Petersen GM, Halling KC, French AJ, Johnson RA, Burgart LJ, Rabe K, Lindor NM, Thibodeau SN (2004) MYH mutations in patients with attenuated and classic polyposis and with young-onset colorectal cancer without polypos. *Gastroenterology* **127**(1): 9–16
- Webb EL, Rudd MF, Houlston RS (2006) Colorectal cancer risk in monoallelic carriers of MYH variants. *Am J Hum Genet* **79**(4): 768–771
- Zhou X-L, Djureinovic T, Werelius B, Lindmark G, Sun XF, Lindblom A (2005) Germline mutations in the MYH gene in Swedish familial and sporadic colorectal cancer. *Genetic Test* **9**: 147–151