

This is a repository copy of *Analysing causal structures with entropy*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/133771/>

Version: Accepted Version

Article:

Weilenmann, Mirjam Sarah and Colbeck, Roger Andrew orcid.org/0000-0003-3591-0576
(2017) *Analysing causal structures with entropy*. Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences. 20170483. ISSN 1364-5021

<https://doi.org/10.1098/rspa.2017.0483>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Analysing causal structures with entropy

Mirjam Weilenmann* and Roger Colbeck†

Department of Mathematics, University of York, Heslington, York, YO10 5DD, UK.

(Dated: 26th September 2017)

A central question for causal inference is to decide whether a set of correlations fit a given causal structure. In general, this decision problem is computationally infeasible and hence several approaches have emerged that look for certificates of compatibility. Here we review several such approaches based on entropy. We bring together the key aspects of these entropic techniques with unified terminology, filling several gaps and establishing new connections regarding their relation, all illustrated with examples. We consider cases where unobserved causes are classical, quantum and post-quantum and discuss what entropic analyses tell us about the difference. This has applications to quantum cryptography, where it can be crucial to eliminate the possibility of classical causes. We discuss the achievements and limitations of the entropic approach in comparison to other techniques and point out the main open problems.

I. INTRODUCTION

Deciding whether a causal explanation is compatible with given statistical data and exploring whether it is the most suitable explanation for the data at hand are central scientific tasks. Sometimes the most reasonable explanation of a set of observations involves unobserved common causes. In the case where the common causes are classical, the well-developed machinery of Bayesian networks can be used [1, 2]. In principle, such networks are well-understood and it is known how to check whether observed correlations are compatible with a given network [3]. In practice, however, testing compatibility for networks that involve unobserved systems is only computationally tractable for small cases [4, 5]. Furthermore, the methodology has to be adapted whenever non-classical common causes are permitted.

Finding good heuristics to help identify correlations that are (in)compatible with a causal structure is currently an active area of research [6–20] and the use of entropy measures is common to many of these [6–13, 17, 18, 20]. Such methods are important in the quantum context, where recent cryptographic protocols rely on the lack of a classical causal explanation for certain quantum correlations in specified causal structures [21–29], an idea that lies behind Bell’s theorem [30] (see also [31]).

In Section II of this article, we review the entropic characterisation of the correlations compatible with causal structures in classical, quantum and more general non-signalling theories. We detail refinements of the approach based on post-selection in Section III. Together, these sections show the current capabilities of entropic techniques, also establishing and clarifying connections between different contributions. Our review is illustrated with several examples to assist its understanding and to make it easily accessible for applications. In Section IV we outline and compare further approaches to the problem, before concluding in Section V with some open questions.

II. ENTROPY VECTOR APPROACH

Characterising the joint distributions of a set of random variables or alternatively considering a multi-party quantum state in terms of its entropy (and of those of its marginals) has a tradition in information theory, dating back to Shannon [32–34]. However, only recently has this approach been extended to account for causal structure [6, 8]. In Sections II A and II B respectively, we review this approach with and without imposing causal constraints. All our considerations are concerned with discrete random variables, for extensions of the approach to continuous random variables (and its limitations) we refer to [8, 35].

*Electronic address: msw518@york.ac.uk

†Electronic address: roger.colbeck@york.ac.uk

A. Classical entropy cone

The entropy cone for a joint distribution of n random variables was introduced in [33]. It is defined in terms of the *Shannon entropy* [32], which for a discrete random variable X taking values $x \in \mathcal{X}$ with probability distribution P_X is defined by

$$H(X) := - \sum_{x \in \mathcal{X}} P_X(x) \log_2 P_X(x),$$

where $0 \log_2(0)$ is taken to be 0^1 .

For a set of $n \geq 2$ jointly distributed random variables, $\Omega := \{X_1, X_2, \dots, X_n\}$, we denote their probability distribution as $P \in \mathcal{P}_n$, where \mathcal{P}_n is the set of all probability mass functions of n jointly distributed random variables. The Shannon entropy maps any subset of Ω to a non-negative real value: $H : \mathcal{P}(\Omega) \rightarrow [0, \infty)$, $X_S \mapsto H(X_S)$, where $\mathcal{P}(\Omega)$ denotes the power set of Ω , and $H(\{\}) = 0$. The entropy of the joint distribution of the random variables Ω and of all its marginals can be expressed as components of a vector in $\mathbb{R}_{\geq 0}^{2^n - 1}$, ordered in the following as²

$$(H(X_1), H(X_2), \dots, H(X_n), H(X_1 X_2), H(X_1 X_3), \dots, H(X_1 X_2 \dots X_n)).$$

We use $\mathbf{H}(P) \in \mathbb{R}_{\geq 0}^{2^n - 1}$ to denote the vector corresponding to a particular distribution $P \in \mathcal{P}_n$. The set of all such vectors is $\Gamma_n^* := \{v \in \mathbb{R}_{\geq 0}^{2^n - 1} \mid \exists P \in \mathcal{P}_n \text{ s.t. } v = \mathbf{H}(P)\}$. Its closure $\overline{\Gamma_n^*}$ includes vectors v for which there exists a sequence of distributions $P_k \in \mathcal{P}_n$ such that $\mathbf{H}(P_k)$ tends to v as $k \rightarrow \infty$. It is known that the *entropy cone* $\overline{\Gamma_n^*}$ is a convex cone for any $n \in \mathbb{N}$ [36]. As such, its boundary may be characterised in terms of (potentially infinitely many) linear inequalities. Because $\overline{\Gamma_n^*}$ is difficult to characterise, we will in the following consider various approximations.

1. Outer approximation: the Shannon cone

The standard outer approximation to $\overline{\Gamma_n^*}$ is the polyhedral cone constrained by the *Shannon inequalities* listed in the following³:

- Monotonicity: For all $X_T, X_S \subseteq \Omega$, $H(X_S \setminus X_T) \leq H(X_S)$.
- Submodularity: For all $X_S, X_T \subseteq \Omega$, $H(X_S \cap X_T) + H(X_S \cup X_T) \leq H(X_S) + H(X_T)$.

These inequalities are always obeyed by the entropies of a set of jointly distributed random variables. They may be concisely rewritten in terms of the following information measures: the *conditional entropy* of two jointly distributed random variables X and Y , $H(X|Y) := H(XY) - H(Y)$, their *mutual information*, $I(X : Y) := H(X) + H(Y) - H(XY)$, and the *conditional mutual information* between two jointly distributed random variables X and Y given a third, Z , denoted $I(X : Y|Z) := H(XZ) + H(YZ) - H(Z) - H(XYZ)$. Hence, the monotonicity constraints correspond to positivity of conditional entropy, $H(X_S \cap X_T | X_S \setminus X_T) \geq 0$, and submodularity is equivalent to positivity of the conditional mutual information, $I(X_S \setminus X_T : X_T \setminus X_S | X_S \cap X_T) \geq 0$. The monotonicity and submodularity constraints can all be generated from a minimal set of $n + n(n-1)2^{n-3}$ inequalities [33]: for the monotonicity constraints it is sufficient to consider the n constraints with $X_S = \Omega$ and $X_T = X_i$ for some $X_i \in \Omega$; for the submodularity constraints it is sufficient to consider those with $X_S \setminus X_T = X_i$ and $X_T \setminus X_S = X_j$ with $i < j$ and where $X_U := X_S \cap X_T$ is any subset of Ω not containing X_i or X_j , i.e., submodularity constraints of the form $I(X_i : X_j | X_U) \geq 0$.

These $n + n(n-1)2^{n-3}$ independent Shannon inequalities can be expressed in terms of a $(n + n(n-1)2^{n-3}) \times (2^n - 1)$ dimensional matrix, which we call M_{SH}^n , such that for any $v \in \Gamma_n^*$, the conditions $M_{\text{SH}}^n \cdot v \geq 0$ hold⁴. More generally, for $v \in \mathbb{R}_{\geq 0}^{2^n - 1}$, a violation of $M_{\text{SH}}^n \cdot v \geq 0$ certifies that there is no distribution $P \in \mathcal{P}_n$ such that $v = \mathbf{H}(P)$. It follows that the *Shannon cone*, $\Gamma_n := \{v \in \mathbb{R}_{\geq 0}^{2^n - 1} \mid M_{\text{SH}}^n \cdot v \geq 0\}$, is an outer approximation of the set of achievable entropy vectors, Γ_n^* [33].

¹ Note $\lim_{p \rightarrow 0^+} p \log_2 p = 0$

² Since the empty set always has zero entropy, we choose to omit it from the entropy vector in this work.

³ It is a matter of convention, whether $H(\{\}) = 0$ is included as a Shannon inequality; we keep this implicit.

⁴ This condition is to be interpreted as the requirement that each component of $M_{\text{SH}}^n \cdot v$ is non-negative.

Example 1. The three variable Shannon cone is $\Gamma_3 = \{v \in \mathbb{R}_{\geq 0}^7 \mid M_{\text{SH}}^3 \cdot v \geq 0\}$, where

$$M_{\text{SH}}^3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 1 \\ 1 & 1 & 0 & -1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & -1 & 0 \\ -1 & 0 & 0 & 1 & 1 & 0 & -1 \\ 0 & -1 & 0 & 1 & 0 & 1 & -1 \\ 0 & 0 & -1 & 0 & 1 & 1 & -1 \end{pmatrix}.$$

The first three rows are monotonicity constraints, the remaining six ensure submodularity.

2. Beyond the Shannon cone

For two variables the Shannon cone coincides with the actual entropy cone, $\Gamma_2 = \Gamma_2^*$, while for three random variables this holds only for the closure of the entropic cone, i.e. $\Gamma_3 = \overline{\Gamma_3^*}$ but $\Gamma_3 \neq \Gamma_3^*$ [36, 37]. For $n \geq 4$ further independent constraints on the set of entropy vectors are needed to fully characterise $\overline{\Gamma_n^*}$, the first of which was discovered in [38].

Proposition 1 (Zhang & Yeung). *For any four discrete random variables T, U, V and W the following inequality holds: $-H(T) - H(U) - \frac{1}{2}H(V) + \frac{3}{2}H(TU) + \frac{3}{2}H(TV) + \frac{1}{2}H(TW) + \frac{3}{2}H(UV) + \frac{1}{2}H(UW) - \frac{1}{2}H(VW) - 2H(TUV) - \frac{1}{2}H(TUW) \geq 0$.*

For $n \geq 4$ the convex cone $\overline{\Gamma_n^*} \not\subseteq \Gamma_n$ is not polyhedral, i.e., it cannot be characterised by finitely many linear inequalities [39]. Nonetheless, many linear entropic inequalities have been discovered [36, 38, 40, 41]. Recently, systematic searches for new entropic inequalities for $n = 4$ have been conducted [42, 43], which recover most of the previously known inequalities; in particular the inequality of Proposition 1 is re-derived and shown to be implied by tighter ones [43]. The systematic search in [43] is based on considering additional random variables that obey certain constraints and then deriving four variable inequalities from the known identities for five or more random variables (see also [38, 39]), an idea that is captured by a so-called copy lemma [38, 43, 44]. In the same article, several rules to generate families of inequalities have been suggested, in the style of techniques introduced by Matúš [39].

For more than four variables, a few additional inequalities are known [38, 40]. Curiously, to our knowledge, all known relevant non Shannon inequalities (i.e., the ones found in [38–43] that are not yet superseded by tighter ones) can be written as a positive linear combination of the *Ingleton quantity*, $I(T : U|V) + I(T : U|W) + I(V : W) - I(T : U)$, and conditional mutual information terms (see also [43]).

3. Inner approximations

For the four variable entropy cone, $\overline{\Gamma_4^*}$, an inner approximation is defined as the region constrained by the Shannon inequalities and the six permutations of the *Ingleton inequality* [45], $I(T : U|V) + I(T : U|W) + I(V : W) - I(T : U) \geq 0$, for random variables T, U, V and W . These inequalities can be concisely written as a matrix $M_I \in \mathbb{R}^6 \times \mathbb{R}^{15}$. The constrained region is called the *Ingleton cone*, $\Gamma^I := \{v \in \mathbb{R}_{\geq 0}^{15} \mid M_{\text{SH}}^4 \cdot v \geq 0 \text{ and } M_I \cdot v \geq 0\}$, and it has the property that $v \in \Gamma^I$ implies $v \in \Gamma_n^*$ [46]. In contrast, there are entropy vectors that violate the Ingleton inequalities, as the following example shows.

Example 2. Let T, U, V and W be four jointly distributed random variables. Let V and W be uniform random bits and let $T = \text{AND}(-V, -W)$ and $U = \text{AND}(V, W)$. This distribution [39] leads to the entropy vector $v \approx (0.81, 0.81, 1, 1, 1.50, 1.50, 1.50, 1.50, 1.50, 2, 2, 2, 2, 2, 2)$, for which $I(T : U|V) + I(T : U|W) + I(V : W) - I(T : U) \approx -0.12$ in violation of the Ingleton inequality.

For five random variables an inner approximation in terms of Shannon, Ingleton and 24 additional inequalities and their permutations is known (including partial extensions to more variables) [47, 48].

B. Entropy cones for causal structures

Causal relations among a set of variables impose constraints on their possible joint distributions, which can be conveniently represented with a causal structure.




Figure 1: (a) Pearl's instrumental scenario. The nodes X , Y and Z are observed, A is unobserved. In the classical case this can be understood in the following way: A random variable X and an unobserved A are used to generate another random variable Z . Then Y is generated from A and the observed output of node Z . In particular, note that no other information can be forwarded from X through the node Z to Y . In the quantum case, the source A shares a quantum system $\rho_A \in \mathcal{S}(\mathcal{H}_A)$, where $\mathcal{H}_A \cong \mathcal{H}_{A_Z} \otimes \mathcal{H}_{A_Y}$. The subsystem A_Z is measured to produce Z and likewise for Y . The subsystems A_Z and A_Y are both considered to be parents of Z (and Y). (b) Bell scenario. The observed variables A and B together with an unobserved system C are used to generate outputs X and Y respectively. In the classical case, C can be modelled as a random variable, in the quantum case it is a quantum state on a Hilbert space $\mathcal{H}_C \cong \mathcal{H}_{C_X} \otimes \mathcal{H}_{C_Y}$. (c) Triangle causal structure. Three observed random variables X , Y and Z share pairwise common causes A , B and C , which in the classical case are modelled by random variables. Some of the valid inequalities such as $2I(X : Y | Z) + I(X : Z | Y) + I(Y : Z | X) - I(X : Y) \geq 0$ can only be recovered using non-Shannon entropic inequalities [50].

Definition 2. A *causal structure* is a set of variables arranged in a directed acyclic graph (DAG), in which a subset of the nodes is assigned as observed.

The directed edges of the graph are intended to represent causation, perhaps by propagation of some influence, and cycles are excluded to avoid the well-known paradoxes associated with causal loops. We will interpret causal structures in different ways depending on the supposed physics of whatever is mediating the causal influence.

One of the simplest causal structures that leads to interesting insights and one of the most thoroughly analysed ones is Pearl's instrumental causal structure, *IC* [49]. It is displayed in Figure 1(a) and will be used as an example throughout this review.

1. Classical causal structures

In the classical case, the causal relations among a set of random variables can be explored by means of the theory of Bayesian networks (see for instance [1, 2] for a complete presentation of this theory).

Definition 3. A *classical causal structure*, C^C , is a causal structure in which each node of the DAG has an associated random variable.

It is common to use the same label for the node and its associated random variable. The DAG encodes which joint distributions of the involved variables are allowed in a causal structure C^C . To explain this we need a little more terminology.

Definition 4. Let X_S , X_T , X_U be three disjoint sets of jointly distributed random variables. Then X_S and X_T are said to be *conditionally independent* given X_U if and only if their joint distribution $P_{X_S X_T X_U}$ can be written as $P_{X_S X_T X_U} = P_{X_S | X_U} P_{X_T | X_U} P_{X_U}$. Conditional independence of X_S and X_T given X_U is denoted as $X_S \perp\!\!\!\perp X_T | X_U$.

Two variables X_S and X_T are (unconditionally) independent if $P_{X_S X_T} = P_{X_S} P_{X_T}$, concisely written $X_S \perp\!\!\!\perp X_T$. With reference to a DAG with a subset of nodes, X , we will use X^\downarrow to denote the ancestors of X and X^\uparrow to denote the descendants of X . The parents of X are represented by $X^{\downarrow 1}$ and the non-descendants are X^\uparrow .

Definition 5. Let C^C be a classical causal structure with nodes $\{X_1, X_2, \dots, X_n\}$. A probability distribution $P_{X_1 X_2 \dots X_n} \in \mathcal{P}_n$ is (Markov) *compatible* with C^C if it can be decomposed as $P_{X_1 X_2 \dots X_n} = \prod_i P_{X_i | X_i^{\downarrow 1}}$.

The compatibility constraint encodes all conditional independences of the random variables in the causal structure C^C . Nonetheless, whether a particular set of variables is conditionally independent of another is more easily read from the DAG, as explained in the following.

Definition 6. Let X , Y and Z be three pairwise disjoint sets of nodes in a DAG G . The sets X and Y are said to be *d-separated*⁵ by Z , if Z blocks any path from any node in X to any node in Y . A path is *blocked* by Z , if the path

⁵ The d in d-separation stands for *directional*.



Figure 2: While in the left causal structure $X \perp\!\!\!\perp Y$, the other three networks share the conditional independence relation $X \perp\!\!\!\perp Y | Z$. This illustrates that the conditional independences are not sufficient to characterise the causal links among a set of random variables.

contains one of the following: $i \rightarrow z \rightarrow j$ or $i \leftarrow z \rightarrow j$ for some nodes i, j and a node $z \in Z$ in that path, or if the path contains $i \rightarrow k \leftarrow j$, where $k \notin Z$.

The d-separation of the nodes in a causal structure is directly related to the conditional independence of its variables. The following proposition corresponds to Theorem 1.2.5 from [1], previously introduced in [51, 52]. It justifies the application of d-separation as a means to identify independent variables.

Proposition 7 (Verma & Pearl). *Let C^C be a classical causal structure and let X, Y and Z be pairwise disjoint subsets of nodes in C^C . If a probability distribution P is compatible with C^C , then the d-separation of X and Y by Z implies the conditional independence $X \perp\!\!\!\perp Y | Z$. Conversely, if for every distribution P compatible with C^C the conditional independence $X \perp\!\!\!\perp Y | Z$ holds, then X is d-separated from Y by Z .*

The compatibility of probability distributions with a classical causal structure is conveniently determined with the following proposition, which has also been called the parental or local Markov condition before (Theorem 1.2.7 in [1]).

Proposition 8 (Pearl). *Let C^C be a classical causal structure. A probability distribution P is compatible with C^C if and only if every variable in C^C is independent of its non-descendants, conditioned on its parents.*

Hence, to establish whether a probability distribution is compatible with a certain classical causal structure, it is enough to check that every variable X is independent of its non-descendants X^\dagger given its parents $X^{\downarrow 1}$, concisely written as $X \perp\!\!\!\perp X^\dagger | X^{\downarrow 1}$, i.e., to check one constraint for each variable. In particular, it is not necessary to explicitly check for all possible sets of nodes whether they obey the independence relations implied by d-separation. Each such constraint can be conveniently expressed as⁶

$$I(X : X^\dagger | X^{\downarrow 1}) = 0. \quad (1)$$

While the conditional independence relations capture some features of the causal structure, they are insufficient to completely capture the causal relations between variables, as illustrated in Figure 2. In this case, the probability distributions themselves are unable to capture the difference between these causal structures: correlations are insufficient to determine causal links between random variables. External interventions allow for the exploration of causal links beyond the conditional independences [1]. However, we do not consider these here.

Let C^C be a classical causal structure involving n random variables $\{X_1, X_2, \dots, X_n\}$. The restricted set of distributions that are compatible with the causal structure C^C is $\mathcal{P}(C^C) := \{P \in \mathcal{P}_n \mid P = \prod_{i=1}^n P_{X_i | X_i^{\downarrow 1}}\}$.

Example 3 (Allowed distributions in the instrumental scenario). The classical instrumental scenario of Figure 1 allows for any four variable distribution in the set $\mathcal{P}(IC^C) = \{P_{AXYZ} \in \mathcal{P}_4 \mid P_{AXYZ} = P_{Y|AZ} P_{Z|AX} P_X P_A\}$.

The restrictions on the allowed distributions also restrict the corresponding entropy cones. Due to Proposition 8 there are at most n independent conditional independence equalities (1) in a causal structure C^C . Their coefficients can be concisely written in terms of a matrix $M_{CI}(C^C)$, where CI stands for conditional independence. For a causal structure C^C , we define the two sets $\Gamma^*(C^C) := \{v \in \Gamma_n^* \mid M_{CI}(C^C) \cdot v = 0\}$ and $\Gamma(C^C) := \{v \in \Gamma_n \mid M_{CI}(C^C) \cdot v = 0\}$, where $\Gamma^*(C^C) \subseteq \Gamma(C^C)$. The following lemma justifies the notation we use for $\Gamma^*(C^C)$; it is the set of achievable entropy vectors in C^C .

Lemma 9. *For a causal structure C^C , $\Gamma^*(C^C) = \{v \in \mathbb{R}_{\geq 0}^{2^n - 1} \mid \exists P = \prod_{i=1}^n P_{X_i | X_i^{\downarrow 1}} \in \mathcal{P}_n \text{ s.t. } v = \mathbf{H}(P)\}$. Furthermore, its topological closure, $\overline{\Gamma^*}(C^C) = \{v \in \overline{\Gamma_n^*} \mid M_{CI}(C^C) \cdot v = 0\}$, is a convex cone.*

⁶ This follows because the relative entropy $D(P\|Q) := \sum_x P_X(x) \log(P_X(x)/Q_X(x))$ satisfies $D(P\|Q) = 0 \iff P = Q$, and because $I(X : X^\dagger | X^{\downarrow 1}) = D(P_{XX^\dagger X^{\downarrow 1}} \| P_{X|X^{\downarrow 1}} P_{X^\dagger | X^{\downarrow 1}} P_{X^{\downarrow 1}})$.

Proof. For the causal structure C^C , let $E(C^C) := \{v \in \mathbb{R}_{\geq 0}^{2^n - 1} \mid \exists P = \prod_{i=1}^n P_{X_i | X_i^{\downarrow 1}} \in \mathcal{P}_n \text{ s.t. } v = \mathbf{H}(P)\}$ be the set of all entropy vectors. Since (1) holds for each variable X_i if and only if $P = \prod_{i=1}^n P_{X_i | X_i^{\downarrow 1}}$ (cf. Proposition 8), $E(C^C) = \{v \in \Gamma_n^* \mid \forall i \in \{1, \dots, n\}, I(X_i : X_i^{\downarrow 1} | X_i^{\downarrow 1}) = 0\}$. Applying the definition of $M_{\text{CI}}(C^C)$ yields $E(C^C) = \Gamma^*(C^C)$. Now, let us consider the set $F(C^C) := \{v \in \overline{\Gamma}_n^* \mid M_{\text{CI}}(C^C) \cdot v = 0\} \subseteq \mathbb{R}^{2^n - 1}$. This is a closed convex set, since $\overline{\Gamma}_n^*$ is known to be closed and convex and since restricting the closed convex cone $\overline{\Gamma}_n^*$ with linear equality constraints retains these properties. More precisely, the set of solutions to the matrix equality $M_{\text{CI}}(C^C) \cdot v = 0$ is also closed and convex. Being the intersection of two closed convex sets, the set $F(C^C)$ is also closed and convex. From this we conclude that $\overline{\Gamma}^*(C^C)$ is convex because $\overline{\Gamma}^*(C^C) = \overline{\{v \in \overline{\Gamma}_n^* \mid M_{\text{CI}}(C^C) \cdot v = 0\}}$ equals $F(C^C)$. (Because $F(C^C)$ is closed, any element $w \in F(C^C)$, in particular any element on its boundary, is the limit of a sequence of elements $\{w_k\}_k$ for $k \rightarrow \infty$, where the w_k lie in the interior of $F(C^C)$ for all k . Hence $w \in \overline{\Gamma}^*(C^C)$.) \square

The convexity of $\overline{\Gamma}^*(C^C)$ is crucial for the considerations of the following sections. Note that in spite of the convexity of $\overline{\Gamma}^*(C^C)$, the set $\mathcal{P}(C^C)$ is generally not convex. This alludes to the fact that significant information about the achievable correlations among the random variables is lost via the mapping from $\mathcal{P}(C^C)$ to the corresponding entropic cone $\overline{\Gamma}^*(C^C)$.

Example 4 (Entropic outer approximation for the instrumental scenario). The instrumental scenario has at most 4 independent conditional independence equalities (1). We find that there are only two, $I(A : X) = 0$ and $I(Y : X \mid AZ) = 0$. This yields $\overline{\Gamma}^*(IC^C) = \{v \in \overline{\Gamma}_4^* \mid M_{\text{CI}}(IC^C) \cdot v = 0\}$ with

$$M_{\text{CI}}(IC^C) = \begin{pmatrix} -1 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 1 \end{pmatrix},$$

where the coordinates are ordered as $(H(A), H(X), H(Y), H(Z), H(AX), H(AY), H(AZ), H(XY), H(XZ), H(YZ), H(AXY), H(AXZ), H(AYZ), H(XYZ), H(AXYZ))$. An outer approximation is given by $\Gamma(IC^C) = \{v \in \Gamma_4 \mid M_{\text{CI}}(IC^C) \cdot v = 0\}$.

In general, the outer approximation to $\overline{\Gamma}^*(C^C)$ can be further tightened by taking non-Shannon inequalities into account. These have led to the derivation of numerous new entropic inequalities for various causal structures [50] (see e.g. the triangle causal structure of Figure 1(c)). For the instrumental scenario, however, such additional inequalities are irrelevant. This can for instance be seen by constructing the following inner approximation to the cone.

Example 5 (Entropic inner approximation for the instrumental scenario [50]). For the instrumental scenario an inner approximation is given in terms of the Ingleton cone and the conditional independence constraints from the previous example, $\Gamma^{\text{I}}(IC^C) = \{v \in \Gamma^{\text{I}} \mid M_{\text{CI}}(IC^C) \cdot v = 0\}$. For this causal structure the Ingleton constraints are implied by the Shannon inequalities and the conditional independence constraints and hence, inner and outer approximation coincide. Consequently, they also coincide with the actual entropy cone, i.e., $\Gamma^{\text{I}}(IC^C) = \Gamma(IC^C) = \overline{\Gamma}^*(IC^C)$. In particular, non-Shannon entropic inequalities cannot improve the outer approximation in this example.

Inner approximations have been considered in [50]. They are particularly useful in cases where identical inner and outer approximations are found, where they identify the actual boundary of the entropy cone. In other cases they can allow parts of the actual boundary to be identified or give clues on how to find better outer approximations.

Arguably all interesting scenarios (such as the previous example) involve unobserved variables that are suspected to cause some of the correlations between the variables we observe. These unobserved variables may yield constraints on the possible joint distributions of the observed variables, a well-known example being a Bell inequality [30]⁷. More generally we would like to infer constraints on the observed variables that follow from the presence of unobserved variables.

For a causal structure on n random variables $\{X_1, X_2, \dots, X_n\}$, the restriction to the set of observed variables is called its *marginal scenario*, denoted \mathcal{M} . Here, we assume w.l.o.g. that the first $k \leq n$ variables are observed and the remaining $n - k$ are not. We are thus interested in the correlations among the first k variables that can be obtained as the marginal of some distribution over all n variables. Without any causal restrictions the set of all probability distributions of the k observed variables is $\mathcal{P}_{\mathcal{M}} := \{P \in \mathcal{P}_k \mid P = \sum_{X_{k+1}, \dots, X_n} P_{X_1 X_2 \dots X_n}\}$, i.e., $\mathcal{P}_{\mathcal{M}} = \mathcal{P}_k$. For a classical causal structure C^C on the set of variables $\{X_1, X_2, \dots, X_n\}$, marginalising all distributions $P \in \mathcal{P}(C^C)$ over the

⁷ For a detailed discussion of the significance of Bell inequality violation on classical causal structures see [31].

$n - k$ unobserved variables leads to the set $\mathcal{P}_{\mathcal{M}}(C^C) := \left\{ P \in \mathcal{P}_k \mid P = \sum_{X_{k+1}, \dots, X_n} \prod_{i=1}^n P_{X_i | X_{i+1}^n} \right\}$. In contrast to the unrestricted case, this set of distributions is in general not recovered by considering a causal structure that involves only k observed random variables, as can be seen in the following example.

Example 6 (Observed distributions in the instrumental scenario). For the instrumental scenario the observed variables are X , Y and Z and their joint distribution is of the form $\mathcal{P}_{\mathcal{M}}(IC^C) = \{P_{XYZ} \in \mathcal{P}_3 \mid P_{XYZ} = \sum_A P_{Y|AZ} P_{Z|AX} P_X P_A\}$.

The first *entropic* inequalities for a marginal scenario were derived in [12], where certificates for the existence of common ancestors of a subset of the observed random variables of at least a certain size were given. One such scenario is the triangle causal structure of Figure 1(c). The systematic entropy vector approach was devised for classical causal structures in [6, 8, 10]. An outer approximation to the entropic cones of a variety of causal structures was given in [11]. In the following we give the details of this approach.

In the entropic picture, marginalisation is performed by eliminating the coordinates that represent entropies of sets of variables containing at least one unobserved variable from the vectors. This corresponds to a projection of a cone in $\mathbb{R}^{2^n - 1}$ to its marginal cone in $\mathbb{R}^{2^k - 1}$ [9]. We will denote this projection $\pi_{\mathcal{M}} : \mathbb{R}^{2^n - 1} \rightarrow \mathbb{R}^{2^k - 1}$. It gives all entropy vectors w of the observed sets of variables, i.e., of the marginal scenario \mathcal{M} , for which there exists at least one entropy vector v in the original scenario with matching entropies on the observed variables.

Starting from the set of all entropy vectors, $\Gamma^*(C^C)$, those relevant for the marginal scenario can be obtained by discarding the appropriate components. For a finitely generated cone such as $\Gamma(C^C)$, its projection can be more efficiently determined from the projection of its extremal rays. In the dual description of the entropic cone in terms of its facets (i.e., its inequality description) the transition to the marginal scenario can be made computationally by eliminating all entropies of sets of variables not contained in \mathcal{M} from the system of inequalities. The standard algorithm that achieves this is Fourier-Motzkin elimination [53], which has been used in this context in [6, 8, 9].

Without any causal restrictions, the entropy cone $\overline{\Gamma}_n^*$ is projected to the marginal cone $\overline{\Gamma}_{\mathcal{M}}^* := \left\{ w \in \mathbb{R}_{\geq 0}^{2^k - 1} \mid \exists v \in \overline{\Gamma}_n^* \text{ s.t. } w = \pi_{\mathcal{M}}(v) \right\}$. Note that if we marginalise $\overline{\Gamma}_n^*$ over $n - k$ variables we recover the entropy cone for k random variables, i.e., $\overline{\Gamma}_{\mathcal{M}}^* = \overline{\Gamma}_k^*$. The same applies to the outer approximations: The n variable Shannon cone Γ_n is projected to the k variable Shannon cone with the mapping $\pi_{\mathcal{M}}$, i.e., $\Gamma_{\mathcal{M}} := \left\{ w \in \mathbb{R}_{\geq 0}^{2^k - 1} \mid \exists v \in \Gamma_n \text{ s.t. } w = \pi_{\mathcal{M}}(v) \right\} = \Gamma_k$. This follows because the n variable Shannon constraints contain the corresponding k variable constraints as a subset, and since any vector in Γ_k can be extended to a vector in Γ_n , for instance by taking $H(X_{k+1}) = H(X_{k+2}) = \dots = H(X_n) = 0$ and $H(X_S \cup X_T) = H(X_S)$ for any $X_T \subseteq \{X_{k+1}, X_{k+2}, \dots, X_n\}$.

For a classical causal structure C^C , we will be interested in the set $\overline{\Gamma}_{\mathcal{M}}^*(C^C) := \left\{ w \in \mathbb{R}_{\geq 0}^{2^k - 1} \mid \exists v \in \overline{\Gamma}^*(C^C) \text{ s.t. } w = \pi_{\mathcal{M}}(v) \right\}$, which is by construction a convex cone, since projection preserves convexity. The following lemma confirms that this is the entropy cone of the marginal scenario \mathcal{M} and thus also formally justifies the method of projecting the sets directly.

Lemma 10. $\overline{\Gamma}_{\mathcal{M}}^*(C^C)$ is equal to the set of entropy vectors compatible with the marginal scenario of the classical causal structure C^C , i.e., $\overline{\Gamma}_{\mathcal{M}}^*(C^C) = \overline{\left\{ w \in \mathbb{R}_{\geq 0}^{2^k - 1} \mid \exists P \in \mathcal{P}_{\mathcal{M}}(C^C) \text{ s.t. } w = \mathbf{H}(P) \right\}}$.

Proof. Let \mathcal{F} denote the set on the rhs of the statement in the lemma. Note that $w \in \overline{\Gamma}_{\mathcal{M}}^*(C^C)$ implies that there exists $v \in \overline{\Gamma}^*(C^C)$ s.t. $w = \pi_{\mathcal{M}}(v)$. Using Lemma 9, we have $v = \mathbf{H}(P)$ for some $P = \prod_i P_{X_i | X_{i+1}^n}$. If we take $P' = \sum_{X_{k+1}, \dots, X_n} P$ then $w = \mathbf{H}(P')$ and hence $w \in \mathcal{F}$.

Conversely, $w \in \mathcal{F}$ implies that there exists $P' \in \mathcal{P}_{\mathcal{M}}(C^C)$ s.t. $w = \mathbf{H}(P')$ and hence there exists $P \in \mathcal{P}(C^C)$ such that $P' = \sum_{X_{k+1}, \dots, X_n} P$. If we take $v = \mathbf{H}(P)$, then $w = \pi_{\mathcal{M}}(v)$ and hence $w \in \overline{\Gamma}_{\mathcal{M}}^*(C^C)$. Taking the topological closure of both sets concludes the proof. \square

An outer approximation to $\overline{\Gamma}_{\mathcal{M}}^*(C^C)$ is $\Gamma_{\mathcal{M}}(C^C) := \left\{ w \in \mathbb{R}_{\geq 0}^{2^k - 1} \mid \exists v \in \Gamma(C^C) \text{ s.t. } w = \pi_{\mathcal{M}}(v) \right\}$, which can be written as $\Gamma_{\mathcal{M}}(C^C) = \left\{ w \in \Gamma_{\mathcal{M}} \mid M_{\mathcal{M}}(C^C) \cdot w \geq 0 \right\}$, where $M_{\mathcal{M}}(C^C)$ is the matrix obtained via Fourier-Motzkin elimination and encodes the set of inequalities on $\mathbb{R}_{\geq 0}^{2^k - 1}$ that are implied by the fact that $M_{\text{CI}}(C^C) \cdot v = 0$ and $M_{\text{SH}}^n \cdot v \geq 0$ hold on $\mathbb{R}_{\geq 0}^{2^n - 1}$ (except for the k -variable Shannon constraints which are already included in $\Gamma_{\mathcal{M}}$).

Example 7 (Entropic outer approximation for the marginal cone of the instrumental scenario [10, 11]). For the instrumental scenario the outer approximation to its marginal cone is found by projecting $\Gamma(IC^C)$ to its three variable scenario and yields $\Gamma_{\mathcal{M}}(IC^C) = \left\{ w \in \Gamma_{\mathcal{M}} \mid M_{\mathcal{M}}(IC^C) \cdot w \geq 0 \right\}$, where $M_{\mathcal{M}}(IC^C) = \begin{pmatrix} -1 & 0 & 1 & 0 & 0 & -1 & 1 \end{pmatrix}$ corresponds to the inequality $I(X : YZ) \leq H(Z)$ from [10, 11]. As $\Gamma(IC^C) = \overline{\Gamma}^*(IC^C)$ holds, we also have $\Gamma_{\mathcal{M}}(IC^C) = \overline{\Gamma}_{\mathcal{M}}^*(IC^C)$.

As mentioned previously, non-Shannon inequalities cannot give any new entropic constraints for IC , as the Shannon approximation is already tight. However, in many causal structures they do. For instance in the triangle scenario of Figure 1(c), non-Shannon inequalities still lead to new entropic constraints, even after marginalisation to the three observed variables [50].

2. Causal structures with unobserved quantum systems

A quantum causal structure differs from its classical counterpart in that unobserved systems correspond to shared quantum states.

Definition 11. A *quantum causal structure*, C^Q , is a causal structure where each observed node has a corresponding random variable, and each unobserved node has an associated quantum system.

In a classical causal structure the edges of the DAG represent the propagation of classical information, and, at a node with incoming edges, the random variable there can be generated by applying an arbitrary function to its parents. We are hence implicitly assuming that all the information about the parents is transmitted to its children (otherwise the set of allowed functions would be restricted). This does not pose a problem since classical information can be copied. In the quantum case, on the other hand, the no-cloning theorem means that the children of a node cannot (in general) all have access to the same information as is present at that node. Furthermore, the analogue of performing arbitrary functions in the classical case is replaced by arbitrary quantum operations. Such a quantum framework that allows for an analysis with entropy vectors was introduced in [13]. In the following we outline this approach. However, for unity of description, our account of quantum causal structures is based upon the viewpoint that is taken for generalised causal structures in [11], which we review in the next section⁸.

Let C^Q be a quantum causal structure. Nodes without input edges correspond to the preparation of a quantum state described by a density operator on a Hilbert space, e.g., $\rho_A \in \mathcal{S}(\mathcal{H}_A)$ for a node A , where for observed nodes this state is required to be classical⁹. For each directed edge in the graph there is a corresponding subsystem with Hilbert space labelled by the edge's input and output nodes. For instance, if Y and Z are the only children of A then there are associated spaces \mathcal{H}_{AY} and \mathcal{H}_{AZ} such that $\mathcal{H}_A = \mathcal{H}_{AY} \otimes \mathcal{H}_{AZ}$ ¹⁰. At an unobserved node, a CPTP map from the joint state of all its input edges to the joint state of its output edges is performed. A node is labelled by its output state. For an observed node the latter is classical. Hence, it corresponds to a random variable that represents the output statistics obtained in a measurement by applying a positive operator valued measure (POVM) to the input states¹¹. If all input edges are classical this can be interpreted as a stochastic map between random variables.

A distribution, P , over the observed nodes of a causal structure C^Q is compatible with C^Q if there exists a quantum state labelling each unobserved node (with subsystems for each unobserved edge) and transformations, i.e., preparations and CPTP maps for each unobserved node as well as POVMs for each observed node, that allow for the generation of P by means of the Born rule. We denote the set of all compatible distributions $\mathcal{P}_{\mathcal{M}}(C^Q)$.

Example 8 (Compatible distributions in the quantum instrumental scenario). For the quantum instrumental scenario (Figure 1(a)), $\mathcal{P}_{\mathcal{M}}(IC^Q) = \{P_{XYZ} \in \mathcal{P}_3 \mid P_{XYZ} = \text{tr}((E_X^Z \otimes F_Z^Y)\rho_A)P_X\}$ is the set of compatible distributions. A state $\rho_A \in \mathcal{S}(\mathcal{H}_{AZ} \otimes \mathcal{H}_{AY})$ is prepared. Depending on the random variable X , a POVM $\{E_X^Z\}_Z$ on \mathcal{H}_{AZ} is applied to generate the output distribution of the observed variable Z . Depending on the latter, another POVM $\{F_Z^Y\}_Y$ is applied to generate the distribution of Y .

The set of entropy vectors of compatible probability distributions over the observed nodes, $\mathcal{P}_{\mathcal{M}}(C^Q)$, is $\Gamma_{\mathcal{M}}^*(C^Q) := \{w \in \mathbb{R}_{\geq 0}^{2^k - 1} \mid \exists P \in \mathcal{P}_{\mathcal{M}}(C^Q) \text{ s.t. } w = \mathbf{H}(P)\}$. Outer approximations $\Gamma_{\mathcal{M}}(C^Q)$ were first derived in [13], a procedure that we outline in the following. For their construction, an entropy is associated to each random variable and to each subsystem of a quantum state (equivalently each edge originating at a quantum node), corresponding to the von

⁸ The difference is as follows: In [13] nodes correspond to quantum systems. All outgoing edges of a node together define a completely positive trace preserving (CPTP) map with output states corresponding to the joint state associated with its child nodes. Similarly, the CPTP map associated to the input edges of a node must map the states of the parent nodes to the node in question. In [11], on the other hand, edges correspond to states whereas the transformations occur at the nodes.

⁹ $\mathcal{S}(\mathcal{H})$ denotes the set of all density operators on a Hilbert space \mathcal{H} .

¹⁰ Note that in the classical case these subsystems may all be taken to be copies of the system itself.

¹¹ Note that preparation and measurement can also be seen as CPTP maps with classical input and output systems respectively, thus allowing for a unified formulation.

Neumann entropy of the respective system. For convenience of exposition, edges and their associated systems share the same label. The *von Neumann entropy* of a density operator $\rho_X \in \mathcal{S}(\mathcal{H}_X)$ is defined as

$$H(X) := -\text{tr}(\rho_X \log_2(\rho_X)).$$

The quantum conditional entropy, mutual- and conditional mutual information are defined as in the classical case and with the von Neumann entropy replacing the Shannon entropy.

Because of the impossibility of cloning, the outcomes and the quantum systems that led to them do not exist simultaneously. Therefore there is in general no joint multi-party quantum state for all subsystems and it does not make sense to talk about the joint entropy of the states and outcomes. More concretely, if a system A is measured to produce Z , then ρ_{AZ} is not defined and hence neither is $H(AZ)$ ¹².

Definition 12. Two subsystems in a quantum causal structure C^Q *coexist* if neither of them is a quantum ancestor of the other. A set of subsystems that mutually coexist is termed *coexisting*.

A quantum causal structure may have several maximal coexisting subsets. Only within such subsets is there a well defined joint quantum state and joint entropy.

Example 9 (Coexisting sets in the quantum instrumental scenario). Consider the quantum version of the instrumental scenario, as illustrated in Figure 1(a). There are three observed variables as well as two edges originating at unobserved (quantum) nodes, hence 5 variables to consider. More precisely, the quantum node A has two associated subsystems A_Z and A_Y . The correlations seen at the two observed nodes Z and Y are formed by measurement on the respective subsystems A_Z and A_Y . The coexisting sets in this causal structure are $\{A_Y, A_Z, X\}$, $\{A_Y, X, Z\}$ and $\{X, Y, Z\}$ and their (non-empty) proper subsets.

Note that without loss of generality we can assume that any initial, i.e., parentless quantum states such as ρ_A above, are pure. This is because any mixed state can be purified, and if the transformations and measurement operators are then taken to act trivially on the purifying systems the same statistics are observed. In the causal structure of Example 9, this implies that ρ_A can be considered to be pure and thus $H(A_Y A_Z) = 0$. The Schmidt decomposition then implies that $H(A_Y) = H(A_Z)$. This is computationally useful as it reduces the number of free parameters in the entropic description of the scenario. Furthermore, by Stinespring's theorem [55], whenever a CPTP map is applied at a node that has at least one quantum child, then one can instead consider an isometry to a larger output system. The additional system that is required for this can be taken to be part of the unobserved quantum output (or one of them in case of several quantum output nodes). Each such case allows for the reduction of the number of variables by one, since the joint entropy of all inputs to such a node must be equal to that of all its outputs.

Quantum states are known to obey submodularity [56] and also obey the condition

- Weak monotonicity [56]: $H(X_S \setminus X_T) + H(X_T \setminus X_S) \leq H(X_S) + H(X_T)$, for all $X_S, X_T \subseteq \Omega$ (recall $H(\{\}) = 0$).

This the dual of submodularity in the sense that the two inequalities can be derived from each other by considering purifications of the corresponding quantum states [57].

Within the context of causal structures, these relations can always be applied between variables in the same coexisting set. In addition, whenever it is impossible for there to be entanglement between the subsystems $X_S \cap X_T$ and $X_S \setminus X_T$ — for instance if these subsystems are in a cq-state — the monotonicity constraint $H(X_S \setminus X_T) \leq H(X_S)$ holds. If it is also impossible for there to be entanglement between $X_S \cap X_T$ and $X_T \setminus X_S$, then the monotonicity relation $H(X_T \setminus X_S) \leq H(X_T)$ holds rendering the weak monotonicity relation stated above redundant.

Altogether, these considerations lead to a set of basic inequalities containing some Shannon and some weak-monotonicity inequalities, which are conveniently expressed in a matrix $M_B(C^Q)$. This way of approximating the entropic cone in the quantum case is inspired by work on the entropic cone for multi-party quantum states [34]. Note also that there are no further inequalities for the von Neumann entropy known to date (contrary to the classical case where a variety of non Shannon inequalities is known) [58–61].

The conditional independence constraints in C^Q cannot be identified by Proposition 8, because variables do not coexist with any quantum parents and hence conditioning a variable on a quantum parent is not meaningful. Nonetheless, among the variables in a coexisting set the conditional independences that are valid for C^C also hold in C^Q . This can be seen as follows. First, the validity of any constraints that involve only observed variables (which are always part of a coexisting set) hold by Proposition 16 below. Secondly, for unobserved systems only their classical ancestors

¹² Attempts to circumvent this have been made, see for example [54].

and none of their descendants can be part of the same coexisting set. An unobserved system is hence independent of any subset of the same coexisting set with which it shares no ancestors. Note that each of the subsystems associated with a quantum node is considered to be a parent of all of the node's children (see Figure 1 for an example).

In addition, suppose X_S and X_T are disjoint subsets of a coexisting set, Ξ , and that the unobserved system A_X is also in Ξ . Then $I(A_X : X_S | X_T) = 0$ if X_T d-separates A_X from X_S (in the full graph including quantum nodes)¹³. The same considerations can be made for sets of unobserved systems. These independence constraints may be assembled in a matrix $M_{\text{QCI}}(C^{\text{Q}})$.

Among the variables that do not coexist, some are obtained from others by means of quantum operations. These variables are thus related by data processing inequalities (DPIs) [62].

Proposition 13 (DPI). *Let $\rho_{X_S X_T} \in \mathcal{S}(\mathcal{H}_{X_S} \otimes \mathcal{H}_{X_T})$ and \mathcal{E} be a completely positive trace preserving (CPTP) map¹⁴ on $\mathcal{S}(\mathcal{H}_{X_T})$ ¹⁵ leading to a state $\rho'_{X_S X_T}$. Then $I(X_S : X_T)_{\rho'_{X_S X_T}} \leq I(X_S : X_T)_{\rho_{X_S X_T}}$.*

The data processing inequalities provide an additional set of entropic constraints, which can be expressed in terms of a matrix inequality $M_{\text{DPI}}(C^{\text{Q}}) \cdot v \geq 0$ ¹⁶. In general, there are a large number of variables for which data processing inequalities hold. It is thus beneficial to derive rules that specify which of the inequalities are needed. First, note that whenever a concatenation of two CPTP maps \mathcal{E}_1 and \mathcal{E}_2 , $\mathcal{E} = \mathcal{E}_2 \circ \mathcal{E}_1$, is applied to a state, then any DPIs for inputs and outputs of \mathcal{E} are implied by the DPIs for \mathcal{E}_1 and \mathcal{E}_2 ¹⁷. Hence, the DPIs for composed maps \mathcal{E} never have to be considered as separate constraints.

Secondly, whenever a state $\rho_{X_S X_T X_R} \in \mathcal{S}(\mathcal{H}_{X_S} \otimes \mathcal{H}_{X_T} \otimes \mathcal{H}_{X_R})$ can be decomposed as $\rho_{X_S X_T X_R} = \rho_{X_S X_T} \otimes \rho_{X_R}$ and a CPTP map \mathcal{E} transforms the state on $\mathcal{S}(\mathcal{H}_{X_S})$. Then any DPIs for $\rho_{X_S X_T X_R}$ are implied by the DPIs for $\rho_{X_S X_T}$ ¹⁸.

Furthermore, whenever a node has classical and quantum inputs, there is not only a CPTP map generating its output state but this map can be extended to a CPTP map that simultaneously retains the classical inputs, as is the content of the following lemma, which also shows that retaining a copy of the classical inputs leads to tighter entropic inequalities.

Lemma 14. *Let Y be a node with classical and quantum inputs X_C and X_Q and \mathcal{E} be a CPTP map that acts at this node, i.e., \mathcal{E} is a map from $\mathcal{S}(\mathcal{H}_{X_C} \otimes \mathcal{H}_{X_Q})$ to $\mathcal{S}(\mathcal{H}_Y)$. Then \mathcal{E} can be extended to a map $\mathcal{E}' : \mathcal{S}(\mathcal{H}_{X_C} \otimes \mathcal{H}_{X_Q}) \rightarrow \mathcal{S}(\mathcal{H}_{X_C} \otimes \mathcal{H}_Y)$ such that $\mathcal{E}' : \rho_{X_C X_Q} \mapsto \rho'_{X_C Y}$ with the property that $\rho'_{X_C Y}$ is classical on \mathcal{H}_{X_C} and $\rho'_{X_C} = \rho_{X_C}$. Furthermore, the DPIs for \mathcal{E}' imply those for \mathcal{E} .*

Proof. The first part of the lemma follows because classical information can be copied, and hence \mathcal{E}' can be decomposed into first copying X_C , and then performing \mathcal{E} ¹⁹.

Suppose $\mathcal{E} : \rho_{X_C} \otimes \rho_{X_Q} \mapsto \rho'_Y$. The second part follows because if $I(X_C X_Q X_S : X_T)_{\rho} \geq I(Y X_S : X_T)_{\rho'}$ is a valid DPI for \mathcal{E} then $I(X_C X_Q X_S : X_T)_{\rho} \geq I(X_C Y X_S : X_T)_{\rho'}$ is valid for \mathcal{E}' . The second of these implies the first by the submodularity relation $I(X_C Y X_S : X_T)_{\rho'} \geq I(Y X_S : X_T)_{\rho'}$. \square

All the above (in)equalities are necessary conditions for a vector to be an entropy vector compatible with the causal structure C^{Q} . They constrain a polyhedral cone in $\mathbb{R}_{\geq 0}^m$, where m is the total number of coexisting sets of C^{Q} ,

$$\Gamma(C^{\text{Q}}) := \{v \in \mathbb{R}_{\geq 0}^m \mid M_{\text{B}}(C^{\text{Q}}) \cdot v \geq 0, M_{\text{QCI}}(C^{\text{Q}}) \cdot v = 0, \text{ and } M_{\text{DPI}}(C^{\text{Q}}) \cdot v \geq 0\}.$$

Example 10 (Entropic constraints for the quantum instrumental scenario). For IC^{Q} , the cone $\Gamma(IC^{\text{Q}}) = \{v \in \mathbb{R}_{\geq 0}^{15} \mid M_{\text{B}}(IC^{\text{Q}}) \cdot v \geq 0, M_{\text{QCI}}(IC^{\text{Q}}) \cdot v = 0 \text{ and } M_{\text{DPI}}(IC^{\text{Q}}) \cdot v \geq 0\}$ involves the matrix $M_{\text{B}}(IC^{\text{Q}})$

¹³ This follows because any quantum states generated from the classical separating variables may be obtained by first producing random variables from the latter (for which the usual d-separation rules hold) and then using these to generate the quantum states in question (potentially after generating other variables in the network), hence retaining conditional independence.

¹⁴ Note that the map from a quantum state to the diagonal state with entries equal to the outcome probabilities of a measurement is a CPTP map and hence also obeys the DPI.

¹⁵ In general \mathcal{E} can be a map between operators on different Hilbert spaces, i.e., $\mathcal{E} : \mathcal{S}(\mathcal{H}'_{X_T}) \rightarrow \mathcal{S}(\mathcal{H}''_{X_T})$. However, as we can consider these operators to act on the same larger Hilbert space we can w.l.o.g. take \mathcal{E} to be a map on this larger space, which we call $\mathcal{S}(\mathcal{H}_{X_T})$.

¹⁶ There are also DPIs for conditional mutual information, e.g., $I(A : B | C)_{\rho'_{ABC}} \leq I(A : B | C)_{\rho_{ABC}}$ for $\rho'_{ABC} = (\mathcal{I} \otimes \mathcal{E} \otimes \mathcal{I})(\rho_{ABC})$, but these are implied by Proposition 13, so they need not be treated separately here.

¹⁷ This follows by deriving the DPIs for input and output states of \mathcal{E}_1 and \mathcal{E}_2 respectively and combining the two.

¹⁸ This follows from $I(X_S : X_T X_R) = I(X_S : X_T)$, $I(X_S X_R : X_T) = I(X_S : X_T)$ and $I(X_S X_T : X_R) = 0$.

¹⁹ Alternatively, we can think of \mathcal{E} as the concatenation of \mathcal{E}' with a partial trace; this allows us to use the same output state ρ' for both maps in the argument below.

that features 29 (independent) inequalities²⁰. In this case a single independence constraint encodes that X is independent of $A_Y A_Z$:

$$M_{\text{QCI}}(IC^{\text{Q}}) = \begin{pmatrix} 0 & 0 & -1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}.$$

Two data processing inequalities are required (cf. Lemma 14), $I(A_Z X : A_Y) \geq I(XZ : A_Y)$ and $I(A_Y Z : X) \geq I(YZ : X)$, which yield a matrix

$$M_{\text{DPI}}(IC^{\text{Q}}) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 & -1 & 1 \end{pmatrix}.$$

The above matrices are all expressed in terms of coefficients of $(H(A_Y), H(A_Z), H(X), H(Y), H(Z), H(A_Y A_Z), H(A_Y X), H(A_Y Z), H(A_Z X), H(XY), H(XZ), H(YZ), H(A_Y A_Z X), H(A_Y XZ), H(XYZ))$. Although the notation suppresses the different states there is no ambiguity because e.g. the entropy of X is the same for all states with subsystem X . The full list of inequalities is provided in the appendix.

From $\Gamma(C^{\text{Q}})$, an outer approximation to the set of compatible entropy vectors $\Gamma_{\mathcal{M}}^*(C^{\text{Q}})$ of the observed scenario \mathcal{M} of C^{Q} can be obtained using Fourier-Motzkin elimination. This leads to $\Gamma_{\mathcal{M}}(C^{\text{Q}}) := \{w \in \mathbb{R}_{\geq 0}^{2^k-1} \mid \exists v \in \Gamma(C^{\text{Q}}) \text{ s.t. } w = \pi_{\mathcal{M}}(v)\}$, which can be written as $\Gamma_{\mathcal{M}}(C^{\text{Q}}) = \{w \in \Gamma_{\mathcal{M}} \mid M_{\mathcal{M}}(C^{\text{Q}}) \cdot w \geq 0\}$. The matrix $M_{\mathcal{M}}(C^{\text{Q}})$ encodes all (in)equalities on $\mathbb{R}_{\geq 0}^{2^k-1}$ implied by $M_{\text{B}}(C^{\text{Q}}) \cdot v \geq 0$, $M_{\text{QCI}}(C^{\text{Q}}) \cdot v = 0$ and $M_{\text{DPI}}(C^{\text{Q}}) \cdot v \geq 0$ (except for the Shannon inequalities, which are already included in $\Gamma_{\mathcal{M}}$). Note that $\Gamma_{\mathcal{M}}(C^{\text{C}}) \subseteq \Gamma_{\mathcal{M}}(C^{\text{Q}}) \subseteq \Gamma_{\mathcal{M}}$, where the first relation holds because all inequalities relevant for quantum states hold in the classical case as well²¹.

Example 11 (Entropic outer approximation for the quantum instrumental scenario). The projection of $\Gamma(IC^{\text{Q}})$ leads to the entropic cone $\Gamma_{\mathcal{M}}(IC^{\text{Q}}) = \{w \in \Gamma_3 \mid M_{\mathcal{M}}(IC^{\text{Q}}) \cdot w \geq 0\}$, for which $M_{\mathcal{M}}(IC^{\text{Q}})$ equals $M_{\mathcal{M}}(IC^{\text{C}})$ from Example 7, thus corresponding to the constraint $I(X : YZ) \leq H(Z)$. Hence, $\overline{\Gamma_{\mathcal{M}}^*}(IC^{\text{Q}})$ coincides with $\overline{\Gamma_{\mathcal{M}}^*}(IC^{\text{C}})$ [50].

This method has been applied to find an outer approximation to the entropy cone of the triangle causal structure in the quantum case (cf. Figure 1(c)) [13]. This approximation did not coincide with the outer approximation to the classical triangle scenario obtained from Shannon inequalities and independence constraints. Whether there are more as yet unknown inequalities in the quantum case remains an open question (as opposed to the classical case where even better outer approximations have already been found [50]). In [13], the method was furthermore combined with the approach reviewed in Section III B, where it was applied to a scenario related to IC (cf. Example 17 below).

3. Causal structures with unobserved systems obeying only no-signalling constraints

The concept of a generalised causal structure was introduced in [11], the idea being to have one framework in which classical, quantum and even more general systems, for instance non-local boxes [63, 64], can be shared by unobserved nodes and where theory independent features of networks and corresponding bounds on our observations may be identified.

Definition 15. A *generalised causal structure* C^{G} is a causal structure which for each observed node has an associated random variable and for each unobserved node has a corresponding non-signalling resource allowed by a generalised probabilistic theory.

Classical and quantum causal structures can be viewed as special cases of generalised causal structures [11, 65]. Generalised probabilistic theories may be conveniently described in the operational-probabilistic framework of [66]. Circuit elements correspond to so-called tests that are connected by wires, which represent propagating systems. In general, such a test has an input system, and two outputs: an output system and an outcome. In case of a trivial

²⁰ Note that the only weak monotonicity relations that are not made redundant by other basic inequalities are $H(A_Y | A_Z X) + H(A_Y) \geq 0$, $H(A_Z | A_Y X) + H(A_Z) \geq 0$, $H(A_Y | A_Z) + H(A_Y | X) \geq 0$ and $H(A_Z | A_Y) + H(A_Z | X) \geq 0$.

²¹ This can be seen by thinking of a classical source as made up of two or more (perfectly correlated) random variables as its subsystems, which are sent to its children and processed there. The Shannon inequalities hold among all of these variables (and also imply any weak monotonicity constraints). The classical independence relations include the quantum ones but may add constraints that involve conditioning on any of the variables' ancestors. These (in)equalities are tighter than the DPIs, which are hence not explicitly considered in the classical case.

input system we are talking about a preparation-test and in case of a trivial output system of an observation-test. In the causal structure framework, a test is associated to each node. However, each such test has only one output: for unobserved nodes this is a general resource state; for observed nodes it is a random variable. Furthermore, resource states do not allow for signalling from the future to the past, i.e., we are considering so-called causal operational-probabilistic theories. This is important for the interpretation of generalised causal structures.

A distribution P over the observed nodes of a generalised causal structure C^G is compatible with C^G if there exists a causal operational-probabilistic theory, a resource for each unobserved edge in that theory and transformations for each node that allow for the generation of P . We denote the set of all compatible distributions $\mathcal{P}_{\mathcal{M}}(C^G)$. As in the quantum case, there is no notion of a joint state of all nodes in the causal structure and of conditioning on an unobserved system. Even more, there is no consensus on the representation of states and their dynamics in general non-signalling theories. To circumvent this, the classical notion of d-separation has been reformulated [11], which enables the following proposition.

Proposition 16 (Henson, Lal & Pusey). *Let C^G be a generalised causal structure and let X , Y and Z be pairwise disjoint subsets of observed nodes in C^G . If a probability distribution P is compatible with C^G , then the d-separation of X and Y by Z implies the conditional independence $X \perp\!\!\!\perp Y | Z$. Conversely, if for every distribution P compatible with C^G the conditional independence $X \perp\!\!\!\perp Y | Z$ holds, then X is d-separated from Y by Z in C^G .*

This allows for the derivation of conditional independence relations among observed variables that hold in any generalised probabilistic theory, which hence restrict a general entropic cone. Furthermore, it rigorously justifies retaining the independence constraints among the (observed) variables in coexisting sets in quantum causal structures (cf. Section II B 2), which can be seen as special cases of generalised causal structures.

In [11], sufficient conditions for identifying causal structures C for which in the classical case, C^C , there are no restrictions on the distribution over observed variables other than those that follow from the d-separation of these variables were derived. Since, by Proposition 16, these conditions also hold in C^Q and C^G , this implies $\mathcal{P}_{\mathcal{M}}(C^C) = \mathcal{P}_{\mathcal{M}}(C^Q) = \mathcal{P}_{\mathcal{M}}(C^G)$. For causal structures with up to six nodes, there are 21 cases (and some that can be reduced to these 21) where such equivalence does not hold and where further relations among the observed variables have to be taken into account [11, 18].

Outer approximations to the entropic cones for causal structures, C^G , based on the observed variables and their independences only were derived in [11]. Moreover, a few additional constraints for certain generalised causal structures were derived there. For example, the entropic constraint $I(X : Y) + I(X : Z) \leq H(X)$ for the triangle causal structure of Figure 1(c) (which had previously been established in the classical case [67]) was found. This constraint does not follow from the observed independences, but nonetheless holds for the triangle causal structure in generalised probabilistic theories.

In spite of this, a systematic entropic procedure, in which the unobserved variables are explicitly modelled and then eliminated from the description, is not available for generalised causal structures. The issue is that we are lacking a generalisation of the Shannon and von Neumann entropy to generalised probabilistic theories that obeys submodularity and for which the conditional entropy can be written as the difference of unconditional entropies [68, 69].

One possible generalised entropy is the *measurement entropy*, which is positive and obeys some of the submodularity constraints (those with $X_S \cap X_T = \{\}$) but not all [68, 69]. Using this, Ref. [70] considered the set of possible entropy vectors for a bipartite state in *box world*, a generalised probabilistic theory that permits all bipartite correlations that are non-signalling [71]. They found no further constraints on the set of possible entropy vectors in this setting (hence, contrary to the quantum case, measurement entropy vectors in box world can violate monotonicity). Other generalised probabilistic theories and multi-party states have, to our knowledge, not been similarly analysed.

4. Other directions for exploring quantum and generalised causal structures

The approaches to quantum and generalised causal structures above are based on adaptations of the theory of Bayesian networks to the respective settings and on retaining the features that remain valid, for instance the relation between d-separation and independence for observed variables [11] (cf. Section II B 3). Other approaches to generalise classical networks to the quantum realm have been pursued [54], where a definition of conditional quantum states, analogous to conditional probability distributions was formulated.

Recent articles have proposed generalisations of Reichenbach's principle [72] to the quantum realm [16, 73, 74]. In Ref. [16] a graph separation rule, q-separation, was introduced, whereas [73, 74] rely on a formulation of quantum networks in terms of quantum channels and their Choi states.

An active area of research is the exploration of frameworks that allow for indefinite causal structures [75–77]. There are several approaches achieving this, such as the process matrix formalism [78], which has led to the derivation of so called causal inequalities and the identification of signalling correlations that are achievable in this framework,

however not with any predefined causal structure [78, 79]. Another framework that is able to describe such scenarios is the theory of quantum combs [80], illustrated by a quantum switch, a quantum bit controlling the circuit structure in a quantum computation. A recent framework with the aim to model cryptographic protocols is also available [81]. Some initial results on the analysis of indefinite causal structures with entropy have recently appeared [82].

In the classical, quantum and generalised causal structures considered above only the observed classical information can be transmitted via a link between two observed variables and, in particular, no additional unobserved system. This understanding of the causal links encodes a Markov condition. In other situations, it can be convenient for the links in the graph to represent a notion of future instead of direct causation, see e.g. [83, 84].

III. ENTROPY VECTOR APPROACH WITH POST-SELECTION

A technique that leads to additional, more fine-grained inequalities is based on post-selecting on the values of parentless classical variables. This technique was pioneered by Braunstein and Caves [85] and has been used to systematically derive numerous entropic inequalities [6–8, 17, 18].

A. Post-selection in classical causal structures

In the following we denote a random variable X post-selected on the event of another random variable, Y , taking a particular value, $Y = y$, as $X_{|Y=y}$. The same notation is used for a set of random variables $S = \{X_1, X_2, \dots, X_n\}$, whose joint distribution is conditioned on $Y = y$, $S_{|Y=y} = \{X_{1|Y=y}, X_{2|Y=y}, \dots, X_{n|Y=y}\}$. The following lemma can be understood as a generalisation of (a part of) Fine's theorem [86, 87].

Lemma 17. *Let C^C be a classical causal structure with a parentless observed node X that takes values $X = 1, 2, \dots, n$ and let P be a joint distribution over all random variables $X \cup X^\dagger \cup X^\ddagger$ in C^C (with P compatible with C^C). Then there exists a joint distribution Q over the $n \cdot |X^\dagger| + |X^\ddagger|$ random variables $\Omega_{|X} := X_{|X=1}^\dagger \cup X_{|X=2}^\dagger \cup \dots \cup X_{|X=n}^\dagger \cup X^\ddagger$ such that $Q(X_{|X=x}^\dagger X^\ddagger) = P(X^\dagger X^\ddagger | X = x)$ for all $x \in \{1, \dots, n\}$.*

Proof. The joint distribution over the random variables $X^\dagger \cup X^\ddagger$ in C^C can be written as $P(X^\dagger X^\ddagger) = \sum_{x=1}^n P(X^\dagger | X^\ddagger X = x) P(X^\ddagger) P(X = x)$. Now take $Q(X_{|X=1}^\dagger \dots X_{|X=n}^\dagger X^\ddagger) = \prod_{x=1}^n P(X^\dagger | X^\ddagger X = x) P(X^\ddagger)$. As required, this distribution has marginals $Q(X_{|X=x}^\dagger X^\ddagger) = P(X^\dagger | X^\ddagger X = x) P(X^\ddagger)$. \square

It is perhaps easiest to think about this lemma in terms of a new causal structure C_X^C on $\Omega_{|X}$ that is related to the original. Roughly speaking the new causal structure is formed by removing X and replacing the descendants of X with several copies each of which have the same causal relations as in the original causal structure (with no mixing between copies). More precisely, if X is a parentless node in C^C we can form a *post-selected causal structure* on $\Omega_{|X}$ (post-selecting on X) as follows: **(1)** For each pair of nodes $A, B \in X^\ddagger$ in C^C , make A a parent of B in C_X^C iff A is a parent of B in C^C . **(2)** For each node $B \in X^\ddagger$ in C^C and for each node $A_{|X=x}$, make B a parent of $A_{|X=x}$ in C_X^C iff B is a parent of A in C^C . **(3)** For each pair of nodes, $A_{|X=x}$ and $B_{|X=x}$, make $B_{|X=x}$ a parent of $A_{|X=x}$ in C_X^C iff B is a parent of A in C^C . (Note that there is no mixing between different values of $X = x$.) See Figures 3 and 5 and Example 12 for illustrations. This view gives us the following corollary of Lemma 17, which is an alternative generalisation of Fine's theorem

Lemma 18. *Let C^C be a classical causal structure with a parentless observed node X that takes values $X = 1, 2, \dots, n$ and let P be a joint distribution over all random variables $X \cup X^\dagger \cup X^\ddagger$ in C^C (with P compatible with C^C). Then there exists a joint distribution Q compatible with the post-selected causal structure C_X^C such that $Q(X_{|X=x}^\dagger X^\ddagger) = P(X^\dagger X^\ddagger | X = x)$ for all $x \in \{1, \dots, n\}$.*

The distributions that are of interest in this new causal structure are the marginals $Q(X_{|X=x}^\dagger X^\ddagger)$ for all x (and their interrelations), as they correspond to distributions in the original scenario. Any constraints on these distributions derived in the post-selected scenario are by construction valid for the (post-selected) distributions compatible with the original causal structure.

Example 12 (Post-selection in the instrumental scenario). Consider the causal structure IC where the parentless variable X takes values 0 or 1. For any P compatible with IC^C , there exists a distribution Q compatible with the post-selected causal structure (Figure 3(a)) such that $Q(Z_{|X=0} Y_{|X=0} A) = P(ZY | AX = 0)P(A)$ and $Q(Z_{|X=1} Y_{|X=1} A) = P(ZY | AX = 1)P(A)$. These marginals and their relations are of interest for the original scenario.




Figure 3: (a) Pearl's instrumental scenario post-selected on binary X . The causal structure is obtained from the IC by removing X and replacing Y and Z with copies, each of which has the same causal relations as in the original causal structure. (b) post-selected Bell scenario with binary inputs A and B .

Note that the above reasoning may be applied recursively. Indeed, the causal structure with variables $\Omega_{|X}$ may be post-selected on the values of one of its parentless nodes. The joint distributions of the nodes $\Omega_{|X}$ and the associated causal structure may be analysed in terms of entropies, as illustrated with the following example.

Example 13 (Entropic constraints for the post-selected Bell scenario [85]). In the Bell scenario with binary inputs A and B (Figure 1(b)), Lemma 18 may be applied first to post-select on the values of A and then of B . This leads to a distribution Q compatible with the post-selected causal structure (on A and B) shown in Figure 3(b), for which $Q(X_{|A=a}Y_{|B=b}) = P(XY | A = a, B = b)$ for $a, b \in \{0, 1\}$ ²². Applying the entropy vector method to the post-selected causal structure and marginalising to vectors of form $(H(X_{|A=0}), H(X_{|A=1}), H(Y_{|B=0}), H(Y_{|B=1}), H(X_{|A=0}Y_{|B=0}), H(X_{|A=0}Y_{|B=1}), H(X_{|A=1}Y_{|B=0}), H(X_{|A=1}Y_{|B=1}))$ yields the inequality $H(Y_1|X_1) + H(X_1|Y_0) + H(X_0|Y_1) - H(X_0|Y_0) \geq 0$ and its permutations [6, 85].²³

The extension of Fine's theorem to more general Bell scenarios [88, 89], i.e., to scenarios involving a number of spacelike separated parties that each choose input values and produce some output random variable (and scenarios that can be reduced to the latter), has been combined with the entropy vector method in [6, 8].

Entropic constraints that are derived in this way provide novel and non-trivial entropic inequalities for the distributions compatible with the original classical causal structure. Ref. [8] introduced this idea and analysed the so-called n -cycle scenario, which is of particular interest in the context of non-contextuality and includes the Bell scenario (with binary inputs and outputs) as a special case²⁴.

In Ref. [6], new entropic inequalities for the bilocality scenario, which is relevant for entanglement swapping [91, 92], as well as quantum violations of the classical constraints on the 4- and 5-cycle scenarios were derived. For the n -cycle scenario, the (polynomial number of) entropic inequalities are sufficient for the detection of any non-local distribution [7]²⁵. In the following we illustrate the method of [6, 8] with a continuation of Example 12.

Example 14 (Entropic approximation for the post-selected instrumental scenario). The entropy vector method from Section II is applied to the 5-variable causal structure of Figure 3(a). The marginalisation is performed to retain all marginals that correspond to distributions in the original causal structure (Figure 1(a)), i.e., any marginals of $P(YZ | X = 0)$ and $P(YZ | X = 1)$. Hence, the 5 variable entropic cone is projected to a cone that restricts vectors of the form $(H(Y_{|X=0}), H(Y_{|X=1}), H(Z_{|X=0}), H(Z_{|X=1}), H(Y_{|X=0}Z_{|X=0}), H(Y_{|X=1}Z_{|X=1}))$. Note that entropies of unobserved marginals such as $H(Y_{|X=0}Z_{|X=1})$ are not included. With this technique, the Shannon constraints for the three components $(H(Y_{|X=0}), H(Z_{|X=0}), H(Y_{|X=0}Z_{|X=0}))$ are recovered (the same holds for $X = 1$); no additional constraints arise here.

It is interesting to compare this to the Bell scenario considered in Example 13. In both causal structures any 4-variable distributions, $P_{Z_{|X=0}Z_{|X=1}Y_{|X=0}Y_{|X=1}}$ and $P_{X_{|A=0}X_{|A=1}Y_{|B=0}Y_{|B=1}}$ respectively, are achievable²⁶. However, the marginal entropy vector in the Bell scenario has more components, leading to additional constraints on the observed variables [6, 85].

In some cases two different causal structures, C_1 and C_2 , can yield the same set of distributions after marginalising, a fact that has been further explored in [93]. When this occurs, either causal structure can be imposed when identifying the set of achievable marginal distributions in either scenario. If the constraints implied by the causal structure C_1

²² In this case the joint distribution is already known to exist by Fine's theorem [86, 87].

²³ Whenever the input nodes take more than two values, the latter may be partitioned into two sets, guaranteeing applicability of these inequalities. Furthermore, [7] showed that these inequalities are sufficient for detecting any behaviour that is not classically reproducible in the Bell scenario where the two parties perform measurements with binary outputs.

²⁴ A full probabilistic characterisation of the n -cycle scenario was given in [90].

²⁵ This is also true of the exponential number of inequalities in the probabilistic case [90].

²⁶ The additional causal links in Figure 3(b) do not affect the set of compatible distributions.




Figure 4: Variations of the instrumental scenario (a), (b) and (c). The causal structure (c) is relevant for the derivation of the information causality inequality where S takes n possible values. (d) and (e) are the causal structures that are effectively analysed when post-selecting on a ternary S in (a) and on a binary S in (c) respectively.

are a subset of those implied by C_2 , then those of C_2 can be used to compute improved outer approximations on the entropic cone for C_1 . Furthermore, valid independence constraints may speed up computations even if they do not lead to any new relations for the observed variables²⁷. Similar considerations also yield a criterion for indistinguishability of causal structures in certain marginal scenarios — if C_1 and C_2 yield the same set of distributions after marginalising then they cannot be distinguished in that marginal scenario.

In examples like the above, where no new constraints follow from post-selection, it may be possible to introduce additional input variables to the causal structure, which may be beneficial for instance if the aim is to certify the presence of quantum nodes in a network. The new parentless nodes may then be used to apply Lemma 17 and the above entropic techniques. Mathematically, introducing further nodes to a causal structure is always possible. However, this is only interesting if experimentally feasible, e.g. if an experimenter has control over certain observed nodes and is able to devise an experiment where he can change their inputs. In the instrumental scenario this may be of interest.

Example 15 (Variations of the instrumental scenario). In this scenario (Figure 1(a)), a measurement on system A_Z is performed depending on X (where in the classical case A_Z can w.l.o.g. be taken to be a copy of the unobserved random variable A). Its outcome Z (in the classical case a function of A) is used to choose another measurement to be performed on A_Y to generate Y (classically another a copy of A). It may often be straightforward for an experimenter to choose between several measurements. In the causal structure this corresponds to introducing an additional observed input S to the second measurement (with the values of S corresponding to different measurements on A_Y). Such an adaptation is displayed in Figure 4(a).²⁸

Alternatively, it may be possible that the first measurement (on A_Z) is chosen depending on a combination of different, independent factors, which each correspond to a random variable X_i . For two variables X_1 and X_2 the corresponding causal structure is displayed in Figure 4(b)²⁹.

Taken together, these two adaptations yield the causal structure of Figure 4(c), relevant in the context of the principle of information causality [94] (see also Example 17 below).

A second approach that relies on very similar ideas (also justified by Lemma 17) is taken in [18]. For a causal structure C^C with nodes $\Omega = X \cup X^\dagger \cup X^\ddagger$, where X is a parentless node, conditioning the joint distribution over all nodes on a particular $X = x$ retains the independences of C^C . In particular, the conditioning does not affect the distribution of the X^\ddagger , i.e., $P(X^\ddagger | X = x) = P(X^\ddagger)$ for all x . The corresponding entropic constraints can be used to derive entropic inequalities without the detour over computing large entropic cones, which may be useful where the latter computations are infeasible. The constraints that are used in [18] are, however, a (diligently but somewhat arbitrarily chosen) subset of the constraints that would go into the entropic technique detailed earlier in this section for the full causal structure. Indeed, when the computations are feasible, applying the full entropy vector method

²⁷ Note that some care has to be taken when identifying valid constraints for scenarios with causal structure [93].

²⁸ Note that for ternary S the outer approximation of the post-selected causal structure of Figure 4(d) with Shannon inequalities does not lead to any interesting constraints (as opposed to the structure of Figure 4(e), which is analysed further in Example 17).

²⁹ This is an example of a causal structure where non-Shannon inequalities among classical variables lead to a strictly tighter outer approximation in the classical and quantum case than the approximations derived using only Shannon and weak-monotonicity constraints (also if there is a causal link from X_1 to X_2) [50].




Figure 5: Causal structures from Example 16. Post-selecting on a binary observed variable C leads to the causal structure (d) in the case of structure (a), whereas both (b) and (c) lead to structure (e). In particular, this shows that the conditional techniques may yield the same results for different causal structures.

to the corresponding post-selected causal structure gives a systematic way to derive constraints, which are in general strictly tighter (cf. Example 16).

So far, the restricted technique has been used in [18] to derive the entropic inequality

$$I(X_{|C=0} : Z) - I(Y_{|C=0} : Z) - I(X_{|C=1} : Z) + I(Y_{|C=1} : Z) \leq H(Z), \quad (2)$$

which is valid for all the classical causal structures of Figure 5 (previously considered in [11]). The inequality was used to certify the existence of classical distributions that respect the conditional independence constraints among the observed variables but that are not achievable in the respective causal structures³⁰. In the following we look at these three causal structures in more detail and illustrate the relation between the two techniques.

Example 16. Applying the post-selection technique for a binary random variable C to the causal structure from Figure 5(a) yields the effective causal structure 5(d). The latter can be analysed with the above entropy vector method, which leads to a cone that is characterised by 14 extremal rays or equivalently in terms of 22 inequalities, both available in the appendix. The inequalities $I(Z : X_{|C=1}) \geq 0$, $I(Z : Y_{|C=0}) \geq 0$, $I(X_{|C=1} : Y_{|C=1} | Z) \geq 0$ and $H(Z | X_{|C=0}) \geq I(X_{|C=1} : Z : Y_{|C=1})$, which are part of this description, imply (2) above. We are not aware of any quantum violations of these inequalities.

Structures (b) and (c) both lead to the causal structure (e) upon post-selecting on a binary C . The latter causal structure turns out to be computationally harder to analyse with the entropy vector method and we have not been able to compute its entropic approximation with the above approach³¹. Hence, the method outlined in [18] is a useful alternative here.

B. Post-selection in quantum and general non-signalling causal structures

In causal structures with quantum and more general non-signalling nodes, Lemma 17 is not valid. For instance, Bell's theorem can be recast as the statement that there are distributions compatible with the quantum Bell scenario for which there is no joint distribution of $X_{|A=0}$, $X_{|A=1}$, $Y_{|B=0}$ and $Y_{|B=1}$ in the post-selected causal structure (on A and B) that has the required marginals (in the sense of Lemma 18).

Nonetheless, the post-selection technique has been generalised to such scenarios [13, 17], i.e., it is still possible to post-select on parentless (and therefore classical) nodes to take specific values. In such scenarios the observed variables can be thought of as obtained from the unobserved resources by means of measurements or tests. If a descendant of the variable that is post-selected on has quantum or general non-signalling nodes as parents, then the different instances of the latter node and of all its descendants do not coexist (even if they are observed, hence classical). This is because such observed variables are generated by measuring a quantum or other non-signalling system. Such a system is altered (or destroyed) in a measurement, hence does not allow for the simultaneous generation of different instances of its children due to the impossibility of cloning.

In the quantum case, this is reflected in the identification of the coexisting sets in the post-selected causal structure³², as is illustrated with the following example.

³⁰ These causal structures may thus also allow for quantum correlations that are not classically achievable.

³¹ We were working with conventional variable elimination software on a desktop computer

³² Different instances of a variable after post-selection have to be seen as alternatives and not as simultaneous descendants of their parent node as the representation of the post-selected causal structure might suggest.

Example 17 (Information causality scenario in the quantum case [13]). The communication scenario used to derive the principle of information causality [94] is based on the variation of the instrumental scenario displayed in Figure 4(c). It has been analysed with the entropy vector method in Ref. [13], an analysis that is presented in the following.

Conditioning on values of the variable S is possible in the classical and quantum cases. However, whereas in the classical case the variables $Y_{|S=s}$ for different S share a joint distribution (cf. Lemma 17), they do not coexist in the quantum case. For binary S , the coexisting sets are $\{X_1, X_2, A_Z, A_Y\}$, $\{X_1, X_2, Z, A_Y\}$, $\{X_1, X_2, Z, Y_{|S=1}\}$ and $\{X_1, X_2, Z, Y_{|S=2}\}$. The only independence constraints in the quantum case are that X_1, X_2 and $A_Y A_Z$ are mutually independent. Marginalising until only entropies of $\{X_1, Y_{|S=1}\}$, $\{X_2, Y_{|S=2}\}$, $\{Z\}$ and their subsets remain, yields only one non-trivial inequality, $\sum_{s=1}^n I(X_s : Y_{|S=s}) \leq H(Z)$, with $n = 2$.³³ The same inequality was previously derived by Pawłowski et al. for general n [94], where the choice of marginals was inspired by the communication task considered. Subsequently, Ref. [13] considered another marginal scenario — the one with with coexisting sets $\{X_1, X_2, Z, Y_{|S=1}\}$, $\{X_1, X_2, Z, Y_{|S=2}\}$ and all of their subsets — which led to additional inequalities.

Similar considerations were applied to causal structures allowing for general non-signalling resources, C^G in [17]. Let $O = X_O^\uparrow \cup X_O^\dagger \cup X$ be the disjoint union of its observed nodes, where X_O^\uparrow are the observed descendants and X_O^\dagger the observed non-descendants of X . If the variable X takes values $x \in \{1, 2, \dots, n\}$, this leads to a joint distribution of $X_O^\uparrow \cup X_O^\dagger$ for each $X = x$, i.e., there is a joint distribution for $P(X_O^\uparrow X_O^\dagger | X = x) = P(X_O^\uparrow | X_O^\dagger X = x)P(X_O^\dagger)$ for all x , denoted $P(X_{O|X=x}^\uparrow X_O^\dagger)$. Because X does not affect the distribution of the independent variables X_O^\dagger , the distributions $P(X_{O|X=x}^\uparrow X_O^\dagger)$ have coinciding marginals on X_O^\dagger , i.e., $P(X_O^\dagger) = \sum_s P(X_{O|X=x}^\uparrow = s, X_O^\dagger)$ for all x , where s runs over the alphabet of X_O^\uparrow . This encodes no-signalling constraints.³⁴

In terms of entropy, there are n entropic cones, one for each $P(X_{O|X=x}^\uparrow X_O^\dagger)$ (which each encode the independences among the observed variables). According to the above, they are required to coincide on the entropies for X_O^\dagger and on all of its subsets. These constraints define a convex polyhedral cone that is an outer approximation to the set of all entropy vectors achievable in the causal structure. Whenever the distributions $P(X_{O|X=x}^\uparrow X_O^\dagger)$ involve fewer than three variables and assuming that all constraints implied by the causal structure and no-signalling have been taken into account³⁵, this approximation is tight because $\overline{\Gamma}_3^* = \Gamma_3$.

Several examples of the use of this technique can be found in Ref. [17], including the original information causality scenario (which we discuss in Example 18) and an entropic analogue of monogamy relations for Bell inequality violations [95, 96].

Example 18 (Information causality scenario in general non-signalling theories). This is related to Example 17 above and reproduces an analysis from [17]. In this marginal scenario we consider the Shannon cones for the three sets $\{X_1, Y_{|S=1}\}$, $\{X_2, Y_{|S=2}\}$ and $\{Z\}$ as well as the constraints $I(X_1 : Y_{|S=1}) \leq H(Z)$ and $I(X_2 : Y_{|S=2}) \leq H(Z)$ which are conjectured to hold [17]. (This conjecture is based on an argument in [97] that covers a special case; we are not aware of a general proof.)

These conditions constrain a polyhedral cone of vectors $(H(X_1), H(X_2), H(Z), H(Y_{|S=1}), H(Y_{|S=2}), H(X_1 Y_{|S=1}), H(X_2 Y_{|S=2}))$, with 8 extremal rays that are all achievable using PR-boxes [63, 64]. Importantly, the stronger constraint $I(X_1 : Y_{|S=1}) + I(X_2 : Y_{|S=2}) \leq H(Z)$, which holds in the quantum case, (cf. Example 17) does not hold here.

IV. ALTERNATIVE TECHNIQUES

Instead of relaxing the problem of characterising the set of probability distributions compatible with a causal structure by considering entropy vectors, other computational techniques are currently being developed. In the following, we give a brief overview of such methods.

³³ Note that this is slightly adapted from [13] where they found $I(X_1 : Y_{|S=1}) + I(X_2 : Y_{|S=2}) \leq H(Z) + I(X_1 : X_2)$, as X_1 and X_2 were not assumed independent there. Furthermore, this is also the only inequality found in the classical case when restricting to this same marginal scenario [17].

³⁴ Note that these are not the only constraints that arise from no-signalling. For instance Example 18 below requires further constraints for each $P(X_{O|X=x}^\uparrow X_O^\dagger)$ that also follow from requiring non-signalling resources. The latter have to be found and added to the description separately.

³⁵ Note that it may not always be obvious how to identify all relevant constraints (cf. the conjectured constraints in Example 18).

In this context, note also that there are methods that allow certification that the only restrictions implied by a causal structure are the conditional independence constraints among the observed variables [11], as well as procedures to show that the opposite is the case [98, 99]. Such methods may (when applicable) indicate whether a causal structure should be analysed further (corresponding techniques are reviewed in [18]).

A. Entropy cones relying on Rényi entropies

Entropy vectors may be computed in terms of other entropy measures, for instance in terms of the α -Rényi entropies [100]. For a quantum state ρ_X , the α -Rényi entropy is $H_\alpha(X) := \frac{1}{1-\alpha} \log \text{tr} \rho_X^\alpha$, for $\alpha \in (0, \infty) \setminus \{1\}$, the cases $\alpha = 0, 1, \infty$ are defined via the relevant limits (note that $H_1(X) = H(X)$).³⁶

One may expect that useful constraints on the compatible distributions can be derived from such entropy vectors. For $0 < \alpha < 1$ and $\alpha > 1$ such constraints were analysed in [101]. In the classical case positivity and monotonicity are the only linear constraints on the corresponding entropy vectors for any $\alpha \neq 0, 1$. For multi-party quantum states monotonicity does not hold for any α , like in the case of the von Neumann entropy. For $0 < \alpha < 1$, there are no constraints on the allowed entropy vectors except for positivity, whereas for $\alpha > 1$ there are constraints, but these are non-linear. The lack of further linear inequalities that generally hold limits the usefulness of entropy vectors using α -Rényi entropies for analysing causal structures. To our knowledge it is not known how or whether non-linear inequalities for Rényi entropies may be employed for this task. The case $\alpha = 0$, where $H_0(X) = \log \text{rank} \rho_X$, has been considered separately in [70], where it was shown that further linear inequalities hold. However, only bi-partitions of the parties were considered and the generalisation to full entropy vectors is still to be explored.

The above considerations do not mention conditional entropy and hence could be taken with the definition $H_\alpha(X|Y) := H_\alpha(XY) - H_\alpha(Y)$. Alternatively, one may consider a definition of the Rényi conditional entropy, for which $H_\alpha(X|YZ) \leq H_\alpha(X|Y)$ [102–106]. With the latter definition, the conditional Rényi entropy cannot be expressed as a difference of unconditional entropies, and so to use entropy vectors we would need to consider the conditional entropies as separate components. Along these lines, one may also think about combining Rényi entropies for different values of α and to use appropriate chain rules [107]. Because of the large increase in the number of variables compared to the number of constraints it is not clear that this will yield useful new conditions.

B. Polynomial inequalities restricting the set of compatible distributions

The probabilistic characterisation of causal structures, depends (in general) on the dimensionality of the observed variables. Computational hardness results suggest that a full characterisation is unlikely to be feasible, except in small cases [108, 109]. Recent progress has been made with the development of procedures to construct polynomial Bell inequalities. A method that resorts to linear programming techniques [15] has led to the derivation of new inequalities for the bilocality scenario (as well as a related four-party scenario). Another, iterative procedure allows for enlarging networks by adding a party to a network in a particular way³⁷. This allows for the constructions of non-linear inequalities for the latter, enlarged network from inequalities that are valid for the former [14].

Furthermore, a recent approach relies on considering enlarged networks, so called inflations, and inferring causal constraints from those [19, 110]. Inflated networks may contain several copies of a variable that each have the same dependencies on ancestors (the latter may also exist in several instances) and which share the same distributions with their originals. Such inflations allow for the derivation of probabilistic inequalities that restrict the set of compatible distributions. These ideas also bear some resemblance to the procedures in [20], in the sense that they employ the idea that certain marginal distributions may be obtained from different networks; they are, however, much more focused on causal structures featuring interesting independence constraints. Inflations allowed the authors of [19] to refute certain distributions as incompatible with the triangle causal structure from Figure 1(c), in particular the so called W-distribution which could neither be proven to be incompatible entropically nor with the covariance matrix approach below.

³⁶ Classical α -Rényi entropies are included in this definition when considering diagonal states.

³⁷ Here, adding a party means adding one observed input and one observed output node as well as an unobserved parent for the output, the latter may causally influence one other output random variable in the network.

C. Semidefinite tests relying on covariance matrices

One may look for mappings of the distribution of a set of observed variables that encode causal structure beyond considering entropies. For causal structures with two generations, i.e., one generation of unobserved variables as ancestors of one generation of observed nodes, a technique has been found using covariance matrices [20]. Each observed variable is mapped to a vector-valued random variable and the covariance matrix of the direct sum of these variables is considered. Due to the law of total expectation, this matrix allows for a certain decomposition depending on the causal structure. For a particular observed distribution and its covariance matrix, the existence of such a decomposition may be tested via semidefinite programming. The relation of this technique to the entropy vector method is not yet well understood. A partial analysis considering several examples is given in Section X of [20].

V. OPEN PROBLEMS

The entropy vector approach has led to many certificates for the incompatibility of correlations with causal structures. However, we are generally still lacking a measure of how well entropic relations can approximate the set of achievable correlations. Firstly, the non-injective mapping from probabilities to entropies is not sufficiently understood and secondly, the current methods employ further approximations, e.g. by restricting the number of non-Shannon inequalities that can be considered at a time. It is as yet unknown whether the entropy vector method (without post-selection) can ever distinguish correlations that arise from classical, quantum and more general non-signalling resources. Such insights may also inform the question of whether there exist novel inequalities for the von Neumann entropy of multi-party quantum states.

The post-selection technique allows for the derivation of additional constraints that may distinguish quantum from classically achievable correlations in the Bell scenario and possibly in other examples. However, the method relies on the causal structure featuring parentless observed nodes, hence it is not always applicable (see e.g. the triangle scenario). In such situations, one may try to combine the entropic techniques reviewed here with the inflation method [19], which might allow for further entropic analysis of several causal structures, e.g., of the triangle scenario.

Criteria to certify whether a set of entropic constraints is able to detect non-classical correlations are currently not available. For many of the established entropic constraints on classical causal structures it is unknown whether or not they are also valid for the corresponding quantum structure. In the case of the Bell scenario this problem has been overcome. It has been shown that the known entropic constraints are even sufficient for detecting any non-classical correlations [7]. However, since the proof is specific to the scenario, finding a systematic tool to analyse the scope of the entropic techniques remains open.

Acknowledgments

We thank Rafael Chaves and Costantino Budroni for confirming details of [17]. RC is supported by the EPSRC's Quantum Communications Hub (grant no. EP/M013472/1) and by an EPSRC First Grant (grant no. EP/P016588/1).

-
- [1] Pearl, J. *Causality: Models, Reasoning, and Inference* (Cambridge University Press, 2009), 2nd edn.
 - [2] Spirtes, P., Glymour, C. N. & Scheines, R. *Causation, Prediction, and Search* (MIT Press, 2000).
 - [3] Geiger, D. & Meek, C. Quantifier elimination for statistical problems. In *Proceedings of the Fifteenth conference on Uncertainty in artificial intelligence*, 226–235 (Morgan Kaufmann Publishers Inc., 1999).
 - [4] Garcia, L. D., Stillman, M. & Sturmfels, B. Algebraic geometry of Bayesian networks. *Journal of Symbolic Computation* **39**, 331–355 (2005).
 - [5] Lee, C. M. & Spekkens, R. W. Causal inference via algebraic geometry: necessary and sufficient conditions for the feasibility of discrete causal models. e-print [arXiv:1506.03880](https://arxiv.org/abs/1506.03880) (2015).
 - [6] Chaves, R. & Fritz, T. Entropic approach to local realism and noncontextuality. *Physical Review A* **85**, 032113 (2012).
 - [7] Chaves, R. Entropic inequalities as a necessary and sufficient condition to noncontextuality and locality. *Physical Review A* **87**, 022102 (2013).
 - [8] Fritz, T. & Chaves, R. Entropic inequalities and marginal problems. *IEEE Transactions on Information Theory* **59**, 803–817 (2013).
 - [9] Chaves, R., Luft, L. & Gross, D. Causal structures from entropic information: geometry and novel scenarios. *New Journal of Physics* **16**, 043001 (2014).

- [10] Chaves, R. *et al.* Inferring latent structures via information inequalities. In *Proceedings of the 30th Conference on Uncertainty in Artificial Intelligence*, 112–121 (AUAI Press, Corvallis, Oregon, 2014).
- [11] Henson, J., Lal, R. & Pusey, M. F. Theory-independent limits on correlations from generalized Bayesian networks. *New Journal of Physics* **16**, 113043 (2014).
- [12] Steudel, B. & Ay, N. Information-theoretic inference of common ancestors. *Entropy* **17**, 2304–2327 (2015).
- [13] Chaves, R., Majenz, C. & Gross, D. Information-theoretic implications of quantum causal structures. *Nature communications* **6**, 5766 (2015).
- [14] Rosset, D. *et al.* Nonlinear Bell inequalities tailored for quantum networks. *Physical Review Letters* **116**, 010403 (2016).
- [15] Chaves, R. Polynomial Bell inequalities. *Physical Review Letters* **116**, 010402 (2016).
- [16] Pienaar, J. & Brukner, Č. A graph-separation theorem for quantum causal models. *New Journal of Physics* **17**, 073020 (2015).
- [17] Chaves, R. & Budroni, C. Entropic nonsignaling correlations. *Phys. Rev. Lett.* **116**, 240501 (2016). URL <https://link.aps.org/doi/10.1103/PhysRevLett.116.240501>.
- [18] Pienaar, J. Which causal structures might support a quantum–classical gap? *New Journal of Physics* **19**, 043021 (2017). URL <http://stacks.iop.org/1367-2630/19/i=4/a=043021>.
- [19] Wolfe, E., Spekkens, R. W. & Fritz, T. The inflation technique for causal inference with latent variables. e-print [arXiv:1609.00672](https://arxiv.org/abs/1609.00672) (2016).
- [20] Kela, A., von Prillwitz, K., Aberg, J., Chaves, R. & Gross, D. Semidefinite tests for latent causal structures. e-print arxiv.org/abs/1701.00652 (2017).
- [21] Ekert, A. K. Quantum cryptography based on Bell’s theorem. *Physical review letters* **67**, 661–663 (1991).
- [22] Mayers, D. & Yao, A. Quantum cryptography with imperfect apparatus. In *Proceedings of the 39th Annual Symposium on Foundations of Computer Science (FOCS-98)*, 503–509 (IEEE Computer Society, Los Alamitos, CA, USA, 1998).
- [23] Barrett, J., Hardy, L. & Kent, A. No signaling and quantum key distribution. *Physical review letters* **95**, 010503 (2005).
- [24] Acín, A., Gisin, N. & Masanes, L. From Bell’s theorem to secure quantum key distribution. *Physical review letters* **97**, 120405 (2006).
- [25] Colbeck, R. *Quantum and Relativistic Protocols For Secure Multi-Party Computation*. Ph.D. thesis, University of Cambridge (2007). Also available as [arXiv:0911.3814](https://arxiv.org/abs/0911.3814).
- [26] Colbeck, R. & Kent, A. Private randomness expansion with untrusted devices. *Journal of Physics A: Mathematical and Theoretical* **44**, 095305 (2011).
- [27] Pironio, S. *et al.* Random numbers certified by Bell’s theorem. *Nature* **464**, 1021–1024 (2010).
- [28] Vazirani, U. & Vidick, T. Fully device-independent quantum key distribution. *Physical review letters* **113**, 140501 (2014).
- [29] Miller, C. A. & Shi, Y. Universal security for randomness expansion from the spot-checking protocol. *SIAM Journal on Computing* **46**, 1304–1335 (2017).
- [30] Bell, J. S. On the Einstein-Podolsky-Rosen paradox. *Physics* **1**, 195–200 (1964).
- [31] Wood, C. J. & Spekkens, R. W. The lesson of causal discovery algorithms for quantum correlations: causal explanations of Bell-inequality violations require fine-tuning. *New Journal of Physics* **17**, 033002 (2015).
- [32] Shannon, C. E. A mathematical theory of communication. *Bell System Technical Journal* **27**, 379–423 (1948).
- [33] Yeung, R. A framework for linear information inequalities. *IEEE Transactions on Information Theory* **43**, 1924–1934 (1997).
- [34] Pippenger, N. The inequalities of quantum information theory. *IEEE Transactions on Information Theory* **49**, 773–789 (2003).
- [35] Chan, T. H. Balanced information inequalities. *IEEE Trans. Inf. Theor.* **49**, 3261–3267 (2003). URL <http://dx.doi.org/10.1109/TIT.2003.820037>.
- [36] Zhang, Z. & Yeung, R. W. A non-Shannon-type conditional inequality of information quantities. *IEEE Transactions on Information Theory* **43**, 1982–1986 (1997).
- [37] Han, T.-S. A uniqueness of Shannon’s information distance and related non-negativity problems. *Journal of Combinatorics, Information & System Sciences* **6**, 320–321 (1981).
- [38] Zhen Zhang & Yeung, R. On characterization of entropy function via information inequalities. *IEEE Transactions on Information Theory* **44**, 1440–1452 (1998). URL <http://ieeexplore.ieee.org/document/681320/>.
- [39] Matus, F. Infinitely many information inequalities. In *2007 IEEE International Symposium on Information Theory*, 41–44 (IEEE, 2007).
- [40] Makarychev, K., Makarychev, Y., Romashchenko, A. & Vereshchagin, N. A new class of non-Shannon-type inequalities for entropies. *Communications in Information and Systems* **2**, 147–166 (2002).
- [41] Dougherty, R., Freiling, C. & Zeger, K. Six new non-Shannon information inequalities. In *2006 IEEE International Symposium on Information Theory*, 233–236 (IEEE, 2006).
- [42] Weidong Xu, Jia Wang & Jun Sun. A projection method for derivation of non-Shannon-type information inequalities. In *2008 IEEE International Symposium on Information Theory*, 2116–2120 (IEEE, 2008).
- [43] Dougherty, R., Freiling, C. & Zeger, K. Non-Shannon information inequalities in four random variables. e-print [arXiv:1104.3602](https://arxiv.org/abs/1104.3602) (2011).
- [44] Kaced, T. Equivalence of two proof techniques for non-Shannon-type inequalities. In *Information Theory Proceedings (ISIT), 2013 IEEE International Symposium on*, 236–240 (IEEE, 2013).
- [45] Ingleton, A. W. Representation of matroids. In Welsh, D. J. A. (ed.) *Combinatorial Mathematics and its applications*, 149–167 (Academic Press, 1971).
- [46] Hammer, D., Romashchenko, A., Shen, A. & Vereshchagin, N. Inequalities for Shannon entropy and Kolmogorov complex-

- ity. *Journal of Computer and System Sciences* **60**, 442–464 (2000). URL <http://linkinghub.elsevier.com/retrieve/pii/S002200009991677X>.
- [47] Dougherty, R., Freiling, C. & Zeger, K. Linear rank inequalities on five or more variables. e-print [arXiv:0910.0284](https://arxiv.org/abs/0910.0284) (2009).
- [48] Dougherty, R. Computations of linear rank inequalities on six variables. In *2014 IEEE International Symposium on Information Theory*, 2819–2823 (IEEE, 2014). URL <http://ieeexplore.ieee.org/lpdocs/epic03/wrapper.htm?arnumber=6875348>.
- [49] Pearl, J. On the testability of causal models with latent and instrumental variables. In *Proceedings of the Eleventh conference on Uncertainty in artificial intelligence*, 435–443 (Morgan Kaufmann Publishers Inc., 1995).
- [50] Weilenmann, M. & Colbeck, R. Non-Shannon inequalities in the entropy vector approach to causal structures. e-print [arXiv:1605.02078](https://arxiv.org/abs/1605.02078) (2016).
- [51] Verma, T. & Pearl, J. Causal networks: Semantics and expressiveness. In *Proceedings of the 4th Workshop on Uncertainty in Artificial Intelligence, Minneapolis, MN, Mountain View, CA*, 352–359 (1988).
- [52] Meek, C. Strong completeness and faithfulness in Bayesian networks. In *Proceedings of the Eleventh conference on Uncertainty in artificial intelligence*, 411–418 (Morgan Kaufmann Publishers Inc., 1995).
- [53] Williams, H. P. Fourier’s method of linear programming and its dual. *The American Mathematical Monthly* **93**, 681–695 (1986).
- [54] Leifer, M. S. & Spekkens, R. W. Towards a formulation of quantum theory as a causally neutral theory of Bayesian inference. *Physical Review A* **88**, 052130 (2013).
- [55] Stinespring, W. F. Positive functions on C^* -algebras. *Proceedings of the American Mathematical Society* **6**, 211–216 (1955).
- [56] Lieb, E. H. & Ruskai, M. B. Proof of the strong subadditivity of quantum-mechanical entropy. *Journal of Mathematical Physics* **14**, 1938 (1973).
- [57] Araki, H. & Lieb, E. H. Entropy inequalities. *Communications in Mathematical Physics* **18**, 160–170 (1970).
- [58] Linden, N. & Winter, A. A new inequality for the von Neumann entropy. *Communications in Mathematical Physics* **259**, 129–138 (2005).
- [59] Cadney, J., Linden, N. & Winter, A. Infinitely many constrained inequalities for the von Neumann entropy. *IEEE Transactions on Information Theory* **58**, 3657–3663 (2012).
- [60] Linden, N., Matus, F., Ruskai, M. B. & Winter, A. The quantum entropy cone of stabiliser states. In *LIPICs - Leibniz International Proceedings in Informatics*, vol. 22, 284 (Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2013).
- [61] Gross, D. & Walter, M. Stabilizer information inequalities from phase space distributions. *Journal of Mathematical Physics* **54**, 082201 (2013).
- [62] Nielsen, M. A. & Chuang, I. L. *Quantum Computation and Quantum Information* (Cambridge University Press, 2000).
- [63] Tsirelson, B. Some results and problems on quantum Bell-type inequalities. *Hadronic Journal Supplement* **8**, 329–345 (1993).
- [64] Popescu, S. & Rohrlich, D. Quantum nonlocality as an axiom. *Foundations of Physics* **24**, 379–385 (1994).
- [65] Fritz, T. Beyond Bell’s theorem II: Scenarios with arbitrary causal structure. *Communications in Mathematical Physics* **341**, 391–434 (2015).
- [66] Chiribella, G., D’Ariano, G. M. & Perinotti, P. Probabilistic theories with purification. *Phys. Rev. A* **81**, 062348 (2010). URL <https://link.aps.org/doi/10.1103/PhysRevA.81.062348>.
- [67] Fritz, T. Beyond Bell’s theorem: correlation scenarios. *New Journal of Physics* **14**, 103001 (2012).
- [68] Short, A. J. & Barrett, J. Strong nonlocality: a trade-off between states and measurements. *New Journal of Physics* **12**, 033034 (2010).
- [69] Barnum, H. *et al.* Entropy and information causality in general probabilistic theories. *New Journal of Physics* **12**, 033024 (2010). URL <http://stacks.iop.org/1367-2630/12/i=3/a=033024>.
- [70] Cadney, J. & Linden, N. Measurement entropy in generalized nonsignalling theory cannot detect bipartite nonlocality. *Phys. Rev. A* **86**, 052103 (2012). URL <https://link.aps.org/doi/10.1103/PhysRevA.86.052103>.
- [71] Barrett, J. Information processing in generalized probabilistic theories. *Physical Review A* **75**, 032304 (2007).
- [72] Reichenbach, H. *The direction of time* (University of California Press, 1956).
- [73] Costa, F. & Shrapnel, S. Quantum causal modelling. e-print [arXiv:1512.07106](https://arxiv.org/abs/1512.07106) (2015).
- [74] Allen, J.-M., Barrett, J., Horsman, D., Lee, C. & Spekkens, R. Quantum common causes and quantum causal models. e-print arxiv.org/abs/1609.09487 (2016).
- [75] Hardy, L. Probability theories with dynamic causal structure: a new framework for quantum gravity. e-print [arXiv:gr-qc/0509120](https://arxiv.org/abs/gr-qc/0509120) (2005).
- [76] Hardy, L. Towards quantum gravity: a framework for probabilistic theories with non-fixed causal structure. *Journal of Physics A: Mathematical and Theoretical* **40**, 3081 (2007). URL <http://stacks.iop.org/1751-8121/40/i=12/a=S12>.
- [77] Hardy, L. Quantum gravity computers: On the theory of computation with indefinite causal structure. In *Quantum Reality, Relativistic Causality, and Closing the Epistemic Circle*, 379–401 (Springer, 2009).
- [78] Oreshkov, O., Costa, F. & Brukner, C. Quantum correlations with no causal order. *Nature communications* **3**, 1092 (2012).
- [79] Baumeler, A. & Wolf, S. Perfect signaling among three parties violating predefined causal order. In *Information Theory (ISIT), 2014 IEEE International Symposium on*, 526–530 (IEEE, 2014).
- [80] Chiribella, G., D’Ariano, G. M., Perinotti, P. & Valiron, B. Quantum computations without definite causal structure. *Phys. Rev. A* **88**, 022318 (2013). URL <https://link.aps.org/doi/10.1103/PhysRevA.88.022318>.

- [81] Portmann, C., Matt, C., Maurer, U., Renner, R. & Tackmann, B. Causal boxes: quantum information-processing systems closed under composition. *IEEE Transactions on Information Theory* **63**, 3277–3305 (2017).
- [82] Miklin, N., Abbott, A. A., Branciard, C., Chaves, R. & Budroni, C. The entropic approach to causal correlations. e-print [arXiv:1706.10270](https://arxiv.org/abs/1706.10270) (2017).
- [83] Colbeck, R. & Renner, R. A system’s wave function is uniquely determined by its underlying physical state. *New Journal of Physics* **19**, 013016 (2017).
- [84] Colbeck, R. & Renner, R. The completeness of quantum theory for predicting measurement outcomes. In Chiribella, G. & Spekkens, R. W. (eds.) *Quantum Theory: Informational Foundations and Foils*, 497–528 (Springer, 2016).
- [85] Braunstein, S. L. & Caves, C. M. Information-theoretic Bell inequalities. *Physical Review Letters* **61**, 662–665 (1988).
- [86] Fine, A. Joint distributions, quantum correlations, and commuting observables. *Journal of Mathematical Physics* **23**, 1306 (1982).
- [87] Fine, A. Hidden variables, joint probability, and the Bell inequalities. *Physical Review Letters* **48**, 291–295 (1982).
- [88] Liang, Y.-C., Spekkens, R. W. & Wiseman, H. M. Specker’s parable of the overprotective seer: A road to contextuality, nonlocality and complementarity. *Physics Reports* **506**, 1 – 39 (2011). URL <http://www.sciencedirect.com/science/article/pii/S0370157311001517>.
- [89] Abramsky, S. & Brandenburger, A. The sheaf-theoretic structure of non-locality and contextuality. *New Journal of Physics* **13**, 113036 (2011). URL <http://stacks.iop.org/1367-2630/13/i=11/a=113036>.
- [90] Araújo, M., Quintino, M. T., Budroni, C., Cunha, M. T. & Cabello, A. All noncontextuality inequalities for the n -cycle scenario. *Phys. Rev. A* **88**, 022118 (2013). URL <https://link.aps.org/doi/10.1103/PhysRevA.88.022118>.
- [91] Branciard, C., Gisin, N. & Pironio, S. Characterizing the nonlocal correlations created via entanglement swapping. *Physical review letters* **104**, 170401 (2010).
- [92] Branciard, C., Rosset, D., Gisin, N. & Pironio, S. Bilocal versus nonbilocal correlations in entanglement-swapping experiments. *Physical Review A* **85**, 032119 (2012).
- [93] Budroni, C., Miklin, N. & Chaves, R. Indistinguishability of causal relations from limited marginals. *Phys. Rev. A* **94**, 042127 (2016). URL <https://link.aps.org/doi/10.1103/PhysRevA.94.042127>.
- [94] Pawłowski, M. *et al.* Information causality as a physical principle. *Nature* **461**, 1101–1104 (2009). URL <http://www.nature.com/doi/10.1038/nature08400>.
- [95] Masanes, L., Acín, A. & Gisin, N. General properties of nonsignaling theories. *Phys. Rev. A* **73**, 012112 (2006). URL <https://link.aps.org/doi/10.1103/PhysRevA.73.012112>.
- [96] Pawłowski, M. & Brukner, Č. Monogamy of Bell’s inequality violations in nonsignaling theories. *Phys. Rev. Lett.* **102**, 030403 (2009). URL <https://link.aps.org/doi/10.1103/PhysRevLett.102.030403>.
- [97] Popescu, S. Nonlocality beyond quantum mechanics. *Nature Physics* **10**, 264–270 (2014).
- [98] Evans, R. J. Graphical methods for inequality constraints in marginalized DAGs. In *Machine Learning for Signal Processing (MLSP), 2012 IEEE International Workshop on*, 1–6 (IEEE, 2012).
- [99] Evans, R. J. Graphs for margins of Bayesian networks. *Scandinavian Journal of Statistics* (2015).
- [100] Rényi, A. On measures of entropy and information. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Vol. 1: Contributions to the Theory of Statistics*, 547–561 (1960).
- [101] Linden, N., Mosonyi, M. & Winter, A. The structure of Rényi entropic inequalities. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **469**, 20120737–20120737 (2013).
- [102] Petz, D. Quasi-entropies for finite quantum systems. *Reports on Mathematical Physics* **23**, 57–65 (1986).
- [103] Tomamichel, M., Colbeck, R. & Renner, R. A fully quantum asymptotic equipartition property. *IEEE Transactions on information theory* **55**, 5840–5847 (2009).
- [104] Müller-Lennert, M., Dupuis, F., Szehr, O., Fehr, S. & Tomamichel, M. On quantum Rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics* **54**, 122203 (2013).
- [105] Frank, R. L. & Lieb, E. H. Monotonicity of a relative Rényi entropy. *Journal of Mathematical Physics* **54**, 122201 (2013).
- [106] Beigi, S. Sandwiched Rényi divergence satisfies data processing inequality. *Journal of Mathematical Physics* **54**, 122202 (2013).
- [107] Dupuis, F. Chain rules for quantum Rényi entropies. *Journal of Mathematical Physics* **56**, 022203 (2015).
- [108] Pitowsky, I. Correlation polytopes: Their geometry and complexity. *Mathematical Programming* **50**, 395–414 (1991).
- [109] Avis, D., Imai, H., Ito, T. & Sasaki, Y. Deriving tight Bell inequalities for 2 parties with many 2-valued observables from facets of cut polytopes. e-print [arXiv:quant-ph/0404014](https://arxiv.org/abs/quant-ph/0404014) (2004).
- [110] Navascués, M. & Wolfe, E. The inflation technique solves completely the classical inference problem. e-print [arXiv:1707.06476](https://arxiv.org/abs/1707.06476) (2017).

Appendix A: Entropy inequalities for Example 10

In the following we provide the basic inequalities for the quantum instrumental scenario, IC^Q , i.e., the constraints making up the matrix $M_B(IC^Q)$.

$$I(A_Y : A_Z) \geq 0 \tag{A1}$$

$$I(A_Y : X) \geq 0 \tag{A2}$$

$$I(A_Z : X) \geq 0 \tag{A3}$$

$$I(A_Y : A_Z|X) \geq 0 \tag{A4}$$

$$I(A_Y : X|A_Z) \geq 0 \tag{A5}$$

$$I(A_Z : X|A_Y) \geq 0 \tag{A6}$$

$$I(A_Y : Z) \geq 0 \tag{A7}$$

$$I(X : Z) \geq 0 \tag{A8}$$

$$I(X : Z|A_Y) \geq 0 \tag{A9}$$

$$I(A_Y : Z|X) \geq 0 \tag{A10}$$

$$I(A_Y : X|Z) \geq 0 \tag{A11}$$

$$I(X : Y) \geq 0 \tag{A12}$$

$$I(Y : Z) \geq 0 \tag{A13}$$

$$I(Y : Z|X) \geq 0 \tag{A14}$$

$$I(X : Z|Y) \geq 0 \tag{A15}$$

$$I(X : Y|Z) \geq 0 \tag{A16}$$

$$H(A_Z|X) \geq 0 \tag{A17}$$

$$H(A_Y A_Z|X) \geq 0 \tag{A18}$$

$$H(X|A_Y A_Z) \geq 0 \tag{A19}$$

$$H(A_Y|XZ) \geq 0 \tag{A20}$$

$$H(X|A_Y Z) \geq 0 \tag{A21}$$

$$H(Z|A_Y X) \geq 0 \tag{A22}$$

$$H(X|YZ) \geq 0 \tag{A23}$$

$$H(Y|XZ) \geq 0 \tag{A24}$$

$$H(Z|XY) \geq 0 \tag{A25}$$

$$H(A_Z|A_Y) + H(A_Z|X) \geq 0 \tag{A26}$$

$$H(A_Y|A_Z) + H(A_Y|X) \geq 0 \tag{A27}$$

$$H(A_Z|A_Y X) + H(A_Z) \geq 0 \tag{A28}$$

$$H(A_Y|A_Z X) + H(A_Y) \geq 0 \tag{A29}$$

Independence constraints and data processing inequalities are provided in the main text. If we include these and remove redundant inequalities we obtain the following set of constraints, which for convenience we give in matrix form (such that $\Gamma(IC^Q) = \{v \in \mathbb{R}_{\geq 0}^{15} \mid M \cdot v \geq 0\}$):




Figure 6: Post-selecting (a) on a binary observed variable C leads to the causal structure (b).

$$M = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

Appendix B: Entropy inequalities for Example 16

The causal structure of Figure 6, previously considered in [11, 18] is analysed entropically by means of the entropic post-selection technique. The outer approximation to the entropic cone of the causal structure of Figure 6 is computed and marginalised to vectors

$$\begin{aligned} & (H(Z), H(X|_{C=0}), H(X|_{C=1}), H(Y|_{C=0}), H(Y|_{C=1}), H(X|_{C=0}Z), H(X|_{C=1}Z), H(Y|_{C=0}Z), \\ & H(Y|_{C=1}Z), H(X|_{C=0}Y|_{C=0}), H(X|_{C=1}Y|_{C=1}), H(X|_{C=0}Y|_{C=0}Z), H(X|_{C=1}Y|_{C=1}Z)). \end{aligned}$$

The following 14 extremal rays are obtained from this computation, where each ray is represented by one particular vector on it. The tip of this pointed polyhedral cone is the zero-vector.

- (1) 0000100010101
- (2) 0001000101010
- (3) 0010001000101
- (4) 0100010001010
- (5) 1111122222222
- (6) 1010112120212
- (7) 1101021212021
- (8) 1000011110011
- (9) 1000111110111
- (10) 1001011111011
- (11) 1010011110111
- (12) 1100011111011
- (13) 1001111111111
- (14) 1110011111111

The corresponding inequality description is given by the 2 equalities and 18 inequalities (or equivalently 22 inequalities if each equality is written as two inequalities).

$$H(X_{|C=0}) + H(Y_{|C=0}) = H(X_{|C=0}Y_{|C=0}) \quad (\text{B1})$$

$$H(X_{|C=1}) + H(Y_{|C=1}) = H(X_{|C=1}Y_{|C=1}) \quad (\text{B2})$$

$$H(X_{|C=1}Y_{|C=1}) \leq H(X_{|C=1}Y_{|C=1}Z) \quad (\text{B3})$$

$$H(X_{|C=0}Y_{|C=0}) \leq H(X_{|C=0}Y_{|C=0}Z) \quad (\text{B4})$$

$$H(Y_{|C=1}Z) \leq H(X_{|C=1}Y_{|C=1}Z) \quad (\text{B5})$$

$$H(Y_{|C=0}Z) \leq H(X_{|C=0}Y_{|C=0}Z) \quad (\text{B6})$$

$$H(X_{|C=1}Z) \leq H(X_{|C=1}Y_{|C=1}Z) \quad (\text{B7})$$

$$H(X_{|C=0}Z) \leq H(X_{|C=0}Y_{|C=0}Z) \quad (\text{B8})$$

$$H(Y_{|C=0}Z) \leq H(Z) + H(Y_{|C=0}) \quad (\text{B9})$$

$$H(Y_{|C=1}Z) \leq H(Z) + H(Y_{|C=1}) \quad (\text{B10})$$

$$H(Y_{|C=1}) + H(X_{|C=1}Z) \leq H(Z) + H(X_{|C=1}Y_{|C=1}) \quad (\text{B11})$$

$$H(Y_{|C=0}) + H(X_{|C=0}Z) \leq H(Z) + H(X_{|C=0}Y_{|C=0}) \quad (\text{B12})$$

$$H(Z) + H(X_{|C=0}Y_{|C=0}Z) \leq H(X_{|C=0}Z) + H(Y_{|C=0}Z) \quad (\text{B13})$$

$$H(Z) + H(X_{|C=1}Y_{|C=1}Z) \leq H(X_{|C=1}Z) + H(Y_{|C=1}Z) \quad (\text{B14})$$

$$H(Y_{|C=1}) + H(X_{|C=1}Z) + H(X_{|C=0}Y_{|C=0}) \leq H(Y_{|C=0}) + H(X_{|C=0}Z) + H(X_{|C=1}Y_{|C=1}Z) \quad (\text{B15})$$

$$H(Y_{|C=1}) + H(Y_{|C=0}Z) + H(X_{|C=0}Y_{|C=0}) \leq H(Y_{|C=0}) + H(Y_{|C=1}Z) + H(X_{|C=0}Y_{|C=0}Z) \quad (\text{B16})$$

$$H(X_{|C=1}Z) + H(Y_{|C=1}Z) + H(X_{|C=0}Y_{|C=0}) \leq H(X_{|C=0}Z) + H(Y_{|C=0}Z) + H(X_{|C=1}Y_{|C=1}Z) \quad (\text{B17})$$

$$H(X_{|C=0}Z) + H(Y_{|C=0}Z) + H(X_{|C=1}Y_{|C=1}) \leq H(X_{|C=1}Z) + H(Y_{|C=1}Z) + H(X_{|C=0}Y_{|C=0}Z) \quad (\text{B18})$$

$$H(Y_{|C=0}) + H(Y_{|C=1}Z) + H(X_{|C=1}Y_{|C=1}) \leq H(Y_{|C=1}) + H(Y_{|C=0}Z) + H(X_{|C=1}Y_{|C=1}Z) \quad (\text{B19})$$

$$H(Y_{|C=0}) + H(X_{|C=0}Z) + H(X_{|C=1}Y_{|C=1}) \leq H(Y_{|C=1}) + H(X_{|C=1}Z) + H(X_{|C=0}Y_{|C=0}Z). \quad (\text{B20})$$