

This is a repository copy of *Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants (Review)*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/133812/>

Version: Published Version

Article:

Gordon, Adrienne, Greenhalgh, Mark and McGuire, William orcid.org/0000-0001-8572-3467 (2018) Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants (Review). Cochrane Database of Systematic Reviews. CD012141. ISSN 1469-493X

<https://doi.org/10.1002/14651858.CD012141.pub2>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Cochrane
Library

Cochrane Database of Systematic Reviews

Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants (Review)

Gordon A, Greenhalgh M, McGuire W

Gordon A, Greenhalgh M, McGuire W.

Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants.

Cochrane Database of Systematic Reviews 2018, Issue 6. Art. No.: CD012141.

DOI: 10.1002/14651858.CD012141.pub2.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	2
OBJECTIVES	4
METHODS	4
RESULTS	6
Figure 1.	7
DISCUSSION	8
AUTHORS' CONCLUSIONS	8
ACKNOWLEDGEMENTS	8
REFERENCES	9
CHARACTERISTICS OF STUDIES	12
CONTRIBUTIONS OF AUTHORS	13
DECLARATIONS OF INTEREST	13
SOURCES OF SUPPORT	13

[Intervention Review]

Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants

Adrienne Gordon¹, Mark Greenhalgh², William McGuire³

¹Neonatology, Royal Prince Alfred Hospital, Sydney, Australia. ²RPA Newborn Care, RPA Women and Babies, Royal Prince Alfred Hospital, Sydney, Australia. ³Centre for Reviews and Dissemination, University of York, York, UK

Contact address: William McGuire, Centre for Reviews and Dissemination, University of York, York, Y010 5DD, UK. William.McGuire@hyms.ac.uk.

Editorial group: Cochrane Neonatal Group.

Publication status and date: New, published in Issue 6, 2018.

Citation: Gordon A, Greenhalgh M, McGuire W. Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants. *Cochrane Database of Systematic Reviews* 2018, Issue 6. Art. No.: CD012141. DOI: 10.1002/14651858.CD012141.pub2.

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Duration of use may be a modifiable risk factor for central venous catheter-associated bloodstream infection in newborn infants. Early planned removal of peripherally inserted central catheters (PICCs) is recommended as a strategy to reduce the incidence of infection and its associated morbidity and mortality.

Objectives

To determine the effectiveness of early planned removal of PICCs (up to two weeks after insertion) compared to an expectant approach or a longer fixed duration in preventing bloodstream infection and other complications in newborn infants.

Search methods

We searched of the Cochrane Central Register of Controlled Trials (CENTRAL; 2018, Issue 4), Ovid MEDLINE, Embase, Maternity & Infant Care Database, and the Cumulative Index to Nursing and Allied Health Literature (CINAHL) (until April 2018), and conference proceedings and previous reviews.

Selection criteria

Randomised and quasi-randomised controlled trials that assessed the effect of early planned removal of umbilical venous catheters (up to two weeks after insertion) compared to an expectant management approach or a longer fixed duration in preventing bloodstream infection and other complications in newborn infants.

Data collection and analysis

Two review authors assessed trial eligibility independently. We planned to analyse any treatment effects in the individual trials and report the risk ratio and risk difference for dichotomous data and mean difference for continuous data, with respective 95% confidence intervals. We planned to use a fixed-effect model in meta-analyses and explore potential causes of heterogeneity in sensitivity analyses. We planned to assess the quality of evidence for the main comparison at the outcome level using “Grading of Recommendations Assessment, Development and Evaluation” (GRADE) methods.

Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants (Review)

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Main results

We did not identify any eligible randomised controlled trials.

Authors' conclusions

There are no trial data to guide practice regarding early planned removal versus expectant management of PICCs in newborn infants. A simple and pragmatic randomised controlled trial is needed to resolve the uncertainty about optimal management in this common and important clinical dilemma.

PLAIN LANGUAGE SUMMARY

Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants

Review question

In newborn infants with a peripherally inserted central catheter in place, does early removal of the catheter reduce the risk of complications, including infection?

Background

Infection in the bloodstream is a frequent and harmful complication for newborn infants who have a peripherally inserted central catheter (a long, narrow, soft and flexible plastic tube inserted through the skin into a vein and advanced several centimetres into the infant's large blood vessels; used as a stable route to deliver drugs and nutrition). Bloodstream infection may cause death and disability. One potential method of reducing the risk of this and other serious complications is to remove the catheter within about two weeks after insertion rather than leaving it for longer until no longer required.

Study characteristics/key results

We did not find any randomised controlled trials that assessed whether removing peripherally inserted central catheters within two weeks prevents infection or other complications in newborn infants.

Conclusions

There are no trial data available to help clinicians to address this common clinical dilemma. Due to the potential for benefit and harm, such a trial may be warranted.

BACKGROUND

Description of the condition

Peripherally inserted central catheters (PICCs), also commonly referred to as percutaneous central venous catheters (CVCs), provide a stable route for the intravenous delivery of drugs or fluids to preterm or sick newborn infants. Typically, PICCs are inserted via a superficial vein in the upper or lower limb, usually avoiding proximal sites such as the femoral or axillary veins, and are then advanced so that the tip of the catheter is sited at the vena caval

junction with the right atrium of the heart (Jain 2013; Wrightson 2013). Peripherally inserted central catheters are less expensive and easier to insert than catheters placed directly or via a subcutaneous tunnel into central veins. Because they are more stable than short peripheral cannulae, their use reduces the risk of extravasation injury from hyperosmolar parenteral nutrition solutions and medications (Ainsworth 2015).

Bloodstream infection is the most common serious complication associated with the use of PICCs in newborn infants. The reported incidence ranges from about 2% to 30% depending on the precise diagnostic criteria and the demographics of the population studied (Chien 2002; Cartwright 2004; van der Zwet 2005;

Garland 2008; Hoang 2008; Olsen 2009; O'Grady 2011; Ohki 2013). Very preterm infants are at the highest risk, but inter-unit variation in the incidence of catheter-associated bloodstream infection is not fully explained by case-mix and may relate to care or infection control practices (Wong 2012). Newborn infants, particularly very preterm infants, with an acquired bloodstream infection have a higher risk of mortality and a range of important morbidities including bronchopulmonary dysplasia, necrotising enterocolitis, retinopathy of prematurity, hepatic dysfunction and prolonged hospitalisation (Saint 2000; Mahieu 2001a; Mahieu 2001b; Chapman 2003; Payne 2004; Adams-Chapman 2006; Hermans 2007; Lahra 2009). Bloodstream infection is associated with higher rates of adverse neurodevelopmental outcomes including cognitive or sensory impairment and cerebral palsy (Stoll 2004; Shah 2008a; Bassler 2009).

The most common causes of PICC-associated bloodstream infections in newborn infants are coagulase-negative staphylococci, other Gram-positive cocci (*Staphylococcus aureus* (*S. aureus*), enterococci), Gram-negative bacilli and fungi (predominantly *Candida* species) (Makhoul 2002; O'Grady 2002; Stoll 2002; Isaacs 2003; Isaacs 2004; Gordon 2006; de Brito 2010). Micro-organisms can gain access through the PICC entry site or via the catheter hub into the PICC lumen or tract. Pathogens adhere to the material of the PICC and secrete an intraluminal or extraluminal biofilm of extracellular polymeric substances (Machado 2009). Bacteria or fungi growing within the biofilm are relatively protected from the host's immune system and circulating antimicrobial agents thus enabling sustained colonisation (Ramirez de Arellano 1994; Stewart 2001). It is often necessary to remove the PICC in order to clear the infection (Benjamin 2001). A thrombus (blood clot) attached to the PICC may be an additional nidus (locus) for infection (Thornburg 2008). Some evidence exists that continuous heparin infusion can reduce the incidence of PICC-occlusion in neonates, but the data are insufficient to determine the effect of this intervention on the risk of infection (Shah 2008b).

Several strategies to prevent PICC-associated bloodstream infections have been developed and adopted, often as multifaceted packages of interventions ('care bundles'). These include strict aseptic precautions when inserting and accessing the PICC, use of needleless intravascular catheter systems and prompt removal when the PICC is no longer needed (O'Grady 2002; Yébenes 2004; Pronovost 2006; Borghesi 2008; Miller 2010; Sannoh 2010; Vanholder 2010; Wirtschafter 2010; Kaplan 2011; O'Grady 2011; Schulman 2011). Care bundles have been shown to reduce bloodstream infection rates in adult, paediatric and neonatal intensive care studies (Pronovost 2006; Miller 2010; Wirtschafter 2010; Kaplan 2011; Schulman 2011; Fisher 2013). Despite these strategies, however, PICC-associated infections remain a major cause of morbidity and mortality in newborn infants and other interventions are required to reduce the infection rates further.

Description of the intervention

Although strong associations between PICC exposure, especially to deliver parenteral nutrition, and the risk of bloodstream infection have been described, observational studies have not provided consistent evidence about the effect of duration of PICC placement ('dwell time') on the risk of invasive infection in newborn infants. Some reports indicate an elevated risk of infection when the PICC has been in place for more than about 10 to 14 days, while others do not show any association until after a dwell time of four to six weeks (Smith 2008; Sengupta 2010; Wong 2012; Milstone 2013; Greenberg 2015). Furthermore, it is not certain to what extent PICC use is an independent risk factor for a bloodstream infection or whether observed associations exist because infants who are smaller, less mature, sicker and receiving more intensive and invasive support are also more likely to have a PICC *in situ*. A Cochrane review of randomised controlled trials of PICCs versus peripheral cannulae for delivering parenteral nutrition to neonates did not show an effect on invasive infection rates. Infants with a PICC *in situ* experienced fewer skin break procedures than infants with a peripheral cannula (or series of cannulae) and this may have balanced the overall risk for acquiring bloodstream infection (Ainsworth 2015).

This review examines the evidence from randomised controlled trials that early planned removal within a pre-specified maximum dwell time versus allowing clinicians to determine when the PICC is removed or replaced affects the risk of bloodstream infection in newborn infants. Limited guidance about this issue is available. The 2011 US Centers for Disease Control and Prevention Healthcare Infection Control Practices Advisory Committee guidelines do not specify a recommended maximum dwell time and advise that PICCs should not be routinely replaced to prevent catheter-related infections (O'Grady 2011). This uncertainty is reflected in surveys of practice that identify wide variation between neonatal care centres with regard to PICC insertion and maintenance (Sharpe 2013; Taylor 2014).

We define the maximum dwell time pragmatically as up to two weeks after insertion based on the typical duration of PICC use for infants in most neonatal units and the minimum time that observational studies have shown this to be associated with a rise in the risk of infection (Smith 2008; Sengupta 2010; Wong 2012; Milstone 2013). The intervention is the pre-specified intent to remove or replace the PICC within this time period, and the control is either (i) any permissive approach that does not pre-specify dwell time but that allows PICC removal or replacement based on clinical criteria (including suspected or confirmed bloodstream infection), or (ii) a longer pre-specified dwell time than the intervention.

How the intervention might work

Pre-specifying a fixed maximum dwell time, with planned removal

rather than an expectant approach, may reduce the risk of a PICC being left *in situ* when not actually in use or needed and reduce the overall length of PICC exposure. These effects could lower the risk of bloodstream infection and its associated complications if the PICC is an independent risk factor for infection. This intervention may also plausibly affect nutrient intake, either reducing receipt of parenteral nutrients, or prompting a more rapid progression to full enteral feeding, or both, with potential consequences for acute morbidity (principally the risk of acute necrotising enterocolitis), growth and development.

Why it is important to do this review

Given the potential for the planned duration of placement of PICCs to affect important outcomes for newborn infants, we undertook a systematic review to identify, appraise and synthesise the available evidence from randomised controlled trials.

Related Cochrane reviews

Other Cochrane reviews assess the effects of other strategies including antimicrobial impregnation or antibiotic locks to prevent PICC-related infection in newborn infants, and early removal versus expectant management of PICC in infants with suspected bloodstream infection (Vasudevan 2011; Balain 2015; Taylor 2015). Another review evaluates the evidence for short-versus longer-term use of umbilical venous catheters for newborn infants (Gordon 2016b).

OBJECTIVES

To determine the effectiveness of early planned removal of PICCs (up to two weeks after insertion), compared to an expectant approach or a longer fixed duration, in preventing infection in newborn infants. We pre-specified subgroup analyses by gestational age at birth (Subgroup analysis and investigation of heterogeneity).

METHODS

Criteria for considering studies for this review

Types of studies

Randomised or quasi-randomised controlled trials, including cluster-randomised controlled trials.

Types of participants

Newborn infants who are to have a PICC placed.

Types of interventions

- Intervention: early planned removal after fixed-term use of PICC (up to two weeks after insertion).
- Control: (i) permissive and expectant approach that allows retention of PICC if needed and in the absence of clinical indicators for removal or replacement, (ii) longer fixed-term use of PICC: at least one week longer than the intervention term.

Trials that assessed the effect of a pre-specified intended duration of PICC placement as part of a package of infection control measures (care bundle) were eligible for inclusion but we planned to analyse them separately from trials of discrete interventions.

We did not intend to include trials where the duration of PICC use was related specifically to the infant's progress in enteral feed intake.

Types of outcome measures

Primary outcomes

- Incidence of laboratory-confirmed bloodstream infection confirmed by culture from blood sampled from peripheral sites (not from indwelling catheters) during hospital admission. 'False-positive' results due to skin contaminants are possible, therefore (where data were available), we planned to exclude cases where infection was attributed to diphtheroids, micrococci, *Propionibacteriaceae* or mixed microbial flora. If sufficient data were available, we planned to examine specific infections with these organisms:
 - coagulase-negative staphylococci;
 - other bacteria (Gram-negative bacilli, *S. aureus*, enterococci);
 - fungi.

Secondary outcomes

- Death before hospital discharge and up one year post-term due to all causes.
- Neurodevelopmental outcomes assessed after 12 months post-term using validated tools: neurological evaluations; developmental scores; and classifications of disability, including auditory and visual disability. We defined neurodevelopmental impairment as the presence of one or more of the following: non-ambulant cerebral palsy; developmental quotient more than two standard deviations below the population mean; and blindness (visual acuity less than 6/60) or deafness (any hearing impairment requiring or unimproved by amplification).

- Death or neurological impairment assessed after 12 months post-term.
- Growth: time (days) to regain birth weight and average rate of weight gain (g/kg/day), linear growth (mm/week), head growth (mm/week) and skinfold thickness growth (mm/week) during hospital admission.
- Extravasation injury: subcutaneous extravasation resulting in skin ulceration; 'deep' extravasation resulting in limb swelling; or 'central' extravasation-infusate in the pleural, peritoneal or pericardial space.
- Number of cannulae or catheters used per infant to administer parenteral fluids until full enteral feeding established.
- Days to full enteral feeding.
- PICC leak, obstruction or breakage necessitating removal of PICC.
- PICC-associated thrombosis necessitating removal of PICC.
- Other morbidity developing after enrolment in the trial until discharge from hospital:
 - bronchopulmonary dysplasia (oxygen supplementation at 36 weeks postmenstrual age);
 - necrotising enterocolitis (Bell stage 2 or 3);
 - retinopathy of prematurity, requiring treatment (medical or surgical).

Search methods for identification of studies

We used the standard search strategy of Cochrane Neonatal (<http://neonatal.cochrane.org/>).

Electronic searches

We searched the Cochrane Central Register of Controlled Trials (CENTRAL, 2018, issue 4), Ovid MEDLINE (1946 to April 2018), OVID Embase (1974 to April 2018), OVID Maternity & Infant Care Database (1971 to April 2018), and CINAHL (1982 to April 2018) using a combination of the text words and MeSH terms described in Appendix 1. We limited the search outputs with the relevant search filters for clinical trials as recommended in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We did not apply any language restrictions. We searched ClinicalTrials.gov and the World Health Organization's International Trials Registry Platform (<http://apps.who.int/trialsearch/Default.aspx>) for completed or ongoing trials.

Searching other resources

We examined reference lists in previous reviews and planned to search the reference lists of any included studies. We searched the proceedings of the annual meetings of the Pediatric Academic Societies (1993 to 2017), the European Society for Pediatric Research (1995 to 2017), the Royal College of Paediatrics and Child Health (2000 to 2018) and the Perinatal Society of Australia and New

Zealand (2000 to 2018). Trials reported only as abstracts were eligible if sufficient information was available from the report, or from contact with the authors, to fulfil the inclusion criteria.

Data collection and analysis

We used the standard methods of Cochrane Neonatal (<http://neonatal.cochrane.org/>).

Selection of studies

We screened the title and abstract of all studies identified by the above search strategy and two review authors (MG and WM) independently assessed the full-text articles for all potentially relevant trials. We excluded those studies that did not meet all of the inclusion criteria and stated the reason for exclusion. We planned to discuss any disagreements with a third author (AG) until consensus was achieved.

Data extraction and management

Two review authors (MG and WM) planned to extract data independently using a data collection form to aid extraction of information on design, methodology, participants, interventions, outcomes and treatment effects from each included study. We planned to discuss any disagreements until we reached a consensus. If data from the trial reports were insufficient, we planned to contact the trialists for further information.

Assessment of risk of bias in included studies

We planned to make explicit judgements about whether studies were at low, high, or unclear risk of bias across the domains suggested in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We planned to assess the likely magnitude and direction of the bias and whether we considered it is likely to impact on the findings. We planned to explore the impact of the level of bias in sensitivity analyses.

Measures of treatment effect

We planned to analyse the treatment effects in the individual trials using Review Manager 5.3 (RevMan 2014) and report risk ratio (RR) and risk difference (RD) for dichotomous data and mean difference (MD) for continuous data, with respective 95% confidence intervals (CIs). We planned to determine the number needed to treat for an additional beneficial outcome (NNTB) or an additional harmful outcome (NNTH) for analyses with a statistically significant difference in the RD. We intended to conduct intention-to-treat analyses.

Unit of analysis issues

The intended unit of analysis was the participating infant in individually-randomised trials and the neonatal unit (or sub-unit) for cluster-randomised trials.

An infant would have been considered only once in an analysis. We planned to exclude infants with multiple enrolments as we would have been unable to address the associated unit of analysis issues.

For cluster-randomised trials, we planned to undertake analyses at the level of the individual while accounting for the clustering in the data using the methods recommended in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011).

Dealing with missing data

We planned to request additional data from the trialists if data on important outcomes were missing or reported unclearly. Where data were still missing, we planned to examine the impact on effect size estimates in sensitivity analyses using the 'best-worst case scenario' technique.

Assessment of heterogeneity

We planned to examine the treatment effects of individual trials and heterogeneity between trial results by inspecting the forest plots. We planned to calculate the I^2 statistic for each RR analysis to quantify inconsistency across studies and describe the percentage of variability in effect estimates that may be due to heterogeneity rather than to sampling error. If we detected moderate or high heterogeneity ($I^2 \geq 50\%$), we planned to explore the possible causes (for example, differences in study design, participants, interventions or completeness of outcome assessments).

Assessment of reporting biases

Had more than 10 trials been included in a meta-analysis, we planned to examine a funnel plot for asymmetry.

Data synthesis

We planned to use the fixed-effect model in [RevMan 2014](#) for meta-analyses (as per Cochrane Neonatal recommendations).

Where substantial heterogeneity existed, we planned to examine the potential causes in subgroup and sensitivity analyses.

Subgroup analysis and investigation of heterogeneity

We planned to undertake the following subgroup analysis:

- very preterm (< 32 weeks) infants (versus infants born > 32 weeks).

Sensitivity analysis

We planned to undertake sensitivity analyses to determine if the findings are affected by including only studies of adequate methodology (low risk of bias), defined as adequate randomisation and allocation concealment, blinding of intervention and measurement, and < 10% loss to follow-up.

'Summary of findings' table


We planned to assess the quality of evidence for the main comparisons at the primary outcomes level using the GRADE approach, as outlined in the GRADE handbook (Guyatt 2011a). Two review authors planned to independently assess the quality of the evidence for outcomes identified as critical or important for clinical decision-making (infection, death). We planned to consider evidence from randomised controlled trials as high quality but downgrade the evidence one level for serious (or two levels for very serious) limitations based upon the following: design (risk of bias), consistency across studies, directness of the evidence, precision of estimates and presence of publication bias (Appendix 2). We planned to use the GRADEpro Guideline Development Tool to create a 'Summary of findings' table to report the quality of the evidence (GRADEpro GDT).

RESULTS

Description of studies

We did not identify any studies or ongoing trials that met our inclusion criteria (Figure 1).

Figure 1. Study flow diagram.



Included studies

We did not identify any eligible trials.

Excluded studies

We excluded one RCT of different durations of placement of umbilical venous catheters in newborn infants (Butler-O'Hara 2006).

Risk of bias in included studies

We did not identify any eligible trials.

Effects of interventions

We did not identify any eligible trials.

DISCUSSION

Summary of main results

Given that deciding the duration of placement of peripherally inserted central catheters (PICCs) is a common and important clinical dilemma that may affect important outcomes for preterm or sick newborn infants, and that substantial uncertainty and variation in practice exists with regard to early planned PICC removal versus expectant management, it is concerning that this question has not yet been addressed in any randomised controlled trials.

Overall completeness and applicability of evidence

In the absence of such trial data, decisions regarding the dwell time of PICCs in infants may continue to rely on the findings of observational studies. These studies, however, have not provided consistent evidence about the effect of duration of PICC placement on the risk of invasive infection and other complications in newborn infants. Any reported associations, moreover, are likely to be confounded because infants who are smaller, less mature, sicker and receiving more intensive and invasive support are both more likely to acquire bloodstream infection and to have a PICC *in situ* (Smith 2008; Sengupta 2010; Wong 2012; Milstone 2013; Greenberg 2015). The variation in policy and practice between neonatal units and clinicians reflects the ongoing uncertainty with which the available observational data are viewed (Sharpe 2013; Taylor 2014).

Potential biases in the review process

Although we conducted a comprehensive search, including conference proceedings, we cannot exclude fully the possibility that other published (but not indexed) or unpublished trials exist.

Agreements and disagreements with other studies or reviews

There appear to be limited data to inform decisions about the optimal duration of placement of PICCs (and other central venous catheters (CVCs)) in other populations of patients. Guidelines published by the Healthcare Infection Control Practices Advisory Committee recommend that PICCs should be removed when no longer essential, but do not recommend *routine* removal and replacement to prevent catheter-related infections, and do not make any recommendations about dwell time (O'Grady 2011).

AUTHORS' CONCLUSIONS

Implications for practice

There are no randomised controlled trials of this intervention to inform policy or practice. Some, but not all, cohort studies suggest that early PICC removal is associated with a lower risk of persistent infection but these findings were not systematically reviewed and should be interpreted with caution because of biases inherent in the study design.

Implications for research

Given the potential for benefit and harm to be associated with the duration of placement of a PICC in a newborn infant, a pragmatic randomised controlled trial of early removal (with replacement if required) versus expectant management may be warranted. Such a trial might primarily address the effect on the risk of catheter-related bloodstream infections (defined using established and validated criteria) in those groups of infants with anticipated prolonged duration of PICC use (such as extremely preterm infants or infants with severe growth-restriction). Trials with sufficient power to detect reliably and precisely effects on the risk of infection and other catheter- and infection-related complications would need to be large, multi-centre and pragmatic in design.

ACKNOWLEDGEMENTS

Ms Kath Wright for developing the electronic search strategy.

REFERENCES

References to studies excluded from this review

Butler-O'Hara 2006 *{published data only}*

Butler-O'Hara M, Buzzard CJ, Reubens L, McDermott MP, DiGrazio W, D'Angio CT. A randomized trial comparing long-term and short-term use of umbilical venous catheters in premature infants with birth weights of less than 1251 grams. *Pediatrics* 2006;**118**(1):e25–35. DOI: 10.1542/peds.2005-1880; PUBMED: 16785289

Additional references

Adams-Chapman 2006

Adams-Chapman I, Stoll BJ. Neonatal infection and long-term neurodevelopmental outcome in the preterm infant. *Current Opinion in Infectious Diseases* 2006;**19**(3):290–7. DOI: 10.1097/01.qco.0000224825.57976.87; PUBMED: 16645492

Ainsworth 2015

Ainsworth SB, McGuire W. Percutaneous central venous catheters versus peripheral cannulae for delivery of parenteral nutrition in neonates. *Cochrane Database of Systematic Reviews* 2015, Issue 10. DOI: 10.1002/14651858.CD004219.pub4

Balain 2015

Balain M, Oddie SJ, McGuire W. Antimicrobial-impregnated central venous catheters for prevention of catheter-related bloodstream infection in newborn infants. *Cochrane Database of Systematic Reviews* 2015, Issue 9. DOI: 10.1002/14651858.CD011078.pub2

Bassler 2009

Bassler D, Stoll BJ, Schmidt B, Asztalos EV, Roberts RS, Robertson CM, et al. Trial of Indomethacin Prophylaxis in Preterms Investigators. Using a count of neonatal morbidities to predict poor outcome in extremely low birth weight infants: added role of neonatal infection. *Pediatrics* 2009;**123**(1):313–8. DOI: 10.1542/peds.2008-0377; PUBMED: 19117897

Benjamin 2001

Benjamin DK Jr, Miller W, Garges H, Benjamin DK, McKinney RE Jr, Cotton M, et al. Bacteremia, central catheters, and neonates: when to pull the line. *Pediatrics* 2001;**107**(6):1272–6. [PUBMED: 11389242]

Borghesi 2008

Borghesi A, Stronati M. Strategies for the prevention of hospital-acquired infections in the neonatal intensive care unit. *Journal of Hospital Infection* 2008;**68**(4):293–300. DOI: 10.1016/j.jhin.2008.01.011; PUBMED: 18329134

Cartwright 2004

Cartwright DW. Central venous lines in neonates: a study of 2186 catheters. *Archives of Disease in Childhood. Fetal*

and Neonatal Edition 2004;**89**(6):F504–8. DOI: 10.1136/adc.2004.049189; PUBMED: 15499142

Chapman 2003

Chapman RL, Faix RG. Persistent bacteremia and outcome in late onset infection among infants in a neonatal intensive care unit. *Pediatric Infectious Disease Journal* 2003;**22**(1):17–21. DOI: 10.1097/01.inf.0000042922.10767.10; PUBMED: 12544403

Chien 2002

Chien LY, Macnab Y, Aziz K, Andrews W, McMillan DD, Lee SK, Canadian Neonatal Network. Variations in central venous catheter-related infection risks among Canadian neonatal intensive care units. *Pediatric Infectious Disease Journal* 2002;**21**(6):505–11. [PUBMED: 12182373]

de Brito 2010

de Brito CS, de Brito DV, Abdallah VO, Gontijo Filho PP. Occurrence of bloodstream infection with different types of central catheter in critically neonates. *Journal of Infection* 2010;**60**(2):128–32. DOI: 10.1016/j.jinf.2009.11.007; PUBMED: 19944717

Fisher 2013

Fisher D, Cochran KM, Provost LP, Patterson J, Bristol T, Metzguer K, et al. Reducing central line-associated bloodstream infections in North Carolina NICUs. *Pediatrics* 2013;**132**(6):e1664–71. DOI: 10.1542/peds.2013-2000; PUBMED: 24249819

Garland 2008

Garland JS, Alex CP, Sevallius JM, Murphy DM, Good MJ, Volberding AM, et al. Cohort study of the pathogenesis and molecular epidemiology of catheter-related bloodstream infection in neonates with peripherally inserted central venous catheters. *Infection Control and Hospital Epidemiology* 2008;**29**(3):243–9. DOI: 10.1086/526439; PUBMED: 18220483

Gordon 2006

Gordon A, Isaacs D. Late onset neonatal Gram-negative bacillary infection in Australia and New Zealand: 1992–2002. *Pediatric Infectious Disease Journal* 2006;**25**(1):25–9. [PUBMED: 16395098]

Gordon 2016b

Gordon A, Greenhalgh M, McGuire W. Early planned removal of umbilical venous catheters to prevent infection in newborn infants. *Cochrane Database of Systematic Reviews* 2016, Issue 4. DOI: 10.1002/14651858.CD012142

GRADEpro GDT [Computer program]

GRADE Working Group, McMaster University. GRADEpro GDT. Version accessed 16 May 2016. Hamilton (ON): GRADE Working Group, McMaster University, 2014.

Greenberg 2015

Greenberg RG, Cochran KM, Smith PB, Edson BS, Schulman J, Lee HC, et al. Effect of catheter dwell time on risk of central line-associated bloodstream infection in infants. *Pediatrics* 2015;**136**(6):1080–6. DOI: 10.1542/peds.2015-0573; PUBMED: 26574587

Guyatt 2011a

Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology* 2011;**64**(4):383–94. [PUBMED: 21195583]

Guyatt 2011b

Guyatt GH, Oxman AD, Vist G, Kunz R, Brozek J, Alonso-Coello P, et al. GRADE guidelines: 4. Rating the quality of evidence—study limitations (risk of bias). *Journal of Clinical Epidemiology* 2011;**64**(4):407–15. [PUBMED: 21247734]

Guyatt 2011c

Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 7. Rating the quality of evidence—inconsistency. *Journal of Clinical Epidemiology* 2011;**64**(12):1294–302. [PUBMED: 21803546]

Guyatt 2011d

Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coello P, Rind D, et al. GRADE guidelines 6. Rating the quality of evidence—imprecision. *Journal of Clinical Epidemiology* 2011;**64**(12):1283–93. [PUBMED: 21839614]

Guyatt 2011e

Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 8. Rating the quality of evidence—indirectness. *Journal of Clinical Epidemiology* 2011;**64**(12):1303–10. [PUBMED: 21802903]

Hermans 2007

Hermans D, Talbotec C, Lacaillie F, Goulet O, Ricour C, Colomb V. Early central catheter infections may contribute to hepatic fibrosis in children receiving long-term parenteral nutrition. *Journal of Pediatric Gastroenterology and Nutrition* 2007;**44**(4):459–63. DOI: 10.1097/MPG.0b013e318031a5c7; PUBMED: 17414144

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Hoang 2008

Hoang V, Sills J, Chandler M, Busalani E, Clifton-Koepfel R, Modanlou HD. Percutaneously inserted central catheter for total parenteral nutrition in neonates: complication rates related to upper versus lower extremity insertion. *Pediatrics* 2008;**121**(5):e1152–9. DOI: 10.1542/peds.2007-1962; PUBMED: 18390957

Isaacs 2003

Isaacs D, The Australasian Study Group For Neonatal Infections. A ten year, multicentre study of coagulase negative staphylococcal infections in Australasian neonatal

units. *Archives of Disease in Childhood. Fetal and Neonatal Edition* 2003;**88**(2):F89-93. [PUBMED: 12598493]

Isaacs 2004

Isaacs D, Fraser S, Hogg G, Li HY. Staphylococcus aureus infections in Australasian neonatal nurseries. *Archives of Disease in Childhood. Fetal and Neonatal Edition* 2004;**89**(4):F331–5. DOI: 10.1136/adc.2002.009480; PUBMED: 15210669

Jain 2013

Jain A, Deshpande P, Shah P. Peripherally inserted central catheter tip position and risk of associated complications in neonates. *Journal of Perinatology* 2013;**33**(4):307–12. DOI: 10.1038/jp.2012.112; PUBMED: 22955288

Kaplan 2011

Kaplan HC, Lannon C, Walsh MC, Donovan EF, Ohio Perinatal Quality Collaborative. Ohio statewide quality-improvement collaborative to reduce late onset sepsis in preterm infants. *Pediatrics* 2011;**127**(3):427–35. DOI: 10.1542/peds.2010-2141; PUBMED: 21339274

Lahra 2009

Lahra MM, Beeby PJ, Jeffery HE. Intrauterine inflammation, neonatal sepsis, and chronic lung disease: a 13-year hospital cohort study. *Pediatrics* 2009;**123**(5):1314–9. DOI: 10.1542/peds.2008-0656; PUBMED: 19403497

Machado 2009

Machado JD, Suen VM, Figueiredo JF, Marchini JS. Biofilms, infection, and parenteral nutrition therapy. *Journal of Parenteral and Enteral Nutrition* 2009;**33**(4):397–403. DOI: 10.1177/0148607108327526; PUBMED: 19401480

Mahieu 2001a

Mahieu LM, De Dooy JJ, De Muynck AO, Van Melckebeke G, Ieven MM, Van Reempts PJ. Microbiology and risk factors for catheter exit-site and hub colonization in neonatal intensive care unit patients. *Infection Control and Hospital Epidemiology* 2001;**22**(6):357–62. DOI: 10.1086/501913; PUBMED: 11519913

Mahieu 2001b

Mahieu LM, Buitenweg N, Beutels P, De Dooy JJ. Additional hospital stay and charges due to hospital-acquired infections in a neonatal intensive care unit. *Journal of Hospital Infection* 2001;**47**(3):223–9. DOI: 10.1053/jhin.2000.0852; PUBMED: 11247683

Makhoul 2002

Makhoul IR, Sujov P, Smolkin T, Lusky A, Reichman B. Epidemiological, clinical, and microbiological characteristics of late-onset sepsis among very low birth weight infants in Israel: a national survey. *Pediatrics* 2002;**109**(1):34–9. [PUBMED: 11773539]

Miller 2010

Miller MR, Griswold M, Harris JM 2nd, Yenokyan G, Huskins WC, Moss M, et al. Decreasing PICU catheter-associated bloodstream infections: NACHRI's quality transformation efforts. *Pediatrics* 2010;**125**(2):206–13. DOI: 10.1542/peds.2009-1382; PUBMED: 20064860

Milstone 2013

Milstone AM, Reich NG, Advani S, Yuan G, Bryant K, Coffin SE, et al. Catheter dwell time and CLABSIs in neonates with PICCs: a multicenter cohort study. *Pediatrics* 2013;**132**(6):e1609–15. DOI: 10.1542/peds.2013-1645; PUBMED: 24218474

O'Grady 2002

O'Grady NP, Alexander M, Dellinger EP, Gerberding JL, Heard SO, Maki DG, et al. Healthcare Infection Control Practices Advisory Committee. Guidelines for the prevention of intravascular catheter-related infections. *Infection Control and Hospital Epidemiology* 2002;**23**(12): 759–69. DOI: 10.1086/502007; PUBMED: 12517020

O'Grady 2011

O'Grady NP, Alexander M, Burns LA, Dellinger EP, Garland J, Heard SO, et al. Healthcare Infection Control Practices Advisory Committee. Guidelines for the prevention of intravascular catheter-related infections. *American Journal of Infection Control* 2011;**39**(4 Suppl 1):S1–34. DOI: 10.1016/j.ajic.2011.01.003; PUBMED: 21511081

Ohki 2013

Ohki Y, Maruyama K, Harigaya A, Kohno M, Arakawa H. Complications of peripherally inserted central venous catheter in Japanese neonatal intensive care units. *Pediatrics International* 2013;**55**(2):185–9. DOI: 10.1111/ped.12033; PUBMED: 23253251

Olsen 2009

Olsen AL, Reinholdt J, Jensen AM, Andersen LP, Jensen ET. Nosocomial infection in a Danish Neonatal Intensive Care Unit: a prospective study. *Acta Paediatrica* 2009;**98**(8):1294–9. DOI: 10.1111/j.1651-2227.2009.01322.x; PUBMED: 19438843

Payne 2004

Payne NR, Carpenter JH, Badger GJ, Horbar JD, Rogowski J. Marginal increase in cost and excess length of stay associated with nosocomial bloodstream infections in surviving very low birth weight infants. *Pediatrics* 2004;**114**(2):348–55. [PUBMED: 15286215]

Pronovost 2006

Pronovost P, Needham D, Berenholtz S, Sinopoli D, Chu H, Cosgrove S, et al. An intervention to decrease catheter-related bloodstream infections in the ICU. *New England Journal of Medicine* 2006;**355**(26):2725–32. DOI: 10.1056/NEJMoa061115; PUBMED: 17192537

Ramirez de Arellano 1994

Ramirez de Arellano E, Pascual A, Martinez-Martinez L, Perea EJ. Activity of eight antibacterial agents on *Staphylococcus epidermidis* attached to Teflon catheters. *Journal of Medical Microbiology* 1994;**40**(1):43–7. DOI: 10.1099/00222615-40-1-43; PUBMED: 8289214

RevMan 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Saint 2000

Saint S, Veenstra DL, Lipsky BA. The clinical and economic consequences of nosocomial central venous catheter-related infection: are antimicrobial catheters useful?. *Infection Control and Hospital Epidemiology* 2000;**21**(6):375–80. DOI: 10.1086/501776; PUBMED: 10879567

Sannoh 2010

Sannoh S, Clones B, Munoz J, Montecalvo M, Parvez B. A multimodal approach to central venous catheter hub care can decrease catheter-related bloodstream infection. *American Journal of Infection Control* 2010;**38**(6):424–9. DOI: 10.1016/j.ajic.2009.07.014; PUBMED: 20137829

Schulman 2011

Schulman J, Stricof R, Stevens TP, Horgan M, Gase K, Holzman IR, et al. New York State Regional Perinatal Care Centers. Statewide NICU central-line-associated bloodstream infection rates decline after bundles and checklists. *Pediatrics* 2011;**127**(3):436–44. DOI: 10.1542/peds.2010-2873; PUBMED: 21339265

Schünemann 2013

Schünemann H, Brożek J, Guyatt G, Oxman A, editors. GRADE Working Group. GRADE Handbook for Grading Quality of Evidence and Strength of Recommendations. Available from <https://gdt.gradepro.org/app/handbook/handbook.html>. Updated October 2013.

Sengupta 2010

Sengupta A, Lehmann C, Diener-West M, Perl TM, Milstone AM. Catheter duration and risk of CLA-BSI in neonates with PICCs. *Pediatrics* 2010;**125**(4):648–53. DOI: 10.1542/peds.2009-2559; PUBMED: 20231192

Shah 2008a

Shah DK, Doyle LW, Anderson PJ, Bear M, Daley AJ, Hunt RW, et al. Adverse neurodevelopment in preterm infants with postnatal sepsis or necrotizing enterocolitis is mediated by white matter abnormalities on magnetic resonance imaging at term. *Journal of Pediatrics* 2008;**153**(2):170–5. DOI: 10.1016/j.jpeds.2008.02.033; PUBMED: 18534228

Shah 2008b

Shah PS, Shah VS. Continuous heparin infusion to prevent thrombosis and catheter occlusion in neonates with peripherally placed percutaneous central venous catheters. *Cochrane Database of Systematic Reviews* 2008, Issue 2. DOI: 10.1002/14651858.CD002772.pub3

Sharpe 2013

Sharpe E, Pettit J, Ellsbury DL. A national survey of neonatal peripherally inserted central catheter (PICC) practices. *Advances in Neonatal Care* 2013;**13**(1):55–74. DOI: 10.1097/ANC.0b013e318278b907; PUBMED: 23360860

Smith 2008

Smith PB, Benjamin DK Jr, Cotten CM, Schultz E, Guo R, Nowell L, et al. Is an increased dwell time of a peripherally inserted catheter associated with an increased risk of bloodstream infection in infants?. *Infection Control and*

- Hospital Epidemiology* 2008;**29**(8):749–53. DOI: 10.1086/589905; PUBMED: 18582196
- Stewart 2001**
Stewart PS, Costerton JW. Antibiotic resistance of bacteria in biofilms. *Lancet* 2001;**358**(9276):135–8. [PUBMED: 11463434]
- Stoll 2002**
Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, et al. Late-onset sepsis in very low birth weight neonates: the experience of the NICHD Neonatal Research Network. *Pediatrics* 2002;**110**(2 Pt 1):285–91. [PUBMED: 12165580]
- Stoll 2004**
Stoll BJ, Hansen NI, Adams-Chapman I, Fanaroff AA, Hintz SR, Vohr B, et al. Neurodevelopmental and growth impairment among extremely low-birth-weight infants with neonatal infection. *JAMA* 2004;**292**(19):2357–65. [PUBMED: 15547163]
- Taylor 2014**
Taylor JE, McDonald SJ, Tan K. A survey of central venous catheter practices in Australian and New Zealand tertiary neonatal units. *Australian Critical Care* 2014;**27**(1):36–42. DOI: 10.1016/j.aucc.2013.11.002; PUBMED: 24315154
- Taylor 2015**
Taylor JE, Tan K, Lai NM, McDonald SJ. Antibiotic lock for the prevention of catheter-related infection in neonates. *Cochrane Database of Systematic Reviews* 2015, Issue 6. DOI: 10.1002/14651858.CD010336.pub2; PUBMED: 26040840
- Thornburg 2008**
Thornburg CD, Smith PB, Smithwick ML, Cotten CM, Benjamin DK Jr. Association between thrombosis and bloodstream infection in neonates with peripherally inserted catheters. *Thrombosis Research* 2008;**122**(6):782–5. DOI: 10.1016/j.thromres.2007.10.001; PUBMED: 17997477
- van der Zwet 2005**
van der Zwet WC, Kaiser AM, van Elburg RM, Berkhof J, Fetter WP, Parlevliet GA, et al. Nosocomial infections in a Dutch neonatal intensive care unit: surveillance study with definitions for infection specifically adapted for neonates. *Journal of Hospital Infection* 2005;**61**(4):300–11. DOI: 10.1016/j.jhin.2005.03.014; PUBMED: 16221510
- Vanholder 2010**
Vanholder R, Canaud B, Fluck R, Jadoul M, Labriola L, Marti-Monros A, et al. Catheter-related blood stream infections (CRBSI): a European view. *Nephrology, Dialysis, Transplantation* 2010;**25**(6):1753–6. DOI: 10.1093/ndt/gfq205; PUBMED: 20466662
- Vasudevan 2011**
Vasudevan C, McGuire W. Early removal versus expectant management of central venous catheters in neonates with bloodstream infection. *Cochrane Database of Systematic Reviews* 2011, Issue 8. DOI: 10.1002/14651858.CD008436.pub2; PUBMED: 21833966
- Wirtschafter 2010**
Wirtschafter DD, Pettit J, Kurtin P, Dalsey M, Chance K, Morrow HW, et al. A state wide quality improvement collaborative to reduce neonatal central line-associated blood stream infections. *Journal of Perinatology* 2010; **30**(3):170–81. DOI: 10.1038/jp.2009.172; PUBMED: 19940855
- Wong 2012**
Wong J, Dow K, Shah PS, Andrews W, Lee S. Percutaneously placed central venous catheter-related sepsis in Canadian neonatal intensive care units. *American Journal of Perinatology* 2012;**29**(8):629–34. DOI: 10.1055/s-0032-1311978; PUBMED: 22566117
- Wrightson 2013**
Wrightson DD. Peripherally inserted central catheter complications in neonates with upper versus lower extremity insertion sites. *Advances in Neonatal Care* 2013;**13**(3): 198–204. DOI: 10.1097/ANC.0b013e31827e1d01; PUBMED: 23722492
- Yébenes 2004**
Yébenes JC, Vidaur L, Serra-Prat M, Sirvent JM, Batlle J, Motje M, et al. Prevention of catheter related blood stream infection in critically ill patients using a disinfectable, needle free connector: a randomised controlled trial. *American Journal of Infection Control* 2004;**32**(5):291–5. DOI: 10.1016/j.ajic.2003.12.004; PUBMED: 15292895

References to other published versions of this review

- Gordon 2016a**
Gordon A, Greenhalgh M, McGuire W. Early planned removal versus expectant management of peripherally inserted central catheters to prevent infection in newborn infants. *Cochrane Database of Systematic Reviews* 2016, Issue 4. DOI: 10.1002/14651858.CD012141
- * Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Butler-O'Hara 2006	RCT of duration of UVC (not PICC) placement

PICC: peripherally inserted central catheter

RCT: randomised controlled trial

UVC: umbilical venous catheter

CONTRIBUTIONS OF AUTHORS

All authors contributed to the development of this review.

DECLARATIONS OF INTEREST

William McGuire is a co-investigator in a UK multi-centre trial of an antimicrobial-impregnated CVC in preterm infants.

Adrienne Gordon and Mark Greenhalgh do not have any conflicts of interest.

SOURCES OF SUPPORT

Internal sources

- University of York, UK.

External sources

- National Institute for Health Research, UK.

This report is independent research funded by a UK National Institute of Health Research Grant (NIHR) Cochrane Programme Grant (16/114/03). The views expressed in this publication are those of the authors and not necessarily those of the NHS, the NIHR, or the UK Department of Health.

- Vermont Oxford Network, Other.