

This is a repository copy of *The impact of environmental change on the use of early pottery by East Asian hunter-gatherers*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/133907/>

Version: Accepted Version

---

**Article:**

Lucquin, Alexandre Jules Andre [orcid.org/0000-0003-4892-6323](https://orcid.org/0000-0003-4892-6323), Robson, Harry Kenneth [orcid.org/0000-0002-4850-692X](https://orcid.org/0000-0002-4850-692X), Eley, Yvette [orcid.org/0000-0001-5434-6830](https://orcid.org/0000-0001-5434-6830) et al. (12 more authors) (2018) The impact of environmental change on the use of early pottery by East Asian hunter-gatherers. *Proceedings of the National Academy of Sciences of the United States of America*. pp. 7931-7936. ISSN 1091-6490

<https://doi.org/10.1073/pnas.1803782115>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# The impact of environmental change on the use of early pottery by East Asian hunter-gatherers

Alexandre Lucquin<sup>1</sup>, Harry K. Robson<sup>1</sup>, Yvette Eley<sup>1</sup>, Shinya Shoda<sup>1</sup>, Dessislava Veltcheva<sup>1</sup>, Kevin Gibbs<sup>2</sup>, Carl Heron<sup>3</sup>, Sven Isaksson<sup>4</sup>, Yastami Nishida<sup>5</sup>, Yasuhiro Taniguchi<sup>6</sup>, Shota Nakajima<sup>6</sup>, Kenichi Kobayashi<sup>7</sup>, Peter Jordan<sup>8</sup>, Simon Kaner<sup>9</sup>, Oliver Craig<sup>1</sup>

<sup>1</sup>University of York, <sup>2</sup>University of California, <sup>3</sup>British Museum, <sup>4</sup>Stockholm University, <sup>5</sup>Niigata Prefectural Museum of History, <sup>6</sup>Kokugakuin University, <sup>7</sup>Chuo University, <sup>8</sup>University of Groningen, <sup>9</sup>Sainsbury Institute for the Study of Japanese Arts and Cultures

Submitted to Proceedings of the National Academy of Sciences of the United States of America

**The invention of pottery was a fundamental technological advancement with far-reaching economic and cultural consequences. Pottery containers first emerged in East Asia during the Late Pleistocene in a wide range of environmental settings but became particularly prominent and much more widely dispersed following climatic warming at the start of the Holocene. Some archaeologists argue that this increasing usage was driven by environmental factors, as warmer climates would have generated a wider range of terrestrial plant and animal resources that required processing in pottery. However, this hypothesis has never been directly tested. Here, in one of the largest studies of its kind, we conducted organic residue analysis of over 800 pottery vessels selected from 46 Late Pleistocene and Early Holocene sites located across the Japanese archipelago to identify their contents. Our results demonstrate that pottery had a strong association with the processing of aquatic resources, irrespective of the ecological setting. Contrary to expectations, this association remained stable even after the onset of Holocene warming, including in more southerly areas where expanding forests provided new opportunities for hunting and gathering. Nevertheless, the results indicate that a broader array of aquatic resources were processed in pottery after the start of the Holocene. We suggest this marks a significant change in the role of pottery of hunter-gatherers, corresponding to an increased volume of production, greater variation in forms and sizes, the rise of intensified fishing, the onset of shellfish exploitation and reduced residential mobility.**

archaeology | early pottery | organic residue analysis | stable isotopes | Jōmon

The production and use of hard, fired earthen containers represents a key technological development in human history. From its prehistoric origins at the end of the last Ice Age, pottery became a fundamental tool for transforming, mixing, storing, and serving foodstuffs almost globally, and was only replaced relatively recently by metal containers. Understanding the motivations for the emergence and wider adoption of pottery is a key question in world prehistory. Ceramic vessels were first invented by hunter-gatherers in East Asia during the Late Pleistocene in Southern China, Japan and the Russian Far East (1–3) during glacial climatic conditions (*ca.* 18,000–16,000 cal BP). With climatic warming in the Early Holocene (*ca.* 11,500 cal BP), pottery was produced in much more substantial quantities and became more widely adopted (4). Organic residue analysis of East Asian early pottery (5–7) is beginning to elucidate the motivations that lay behind early pottery innovation and its more widespread adoption. However, so far there has been no systematic investigation of pottery use across the transition from the Pleistocene to the Holocene.

One of the best areas to investigate the development of ceramic technology is the Japanese archipelago due to the intensively studied sequence of hunter-gatherer pottery, known as Jōmon (meaning cord marked). The Jōmon ceramic sequences not only offer the chance to study potential continuity or change

in pottery function across the Pleistocene-Holocene transition (See SI Appendix, Figs. S1 and S2) but also offer scope to explore this process in a wide range of ecological settings (Fig. 1) because the main Japanese islands span a large latitudinal range (30°N–46°N), which ranged from steppe-tundra in the north to warm evergreen broadleaf forest in the south (Fig. 1). The transition from the Pleistocene to Holocene is clearly apparent in changes to the composition (see SI Appendix, Fig. S1) and extent of the pottery assemblages, although changes in volumes and sizes are more difficult to assess due to their highly fragmented nature. Firstly and most noticeably there is a substantial, 100-fold increase in the number of sherds recovered on early Initial Jōmon (Stage 4) sites compared to Final Incipient sites (Stage 3) across the archipelago (4). This cannot simply be explained by a greater intensity of occupation through time. Even large Incipient sites, such as Kuzuharazawa IV in Shizuoka, have less than a thousand sherds where similarly sized Initial Jōmon sites, such as Nakano B in Hokkaido or Jozuka in Kyushu have yielded tens to hundreds of thousands with the ratio of potsherd to other artefacts also dramatically increasing (8). Secondly, clearly defined regional styles and manufacturing techniques emerge in the Early Holocene that are thought to reflect a greater integration of production and use.


The emergence of regional pottery styles and greater scale of production corresponds to transformation of the local environment in many areas. These include the expansion of broadleaf forests, particularly in Southern Japan, (Fig. 1) with increased

## Significance

**The motivations for the widespread adoption of pottery is a key theme in world prehistory and is often linked with climate warming at the start of the Holocene. Through organic residue analysis, we investigated the contents of >800 ceramic samples from across the Japanese archipelago, a unique assemblage that transcends the Pleistocene-Holocene boundary. Against our expectations we found that pottery use did not fundamentally change in the Early Holocene. Instead aquatic resources dominated in both periods regardless of the environmental setting. Nevertheless, we found that a broader range of aquatic foods were processed in Early Holocene vessels corresponding to increased ceramic production, reduced mobility, intensified fishing and the start of significant shellfish gathering at this time.**

## Reserved for Publication Footnotes

137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204




**Fig. 1.** Locations of the sampling sites, distributions of the aquatic biomarkers/phytanic acid SRR ratios (from Table 1) and change in vegetation cover (9, 10) across the Japanese archipelago from the Late Pleistocene/Incipient Jōmon (A) to Early Holocene/Initial Jōmon (B). The maps account for changes in sea level across these periods (44). Key: Dark red, complete suite of aquatic biomarkers and/or phytanic acid SRR ratio >75.5%; Red, partial suite of aquatic biomarkers and/or phytanic acid SRR ratio <75.5%; Open, absence of aquatic biomarkers and/or phytanic acid SRR ratio <75.5%; T/BF, tundra/boreal forest; CTDF, cool temperature deciduous forest; WTDF, warm temperature deciduous forest; LF, lucidophyllous forest.

**Table 1.** The frequency of aquatic derived residues associated with Incipient and Initial Jōmon pottery from Japan.

Period	# Samples (with lipid)	Full suite of aquatic biomarkers* % (n)	Partial suite of aquatic biomarkers** % (n)	Phytanic acid % (n)	>75.5% SSR-phytanic % (n)	Minimum number of aquatic vessels† % (n)
Incipient (ca. 14,460-11,310 cal BP)	179 (156)	30.8% (48)	7.1% (11)	93.6% (146)	43.6% (68)	46.8% (73)
Initial (ca. 11,500-8,000 cal BP)	622 (566)	10.8% (61)	6.9% (39)	77.0% (436)	42.0% (238)	45.2% (256)
Total	801 (722)	15.1% (109)	6.9% (50)	80.6% (582)	42.4% (306)	45.6% (329)

\* Presence of C<sub>18</sub> and C<sub>20</sub> APAAs together with one of three isoprenoid fatty acids.  
 \*\* Presence of C<sub>18</sub> APAAs and TMTD.  
 † Having either phytanic acid SSR ratio >75.5% or containing aquatic biomarkers.



**Fig. 2.** Relative treatment effects (RTE) of different geographical and temporal variables on the presence of aquatic derived lipids in Incipient and Initial pottery using a non-parametric multivariate test. RTE of treatment "k" is defined as the probability that a randomly chosen subject from treatment "k" displays a higher response than a subject that is randomly chosen from any of the treatment groups, including treatment "k". The range of possible effect is 0.27 to 0.73.


opportunities for the exploitation of terrestrial resources, such as forest game, acorns and chestnuts, (9, 10) but also greater access to marine resources through expansion of the coastal shelf (11). Increased pottery production at this time is often seen as a response to the need for processing these newly available resources, as well as intensification and increased sedentism (12) in response to the ameliorated climate and changing coastline.

There is, however, little direct evidence to support this view. The analysis of animal and plant remains tentatively show a broadening of the available resources exploited in the Holocene (13) but the data are severely constrained due to generally poor organic preservation in Japan's prevailing acidic soils (14). Some of the best palaeoeconomic data derives from coastal and lacustrine shell middens that commence during the Initial Jōmon period and while these point to a broad economic base, with terrestrial plant and animal remains well represented in addition to fish remains and shell (e.g. (15, 16), it is unknown whether pottery use also broadened at this point.


Organic residue analysis provides the only approach for directly examining the contents of pottery vessels, and in the absence of quantifiable numbers of faunal and floral remains at the majority of sites (17), it is also a valuable tool for examining palaeoeconomic change through this critical period in East Asian prehistory. Previous studies have already shown that Incipient Jōmon vessels dating to the Late Pleistocene (ca. 15,000–11,500 cal BP) were predominantly used for processing aquatic species, particularly seasonally abundant marine and anadromous fish (5, 6). Produced in low numbers compared to other artefacts (4), it has been suggested that pottery did not have a major economic function at this time and may have been prestige items associated with the collective procurement of aquatic foods during periods of sedentism by otherwise largely mobile Pleistocene hunter-gatherer groups (5, 6, 18). In contrast, the only organic residue analysis of Initial Jōmon pottery is limited to a small number of sherds from the site of Torihama in Western Japan (5). It is therefore not known whether the function of pottery fundamentally changed in the Early Holocene, as a consequence of

205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272

273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340



**Fig. 3.** Bulk and single compound stable isotope data from Late Pleistocene/Incipient Jōmon (blue) and Early Holocene/Initial Jōmon (red) ceramic vessels.  $\delta^{13}\text{C}$  values of  $\text{C}_{16:0}$  and  $\text{C}_{18:0}$  n-alkanoic acids extracted from Late Pleistocene/Incipient (A) and Early Holocene/Initial (B) Jōmon pottery, which show a broadening of aquatic resources. The 95% confidence ellipses are based on modern Japanese authentic reference fats (5, 6, 21–23). Bulk  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  stable isotope data (C) obtained from carbonised residues adhering to Incipient and Initial Jōmon vessels (some data previously reported in (5, 6, 40, 41)). (D) Box plot of the  $\delta^{15}\text{N}$  values, which also demonstrate a broadening of aquatic resources. Key: Filled circle, sample with aquatic biomarkers and/or phytanic acid SRR ratio >75.5%; Open circle, absence of aquatic biomarkers and/or phytanic acid SRR ratio <75.5%.



**Fig. 4.** Estimated percentage contributions of lipid from different food sources to (A) Late Pleistocene/Incipient and (B) Early Holocene/Initial Jōmon pottery using a concentration dependent mixing model. The model parameters are described in the SI Appendix. Box plots show the range of median % contributions estimated from each pot for each food source. The summed probability density distributions (grey) shows the relative likelihood of the contribution of each food resource summed across the two samples groups and normalised to account for differences in samples size.

the ameliorating climates, nor how responses varied across the archipelago.

To investigate further, here we present new chemical and isotopic analysis of 638 sherds and 77 charred deposits from 39 Incipient (*ca.* 14,460–11,310 cal BP) and Initial Jōmon sites (*ca.* 11,500–8,000 cal BP). The sites were chosen to examine variability over an ecological transect through Japan (see SI Appendix, Datasets S1 and S2), including inland and coastal localities (Fig. 1) at variable elevations (0–1,500 m). The majority of Incipient Jōmon sherds have cord-marked decoration corresponding to Phase 3a (*ca.* 14,460–12,000 cal BP) and 3b (*ca.* 12,030–11,310 cal BP) (see SI Appendix, Fig. S2) as defined by Taniguchi (12), with the majority corresponding to the Younger Dryas chronozone. When combined with previous data (5, 6), we have a comprehensive corpus of over 800 samples from 46 sites making this one of the largest studies of its kind. We hypothesised that en-

vironmental factors (e.g. site location, ecological zone, elevation) would largely determine the use of pottery with an increase in the processing of aquatic organisms in the cooler northern regions where terrestrial resources were less available. Further, we may expect to see a clear increase in oil rich plant products, such as nuts and seeds, and ruminant products, such as sika deer (*Cervus nippon*), and a shift away from aquatic resources in all but coastal sites at the start of the Holocene associated with climate amelioration.

### Results and Discussion

Overall, interpretable amounts of lipids were readily extractable from fragments of Jōmon pottery and adhering charred deposits using an acid/methanol extraction procedure (see Methods Summary). In total, 94% (611/652) of the potsherds and 74% (111/149) of the carbonised deposits yielded appreciable quantities of lipids i.e. that were either above the minimum amount required for interpretation (>5 $\mu\text{g g}^{-1}$  for potsherds and >100 $\mu\text{g g}^{-1}$  for charred deposits) (6, 19) or contained distinctive lipids traceable to a specific source.

#### Evidence for the processing of aquatic foods

Although the procedure deployed is suitable for identifying fats, oils and waxes from a wide range of plant and animal products (20), a distinctive feature of many of the Jōmon sherds analysed was the presence of aquatic derived lipids. In total, 15.1% (109/722; Table 1) of the samples analysed satisfy the established criteria for the presence of ‘aquatic biomarkers’ in pottery (5, 20), which includes the presence of  $\omega$ -(*o*-alkylphenyl) alkanolic acids (APAAs) with  $\text{C}_{18}$  and  $\text{C}_{20}$  carbon atoms and isoprenoid fatty acids (either phytanic, pristanic or 4,8,12-trimethyl tridecanoic acid (TMTD)). Notably the  $\text{C}_{20}$  APAAs are formed during the protracted heating of the  $\text{C}_{20:\text{x}}$  mono- and polyunsaturated fatty acids which are only found in appreciable concentrations in freshwater and marine animals (21, 22). The presence of APAAs implies that the pottery vessels were subjected to prolonged heating (typically >270 $^{\circ}\text{C}$ , >17h; (21, 22)), easily achieved through boiling or roasting of their contents, which is consistent with the presence of charred ‘foodcrusts’ on many vessels. Multi-branched isoprenoid fatty acids originate from the breakdown of phytol, a constituent of chlorophyll, but only accumulate at high concentrations in ruminant and aquatic animal tissues. In particular, TMTD is considered more of a characteristic of aquatic oils (23).

341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408

409 These 'aquatic biomarker' estimates should be considered  
410 as a minimum percentage, since APAAs are not always formed  
411 during food preparation and both APAAs and isoprenoids may  
412 be lost in the burial environment relative to other lipid molecules  
413 with higher relative concentrations. A further 6.9% (50/722) have  
414 C<sub>18</sub> APAAs and TMTD which are most likely aquatic in origin (i.e.  
415 'partial aquatic biomarkers'; Table 1), while the majority of samples  
416 (81%; 582/722) contained phytanic acid, the most frequent  
417 isoprenoid acid. Among the resources available to Japanese Pleis-  
418 tocene and Holocene hunter-gatherers, wild ruminants such as  
419 sika deer offer the only other major source of phytanic acid other  
420 than aquatic oils (7, 24). To distinguish these, we examined the  
421 ratio of the two naturally occurring configurations, or diastere-  
422 omers, of phytanic acid (3S,7R,11R,15-phytanic acid ~ SRR, and  
423 3R,7R,11R,15-phytanic acid ~ RRR (see Methods Summary)).  
424 Despite considerable overlap, the SRR isomer tends to dominate  
425 in aquatic oils compared to ruminant fats (7, 24) and a SRR%  
426 above 75.5% can be assigned to this source, using a conservative  
427 limit (95% confidence). Over 52% (306/582) of the samples with  
428 phytanic acid met this criteria, for the remainder the source of  
429 phytanic acid is uncertain as it fell within both the aquatic and  
430 ruminant range.

431 Using the SRR ratio and presence of aquatic biomarkers, we  
432 conservatively assigned a minimum number of vessels analysed that  
433 were used to process aquatic foods across the Pleistocene-  
434 Holocene transition (Table 1). Overall, there was a striking  
435 consistency in the use of pottery throughout the Japanese archipelago  
436 regardless of period or environmental setting (Fig. 1). A non-  
437 parametric multivariate inference test (25) showed significant  
438 ( $p < 0.001$ ) effects of period, latitude, longitude, elevation, distance  
439 from the coast, precipitation, temperature and vegetation  
440 cover on the frequency of aquatic resources in the vessels (Fig.  
441 2; see SI Appendix). However, when the relative effects were  
442 quantified (Fig. 2) the site's distance from the coast and its  
443 elevation had the greatest effect but even this effect was not  
444 strong (i.e. the relative effect value does not approach the minimum  
445 or maximum effect). The other environmental variables  
446 and the period classification of the vessels had no or very weak  
447 effects (0.43-0.54) on the presence/absence of aquatic derived  
448 lipids. Pots were used to process aquatic resources at an equally  
449 high frequency throughout the archipelago from Hokkaido and  
450 Northern Honshu to Kyushu.

451 There was only a slight decrease in pottery used for process-  
452 ing aquatic resources between the Incipient (47%, 73/156) and  
453 Initial (45%, 256/566; Table 1) Jōmon. These results refute the  
454 expectation of a dramatic change in the function of pottery at start  
455 of the Holocene when terrestrial resources were more available,  
456 even accounting for potential biases in site location between  
457 periods (Fig. 2). These data corroborate what we have suggested  
458 previously (5, 20) that pottery production for the exploitation of  
459 aquatic resources was embedded as a cultural norm in the social  
460 memories of these foragers.

461 Pottery from sites more distant from the palaeocoastline  
462 tended to have fewer aquatic derived lipids but the effect was  
463 marginal and co-varied with site elevation. Indeed, aquatic prod-  
464 ucts were frequently identified in ceramics from inland riverine  
465 and lacustrine sites (Fig. 1) most likely pointing to the exploita-  
466 tion of freshwater resources and/or migratory species, such as  
467 salmonids. To distinguish the source of these residues further we  
468 examined the carbon isotope ( $\delta^{13}\text{C}$ ) values of the major saturated  
469 fatty acids (C<sub>16:0</sub> and C<sub>18:0</sub>) extracted from the sherds, and bulk  
470 carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) stable isotope values of any  
471 adhering carbonised residues (see SI Appendix, Dataset S2).  
472 In Fig. 3, the fatty acid data are compared with  $\delta^{13}\text{C}$  values  
473 obtained from modern authentic Japanese plants and animals  
474 (5, 6, 26–28). These generally support the lipid biomarker data  
475 with many vessels plotting in the reference ranges for aquatic

477 resources. Interestingly, there was only a weak negative corre-  
478 lation between distance from the coast and the  $\delta^{13}\text{C}_{16:0}$  value  
479 (Spearman  $\rho(560) = -0.25$   $p < 0.001$ ) and no correlation with the  
480 bulk  $\delta^{13}\text{C}$  value (Spearman  $\rho(190) = 0.03$   $p = 0.6667$ ) as may have  
481 been expected if marine resources were preferentially processed  
482 at coastal sites compare to inland localities. This may be explained  
483 by the exploitation of migratory fish, such as salmonids, which  
484 have  $\delta^{13}\text{C}$  values that approach the marine range (Fig. 3).

485 Fully marine species, beyond the isotopic range of reference  
486 salmonids (Fig. 3; see SI Appendix, Dataset S3) were identified in  
487 pottery from sites located >15 km from the prehistoric coastline,  
488 the maximum logistical walking distance for a logistical day trip  
489 (29), which suggests that aquatic resources were not only acquired  
490 locally for direct consumption but could also have been preserved  
491 (e.g. dried) and transported. These include an Incipient vessel  
492 from Taisho 3 and 13 Initial vessels from Haizuka, Higashimyou,  
493 Nishinojo, Nisshin 3 and Taisho 3. Although site elevation has  
494 the greatest effect on the presence/absence of aquatic derived  
495 lipids, even pottery from remote mountainous areas were also  
496 used to process aquatic foods. At Yukura Cave (at elevation  
497 of ca. 1534 m) almost half the vessels had residues typical of  
498 salmonids with the nearest source, the Shinano river (30), located  
499 ca. 15 km away. Conversely at these remote hunting sites and  
500 more broadly in Japan's warmer forested areas, a surprisingly low  
501 number of residues could be attributed to forest game species,  
502 such as sika deer and wild boar (*Sus scrofa*) (Fig. 3), implying they  
503 were processed in other ways.

504 There is also surprisingly limited data to suggest that plant  
505 foods were processed in Incipient or Initial Jōmon pottery across  
506 Japan. Low to trace amounts of leafy plant-derived lipids, includ-  
507 ing phytosterol, long-chain even-numbered fatty acids or long-  
508 chain odd-numbered alkanes, were present in some samples,  
509 most notably at Kenshojo in Southern Kyushu (see SI Appendix,  
510 Dataset S2). Isotopic analysis of the foodcrusts adhering to the  
511 potsherds also generally had lower (<22) C:N atomic ratios  
512 (mean =  $12.0 \pm 5.1$ ) more typical of carbonised terrestrial animal  
513 and marine tissues than plant remains (27) (See SI Appendix,  
514 Dataset S2). Plant resources, particularly acorns and chestnuts,  
515 and artefacts associated with plant processing are frequently  
516 found on Incipient and Initial sites (9, 31–33) suggesting they were  
517 an important feature of the Jōmon economy. While the organic  
518 residue evidence cannot rule out the presence of plants in pottery  
519 entirely, the data clearly show that Incipient and Initial Jōmon  
520 vessels were not extensively used for this purpose. Rather we  
521 contend that Incipient and Initial Jōmon hunter-gatherers had  
522 a clear preference for preparing aquatic resources over terres-  
523 trial animal and plant products in pottery. Moreover, we assert  
524 that this cooking practice was pervasive over a wide range of  
525 environmental settings and persistent through time and through  
526 significant climate change.

#### 527 *Holocene pottery used for processing of a wider array of aquatic* 528 *resources*

529 Although there is strong evidence that aquatic resources were  
530 exploited in both periods we found evidence across the Japanese  
531 archipelago of diversification in the types of aquatic foods pro-  
532 cessed in the pottery at the start of the Holocene. A much  
533 narrower range of  $\delta^{13}\text{C}_{16:0}$  and  $\delta^{13}\text{C}_{18:0}$  values were obtained from  
534 Late Pleistocene (Incipient Jōmon) pottery when compared to  
535 those from the Early Holocene (Initial Jōmon) (Figs. 2A and  
536 2B). During the Incipient Jōmon,  $\delta^{13}\text{C}_{16:0}$  and  $\delta^{13}\text{C}_{18:0}$  values  
537 are relatively homogenous i.e. have low variances ( $\sigma^2 = 3.5$ ,  $n =$   
538 119, mean =  $-26.3\%$ ). Regardless of the geographic setting, the  
539 majority of values fall within the ranges established from the  
540 analysis of modern marine organisms and salmonids (Fig. 3A),  
541 which was corroborated by the presence of aquatic biomarkers  
542 in many of the samples. These data support the general model  
543 proposed previously (6, 34, 35) that the earliest phases of pottery  
544

use are highly specialised and focused on seasonally available aquatic resources.

In contrast, the variance of  $\delta^{13}\text{C}_{16:0}$  values significantly increased (Brown–Forsythe test  $F(1,558)=10.42$   $p<0.005$ ) during the Initial Jōmon ( $\sigma^2=6.9$ ,  $n=441$ , mean =  $-27.0\text{‰}$  for  $\text{C}_{16:0}$ ; Fig. 3B). This most likely reflects a broadening of the aquatic foods processed to encompass a greater range of both marine and freshwater species (Fig. 3B). The high frequency of the other aquatic derived lipids on Initial Jōmon sherds supports this contention but mixing with terrestrial animal and even plant resources, also relatively depleted in  $^{13}\text{C}$ , cannot be ruled out entirely. In order to investigate the effects of mixing different resources in the vessels we applied a concentration dependent Bayesian mixing model (36) that used the  $\delta^{13}\text{C}_{16:0}$ ,  $\delta^{13}\text{C}_{18:0}$  and SRR% values as proxies, with priors based on the presence of isoprenoid and APAAs (see SI Appendix). This model was used to examine the likely probability of different proportions of lipids derived from plants (acorns and chestnuts), freshwater organisms (fish), wild boar, wild ruminants (sika deer) and marine organisms/salmonids to each pot. By summing the probabilities for each period and examining their densities (Fig. 4), only aquatic organisms can be reliably considered to have made a substantial contribution (i.e.  $>25\%$  of total lipid) in either periods (Fig. 4) based on the assumptions used in the model. Noticeably, however, the percentage contribution of lipid from freshwater organisms is predicted to increase from the Incipient to Initial Jōmon (Fig. 4) consistent with a broadening of pottery use at this time.

Surprisingly few vessels contained substantial amounts of non-aquatic products. Ruminant, wild boar and acorn/chestnut were estimated by the model to have made a contribution of  $>25\%$  in 21, 1 and 7 samples respectively. It should be noted that their contribution to the remaining vessels cannot be ruled out entirely; between 0 to 25% lipid contributions from non-aquatic sources were most likely, although depending on their lipid content, these could have had a greater relative contribution by total weight. Further distinction is not possible using the isotope approach and SRR% alone. Even where prior information from the biomarker evidence is deployed there is a high degree of equifinality regarding the source contributions.

A broadening of the range of aquatic resources processed in pottery during the Holocene and across the Japanese archipelago can also be seen from the bulk nitrogen ( $\delta^{15}\text{N}$ ) stable isotope values of carbonised residues adhering to pottery (Fig. 3C and 3D). Nitrogen stable isotope values of charred deposits are often used to crudely distinguish between high trophic level aquatic resources and lower trophic level terrestrial organisms (37), although  $^{15}\text{N}$  enrichment due to charring also needs to be accounted for (38, 39). In total,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values were obtained on 157 samples from 21 sites (Fig. 3C), which were complemented with previously published data undertaken as part of AMS radiocarbon ( $^{14}\text{C}$ ) dating programs (40, 41). As with  $\delta^{13}\text{C}_{16:0}$ , a broader range of nitrogen isotope values were obtained from the Initial Jōmon pottery (Incipient variance,  $\sigma^2=5.6$ ,  $n=119$ ; Initial variance,  $\sigma^2=10.9$ ,  $n=71$ ; Brown–Forsythe test  $F(1,188)=13.49$   $p<0.005$ ). A decrease in  $\delta^{15}\text{N}$  values (Fig. 3D) was also observed between the Incipient (median =  $11.5\text{‰}$ ) and Initial Jōmon (median =  $8.8\text{‰}$ ) and overall the distributions of the  $\delta^{15}\text{N}$  values were significantly different (Mann–Whitney  $U$  test;  $U=6024$ ;  $p<0.005$ ). In contrast, the range of  $\delta^{13}\text{C}$  values is similar ( $U=4128$ ;  $p=0.79$ ) between the Incipient ( $-27$  to  $-20\text{‰}$ , median =  $-24\text{‰}$ ) and Initial Jōmon ( $-26$  to  $-19\text{‰}$ , median =  $-24\text{‰}$ ).

Interestingly, aquatic biomarkers were frequently observed in charred deposits with lower  $\delta^{15}\text{N}$  values (Fig. 3C) ruling out predominantly terrestrial input. Although outside the range of

marine finfish and marine mammals ( $>12\text{‰}$ , 17, 40), these  $\delta^{15}\text{N}$  values are within the range of values obtained on lower trophic level freshwater fish and marine/freshwater shellfish (17, 40), accounting for a  $1\text{‰}$  increase with charring (38, 40). Therefore, a more likely explanation is that the observed broadening and decrease in  $\delta^{15}\text{N}$  values of charred deposits observed in the Holocene (Fig. 3D) is due to a diversification of aquatic resource exploitation to encompass freshwater fishing and/or shellfish collection. This explanation is also consistent with the establishment of shell middens in Japan at this time (16) but currently we are unable to unequivocally distinguish shellfish derived residues with the methods at our disposal.

### Conclusion

There is a dramatic increase in the scale of pottery use across Japan after the onset of Early Holocene warming. We have investigated the extent to which these environmental changes drove diversification in pottery function as a broader range of resources became readily available. The earliest pottery in Japan was used to process aquatic resources, but contrary to expectations, we found no evidence that its function expanded in the Early Holocene to include the processing of terrestrial animal and plant resources. Instead, our results show remarkable continuity and consistency in the function of pottery across the Pleistocene–Holocene transition, pointing towards a strong cultural association between pottery and the processing of aquatic resources. This pattern also holds throughout the different ecological zones of the Japanese archipelago. As a result, we suggest that after its first invention, pottery developed particular cultural associations linked to processing aquatic resources, and that these were robust enough to withstand the effects of major climatic and environmental transformations at the Pleistocene–Holocene transition. Moreover, these ‘culinary’ preferences persisted across Japan, even in warmer southern areas where abundant nut and plant resources were increasingly available. Our earlier study from the Torihama shell midden site (5–7) in Japan indicates that this cognitive association persisted until at least the Middle Holocene, and may only have been truncated by the arrival of rice and millet agriculture ca. 2,500 cal BP. A similar association between early pottery and aquatic resources has also been identified in adjacent regions of East Asia such as Sakhalin Island (6, 34, 35) and the Korean Peninsula (5–7). Here, pottery appears in the Early and Middle Holocene and from the outset demonstrates close association with processing of marine foods.

Our current research also identified an important new pattern, which is that pottery was used to process a broader spectrum of aquatic foods in the Early Holocene, including shellfish, freshwater fish and a greater range of marine taxa. This corresponds to significant climate warming ca. 11,500 years ago which may have reduced salmonid stocks in Northern Japan (42, 43) prompting a switch to other aquatic species, but also created greater opportunities for inshore fishing and shellfish gathering through the expansion of the marine shelf (11). Also at this time, pottery traditions began to flourish, with greater variation in forms and volumes reflecting intensified usage. We suggest that this represents an important change away from the small-scale and specialised use of pottery in the Late Pleistocene to a greater utilitarian function in the Early Holocene as fishing and shellfish gathering intensified. Whether this change served as a driving force for the wider-range dispersal of hunter-gatherer pottery from East Asia into surrounding areas along aquatic ecotones (2), needs to be tested through further organic residue analysis and greater AMS radiocarbon dating of early pottery sites.

Finally, we are unable to explain either the invention of pottery in the Late Pleistocene or its more varied and intensified use in the Holocene in purely functional terms. Indeed, aquatic foods were undoubtedly exploited by maritime East Asian hunter-gatherers well before pottery appeared (45). Social and demo-

681 graphic factors, indirectly linked to economic change, provide  
682 a more compelling argument. We suggest that pottery was initially  
683 developed as a novel, prestige technology during periods  
684 of seasonal population aggregation focused on cooperative har-  
685 vesting of migratory fish, such as salmonids. From the start of  
686 the Holocene, however, it was produced in significantly larger  
687 quantities, associated with intensification of aquatic resource  
688 exploitation and increasing sedentism.

#### 689 Methods summary

690 We obtained 652 ceramic sherds and 172 adhering carbonised  
691 residues from 46 archaeological sites throughout the Japanese  
692 archipelago. Assignment to the Incipient or Initial Jōmon was  
693 based on pottery typology or independently through the AMS  
694 radiocarbon ( $^{14}\text{C}$ ) dating of associated organic materials.

695 **Organic residue analysis:** Lipids were directly extracted and  
696 methylated with acidified methanol according to established  
697 methods (6, 7). Briefly, methanol (1 or 4 mL) was added to  
698 homogenised carbonised residues (10–20 mg) or ceramic powders  
699 (0.5–1.0 g) drilled (2–5 mm depth) from the interior or exterior  
700 surface of the sherd. The sample was sonicated in a water bath for

701 15 min, and acidified with concentrated sulphuric acid (200 or 800  
702  $\mu\text{L}$ ). The acidified suspension was heated in a block for 4 h at 70  
703  $^{\circ}\text{C}$ . Lipids were extracted *n*-hexane (3  $\times$  2 ml), and subsequently  
704 analysed by GC-MS and GC-c-IRMS (see SI Appendix). Interior  
705 foodcrusts or exterior carbonised residues were also analysed  
706 by Elemental Analysis-Isotope Ratio Mass Spectrometry (see SI  
707 Appendix) using previously reported protocols (5, 6).

708 **Statistical and GIS:** All statistical tests were performed using  
709 *R studio* (version 1.0.136) and *Past* (version 3.18). Mapping was  
710 undertaken with QGIS (version 2.18.9).

#### 711 Acknowledgments

712 This work was supported by the Arts and Humanities Research Council  
713 (The Innovation and Development of Pottery in East Asia project, No.  
714 AH/L00691X/1), a Marie Curie International Incoming Fellowship (IIF-624467)  
715 to SS and a Wenner-Gren Post PhD Research Grant (8565) to KG. HKR  
716 acknowledges the support of the British Academy. We should like to thank  
717 Matthew Von Tersch (University of York) and Marise Gorton (University of  
718 Bradford) for undertaking the bulk stable isotope analyses. We are grateful  
719 to Yuichiro Kudo (National Museum of Japanese History) for assistance SI  
720 Appendix, Fig. S2, and to the three anonymous referees for their helpful  
721 comments that improved the text.

- 722 1. Kuzmin YV (2017) The origins of pottery in East Asia and neighboring regions: An analysis  
723 based on radiocarbon data. *Quat Int* 441:29–35.
- 724 2. Jordan P, Gibbs K, Hommel P, Piezonka H, Silva F (2016) Modelling the diffusion of pottery  
725 technologies across Afro-Eurasia: emerging insights and future research. *Antiquity* 90(351):  
726 590–603.
- 727 3. Yanshina OV (2017) The earliest pottery of the eastern part of Asia: Similarities and  
728 differences. *Quat Int* 441:69–80.
- 729 4. Taniguchi Y (2017) The Beginning of Pottery Technology in Japan: The Dating and Function  
730 of Incipient Jomon Pottery. *The Emergence of Pottery in West Asia*, eds Tsuneki A, Nieuwen-  
731 huyse O, Campbell S (Oxbow Books, Oxford).
- 732 5. Lucquin A, et al. (2016) Ancient lipids document continuity in the use of early  
733 hunter-gatherer pottery through 9,000 years of Japanese prehistory. *Proc Natl Acad Sci USA*  
734 113(15):3991–3996.
- 735 6. Craig OE, et al. (2013) Earliest evidence for the use of pottery. *Nature* 496(7445):351–354.
- 736 7. Shoda S, Lucquin A, Ahn J-H, Hwang C-J, Craig OE (2017) Pottery use by early Holocene  
737 hunter-gatherers of the Korean peninsula closely linked with the exploitation of marine  
738 resources. *Quat Sci Rev* 170:164–173.
- 739 8. Aira Town Educational Board (2005) Excavation Report: The Kenshojo Site (in Japanese).
- 740 9. Sakaguchi T (2009) Storage adaptations among hunter-gatherers: A quantitative approach  
741 to the Jomon period. *J Anthropol Archaeol* 28(3):290–303.
- 742 10. Melvin Aikens C, Akazawa T (1996) The Pleistocene–Holocene Transition in Japan and  
743 Adjacent Northeast Asia. *Humans at the End of the Ice Age, Interdisciplinary Contributions to*  
744 *Archaeology*, eds Straus LG, Eriksen BV, Erlandson JM, Yesner DR (Springer, New York,  
745 NY).
- 746 11. Sato H, Izuho M, Morisaki K (2011) Human cultures and environmental changes in the  
747 Pleistocene–Holocene transition in the Japanese Archipelago. *Quat Int* 237(1):93–102.
- 748 12. Kaner S, Taniguchi Y (2017) The Development of Pottery and Associated Technological  
749 Developments in Japan, Korea, and the Russian Far East. *Handbook of East and Southeast*  
750 *Asian Archaeology*, eds Habu J, Lape P, Olsen J (Springer, New York, NY).
- 751 13. Matsui A, Kanehara M (2006) The question of prehistoric plant husbandry during the Jomon  
752 period in Japan. *World Archaeol* 38(2):2590273.
- 753 14. Matsui A (2014) Shuryou no Taisho (Hunting game). *Jomon Jidai (Ge) (Jomon Period (Second*  
754 *Volume), The Archaeology of Japan Lecture* (Aoki Shoten, Tokyo), pp 3–35.
- 755 15. Saga City Educational Board (2016) The Higashimyou site cluster synthesis report (in  
756 Japanese).
- 757 16. Habu J, Matsui A, Yamamoto N, Kanno T (2011) Shell midden archaeology in Japan: Aquatic  
758 food acquisition and long-term change in the Jomon culture. *Quat Int* 239(1–2):19–27.
- 759 17. Yoneda M, et al. (2004) Isotopic evidence of inland-water fishing by a Jomon population  
760 excavated from the Boji site, Nagano, Japan. *J Archaeol Sci* 31(1):97–107.
- 761 18. Hayden B (1995) The emergence of prestige technologies and pottery. *The Emergence*  
762 *of Pottery: Technology and Innovation in Ancient Societies* (Smithsonian Institution Press,  
763 Washington, DC), pp 257–266.
- 764 19. Evershed RP (2008) Experimental approaches to the interpretation of absorbed organic  
765 residues in archaeological ceramics. *World Archaeol* 40(1):26–47.
- 766 20. Evershed RP (2008) Organic residue analysis in archaeology: the archaeological biomarker  
767 revolution. *Archaeometry* 50(6):895–924.
- 768 21. Hansel FA, Copley MS, Madureira LAS, Evershed RP (2004) Thermally produced  $\omega$ -  
769 (o-alkylphenyl) alkanolic acids provide evidence for the processing of marine products in  
770 archaeological pottery vessels. *Tetrahedron Lett* 45(14):2999–3002.
- 771 22. Evershed RP, Copley MS, Dickson L, Hansel FA (2008) Experimental evidence for the  
772 processing of marine animal products and other commodities containing polyunsaturated  
773 fatty acids in pottery vessels. *Archaeometry* 50(1):101–113.
- 774 23. Ackman RG, Hooper SN (1968) Examination of isoprenoid fatty acids as distinguishing  
775 characteristics of specific marine oils with particular reference to whale oils. *Comp Biochem*  
776 *Physiol* 24:549–565.
- 777 24. Lucquin A, Colonese AC, Farrell FFG, Craig OE (2016) Utilising phytanic acid diastere-  
778 omers for the characterisation of archaeological lipid residues in pottery samples. *Tetrahedron*  
779 *Lett* 57(6):703–707.
- 780 25. Burchett W, Ellis A, Harrar S, Bathke A (2017) Nonparametric Inference for Multivariate  
781 Data: The R Package nprmv. *J Stat Softw* 76(4):1–18.
- 782 26. Horiuchi A, Miyata Y, Kamijo N, Cramp L, Evershed RP (2015) A Dietary Study of the  
783 Kamegaoka Culture Population during the Final Jomon Period, Japan, Using Stable Isotope  
784 and Lipid Analyses of Ceramic Residues. *Radiocarbon* 57(4):721–736.
- 785 27. Heron C, et al. (2016) Molecular and isotopic investigations of pottery and “charred remains”  
786 from Sannai Maruyama and Sannai Maruyama No. 9, Aomori Prefecture. *Japanese Journal*  
787 *of Archaeology* 4(1):29–52.
- 788 28. Miyata Y, Horiuchi A, Takada H, Nakamura T (2015) Evaluation of sea mammals as marine  
789 resource by lipid analysis in pottery excavated from Mawaki archaeological site, Ishikawa,  
790 Japan. *The 32nd Annual Meeting of Japan Society for Scientific Studies on Cultural Property*  
791 *Abstracts*: 40–41.
- 792 29. Kelly RL (2013) *The Lifeways of Hunter-Gatherers: The Foraging Spectrum* (Cambridge  
793 University Press, Cambridge).
- 794 30. Kaner S (2009) Long-term innovation: The appearance and spread of pottery in the Japanese  
795 Archipelago. *Ceramics Before Farming: The Dispersal of Pottery Among Prehistoric Eurasian*  
796 *Hunter-Gatherers*, eds Jordan P, Zvelebil M (Left Coast Press, Walnut Creek, CA).
- 797 31. Crawford GW (2011) Advances in Understanding Early Agriculture in Japan. *Curr Anthropol*  
798 52(S4):S331–S345.
- 799 32. Noshiro S, Sasaki Y (2014) Pre-agricultural management of plant resources during the  
800 Jomon period in Japan—a sophisticated subsistence system on plant resources. *J Archaeol*  
801 *Sci* 42:93–106.
- 802 33. Shitubani A (2009) Late Pleistocene to Early Holocene Plant Movements in Southern  
803 Kyushu, Japan. *Arch* 5(1):124–133.
- 804 34. Taché K, Craig OE (2015) Cooperative harvesting of aquatic resources and the beginning of  
805 pottery production in north-eastern North America. *Antiquity* 89(343):177–190.
- 806 35. Gibbs K, et al. (2017) Exploring the emergence of an “Aquatic” Neolithic in the Russian Far  
807 East: organic residue analysis of early hunter-gatherer pottery from Sakhalin Island. *Antiquity*  
808 91(360):1484–1500.
- 809 36. Fernandes R, et al. (2018) Reconstruction of prehistoric pottery use from fatty acid carbon  
810 isotope signatures using Bayesian inference. *Org Geochem* 117:31–42.
- 811 37. Heron C, Craig OE (2015) Aquatic Resources in Foodcrusts: Identification and Implication.  
812 *Radiocarbon* 57(4): 707–719.
- 813 38. Nitsch EK, Charles M, Bogaard A (2015) Calculating a statistically robust  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$   
814 offset for charred cereal and pulse seeds. *STAR: Science & Technology of Archaeological*  
815 *Research* 1(1):1–8.
- 816 39. Fraser RA, et al. (2013) Assessing natural variation and the effects of charring, burial and  
817 pre-treatment on the stable carbon and nitrogen isotope values of archaeobotanical cereals  
818 and pulses. *J Archaeol Sci* 40(12):4754–4766.
- 819 40. Yoshida K, et al. (2013) Dating and stable isotope analysis of charred residues on the Incipient  
820 Jomon pottery (Japan). *Radiocarbon* 55(2–3):1322–1333.
- 821 41. Kunikita D, et al. (2013) Dating Charred Remains on Pottery and Analyzing Food Habits in  
822 the Early Neolithic Period in Northeast Asia. *Radiocarbon* 55(2–3):1334–1340.
- 823 42. Ishida Y, et al. (2001) Archeological evidence of Pacific salmon distribution in northern Japan  
824 and implications for future global warming. *Prog Oceanogr* 49(1):539–550.
- 825 43. Ishida Y, Yamada A, Nagasawa K (2016) Future Climate-Related Changes in Fish Species  
826 Composition Including Chum Salmon (*Oncorhynchus keta*) in Northern Japanese Waters,  
827 Inferred from Archaeological Evidence. *North Pacific Anadromous Fish Commission Bulletin*  
828 6: 243–258.
- 829 44. Lambeck K, Rouby H, Purcell A, Sun Y, Sambridge M (2014) Sea level and global  
830 ice volumes from the Last Glacial Maximum to the Holocene. *Proc Natl Acad Sci USA*  
831 111(43):15296–15303.
- 832 45. O’Connor S, Ono R, Clarkson C (2011) Pelagic fishing at 42,000 years before the present and  
833 the maritime skills of modern humans. *Science* 334(6059):1117–1121.