

This is a repository copy of *Surfaces away from horizons are not thermodynamic*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/133973/>

Version: Accepted Version

Article:

Wang, Zhi-Wei and Braunstein, Samuel Leon orcid.org/0000-0003-4790-136X (2018)
Surfaces away from horizons are not thermodynamic. *Nature Communications*. 2977.
ISSN 2041-1723

<https://doi.org/10.1038/s41467-018-05433-9>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Surfaces away from horizons are not thermodynamic

Zhi-Wei Wang^{1,2} and Samuel L. Braunstein²

¹*College of Physics, Jilin University, Changchun, 130012, People's Republic of China and*

²*Computer Science, University of York, York YO10 5GH, United Kingdom **

Since the 1970's it has been known that black hole (and other) horizons are truly thermodynamic. More generally, surfaces which are not horizons have also been conjectured to behave thermodynamically. Initially, for surfaces microscopically expanded from a horizon to so-called stretched horizons, and more recently, for more general ordinary surfaces in the emergent gravity program. To test these conjectures we ask whether such surfaces satisfy an analogue to the first law of thermodynamics (as do horizons). For static asymptotically-flat spacetimes we find that such a first law holds on horizons. We prove that this law remains an excellent approximation for stretched horizons, but counter-intuitively this result illustrates the insufficiency of the laws of black hole mechanics alone from implying truly thermodynamic behavior. For surfaces away from horizons in the emergent gravity program the first law fails (except for spherically-symmetric scenarios) thus undermining the key thermodynamic assumption of this program.

In 1973, Bardeen, Carter, and Hawking derived the laws of black hole mechanics which are in direct analogy with the laws of thermodynamics¹. Together with the discovery of Hawking radiation², the truly thermodynamic behavior of black hole horizons became well established. Indeed such thermodynamic behavior is now well accepted for all spacetime horizons, including those due to accelerated observers^{3,4} and cosmological horizons⁵.

Later, other surfaces were also attributed with thermodynamic properties. Firstly, stretched horizons were claimed to be thermodynamic, effectively acting as radiating black bodies⁶ with a temperature $T = \kappa/(2\pi)$ determined by their local surface gravity κ and an entropy (a 'state variable') associated with a statistical mechanical interpretation of black hole entropy^{6,7}. An explicit re-derivation of the laws of black hole mechanics has not been previously carried out for stretched horizons. More recently, a class of ordinary surfaces has been conjectured to behave thermodynamically, forming the key assumption in the emergent gravity program⁸. This thermodynamic attribution was justified in part by using it in a heuristic derivation of the full Einstein field equations in static asymptotically-flat spacetime⁸.

Here we ask whether canonical General Relativity is consistent with the assumption that such ordinary surfaces can be rigorously seen to behave thermodynamically. We attack this question by focusing on the analogue to the first law of thermodynamics. Originally this law was derived in an analysis that was specialized to the behavior of horizons¹. We remove this specialization to reveal the behavior of ordinary surfaces in an analysis of the first law. Here we report that the first law holds to an excellent approximation for stretched horizons. Finally, with the exception of fully spherically-symmetric scenarios, we find that for ordinary surfaces in the emergent gravity program that the first law fails to hold.

Results

Energy conservation. For a static asymptotically-

flat spacetime with timelike Killing vector K^μ one may derive the total gravitating mass M as an integral over a spacelike hypersurface Σ that is truncated (or bounded) internally by an ordinary 2-surface $\partial\Sigma_{\text{in}}$ (see Fig. 1)

$$M = \frac{1}{4\pi} \int_{\Sigma} R_{\mu\nu} K^\mu \hat{T}^\nu \sqrt{|\gamma^{(\Sigma)}|} d^3x + \frac{1}{4\pi} \int_{\partial\Sigma_{\text{in}}} \kappa dA. \quad (1)$$

(See the Supplementary Methods for a detailed derivation and definition of each term.) This expression is a straightforward extension of that used in 1973 by Bardeen et al.¹ in their derivation of the first law of thermodynamics for black holes, though there the internal boundary was a horizon. Here κ is a natural extension of the surface gravity for non-rotating spacetimes.




FIG. 1: Schematic of the spacelike three-dimensional hypersurface of interest, Σ , with an inner boundary $\partial\Sigma_{\text{in}}$ and a boundary at infinity $\partial\Sigma_{\infty}$. Here \hat{N}^μ is the spacelike 4-vector normal to the boundaries of Σ (note the direction convention on the inner boundary). We assume a general mass distribution within the inner boundary and no matter outside it.

Local temperature. Following the results for horizons¹, it is tempting to seek to interpret $\kappa/(2\pi)$ from Eq. (1) as the local temperature at any point along an arbitrary 2-surface $\partial\Sigma_{\text{in}}$. However, this would be unsatisfactory if true for arbitrary surfaces, since this local

temperature would not be in thermal equilibrium with an actual physical screen held fixed at the same location; the temperature now coming from the Unruh effect³ and the local proper acceleration required to keep each portion of the screen stationary. Only for surfaces of constant Newtonian gravitational potential ϕ , where the proper acceleration of a stationary observer and the local normal to the surface are parallel, is such thermal equilibrium possible (see Supplementary Methods). Thus the temptation of such a thermodynamic interpretation should be restricted to the family of ordinary surfaces satisfying $\phi = \text{constant}$.

Indeed, this restricted temptation appears to have been satisfied in the emergent gravity program, where for static asymptotically-flat spacetimes, ordinary surfaces of constant ϕ are dubbed holographic screens and are claimed to have a local temperature⁸ given by $T = \kappa/(2\pi)$ and even to possess a ‘state variable’ quantifying the number of ‘bits’ on the screen. These putative thermodynamic properties are then used in a heuristic derivation of the full Einstein field equations⁸. If correct, such a claim would mean that the emergent gravity program would already subsume many decades of results associated with full General Relativity in this setting.

First law of thermodynamics. Here we test this thermodynamic assumption by asking whether perturbations of Eq. (1) reproduce the first law of thermodynamics. After all, thermodynamics is primarily a theory about how energy transforms under change, and this aspect of the theory is embodied in the first law. In the simplest case, where the hypersurface Σ is empty of matter, this law should read

$$\delta M = \frac{1}{8\pi} \int_{\partial\Sigma_{\text{in}}} \kappa \delta(dA). \quad (2)$$

We start by following Bardeen et al.’s original analysis¹, generalizing it where necessary to deal with a boundary $\partial\Sigma_{\text{in}}$ which is an arbitrary ordinary surface instead of a horizon. Under diffeomorphic metric perturbations we find (see Supplementary Methods)

$$\delta M = \frac{-1}{32\pi} \int_{\partial\Sigma_{\text{in}}} [4\delta\theta^{(l)} + (k_3 + k_6)\theta^{(l)} + (k_3 - k_6)\sigma_+^{(l)} + (k_4 + k_5)\sigma_\times^{(l)}] \mathcal{N} dA + \frac{1}{8\pi} \int_{\partial\Sigma_{\text{in}}} \kappa \delta(dA). \quad (3)$$

Here $\theta^{(l)}$ and $\sigma_j^{(l)}$ ($j = +, \times$) are the expansion and shears of null normal congruences of geodesics, the change in the expansion under the diffeomorphism is given by

$$\delta\theta^{(l)} = -\frac{k_2}{2}\theta^{(l)} + \frac{1}{2}(k_3 + k_6)_{;\rho} \hat{N}^\rho, \quad (4)$$

and the k_j are functions corresponding to independent components of the metric perturbation. As the expansion and shears vanish identically on the horizon⁹, we see that Eq. (3) trivially reduces to the first law, Eq. (2), thus reproducing the famous 1973 result¹. Similarly, it follows

straightforwardly that for surfaces sufficiently close to the horizon (so-called stretched horizons), the corrections to the first law will be negligible.

Surfaces away from horizons. So far we have assumed that the inner boundaries before and after the diffeomorphic perturbation are arbitrary. But could the perturbed boundary be chosen in a specific manner so as to cause the unwanted terms in Eq. (3) to vanish? As already noted, holographic screens correspond to surfaces of constant Newtonian potential $\phi = \text{constant}$. Thus, the perturbed screen relies on a specification of the constant $\delta\phi$. It is easy to show that $\delta\phi = \frac{1}{2}k_1$ (see Supplementary Methods), where k_1 is a metric perturbation of which the unwanted terms in Eq. (3) are wholly independent. Thus, the ordinary surfaces used within the emergent gravity program cannot generally satisfy the first law, Eq. (2).

One caveat to this claim comes when we consider a fully spherically symmetric scenario; where both the initial spacetime and screen are spherically symmetric, so the initial shears $\sigma_j^{(l)}$ vanish, and also the final spacetime and screen are spherically symmetric, placing further constraints on the k_j . In this case, Birkhoff’s theorem¹⁰ for spherically symmetric metrics imposes extra constraints between the metric components so that a perturbed screen may always be chosen so as to satisfy the form of the first law¹¹. However, as noted above, this form will not be preserved under arbitrary metric perturbations.

Discussion

The implications of our results are now described for (i) stretched horizons, and (ii) ordinary surfaces.

(i) Stretched horizons have long been considered to act as black bodies⁶, effectively radiating with a temperature $\kappa/(2\pi)$. Thus, our demonstration that they also satisfy the first law to an excellent approximation hardly seems surprising. Nevertheless, we do not believe that our result here should be interpreted as implying that the surfaces corresponding to stretched horizons themselves should be imbued with actual thermodynamic properties.

In particular, we may consider an alternative spacetime, identical from the stretched horizon outward, but instead of a horizon, we consider an infinitesimal shell of matter just outside what would correspond to its Schwarzschild radius were the shell to collapse further, yet still within the ‘stretched horizon’. In this latter spacetime, there is no horizon and hence no Hawking radiation. Notwithstanding this, our work proves that the ‘stretched horizon’ still closely satisfies the first law.

We conclude from this that the laws of black hole mechanics are not sufficient in themselves to guarantee whether any particular surface is truly thermodynamic in nature. For stretched horizons, we interpret this reasoning to imply that their full thermodynamic behavior is only inherited from the presence of an underlying horizon, but is not intrinsic to stretched horizons themselves.

This conclusion appears to mimic the initial reluctance of general relativists¹ from accepting black hole horizons as truly thermodynamic despite the deep analogy to thermodynamics uncovered in the laws of black hole mechanics. By contrast, these laws should still be considered a necessary condition.

(ii) Our analysis further rigorously shows that the family of ordinary surfaces called holographic screens will generally not obey a first law of thermodynamics, in contrast to the long-standing result for horizons¹. (Other families would not even be in thermal equilibrium with a physical surface at the same location.) Recall that the first law is more general than thermodynamics: the ‘temperature’ is merely an integrating factor relating changes in energy to changes in some state variable (entropy in the case of thermodynamics). Failure of the first law means that the putative state variable is not a variable of state at all. Therefore, even in static asymptotically-flat spacetimes, where the emergent gravity program claims to derive the full Einstein field equations, our results show that the key assumption of this program is actually inconsistent with General Relativity.

Methods

Energy conservation under diffeomorphisms. In order to attempt to derive a first law for ordinary surfaces we closely follow in the footsteps of Bardeen, Carter and Hawking’s 1973 classic paper¹. The first step is to obtain an integral equation for the net energy in a static system,

Eq. (1), where instead of an inner boundary located at a black hole horizon, this boundary is an ordinary surface. Next, we consider small ‘changes’ in the net energy corresponding to shifting to a parametrically nearby solution to the Einstein field equations. This ‘differential’ version is determined by studying the behavior of the net energy under spacetime diffeomorphisms of the initial metric¹. As in Bardeen et al., “gauge” freedom in the choice of coordinates is used to ensure that the hypersurfaces before and after the diffeomorphism are covered by identical sets of coordinates.

Study assumptions. Our analysis is limited to static asymptotically-flat solutions, with zero shift vector, $\beta^\mu = 0$. For simplicity, we assume that the spacetime of interest is non-rotating, and that there is no matter exterior to the holographic screen ($T^{\mu\nu} = 0$). We work throughout in natural units where $G = c = \hbar = k_B = 1$. Full and extensive details of the analysis are provided in the Supplementary Methods.

Data availability. The authors declare that all relevant data of this study are contained in the article and its Supplementary Information

Acknowledgements.

Z-WW acknowledges the support from Jilin University and China Scholarships Council. Z-WW is grateful to Tang Aoqing Honors Program in Science.

Author contributions.

The authors are equally responsible for all calculations and for writing and editing the manuscript.

The authors declare no competing interests.

* Electronic address: phyjasonwang@gmail.com; Electronic address: sam.braunstein@york.ac.uk

¹ Bardeen, J. M., Carter, B. & Hawking, S. W. The four laws of black hole mechanics. *Communications in Mathematical Physics* **31**, 161–170 (1973).

² Hawking, S. W. Particle creation by black holes. *Communications in mathematical physics* **43**, 199–220 (1975).

³ Unruh, W. G. Notes on black-hole evaporation *Physical Review D* **14**, 870 (1976).

⁴ Jacobson, T. Thermodynamics of spacetime: The Einstein equation of state. *Physical Review Letters* **75**, 1260 (1995).

⁵ Gibbons, G. W. & Hawking, S. W. Cosmological event horizons, thermodynamics, and particle creation. *Physical Review D* **15**, 2738 (1977).

⁶ Thorne, K. S., Price, R. H. & Macdonald, D. A. *Black Holes: The Membrane Paradigm* (Yale University Press,

Yale, 1986) pp.35-37.

⁷ Zurek, W. H. & Thorne, K. S. Statistical mechanical origin of the entropy of a rotating, charged black hole. *Physical review letters* **54**, 2171 (1985).

⁸ Verlinde, E. On the Origin of Gravity and the Laws of Newton. *Journal of High Energy Physics* **2011**, 29 (2011).

⁹ Hawking, S. W. & Ellis, G. F. R. *The large scale structure of space-time.* (Cambridge University Press, 1973), pp82-83, p334,.

¹⁰ Birkhoff, G. D. & Langer, R. E. *Relativity and modern physics.* (Cambridge, Harvard University Press, 1923).

¹¹ Chen, Y.-X. & Li, J.-L. First law of thermodynamics on holographic screens in entropic force frame. *Physics Letters B* **700**, 380–384 (2011).