

This is a repository copy of *Non-equilibrium particle dynamics with unbounded number of interacting neighbors*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/135626/>

Version: Accepted Version

Article:

Finkelshtein, Dimitri and Daletskii, Alexei orcid.org/0000-0003-3185-9806 (2018) Non-equilibrium particle dynamics with unbounded number of interacting neighbors. *Journal of Statistical Physics*. pp. 1-21. ISSN 1572-9613

<https://doi.org/10.1007/s10955-018-2159-x>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Non-equilibrium particle dynamics with unbounded number of interacting neighbors

Alexei Daletskii¹ Dmitri Finkelshtein²

September 12, 2018

Abstract

We consider an infinite system of first order differential equations in \mathbb{R}^ν , parameterized by elements x of a fixed countable set $\gamma \subset \mathbb{R}^d$, where the right-hand side of each x -equation depends on a finite but in general unbounded number n_x of variables (a row-finite system).

Such systems describe in particular (non-equilibrium) dynamics of spins $q_x \in \mathbb{R}^\nu$ of a collection of particles labelled by points $x \in \gamma$. Two spins q_x and q_y interact via a pair potential if the distance between x and y is no more than a fixed interaction radius. In contrast to the case where γ is a regular graph, e.g. \mathbb{Z}^d , the number n_x of particles interacting with particle x can be unbounded in x . Our main example of a “growing” configuration γ is a typical realization of a Poisson (or Gibbs) point process.

Under certain dissipativity-type condition on the right-hand side of our system and a bound on growth of n_x , we prove the existence and (under additional assumptions) uniqueness of infinite lifetime solutions with explicit estimates of growth in parameter x and time t .

For this, we obtain uniform estimates of solutions to approximating finite systems using a version of Ovsyannikov’s method for linear systems in a scale of Banach spaces. As a by-product, we develop an infinite-time generalization of the Ovsyannikov method.

Keywords: interacting particle systems, row-finite systems, scale of Banach spaces, Hamiltonian dynamics, gradient diffusion, self-organised systems, Ovsyannikov’s method, dissipativity

2010 Mathematics Subject Classification: 82C22, 34A12, 34A30

¹Department of Mathematics, The University of York, York YO1 5DD, U.K. (ad557@york.ac.uk).

²Department of Mathematics, Swansea University, Singleton Park, Swansea SA2 8PP, U.K. (d.l.finkelshtein@swansea.ac.uk).

1 Introduction

In recent decades, there has been an increasing interest to the study of countable systems of particles randomly distributed in the Euclidean space \mathbb{R}^d , which appear, in particular, in modeling of non-crystalline (amorphous) substances, e.g. ferrofluids and amorphous magnets, see e.g. [14], [13, Section 11], [2] and [4, 5]. Each particle is characterized by its position $x \in \mathbb{R}^d$ and an internal parameter (spin) $q_x \in S = \mathbb{R}^\nu$. For a given fixed (“quenched”) configuration γ of particle positions, which is a locally finite subset of \mathbb{R}^d , one considers a system of differential equations describing (non-equilibrium) dynamics of spins q_x , $x \in \gamma$.

Two spins q_x and q_y are allowed to interact via a pair potential if the distance between x and y is no more than a fixed interaction radius $r > 0$, that is, they are neighbors in the geometric graph defined by γ and r . The case where the vertex degrees of the graph are globally bounded (in particular, if γ has a regular structure, e.g. $\gamma = \mathbb{Z}^d$) has been well-studied (even in the stochastic case), see [11], [6] and modern developments in [9], and references therein. However, aforementioned applications to non-crystalline substances require to deal with unbounded vertex degrees. For example, if configuration γ is distributed according to a Poisson or, more generally, Gibbs measure with a superstable low regular interaction energy, the typical number of “neighbors” of a particle located at $x \in \mathbb{R}^d$ is proportional to $\log|x|$ (see Example 2.10 below).

More generally, we consider, for a fixed countable $\gamma \subset \mathbb{R}^d$, the system

$$\frac{d}{dt}q_x(t) = F_x(\bar{q}(t)), \quad q_x(0) = q_{x,0} \in S, \quad x \in \gamma, \quad (1.1)$$

where $\bar{q}(t) = (q_x(t))_{x \in \gamma} \in S^\gamma$, $t \in \mathbb{R}_+ := [0, \infty)$ and $F_x : S^\gamma \rightarrow S$ for each $x \in \gamma$. We suppose that the system is row-finite, that is, F_x depends only on a finite number n_x of components of the vector \bar{q} , which may be unbounded in $x \in \gamma$.

In our main motivating example, the right-hand side of (1.1) has the form

$$F_x(\bar{q}) = \sum_{y \in \gamma} W_{xy}(q_x, q_y) - \nabla U_x(q_x), \quad (1.2)$$

where $W_{xy} = 0$ if $|x-y| > r$. In this case, (1.1) describes general gradient-type spin dynamics for particle configuration γ and interaction radius r . A slight modification of this framework covers also the case of the infinite anharmonic (Hamiltonian) system on γ :

$$\frac{d}{dt}q_x(t) = p_x(t), \quad \frac{d}{dt}p_x(t) = F_x(\bar{q}(t)). \quad (1.3)$$

A natural approach to the study of system (1.1) would be to realize it as an evolution equation (with “good” coefficients) in a Banach space of sequences. However, in many cases, this proves to be impossible even for linear systems, as it happens for instance on unbounded vertex degree graphs. Beyond the framework of a fixed Banach space, linear row-finite systems are always solvable, although the corresponding solutions might show an uncontrolled growth in $x \in \gamma$, see e.g. in [7, § 6].

The aim of this work is to show the existence of solutions $q_x(t)$, $x \in \gamma$, to system (1.1) keeping control over the growth of the solution in $x \in \gamma$. In order to do this, we introduce an increasing scale of Banach spaces S_α^γ , $\alpha \geq 0$, consisting of sequences $\bar{q} = (q_x)_{x \in \gamma}$ with $|q_x| \leq \text{const } w(|x|)^\alpha$ for a fixed weight function w . Our main result (Theorem 2.4) states that for each $0 < \alpha < \beta$ and any initial condition $\bar{q}_0 \in S_\alpha^\gamma$ there exists a solution $\bar{q}(t) \in S_\beta^\gamma$ of system (1.1) on infinite time-interval $[0, \infty)$, subject to certain dissipativity-type conditions on F_x and a bound on the order of growth of n_x in $x \in \gamma$.

Effectively, the aforementioned logarithmic in $|x|$ growth of n_x (corresponding to a Gibbsian configuration) appears maximally allowed one, see Example 2.10. The corresponding weight is then exponential, $w(|x|) = e^{|x|}$; in other words, given an initial condition $\bar{q}(0) = (q_x(0))_{x \in \gamma}$ with at most exponential growth in x , we show that $\bar{q}(t)$ is of the same type for all $t > 0$. An informal balance condition between n_x and $w(|x|)$ is given by $w(|x|) \sim \exp(\exp(n_x))$, see Subsection 2.2 for details.

For the proof of the existence result, we approximate (1.1) by finite systems. Uniform estimates of the corresponding solutions are obtained using a version of the so-called Ovsyannikov method for linear systems in a scale of Banach spaces. In contrast to the classical Ovsyannikov method (see e.g. [7]), our modified version gives the existence of solutions with infinite lifetime (Theorem 3.1).

We also prove the uniqueness of the solution to (1.1), which requires however stronger conditions on F_x (Theorem 5.1).

We present three types of examples. In Subsection 6.1, we consider a general gradient-type system of the form (1.1), (1.2). The existence of the solutions is proved for polynomially bounded pair potentials W , properly dominated by the self-interaction potential U , see Proposition 6.1. The uniqueness is shown under stronger assumptions (cf. Proposition 6.2), which essentially boil down to either linear growth of W (the harmonic case), or slow (log log) growth of n_x together with linear bound on the initial condition.

In Subsection 6.2, we replace $(q_x)_{x \in \gamma}$ in the right hand side of (1.1) by the pairs $(p_x, q_x)_{x \in \gamma}$ with $\frac{d}{dt} p_x = q_x$ to study infinite anharmonic (Hamiltonian) systems (1.3) on γ . Note that the latter example, in the case of the regular $\gamma = \mathbb{Z}^d$, was studied in [11]. Finally, in Subsection 6.3, we consider an example

of multiparticle dynamics of somewhat different type, motivated by the study of self-organized systems, see review in [12].

2 Problem description and the existence result

2.1 The setup and main result

We start with more precise description of our model. We suppose that γ is a locally finite subset of the space $X = \mathbb{R}^d$, $d \geq 1$, i.e. the set $\gamma \cap \Lambda$ is finite for any compact $\Lambda \subset X$. We denote by S^η the vector space of elements of the form $(q_x)_{x \in \eta}$ with $q_x \in S$ for $x \in \eta \subseteq \gamma$.

Let us fix a number $r > 0$ and introduce the family of *finite* sets

$$\gamma_{x,r} := \{y \in \gamma \mid |x - y| \leq r\}, \quad x \in \gamma.$$

Consider the space $S^{\gamma_{x,r}}$ endowed with the (finite) Cartesian product topology and introduce the notation $\bar{q}_{x,r} := (q_u)_{u \in \gamma_{x,r}}$.

We suppose that the following condition holds, which reflects the fact that the system (1.1) is row-finite.

Condition 1. *For each $x \in \gamma$ there exists $f_x \in C^1(S^{\gamma_{x,r}}, S)$ such that*

$$F_x(\bar{q}) = f_x(\bar{q}_{x,r}), \quad \bar{q} \in S^\gamma.$$

Throughout the paper, we will understand solutions of (1.1) in the following sense.

Definition 2.1. We call a map $\bar{q} : [0, T) \rightarrow S^\gamma$ a (pointwise) solution of system (1.1) if the function $t \mapsto q_x(t) \in S$ is continuous (resp. continuously differentiable) on the interval $[0, T)$ (resp. $(0, T)$) and satisfies (1.1) for each $x \in \gamma$.

In what follows, we assume that the family of mappings $(F_x)_{x \in \gamma}$ is in certain sense dissipative. To this end, let $U_x \in C^2(S, \mathbb{R}_+)$, $x \in \gamma$, be a family of functions such that

$$U_x(q) \geq C_1 |q|, \quad q \in S, \quad |q| \geq 1, \quad (2.1)$$

$$U_x(q) \leq C_2(|q|^j + 1), \quad q \in S, \quad (2.2)$$

for some $C_1, C_2 > 0$ and $j \in \mathbb{N}$. Next, we introduce the following notation:

$$\text{for } x \in \gamma, \quad y \sim x \text{ means that } y \in \gamma_{x,r}.$$

Let also $n_x = n_{x,r}(\gamma) \geq 1$ denote the number of points in $\gamma_{x,r}$, $x \in \gamma$.

Condition 2. Let $U_x \in C^2(S, \mathbb{R}_+)$, $x \in \gamma$, satisfy (2.1)–(2.2), and there exist $C > 0$ and $m \in \mathbb{N}$ such that

$$F_x(\bar{q}) \cdot \nabla U_x(q_x) \leq C \sum_{y \sim x} (n_x n_y)^m U_y(q_y), \quad \bar{q} \in S^\gamma, \quad x \in \gamma. \quad (2.3)$$

Here the dot \cdot denotes Euclidean inner product in S . Examples of the families $(F_x)_{x \in \gamma}$ satisfying Conditions 1–2 will be given in Section 6.

Let $w : \mathbb{R}_+ \rightarrow [1, \infty)$ be a non-decreasing function. We define the family of Banach spaces

$$S_\alpha^\gamma := \left\{ \bar{q} \in S^\gamma \mid \|\bar{q}\|_\alpha := \sup_{x \in \gamma} \frac{|q_x|}{w(|x|)^\alpha} < \infty \right\}, \quad \alpha > 0. \quad (2.4)$$

Here and below, with an abuse of notations, $|x|$ means the Euclidean norm in $X = \mathbb{R}^d$, whereas $|q_x|$ means the Euclidean norm in $S = \mathbb{R}^\nu$. Clearly, the family (2.4) is increasing in α , i.e. $S_\alpha^\gamma \subset S_\beta^\gamma$ for $0 < \alpha < \beta$.

We introduce now certain balance condition on the growth of n_x and $w(|x|)$ as $|x| \rightarrow \infty$.

Definition 2.2. Let \mathcal{R} denote the class of non-decreasing functions $f : \mathbb{R}_+ \rightarrow [1, \infty)$ such that

$$f_\tau := \sup_{s \in \mathbb{R}_+} \frac{f(\tau + s)}{f(s)} < \infty \quad (2.5)$$

for any $\tau > 0$.

Examples of functions from \mathcal{R} are considered in Subsection 2.2 below.

Definition 2.3. Let $w, z \in \mathcal{R}$. We call the pair (w, z) admissible if for any $\beta, \mu > 0$ and $\alpha \in (0, \beta)$ there exists $D = D(\beta, \mu) \geq 1$ such that

$$\sup_{s \in \mathbb{R}_+} z(s)^\mu w(s)^{-\alpha} \leq \frac{D}{\alpha}. \quad (2.6)$$

We formulate now the balance condition. Let w be the weight function defining the scale of spaces S_α^γ in (2.4).

Condition 3. There exists $z \in \mathcal{R}$ such that

$$n_x \leq z(|x|), \quad x \in \gamma, \quad (2.7)$$

and the pair (w, z) is admissible.

The following is our main existence result.

Theorem 2.4. *Let Conditions 1–3 hold. Then, for each $\alpha \geq 0$ and any initial condition $\bar{q}_0 = (q_{x,0})_{x \in \gamma} \in S_\alpha^\gamma$, there exists a pointwise solution $\bar{q}(t)$ of system (1.1) on infinite time-interval $[0, \infty)$. Moreover, for $j \in \mathbb{N}$ from (2.2) we have the inclusion*

$$\bar{q}(t) \in \bigcap_{\beta > j\alpha} S_\beta^\gamma, \quad t \in [0, \infty),$$

and the estimate

$$\|\bar{q}(t)\|_\beta \leq C_p(\alpha, \beta; t) (\|\bar{q}_0\|_\alpha^j + 1) \quad (2.8)$$

holds true for any $\beta > j\alpha$ and $p > 1$. Here $C_p(\alpha, \beta) : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is an entire function of order $\rho = \frac{p}{p-1} > 1$ and type $\sigma = B^\rho e^{\rho(\beta r + \beta - j\alpha + 1)} (e\rho)^{-1} (\beta - j\alpha)^{-\frac{1}{p-1}} > 0$ with some $B = B(p) > 0$.

Remark 2.5. Clearly, by choosing $p > 1$ large enough, one can make the order $\rho > 1$ arbitrary close to 1.

Remark 2.6. The order and type of C_p are positive and finite. Thus, for any $\varepsilon > 0$, there exists $T_\varepsilon > 0$ such that

$$\sup_{t \in [0, T]} \|\bar{q}(t)\|_\beta \leq e^{(\sigma + \varepsilon)T^{\rho + \varepsilon}} (\|\bar{q}_0\|_\alpha^j + 1), \quad T > T_\varepsilon.$$

2.2 Discussion of Condition 3 and examples

Our next goal is to explain in more detail the definition of admissible pairs.

We start with the discussion of the class \mathcal{R} . Let $f : \mathbb{R}_+ \rightarrow [1, \infty)$ be a non-decreasing function. Then, for each $\tau > 0$, the function $\frac{f(\tau+s)}{f(s)}$ is bounded in s on any closed subinterval of \mathbb{R}_+ . Thus a sufficient condition for (2.5) is that $\lim_{s \rightarrow \infty} \frac{f(\tau+s)}{f(s)} < \infty$, $\tau > 0$. The latter, by e.g. [1, Theorem 1.4.1] applied to the function $g(s) := f(\log s)$, $s > 1$, is equivalent to the requirement that g has regular variation of index $\rho \in \mathbb{R}$, i.e. $\lim_{s \rightarrow \infty} \frac{g(\tau s)}{g(s)} = \tau^\rho$, $\tau > 0$. Combining [1, Theorem 1.4.1] and the examples of [1, p. 16], we conclude that a function $g(s) = s^\rho h(s)$, $\rho \in \mathbb{R}$, has regular variation of index ρ , if e.g. $h(s)$ is a product of non-negative powers of $\log s$, $\log \log s$ (and so on), $e^{(\log s)^\nu}$, $\nu \in (0, 1)$ etc. Considering $f(s) := g(e^s)$, $s \in \mathbb{R}_+$, we get that e.g. the functions

$$f(s) = (\log s)^\kappa s^\nu e^{\rho s^\mu}, \quad \rho, \kappa, \nu \geq 0, \quad \mu \in (0, 1]$$

belong to the set \mathcal{R} , as well as the functions $\log f(s)$, $\log \log f(s)$ and so on.

The following Lemma describes a simple way to generate admissible pairs.

Lemma 2.7. (1) Let $w \in \mathcal{R}$, $w \geq e^e$, and z be given by the formula

$$z(s) = v \log(\log w(s)), \quad v > 0.$$

Then the pair (w, z) is admissible.

(2) Let (w, z) be an admissible pair. Then the pair $(w + c_1, z + c_2)$ is admissible for any $c_1, c_2 \geq 0$.

(3) Let (w, z) be an admissible pair, $v \in \mathcal{R}$ and $v(s) \geq w(s)$, $s \in \mathbb{R}_+$. Then (v, z) is an admissible pair.

Proof. As we already pointed out, $z \in \mathcal{R}$. It is straightforward to check that, for any $\mu > 0$, there exists $d_\mu > 0$, such that $(\log \tau)^\mu \leq d_\mu \tau$ for all $\tau > 0$. Then, denoting $v(s) := \log w(s) \geq 0$, $s \in \mathbb{R}_+$, we obtain

$$z(s)^\mu w(s)^{-\alpha} \leq v^\mu d_\mu (\log w(s)) w(s)^{-\alpha} = v^\mu d_\mu v(s) e^{-\alpha v(s)} \leq \frac{v^\mu d_\mu}{e\alpha},$$

which completes the proof of part (1). Parts (2) and (3) are obvious. \square

Remark 2.8. In (1), it is sufficient to assume that $w(s) \geq e^e$, $s \geq s_0$ for some $s_0 > 0$ only, and set e.g. $w(s) = 1$ for $s \in [0, s_0)$. Then one can choose $z(s) = v \log(\log(w(s)))$ for $s \geq s_0$ and $z(s) = 1$ otherwise.

Let us note that the structure of the given underlying set γ dictates the choice of the function z , which in turn determines suitable weight function w and, ultimately, the conditions on the family F_x , $x \in \gamma$. Below are three examples of admissible pairs associated with different type of γ .

Example 2.9 (Minimal growth). Assume that the number of elements in $\gamma_{x,r}$ is globally bounded, that is, there exists a constant $z_0 \geq 1$ such that $n_x \leq z_0$ for all $x \in \gamma$. An important example of such γ is given by the integer lattice \mathbb{Z}^d . Then we can set $z(s) \equiv z_0$ and choose *an arbitrary* non-decreasing function $w : \mathbb{R}_+ \rightarrow [1, \infty)$ such that $w(s) \geq e^e$ for $s \geq s_0$, cf. Remark 2.8. The choice of w is dictated by the growth of the initial condition $\bar{q}_0 = (q_{x,0})_{x \in \gamma} \in S_\alpha^\gamma$, cf. Theorem 2.4 (i.e. by the growth of $|q_{x,0}|$ in $x \in \gamma$). Note also that our main technical tool, the modified Ovsiannikov theorem (cf. Theorem 3.1 and Proposition 4.2), becomes then redundant, as the corresponding linear operator A defined by formula (2.13) will be bounded in any such (fixed) S_α^γ , $\alpha > 0$ (i.e. the dynamics evolves in one Banach space). This case is well-studied, see [11], where $w(s) = e^s$ (and $\alpha = 1$) was considered. In particular, if $(|q_{x,0}|)_{x \in \gamma}$ is bounded, one can choose $w(s) \equiv e^e$; then all the spaces S_α^γ , $\alpha > 0$ will coincide with the space $l_\infty(S)$ of bounded sequences (and A is bounded there).

Example 2.10 (Maximal growth). Assume that the number of elements in $\gamma_{x,r}$ grows logarithmically, that is, there exists $a(r) > 0$ such that

$$n_{x,r}(\gamma) \leq z(|x|), \quad z(s) := a(r)(1 + \log(1 + s)) \quad (2.9)$$

for all $x \in X$ and $r > 0$. This bound holds for a typical random configuration distributed according to a Ruelle measure on X , see e.g. [15] and [10, p. 1047]. Then we can set $w(s) = e^s$, $s \geq e$, cf. Remark 2.8. It follows from Lemma 2.7 that the pair (w, z) is admissible. This is our most important motivating case, see Section 6 and Example 6.4.

Example 2.11 (Medium growth). Let, for some $v > 0$,

$$z(s) = v(1 + \log \log(e + s)). \quad (2.10)$$

Then, according to Lemma 2.7, we can set $w(s) = 1 + s$. In this case, more comprehensive study of the solutions of system (1.1) can be accomplished, at least in the framework of systems with pair interaction, which we discuss in Section 6 (see Example 6.5). Indeed, slow growth of the weight function w allows to show the uniqueness of the solution living in S_β^γ with any (fixed) $\beta > 0$.

2.3 Scheme of the proof

To explain the scheme of the proof of Theorem 2.4, we start with the following simple observation. Let $\bar{q}(t)$ solve (1.1), and define

$$\mathcal{L}_x(t) := U_x(q_x(t)), \quad x \in \gamma, \quad t \geq 0. \quad (2.11)$$

Then Condition 2 yields

$$\begin{aligned} \frac{d}{dt} \mathcal{L}_x(t) &= \frac{d}{dt} q_x(t) \cdot \nabla U_x(q_x(t)) = F_x(\bar{q}(t)) \cdot \nabla U_x(q_x(t)) \\ &\leq C \sum_{y \sim x} (n_x n_y)^m U_y(q_y) = \sum_{y \in \gamma} A_{x,y} \mathcal{L}_y(t), \end{aligned} \quad (2.12)$$

where

$$A_{x,y} := \begin{cases} C(n_x n_y)^m, & y \sim x, \\ 0, & \text{otherwise.} \end{cases} \quad (2.13)$$

Introduce the infinite matrix $A = (A_{x,y})_{x,y \in \gamma}$. Relation (2.12) implies the following differential inequality in S^γ :

$$\frac{d}{dt} \mathcal{L}(t) \leq A \mathcal{L}(t), \quad \mathcal{L}(t) = (\mathcal{L}_x(t))_{x \in \gamma}. \quad (2.14)$$

Now we can proceed as follows. First we will prove that, for any $0 < \alpha < \beta$, the differential equation

$$\frac{d}{dt}\Psi(t) = A\Psi(t), \quad \Psi(0) \in S_\alpha^\gamma, \quad (2.15)$$

which corresponds to the inequality (2.14), has a classical solution with infinite lifetime in S_β^γ (Proposition 4.2). To this end, we use a modified version of Ovsyannikov's method (proved in Theorem 3.1). Informally, we will have then that $\mathcal{L}_x(t) \leq \Psi_x(t)$, $x \in \gamma$, $t \geq 0$ (here and below we understand inequalities between elements of $S = \mathbb{R}^\nu$ in the coordinate-wise sense). Next, we will approximate (1.1) by finite volume cut-off systems and use the corresponding finite-dimensional versions of (2.14) and (2.15) to find (uniform in volume $\Lambda \subset X$) estimates, which will allow us to pass to a limit as $\Lambda \rightarrow X$.

3 Linear equations in a scale of Banach spaces

In this section we prove a general result on the existence of (infinite-time) solutions for a special class of linear differential equations, which extends the so-called Ovsyannikov method, see e.g. [7].

Let us consider a family of Banach spaces \mathbb{B}_α indexed by $\alpha \in (0, \beta]$ with $\beta < \infty$ fixed, and denote by $\|\cdot\|_\alpha$ the corresponding norms. We assume that

$$\mathbb{B}_{\alpha'} \subset \mathbb{B}_{\alpha''} \text{ and } \|u\|_{\alpha''} \leq \|u\|_{\alpha'} \text{ if } \alpha' < \alpha'', \quad u \in \mathbb{B}_{\alpha''}, \quad (3.1)$$

where the embedding means that $\mathbb{B}_{\alpha'}$ is a vector subspace of $\mathbb{B}_{\alpha''}$. For any $\delta \in (0, \beta]$, we set $\mathbb{B}_{\delta-} := \bigcup_{0 < \alpha < \delta} \mathbb{B}_\alpha$.

Let $A : \mathbb{B}_{\beta-} \rightarrow \mathbb{B}_{\beta-}$ be a linear operator. We assume that A is a bounded operator acting from $\mathbb{B}_{\alpha'}$ to $\mathbb{B}_{\alpha''}$ for any $0 < \alpha' < \alpha'' \leq \beta$, and

$$\|Ax\|_{\alpha''} \leq c(\alpha'' - \alpha')^{-q} \|x\|_{\alpha'} \quad (3.2)$$

for all $x \in \mathbb{B}_{\alpha'}$ and some constants $c = c(\beta) > 0$ and $q \in (0, 1)$ (both independent of α' and α'').

Theorem 3.1. *Assume that the bound (3.2) holds. Then, for any $\alpha \in (0, \beta)$ and $u_0 \in \mathbb{B}_\alpha$, there exists a continuous function $u : [0, \infty) \rightarrow \mathbb{B}_\beta$ with $u(0) = u_0$ such that:*

- 1) u is continuously differentiable on $(0, \infty)$;
- 2) $Au(t) \in \mathbb{B}_\beta$ for all $t \in (0, \infty)$;

3) u solves the differential equation

$$\frac{d}{dt}u(t) = Au(t), \quad t \in (0, \infty). \quad (3.3)$$

Moreover,

$$\|u(t)\|_{\beta} \leq a(t)\|u_0\|_{\alpha}, \quad t > 0, \quad (3.4)$$

where $a(t)$ is an entire function of order $\rho = \frac{1}{1-q}$ and type $\sigma = (ce)^{\rho}(e\rho)^{-1}(\beta - \alpha)^{-q\rho}$.

Proof. Let us first observe that, by (3.2), A is a well-defined operator in any $\mathbb{B}_{\delta-}$, $\delta \in (\alpha, \beta)$. Thus, for any $n \geq 1$, one can consider $A^n : \mathbb{B}_{\delta-} \rightarrow \mathbb{B}_{\delta-}$. Embeddings $\mathbb{B}_{\alpha} \subset \mathbb{B}_{\delta-} \subset \mathbb{B}_{\delta}$ imply that $A^n : \mathbb{B}_{\alpha} \rightarrow \mathbb{B}_{\delta}$ is a well-defined linear operator.

Fix $u_0 \in \mathbb{B}_{\alpha}$ and define the sequence $u_k = A^k u_0 (= Au_{k-1})$, $k = 1, 2, \dots$. Then, for any $n \geq 1$ and $\alpha_1, \dots, \alpha_n$ such that $\alpha < \alpha_1 < \dots < \alpha_n = \delta$, we have $u_k \in \mathbb{B}_{\alpha_k-} \subset \mathbb{B}_{\alpha_k}$ and

$$\|u_k\|_{\alpha_k} \leq c(\alpha_k - \alpha_{k-1})^{-q} \|u_{k-1}\|_{\alpha_{k-1}}.$$

Setting $\alpha_k := \alpha + k\frac{\delta-\alpha}{n}$, we obtain the estimate

$$\|A^n u_0\|_{\delta} \leq D^n n^{qn} \|u_0\|_{\alpha}, \quad D := c(\delta - \alpha)^{-q}, \quad n \geq 1.$$

Therefore the series

$$u(t) := \sum_{n=0}^{\infty} \frac{t^n}{n!} A^n u_0 \quad (3.5)$$

is uniformly convergent in \mathbb{B}_{δ} on the interval $[0, T]$ for each $T > 0$. Indeed, the inequality $n! \geq \left(\frac{n}{e}\right)^n$ implies that

$$\|u(t)\|_{\delta} \leq \|u_0\|_{\alpha} \sum_{n=0}^{\infty} \frac{(De)^n}{n^{(1-q)n}} t^n < \infty \quad (3.6)$$

for any $t > 0$, and the series in the r.h.s. of (3.6) defines an entire function.

Similarly, the series

$$v(t) := \sum_{n=0}^{\infty} \frac{t^n}{n!} A^{n+1} u_0$$

is uniformly convergent in \mathbb{B}_{δ} on the interval $[0, T]$. As a result,

$$v(t) = \frac{d}{dt}u(t) \quad (3.7)$$

in \mathbb{B}_δ . Since the norm in \mathbb{B}_δ is stronger than in \mathbb{B}_β , the function $[0, T] \ni t \mapsto u(t)$ is differentiable in \mathbb{B}_β and (3.7) holds in \mathbb{B}_β , too. Observe that A maps $\mathbb{B}_\delta \rightarrow \mathbb{B}_\beta$ continuously. Therefore, we can apply A to the right hand side of (3.5) and obtain $Au(t) = v(t) \in \mathbb{B}_\beta$. This proves parts 1) – 3) of the statement.

Next, fix any $\delta_0 \in (\alpha, \beta)$. The series in the r.h.s. of (3.6) converges uniformly in $\delta \in [\delta_0, \beta]$ (recall that D depends on δ) to a continuous decreasing function of δ . Therefore, one can pass to the limit as $\delta \rightarrow \beta$ in (3.6), to get

$$\|u(t)\|_\beta \leq \|u_0\|_\alpha \sum_{n=0}^{\infty} \frac{(Be)^n}{n^{(1-q)n}} t^n < \infty, \quad B := c(\beta - \alpha)^{-q}. \quad (3.8)$$

The sum of the series in the r.h.s. of (3.8) is an entire function of order

$$\rho := \limsup_{n \rightarrow \infty} \frac{n \ln n}{-\ln \frac{(Be)^n}{n^{(1-q)n}}} = \frac{1}{1-q}.$$

The type σ of this entire function satisfies the equality

$$(\sigma e \rho)^{\frac{1}{\rho}} = \limsup_{n \rightarrow \infty} n^{\frac{1}{\rho}} \left(\frac{(Be)^n}{n^{(1-q)n}} \right)^{\frac{1}{n}} = Be,$$

i.e.

$$\sigma = \frac{(Be)^\rho}{e\rho} = \frac{(ce)^\rho}{e\rho(\beta - \alpha)^{q\rho}},$$

which completes the proof. \square

Remark 3.2. The order and type of $a(t)$ are positive and finite. Thus, for any $\varepsilon > 0$, there exists $T_\varepsilon > 0$ such that

$$\sup_{t \in [0, T]} \|u(t)\|_\beta \leq e^{(\sigma + \varepsilon)T^{\rho + \varepsilon}} \|u_0\|_\alpha$$

for all $T > T_\varepsilon$.

Remark 3.3. The function u from Theorem 3.1 is unique. The proof can be obtained in a similar way to the “standard” Ovsyannikov method, see e.g. [8, Theorem 2.4]. Note that we do not use this fact in the present paper.

4 Proof of the existence result

4.1 Row-finite linear systems

We start with the study of the linear system (2.15). First, we show that the matrix (2.13) generates a bounded operator in the scale $(S_\alpha^\gamma)_{\alpha > 0}$.

Lemma 4.1. *Let Condition 3 hold. Then the matrix $(A_{x,y})_{x,y \in \gamma}$, given by (2.13), generates a bounded linear operator $A : S_{\alpha'}^\gamma \rightarrow S_{\alpha''}^\gamma$ for any $0 < \alpha' < \alpha'' < \beta$. The corresponding norm of A satisfies the estimate*

$$\|A\|_{\alpha', \alpha''} \leq Bw_r^{\alpha'} (\alpha'' - \alpha')^{-1/p} \quad (4.1)$$

with an arbitrary $p > 1$ and some $B = B(p) > 0$.

Proof. Fix arbitrary $0 < \alpha' < \alpha''$. For any $\bar{q} \in S_{\alpha'}^\gamma$, we have

$$\|A\bar{q}\|_{\alpha''} \leq C \sup_{x \in \gamma} \sum_{y \sim x} (n_x n_y)^m |q_y| w(|x|)^{-(\alpha'' - \alpha')} w(|x|)^{-\alpha'},$$

by (2.4), (2.13). Recall that $y \sim x$ implies $|y - x| \leq r$, and hence

$$|y| \leq r + |x|. \quad (4.2)$$

Then, since w is and satisfies (2.5), we have

$$w(|x|)^{-\alpha'} \leq \left(\frac{w(r + |x|)}{w(|x|)} \right)^{\alpha'} w(|y|)^{-\alpha'} \leq w_r^{\alpha'} w(|y|)^{-\alpha'},$$

where w_r is as in (2.5), so that

$$\begin{aligned} \|A\bar{q}\|_{\alpha''} &\leq Cw_r^{\alpha'} \sup_{x \in \gamma} w(|x|)^{-(\alpha'' - \alpha')} \sum_{y \sim x} (n_x n_y)^m |q_y| w(|y|)^{-\alpha'} \\ &\leq w_r^{\alpha'} \|\bar{q}\|_{\alpha'} \sup_{x \in \gamma} \left(A_x w(|x|)^{-(\alpha'' - \alpha')} \right), \end{aligned} \quad (4.3)$$

where

$$A_x := C \sum_{y \sim x} (n_x n_y)^m, \quad x \in \gamma. \quad (4.4)$$

By (2.7), (4.2) and (2.5), we have

$$n_y \leq z(|y|) \leq \frac{z(r + |x|)}{z(x)} z(|x|) \leq z_r z(|x|), \quad y \sim x, \quad x \in \gamma.$$

Therefore,

$$A_x \leq Cn_x^{m+1} z_r^m z(|x|)^m = Mz(|x|)^{2m+1},$$

where $M := Cz_r^m$.

Fix an arbitrary $p > 1$. By (4.3) and (2.6) we obtain the bound

$$\begin{aligned} \|A\bar{q}\|_{\alpha''} &\leq Mw_r^{\alpha'} \|\bar{q}\|_{\alpha'} \sup_{x \in \gamma} \left(z(|x|)^{p(2m+1)} w(|x|)^{-p(\alpha'' - \alpha')} \right)^{\frac{1}{p}} \\ &\leq Mw_r^{\alpha'} \|\bar{q}\|_{\alpha'} \left(\frac{D(p\beta, p(2m+1))}{p(\alpha'' - \alpha')} \right)^{\frac{1}{p}} = Bw_r^{\alpha'} (\alpha'' - \alpha')^{-1/p} \|\bar{q}\|_{\alpha'}, \end{aligned}$$

where $B = M \left(\frac{1}{p} D(p\beta, p(2m+1)) \right)^{\frac{1}{p}} > 0$. This completes the proof. \square

Now we can use Theorem 3.1 and prove the existence of solutions of equation (2.15).

Proposition 4.2. *Let Condition 3 hold and let $A = (A_{x,y})_{x,y \in \gamma}$ be given by (2.13). Then, for any $0 < \alpha < \beta$, equation (2.15) with $\Psi(0) \in S_\alpha^\gamma$ has a classical solution $\Psi(t) \in S_\beta^\gamma$, $t \geq 0$. Moreover, the estimate*

$$\|\Psi(t)\|_\beta \leq A_p(\alpha, \beta; t) \|\Psi(0)\|_\alpha, \quad t \geq 0, \quad (4.5)$$

holds true for each $p > 1$, with an entire function $A_p(\alpha, \beta; \cdot) : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ of order $\rho = \frac{p}{p-1} > 1$ and type $\sigma = B^\rho w_r^{\beta\rho} e^{\rho-1} \rho^{-1} (\beta - \alpha)^{-\frac{1}{p-1}} > 0$.

Proof. Take any $p > 1$. By Lemma 4.1, for any $0 < \alpha < \alpha' < \alpha'' \leq \beta$, the estimate (3.2) (for the scale of spaces $\mathbb{B}_\alpha = S_\alpha^\gamma$) holds with $q = \frac{1}{p} \in (0, 1)$ and $c = B e^{\beta r + \beta - \alpha} > B e^{\alpha' r + \alpha'' - \alpha'} > 0$. Then the result follows directly from Theorem 3.1. \square

4.2 Finite-dimensional approximations

For a compact set $\Lambda \subset X$, consider the collection $\bar{q}^\Lambda(t) = (q_x^\Lambda(t))_{x \in \gamma}$ of functions $q_x^\Lambda : \mathbb{R}_+ \rightarrow S$ such that

$$\begin{cases} \frac{d}{dt} q_x^\Lambda(t) = F_x(\bar{q}^\Lambda(t)), & t > 0, x \in \gamma_\Lambda, \\ q_x^\Lambda(t) = q_x(0), & t > 0, x \in \gamma_{\Lambda^c}, \\ q_x^\Lambda(0) = q_x(0), & x \in \gamma, \end{cases} \quad (4.6)$$

where $\gamma_\Lambda = \gamma \cap \Lambda$, $\gamma_{\Lambda^c} = \gamma \cap \Lambda^c$, $\Lambda^c := X \setminus \Lambda$.

Proposition 4.3. *Let Conditions 1–3 hold. Then, for any compact $\Lambda \subset X$ and $\alpha > 0$, the system (4.6) with an initial condition $\bar{q}_0 = (q_x(0))_{x \in \gamma} \in S_\alpha^\gamma$ has a unique solution $\bar{q}^\Lambda(t)$, $t \geq 0$. Moreover, for each $x \in \gamma$, there exists an entire function $Q_x : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, such that the estimate*

$$|q_x^\Lambda(t)| \leq Q_x(t), \quad t \geq 0, \quad (4.7)$$

holds for all compact sets $\Lambda \subset X$.

Proof. For an arbitrary compact $\Lambda \subset X$, the system (4.6) is essentially finite-dimensional with continuously differentiable coefficients. Thus there exists a unique solution of (4.6) with lifetime $T_\Lambda \leq \infty$. Moreover, $T_\Lambda < \infty$ implies that $|q_x^\Lambda(t)| \rightarrow \infty$ as $t \nearrow T_\Lambda$ for some $x \in \gamma_\Lambda$.

Similarly to (2.11), define

$$\mathcal{L}_x^\Lambda(t) := U_x(q_x^\Lambda(t)), \quad x \in \gamma, \quad t \in [0, T_\Lambda], \quad (4.8)$$

and note that $\mathcal{L}_x^\Lambda(0) = \mathcal{L}_x(0) = U_x(q_x(0))$. Observe that $\frac{d}{dt}\mathcal{L}_x^\Lambda(t) = 0$ for $x \in \gamma_{\Lambda_c}$, $t > 0$. Thus, similarly to (2.12), we have the inequality

$$\frac{d}{dt}\mathcal{L}_x^\Lambda(t) \leq \sum_{y \in \gamma} a_{x,y}^\Lambda \mathcal{L}_y^\Lambda(t), \quad x \in \gamma, \quad t \in [0, T_\Lambda], \quad (4.9)$$

where

$$a_{x,y}^\Lambda := \begin{cases} A_{x,y}, & x \in \Lambda, \\ 0, & \text{otherwise} \end{cases} \quad (4.10)$$

and $A_{x,y}$ is given by (2.13). Denote

$$\Lambda_r := \{x \in X \mid \text{dist}(x, \Lambda) \leq r\}.$$

We can replace γ with $\gamma_{\Lambda_r} = \gamma \cap \Lambda_r$ in the r.h.s. of (2.13), because $a_{x,y}^\Lambda = 0$ if $y \notin \Lambda_r$. Also, the relation $\frac{d}{dt}\mathcal{L}_x^\Lambda(t) = 0 \leq \mathcal{L}_x^\Lambda(t)$, $x \in \gamma \cap (\Lambda_r \setminus \Lambda)$, implies that (2.13) always holds for $x \notin \Lambda_r$. Thus the system of differential inequalities (4.9) is also essentially finite.

The classical comparison theorem (see e.g. [16]) implies that

$$0 \leq \mathcal{L}_x^\Lambda(t) \leq \Psi_x^{\Lambda_r}(t), \quad x \in \gamma_{\Lambda_r}, \quad (4.11)$$

where the collection of functions $(\Psi_x^{\Lambda_r}(t))_{x \in \gamma_{\Lambda_r}}$ satisfies the following system of equations

$$\frac{d}{dt}\Psi_x^{\Lambda_r}(t) = \sum_{y \in \gamma_{\Lambda_r}} a_{x,y}^\Lambda \Psi_y^{\Lambda_r}(t), \quad \Psi_x^{\Lambda_r}(0) = \mathcal{L}_x^\Lambda(0), \quad x \in \gamma_{\Lambda_r}.$$

The latter system can be considered as a single equation with the cut-off matrix $A^\Lambda = (a_{x,y}^\Lambda)_{x,y \in \gamma}$:

$$\dot{\Psi}^{\Lambda_r}(t) = A^\Lambda \Psi^{\Lambda_r}(t), \quad \Psi_x^{\Lambda_r}(0) = \mathcal{L}_x^\Lambda(0), \quad x \in \gamma, \quad (4.12)$$

with the ‘trivial’ extension $\Psi_x^{\Lambda_r}(t) = \mathcal{L}_x^\Lambda(t)$, $t \geq 0$, for all $x \in X \setminus \Lambda_r$.

By (2.11) and (2.2) we have

$$|\mathcal{L}_x(0)| \leq C_2(|q_x(0)|^j + 1). \quad (4.13)$$

Therefore $\bar{q}_0 \in S_\alpha^\gamma$ yields $(\mathcal{L}_x^\Lambda(0))_{x \in \gamma} = (\mathcal{L}_x(0))_{x \in \gamma} =: \mathcal{L}(0) \in S_{j\alpha}^\gamma$ and

$$\|\mathcal{L}(0)\|_{j\alpha} \leq C_3 (\|\bar{q}_0\|_\alpha^j + 1) \quad (4.14)$$

for some constant C_3 . It is evident that estimate (4.1) holds for the operator A^Λ instead of A , with the same constant $B > 0$. Therefore, using the arguments similar to those in the proof of Proposition 4.2, we can show that the solution $\Psi^{\Lambda r}(t)$ to the system (4.12) exists for all $t \geq 0$ and satisfies the estimate

$$\|\Psi^{\Lambda r}(t)\|_{j\beta} \leq A_p(j\alpha, j\beta; t) \|\mathcal{L}(0)\|_{j\alpha}, \quad \beta > \alpha, \quad t \geq 0, \quad (4.15)$$

with the same entire function A_p as in (4.5).

Next, using (2.1), (4.8), (4.11) and (4.15), we can write, for each $t \geq 0$,

$$\begin{aligned} |q_x^\Lambda(t)| &\leq \frac{1}{C_1} U_x(q_x^\Lambda(t)) + 1 = \frac{1}{C_1} \mathcal{L}_x^\Lambda(t) + 1 \leq \frac{1}{C_1} \Psi_x^{\Lambda r}(t) + 1 \\ &\leq \frac{1}{C_1} A_p(j\alpha, j\beta; t) \|\mathcal{L}(0)\|_{j\alpha} w(x)^{j\beta} + 1 =: Q_x(t) < \infty. \end{aligned} \quad (4.16)$$

This bound holds for any $x \in \gamma$, which implies that $T_\Lambda = \infty$ and (4.7) holds. The proof is complete. \square

Corollary 4.4. *Estimates (4.16) and (4.14) imply that*

$$\|\bar{q}^\Lambda(t)\|_\beta \leq C_p(\alpha, \beta; t) (\|\bar{q}_0\|_\alpha^j + 1), \quad t \geq 0, \quad (4.17)$$

for any $\beta > j\alpha$. Here $C_p(\alpha, \beta; \cdot) : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is an entire function of order $\rho = \frac{p}{p-1} > 1$ and type $\sigma = B^\rho e^{\rho(\beta r + \beta - j\alpha + 1)} (e\rho)^{-1} (\beta - j\alpha)^{-\frac{1}{p-1}} > 0$ with some $B = B(p) > 0$.

Now we are ready to prove Theorem 2.4. The proof is essentially similar to that in [11].

Proof of Theorem 2.4. Fix $\alpha > 0$, $\beta > j\alpha$ and $\bar{q}_0 \in S_\alpha^\gamma$. Choose any sequence Λ^0 of compacts Λ exhausting X . Let $\bar{q}^\Lambda(t)$ solve the corresponding system (4.6). Observe that both sides of inequality (4.7) are continuous in t , which implies that

$$\sup_{t \in [0, T]} |q_x^\Lambda(t)| \leq \sup_{t \in [0, T]} Q_x(t) < \infty,$$

for each $x \in \gamma$ and any $T > 0$. The equality $\dot{q}_x^\Lambda = F_x(\bar{q}^\Lambda)$ together with continuity of F_x on $S^{\gamma_{x,r}}$ imply then that $\sup_{t \in [0, T]} |\dot{q}_x^\Lambda(t)|$, $x \in \gamma$, is also bounded uniformly in Λ .

Let us fix an arbitrary indexation of γ , so that $\gamma = (x_k)_{k=1}^\infty$. The Arzelà-Ascoli theorem implies that there exists a subsequence $\Lambda^{(1)} = (\Lambda_n^{(1)})$ of $\Lambda^{(0)}$ and some $q_{x_1}(t) \in S$ such that $q_{x_1}^{\Lambda_n^{(1)}}(t) \rightrightarrows q_{x_1}(t)$ as $n \rightarrow \infty$, where \rightrightarrows denotes

the uniform convergence in $t \in [0, T]$. Repeating these arguments, we can see that for any $k \in \mathbb{N}$ there exists a subsequence $\mathbf{\Lambda}^{(k)} = (\Lambda_n^{(k)})$ of $\mathbf{\Lambda}^{(k-1)}$ and $q_{x_k}(t) \in S$ such that $q_{x_k}^{\Lambda_n^{(k)}}(t) \rightrightarrows q_{x_k}(t)$, $n \rightarrow \infty$. Then of course $q_{x_m}^{\Lambda_n^{(k)}}(t) \rightrightarrows q_{x_m}(t)$, $n \rightarrow \infty$, for all $m \leq k$.

Consider now the ‘‘diagonal’’ sequence $\mathbf{\Lambda}$ with elements $\Lambda_n := \Lambda_n^{(n)}$. It is clear that $q_{x_k}^{\Lambda_n}(t) \rightrightarrows q_{x_k}(t)$, $n \rightarrow \infty$, for all $k \in \mathbb{N}$.

The limit transition in both sides of the equality

$$q_x^{\Lambda_n}(t) = q_{x,0}^{\Lambda_n} + \int_0^t F_x(q_x^{\Lambda_n}(s)) ds, \quad x \in \gamma,$$

shows that the functions $q_x(t) = \lim q_x^{\Lambda_n}(t)$, $x \in \gamma$, solve the system (1.1). The inclusion $\bar{q}(t) \in \bigcap_{\beta > j\alpha} S_\beta^\gamma$ and bounds (2.8) for all $\beta > j\alpha$ follow from (4.17). The proof is complete. \square

5 The uniqueness

In this section, we will discuss conditions that guarantee the uniqueness of the solution $\bar{q}(t) \in S_\beta^\gamma$. We fix $\beta > 0$ and let $\Delta_{\beta,R}$ be the ball of radius $R > 0$ in S_β^γ centered at $0 = (0)_{x \in \gamma} \in S_\beta^\gamma$.

For $x, y \in \gamma$ denote by $\frac{\partial F_x(\bar{q})}{\partial q_y}$ the Jacobian matrix of the mapping F_x w.r.t. the variable q_y . By Condition 1, this Jacobian is the zero matrix if $y \notin \gamma_{x,r}$. We also define the corresponding gradient as the following vector with matrix components:

$$\nabla F_x(\bar{q}) = \left(\frac{\partial F_x(\bar{q})}{\partial q_y} \right)_{y \in \gamma}, \quad x \in \gamma.$$

Since all but finite number of the components of the gradient vector are zero matrices, we can define its norm

$$|\nabla F_x(\bar{q})| := \sum_{y \in \gamma} \left\| \frac{\partial F_x(\bar{q})}{\partial q_y} \right\| < \infty, \quad x \in \gamma,$$

where $\|\cdot\|$ denotes the standard norm of a linear operator in S .

To ensure the uniqueness, we assume the following:

Condition 4. For any $R > 0$, there exists a constant $C_R > 0$ such that

$$\sup_{\bar{q} \in \Delta_{\beta,R}} |\nabla F_x(\bar{q})| \leq C_R (|x| + 1), \quad x \in \gamma.$$

Theorem 5.1. *Let $\beta > 0$, and assume that Condition 4 holds. Suppose that the weight sequence $w(s)$ in (2.4) satisfies the bound $w(s) \leq e^{\nu s}$, $s \in \mathbb{R}_+$ for some $\nu > 0$. Let $\bar{q}^{(1)}(t), \bar{q}^{(2)}(t) \in S_\beta^\gamma$ be two pointwise solutions of (1.1) on $[0, T]$, and let $\bar{q}^{(1)}(0) = \bar{q}^{(2)}(0)$. Then $\bar{q}^{(1)}(t) = \bar{q}^{(2)}(t)$, $t \in [0, T]$.*

Proof. We start by observing that, because of Condition 1, we have the bound

$$\begin{aligned} & |F_x(\bar{q}^{(1)}) - F_x(\bar{q}^{(2)})| \\ & \leq \sum_{y \in \gamma_{x,r}} |q_y^{(1)} - q_y^{(2)}| \sup_{\tau \in [0,1]} \left\| \frac{\partial F_x}{\partial q_y} (\tau \bar{q}^{(1)} + (1-\tau) \bar{q}^{(2)}) \right\|, \end{aligned} \quad (5.1)$$

which holds for any $\bar{q}^{(l)} = (q_x^{(l)})_{x \in \gamma}$, $l = 1, 2$.

Assume now that $\bar{q}^{(1)}, \bar{q}^{(2)} \in \Delta_{\beta,R}$, $R > 0$. Then $\tau \bar{q}^{(1)} + (1-\tau) \bar{q}^{(2)} \in \Delta_{\beta,R}$, $\tau \in [0, 1]$. Let $n \geq 1$ and $x \in \gamma$ be such that $|x| \leq nr$. Then $|y| \leq (n+1)r$ for any $y \in \gamma_{x,r}$, cf. (4.3). Thus (5.1) implies the following estimate:

$$|F_x(\bar{q}^{(1)}) - F_x(\bar{q}^{(2)})| \leq \sup_{|y| \leq (n+1)r} |q_y^{(1)} - q_y^{(2)}| \sup_{\bar{q} \in \Delta_{\beta,R}} |\nabla F_x(\bar{q})|. \quad (5.2)$$

Let us now $\bar{q}^{(1)}(t), \bar{q}^{(2)}(t) \in S_\beta^\gamma$ be two solutions of (1.1) on $t \in [0, T]$. Set $R := \max_{t \in [0, T], l=1,2} \|\bar{q}^{(l)}(t)\|_\beta$, so that $\bar{q}^{(l)}(t) \in \Delta_{\beta,R}$, and let $C_R > 0$ be such that Condition 4 holds. Denote

$$\delta_n(t) := \sup_{|x| \leq nr} |q_x^{(1)}(t) - q_x^{(2)}(t)|, \quad n \geq 1, t \in [0, T]. \quad (5.3)$$

For any $x \in \gamma$ such that $|x| \leq nr$, it follows from the integral form of (1.1), inequality (5.2) and Condition 4 that

$$\begin{aligned} |q_x^{(1)}(t) - q_x^{(2)}(t)| & \leq \int_0^t |F_x(\bar{q}^{(1)}(s)) - F_x(\bar{q}^{(2)}(s))| ds \\ & \leq \int_0^t \delta_{n+1}(s) \sup_{\bar{q} \in \Delta_{\beta,R}} |\nabla F_x(\bar{q})| ds \\ & \leq C_R (|x| + 1) \int_0^t \delta_{n+1}(s) ds. \end{aligned}$$

Therefore,

$$\delta_n(t) \leq C_R (nr + 1) \int_0^t \delta_{n+1}(s) ds \leq \mu nr \int_0^t \delta_{n+1}(s) ds,$$

where $\mu := C_R(1 + \frac{1}{r}) > 0$. The N -th iteration of this estimate gives

$$\delta_n(t) \leq \frac{(t\mu r)^N}{N!} n(n+1)\dots(n+N-1) \sup_{s \leq t} \delta_{n+N}(s). \quad (5.4)$$

A direct computation using (2.4), (5.3) and the inclusion $\bar{q}^{(1)}(t), \bar{q}^{(2)}(t) \in \Delta_{\beta, R}$ shows that $\delta_{n+N}(s) \leq 2Rw((n+N)r)^\beta$, $s \geq 0$. Using the assumption $w(s) \leq e^{\nu s}$, $s \in \mathbb{R}_+$, we get that (5.4) implies the bound

$$\begin{aligned} \delta_n(t) &\leq 2Re^{\beta\nu(n+N)r} (t\mu r)^N \binom{n+N-1}{N} \\ &\leq 2Re^{\beta\nu nr} \left(t\mu r e^{\beta\nu r+1} \frac{n+N-1}{N} \right)^N, \end{aligned}$$

where we used the well-known inequality $\binom{M}{N} \leq (\frac{M\epsilon}{N})^N$, $1 \leq N \leq M$. Therefore, for all $n \geq 1$ and $N > n-1$ we have

$$\delta_n(t) < 2Re^{\beta\nu nr} (2t\mu r e^{\beta\nu r+1})^N, \quad t \geq 0. \quad (5.5)$$

Observe that for $t < t_0 := (2\mu r e^{\beta\nu r+1})^{-1}$ the r.h.s. of inequality (5.5) converges to zero as $N \rightarrow \infty$, which in turn implies that $\delta_n(t) = 0$ for all $n \geq 1$. Thus $\bar{q}^{(1)}(t) = \bar{q}^{(2)}(t)$ for $t \in [0, t_0)$.

These arguments can be repeated on each of the time intervals $[t_k, t_{k+1})$ with $t_k := kt_0$, $k = 1, 2, \dots$, which shows that $\bar{q}^{(1)}(t) = \bar{q}^{(2)}(t)$, $t \in [0, T)$, for an arbitrary $T > 0$. \square

6 Dynamics of interacting particle systems

Our main example is motivated by the study of (deterministic) dynamics of interacting particle systems. In this case, γ represents a collection of particles indexed by elements x of X , may be interpreted as particle positions. A particle with position $x \in \gamma$ carries an internal parameter (spin) $\sigma_x \in S$. While the positions of our particles are fixed, the spins evolve according to system (1.1). Here we will consider two types of the time-evolution with pair spin-spin interaction- general gradient-type dynamics and the Hamiltonian anharmonic dynamics. In the last subsection, we give an example of somewhat different type, motivated by the study of self-organized systems.

6.1 Gradient-type dynamics

Consider the following example of the family $(F_x)_{x \in \gamma}$ that fulfills Condition 1:

$$F_x(\bar{q}(t)) = R_x(\bar{q}_{x,r}(t)) - \nabla U_x(q_x(t)), \quad x \in \gamma, \quad (6.1)$$

with

$$R_x(\bar{q}) = \sum_{y \sim x, y \neq x} W_{xy}(q_x, q_y), \quad x \in \gamma. \quad (6.2)$$

Here, for each $x \in \gamma$, we denote $\bar{q}_{x,r}(t) := (q_y(t))_{y \in \gamma_{x,r}}$ and assume that $U_x \in C^2(S, \mathbb{R}_+)$, and hence $\nabla U_x \in C^1(S, S)$. Next, for each $\{x, y\} \subset \gamma$, $x \sim y$, we assume that $W_{xy} \in C^2(S^2, S)$.

Proposition 6.1. *Let, for some $k \geq 1$, $j \geq 2k$, $J_U > 0$, condition (2.2) holds, and condition (2.1) is reinforced to*

$$U_x(q) \geq J_U |q|^{2k}, \quad q \in S, x \in \gamma. \quad (6.3)$$

Let also, for some $J_W > 0$,

$$|W_{xy}(q_1, q_2)| \leq J_W (|q_1|^k + |q_2|^k + 1), \quad q_1, q_2 \in S, x, y \in \gamma. \quad (6.4)$$

Then there exists $C > 0$ such that Condition 2 holds with $m = 1$.

Proof. First we note that for F of the form (6.1) condition (2.3) reads as

$$R_x(\bar{q}) \cdot \nabla U_x(q_x) - |\nabla U_x(q_x)|^2 \leq C n_x^m \sum_{y \sim x} n_y^m U_y(q_y) \quad (6.5)$$

for any $\bar{q} \in S^{\gamma_{x,r}}$ and $x \in \gamma$. The inequality $R_x \cdot \nabla U_x \leq |\nabla U_x|^2 + |R_x|^2$ implies that the following bound is sufficient for (6.5) to hold:

$$|R_x(\bar{q})|^2 \leq C n_x^m \sum_{y \sim x} n_y^m U_y(q_y), \quad \bar{q} \in S^\gamma, x \in \gamma. \quad (6.6)$$

We are going to check now (6.6) with $m = 1$, for R_x given by (6.2) under the conditions above. By (6.2) and (6.4), there exist $J_1, J_2, J_3 > 0$, such that

$$|R_x(\bar{q})|^2 \leq J_1 n_x \sum_{y \sim x, y \neq x} |q_y|^{2k} + J_2 n_x^2 |q_x|^{2k} + J_3 n_x.$$

On the other hand, (6.3) yields

$$\begin{aligned} n_x \sum_{y \sim x} n_y U_y(q_y) &\geq n_x \sum_{y \sim x, y \neq x} J_U (|q_y|^{2k} + 1) + n_x^2 J_U (|q_x|^{2k} + 1) \\ &= J_U n_x \sum_{y \sim x, y \neq x} |q_y|^{2k} + J_U n_x^2 |q_x|^{2k} + 2J_U n_x^2 - J_U n_x. \end{aligned}$$

Thus (6.6) holds with $m = 1$ and $C = \max\{J_1, J_2, J_3\} / J_U$ (we used also that $n_x \geq 1$ as $x \in \gamma_{x,r}$). \square

Proposition 6.2. *Assume that the interaction potentials W and U satisfy the bounds*

$$\|\nabla^2 U_y(q)\| \leq K_U(|q|^{j-2} + 1), \quad q \in S, x \in \gamma, \quad (6.7)$$

$$\left\| \frac{\partial}{\partial q_2} W_{xy}(q_1, q_2) \right\| \leq K_W(|q_1|^{k-1} + |q_2|^{k-1} + 1), \quad q_1, q_2 \in S, x, y \in \gamma, \quad (6.8)$$

for some $k \geq 1$, $j \geq 2k$ and $K_U, K_W > 0$. Let the pair (w, z) satisfy Condition 3 and, moreover, the inequality

$$z(s)w(s)^{\beta(k-1)} + w(s)^{\beta(j-2)} \leq K_1 s + K_2, \quad s \in \mathbb{R}_+. \quad (6.9)$$

holds for some $K_1, K_2 > 0$. Then Condition 4 holds.

Proof. Again, we note that for F as in (6.1) Condition 4 obtains the following form: for any $R > 0$ there exists a constant $C_R > 0$ such that

$$\sup_{\bar{q} \in \Delta_{\beta, R}} \sum_{y \in \gamma_{x, r}} \left\| \frac{\partial R_x(\bar{q}_{x, r})}{\partial q_y} - \mathbb{1}_{\{y=x\}} \nabla^2 U_x(q_x) \right\| \leq C_R(|x| + 1), \quad x \in \gamma,$$

where ∇^2 is the Hessian matrix. Therefore, it is sufficient to assume that

$$\sup_{\bar{q} \in \Delta_{\beta, R}} \left(\sum_{y \in \gamma_{x, r}} \left\| \frac{\partial R_x(\bar{q}_{x, r})}{\partial q_y} \right\| + \|\nabla^2 U_x(q_x)\| \right) \leq C_R(|x| + 1), \quad x \in \gamma. \quad (6.10)$$

Let now R_x be given by (6.2). Then, $\frac{\partial R_x(\bar{q}_{x, r})}{\partial q_x}$ is the zero matrix, and (6.10) can be rewritten, for each $x \in \gamma$, as follows

$$\sup_{\bar{q} \in \Delta_{\beta, R}} \left(\sum_{y \sim x, y \neq x} \left\| \frac{\partial}{\partial q_y} W_{xy}(q_x, q_y) \right\| + \|\nabla^2 U_x(q_x)\| \right) \leq C_R(|x| + 1). \quad (6.11)$$

Recall that $\bar{q} \in \Delta_{\beta, R}$ implies

$$|q_x| \leq R w(|x|)^\beta, \quad x \in \gamma.$$

Note that one can assume $R \geq 1$. Then, by (6.8), (2.5), (2.7), we have

$$\begin{aligned} \sum_{y \sim x, y \neq x} \left\| \frac{\partial}{\partial q_y} W_{xy}(q_x, q_y) \right\| &\leq K_W R^k \sum_{y \sim x, y \neq x} (w(|x|)^{\beta(k-1)} + w(|y|)^{\beta(k-1)} + 1) \\ &\leq K_W R^k (n_x w(|x|)^{\beta(k-1)} + n_x w(|x| + r)^{\beta(k-1)} + 1) \\ &\leq K_W R^k (2 + w_r^{\beta(k-1)}) z(|x|) w(|x|)^{\beta(k-1)}. \end{aligned}$$

Next, by (6.7),

$$\|\nabla^2 U_x(q_x)\| \leq 2K_U R^{j-2} w(|x|)^{\beta(j-2)}.$$

As a result, by using (6.9), we get (6.11). \square

Remark 6.3. A typical example of the pair interaction is

$$W_{xy}(q_x, q_y) = V(q_x - q_y), \quad V \in C^2(S, S).$$

Then, to ensure (6.4), it is enough to assume that

$$|V(q)| \leq J_V(|q|^k + 1), \quad q \in S$$

for some $J_V > 0$. Moreover, $\left\| \frac{\partial}{\partial q} V(q) \right\| = \left\| \frac{\partial}{\partial q_2} W(q_1, q_2) \right\|$ for $q = q_1 - q_2$, and hence (6.8) holds if only, for some $K_V > 0$,

$$\left\| \frac{\partial}{\partial q} V(q) \right\| \leq K_V(|q|^{k-1} + 1), \quad q \in S.$$

Now we will revisit our examples of admissible pairs and study the corresponding conditions of the existence and uniqueness of the corresponding dynamics. Recall that the exponents j and k in (2.2), (6.4), (6.7), (6.8) are always related by the assumption $j \geq 2k$, $k \geq 1$.

Example 6.4 (Example 2.10 revisited). Let (2.9) hold. Then Theorem 2.4 holds with $w(s) = e^{s+e}$. Thus a solution with initial value

$$|q_x(0)| \leq c \exp(\alpha|x|), \quad x \in \gamma,$$

will live in each of the spaces S_β^γ and thus satisfy the bound

$$|q_x(t)| \leq c(t) \exp(\beta|x|), \quad x \in \gamma, t > 0,$$

for all $\beta > j\alpha$ (here $c(t) = c(\alpha, \beta, t)$). It is clear that the uniqueness condition (6.9) of Proposition 6.2 holds only for $k = 1$, $j = 2$.

Example 6.5 (Example 2.11 revisited). Let (2.7) hold, z satisfy (2.10), and $w(s) = 1 + s$. Assume without loss of generality that $j > 2$. Then for $1 \leq k \leq \frac{j}{2}$ we have

$$z(s)w(s)^{\beta(k-1)} + w(s)^{\beta(j-2)} \leq z(s)(1+s)^{\frac{1}{2}} + (1+s),$$

so that (6.9) holds in the corresponding S_β^γ with any $\beta \leq (j-2)^{-1}$. Thus we can apply both Theorems 2.4 and 5.1 with arbitrary $\alpha < j^{-1}(j-2)^{-1}$ and $\beta \in (j\alpha, (j-2)^{-1}]$. As a result, for any initial data

$$|q_x(0)| \leq c(1 + |x|)^{\frac{1}{j(j-2)}}, \quad x \in \gamma,$$

there exists the unique solution satisfying the estimate

$$|q_x(t)| \leq c(t)(1 + |x|)^{\frac{1}{j-2}}, \quad x \in \gamma.$$

Moreover, the stronger bound

$$|q_x(t)| \leq c(t)(1 + |x|)^{\frac{1}{\beta}}, \quad x \in \gamma.$$

will hold for all $\beta \in (j\alpha, (j-2)^{-1}]$.

Example 6.6. A simple example of interactions satisfying assumptions (6.3)–(6.8) is given by

$$W_{xy}(q_x, q_y) = Jq_y, \quad J \in \mathbb{R},$$

and

$$U_y(q_y) = aq_y^2 - b, \quad a, b > 0.$$

In this case (1.1) is just the linear system

$$\dot{q}_x = -aq_x + J \sum_{y \sim x} q_y.$$

The existence and uniqueness of solutions in any S_β^γ , which follows from Theorem 2.4, can be proved by a direct application of Theorem 3.1.

6.2 Infinite Anharmonic Systems

Consider the Hamiltonian system

$$\dot{q}_x = p_x, \quad \dot{p}_x = R_x(\bar{q}) - \nabla U_x(q_x), \quad x \in \gamma,$$

with R_x and U_x as in the previous section. It fits the framework of Section 2 with the “double” state space $S \times S \ni (\bar{q}, \bar{p})$ in place of S and

$$F_x(\bar{q}, \bar{p}) = (\bar{p}_x, (R_x(\bar{q}) - \nabla U_x(q_x))) \in S \times S, \quad x \in \gamma,$$

and the single-particle Hamiltonian

$$H_x(q, p) = \frac{1}{2} |p|^2 + U_x(q)$$

replacing U_x in Condition 2.

Proposition 6.7. *Assume that R and U satisfy (6.6) resp. (6.10). Then the modified version of Condition 2 resp. Condition 4 holds.*

Proof. We clearly have

$$F_x(\bar{q}, \bar{p}) \cdot \nabla H_x(q_x, p_x) = p_x \cdot R_x(\bar{q}) \leq \frac{1}{2} (|p_x|^2 + |R_x(\bar{q})|^2).$$

Inequality (6.6) implies now that

$$\begin{aligned} F_x(\bar{q}, \bar{p}) \cdot \nabla H_x(q_x, p_x) &\leq \frac{1}{2} |p_x|^2 + \frac{1}{2} C n_x^m \sum_{y \sim x} n_y^m U_y(q_y) \\ &\leq \frac{1}{2} C n_x^m \sum_{y \sim x} n_y^m H_y(q_y, p_y), \end{aligned}$$

and the modified Condition 2 holds. A direct calculation shows that (6.10) implies Condition 4. \square

6.3 Self-alignment dynamics

This example is motivated by various models of self-organized dynamics based on the alignment. We refer to the review in [12] and references therein for the theory and applications of these models in biological, physical and social sciences. In this framework, equation (1.1) takes the form

$$\frac{d}{dt} q_x(t) = \sum_{y \sim x} g_{x,y}(\bar{q}_{x,r}(t)) (q_y(t) - q_x(t)), \quad (6.12)$$

where $g_{x,y} \in C^1(S^{\gamma_{x,r}}, \mathbb{R})$, $x, y \in \gamma$, is a family of non-negative uniformly bounded functions.

Note that in [12] the underlying set γ is supposed to be finite and either fixed or allowed to evolve. Thus (6.12) can be considered as the infinite-particle (“quenched”) version of these models.

We consider two examples inspired by the so-called opinion dynamics, in which functions $g_{x,y}$ are given by either

$$g_{x,y}(\bar{q}_{x,r}) = \frac{\phi(|q_y - q_x|)}{n_x}, \quad x \in \gamma, y \sim x, \quad (6.13)$$

or

$$g_{x,y}(\bar{q}_{x,r}) = \frac{\phi(|q_y - q_x|)}{\sum_{z \sim x} \phi(|q_z - q_x|)}, \quad x \in \gamma, y \sim x, \quad (6.14)$$

with $0 < \phi(s) \leq \phi_0$, $s > 0$ for some $\phi_0 > 0$.

Let us prove that Condition 2 holds with $U_x(q_x) := \frac{1}{2} |q_x|^2$, $x \in \gamma$. Indeed,

$$\begin{aligned} F_x(\bar{q}) \cdot \nabla U_x(q_x) &= \sum_{y \sim x} g_{x,y}(\bar{q}_{x,r}) (q_y \cdot q_x - |q_x|^2) \\ &\leq \sum_{y \sim x} g_{x,y}(\bar{q}_{x,r}) \frac{1}{2} (|q_y|^2 - |q_x|^2) \leq G \sum_{y \sim x} U_y(q_y), \end{aligned}$$

for some constant $G > 0$, and hence (2.3) holds with $m = 1$ and an arbitrary $n_x \geq 1$, $x \in \gamma$. In particular, there are no further restrictions on the function z in Condition 3.

Next, to check Condition 4, we note that

$$\frac{\partial}{\partial q_y} F_x(\bar{q}) = \sum_{y \sim x} g_{x,y}(\bar{q}_{x,r}) \mathbb{1} + \sum_{y \sim x} \frac{\partial}{\partial q_y} g_{x,y}(\bar{q}_{x,r}) \times q_y,$$

where $u \times v$ denotes (for $u, v \in S = \mathbb{R}^\nu$) the matrix $(u_i v_j)_{1 \leq i, j \leq \nu}$, and $\mathbb{1}$ is the identity matrix on S . We consider $g_{x,y}$ given by (6.13) and (6.14), respectively.

In the case of (6.13) we have

$$\frac{\partial}{\partial q_y} g_{x,y}(\bar{q}_{x,r}) = \frac{\phi'(|q_y - q_x|)}{n_x} \frac{q_y}{|q_y|}.$$

Observe that $\|q_y \times q_y\| \leq c|q_y|^2$, where the constant $c > 0$ depends only on the choice of norm on $S = \mathbb{R}^\nu$. Thus we obtain the inequality

$$\left\| \frac{\partial}{\partial q_y} F_x(\bar{q}) \right\| \leq \phi_0 + \sup_{s \geq 0} |\phi'(s)| \sum_{y \sim x} \frac{c|q_y|}{n_x}.$$

Therefore, assuming that $|\phi'(s)| \leq \phi_1$ for $s \geq 0$, we see that

$$|\nabla F_x(\bar{q})| \leq \phi_0 n_x + \frac{c\phi_1}{n_x} \|\bar{q}\|_\beta n_x w(|x| + r)^\beta,$$

and hence Condition 4 holds provided

$$z(s) + w(s)^\beta \leq K_1 s + K_2, \quad s \geq 0,$$

for some $K_1, K_2 > 0$.

In the case of $g_{x,y}$ given by (6.14) we obtain

$$\frac{\partial}{\partial q_y} g_{x,y}(\bar{q}_{x,r}) = \frac{\phi'(|q_y - q_x|) \sum_{z \sim x, z \neq y} \phi(|q_z - q_x|)}{\left(\sum_{z \sim x} \phi(|q_z - q_x|) \right)^2} \frac{q_y}{|q_y|}$$

and proceed in a completely similar way provided the condition of boundedness of $|\phi'(s)|$ is replaced by the bound $|\phi'(s)| \leq \phi_1 \phi(s)$, $s \geq 0$.

Finally, one can consider the modification of (6.12), which corresponds to the so-called flocking dynamics [12]:

$$\frac{d}{dt} q_x(t) = \sum_{y \sim x} g_{x,y}(\bar{p}_{x,r}(t)) (q_y(t) - q_x(t)), \quad \frac{d}{dt} p_x(t) = q_x(t). \quad (6.15)$$

In particular, for $g_{x,y}$ given by (6.13), we will get an infinite-particle counterpart of the well-known Cucker–Smale dynamics, see e.g. [3, 12]. It is straightforward to check that

$$U_x(q_x, p_x) := \frac{1}{2}|q_x|^2 + \frac{1}{2}|p_x|^2$$

fulfills Condition 2 and Condition 4 holds for $g_{x,y}$ as in (6.13) or (6.14).

Acknowledgements

We would like to thank Sergio Albeverio, Yuri Kondratiev, Oles Kutovyi and Tanja Pasurek for their interest to this work and stimulating discussions. Financial support of the Alexander von Humboldt Stiftung, the European Commission under the project STREVCOMS PIRSES-2013-612669 and DFG through SFB 1283 “Taming uncertainty and profiting from randomness and low regularity in analysis, stochastics and their applications” is gratefully appreciated.

References

- [1] N. H. Bingham, C. M. Goldie, J. L. Teugels, Regular variation, *Encyclopedia of Mathematics and its Applications* **27**, Cambridge University Press, Cambridge 1989.
- [2] A. Bovier, *Statistical Mechanics of Disordered Systems. A Mathematical Perspective* (Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 2006).
- [3] J. A. Canizo, J. A. Carrillo, J. Rosado, Collective behavior of animals: Swarming and complex patterns, *Arbor* **186** (2010), 1035–1049.
- [4] A. Daletskii, Yu. Kondratiev, Yu. Kozitsky, T. Pasurek, Gibbs states on random configurations, *J. Math. Phys.* **55** (2014), 083513.
- [5] A. Daletskii, Yu. Kondratiev, Yu. Kozitsky, T. Pasurek, Phase Transitions in a quenched amorphous ferromagnet, *J. Stat. Phys.* **156**(2014), 156–176.
- [6] G. Da Prato, J. Zabczyk, Ergodicity for Infinite Dimensional Systems, *London Mathematical Society Lecture Note Series* **229**, University Press, Cambridge, 1996.
- [7] K. Deimling, Ordinary differential equations in Banach spaces, *Lecture Notes in Mathematics* **596**, Springer 1977.

- [8] D. Finkelshtein, Around Ovsiannikov's method, *Meth. Funct. Anal. Topology* **21** (2015), No. 2, 134–150.
- [9] J. Inglis, M. Neklyudov, B. Zegarliński, Ergodicity for infinite particle systems with locally conserved quantities, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **15** (2012), No. 1, 1250005.
- [10] D. Klein and W. S. Yang, A characterization of first order phase transitions for superstable interactions in classical statistical mechanics. *Journal of Statistical Physics* **71** (1993), 1043–1062.
- [11] E. Lanford, J. Lebowitz, E. Lieb, Time Evolution of Infinite Anharmonic Systems, *Journal of Statistical Physics* **16** (1977), No. 6, 453–461.
- [12] S. Motsch, E. Tadmor, Heterophilous Dynamics Enhances Consensus, *SIAM Review* **56** (2014), No. 4, 577–621.
- [13] R. C. O'Handley, *Modern Magnetic Materials: Principles and Applications*, (Wiley, 2000).
- [14] S. Romano and V. A. Zagrebnov, Orientational ordering transition in a continuous-spin ferrofluid, *Phys. A* **253** (1998), 483–497.
- [15] D. Ruelle, Superstable interactions in classical statistical mechanics, *Commun. Math. Phys.* **18** (1970), 127–159.
- [16] W. Walter, *Differential and integral inequalities*, Springer 1970.