

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is an author's version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/35987>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

Diagrammatic proof of the BCFW recursion relation for gluon amplitudes in QCD.

**Petros D. Draggiotis^{(a,c)1}, Ronald H.P. Kleiss^{(b)2}, Achilleas Lazopoulos^{(b)3}
and Costas G. Papadopoulos^{(c)4}**

^(a)**University of Athens, Physics Department, Nuclear & Particle Physics
Section, GR – 15771, Athens, Greece**

^(b)**IMAPP, Institute of Mathematics, Astrophysics and Particle Physics,
Radboud University, Nijmegen, the Netherlands**

^(c)**Institute of Nuclear Physics, NCSR "Demokritos", 15310, Athens, Greece**

Abstract

We present a proof of the Britto-Cachazo-Feng-Witten tree-level recursion relation for gluon amplitudes in QCD, based on a direct equivalence between BCFW decompositions and Feynman diagrams. We demonstrate that this equivalence can be made explicit when working in a convenient gauge. We exhibit that gauge invariance and the particular structure of Yang-Mills vertices guarantees the validity of the BCFW construction.

1 Introduction

During the last year much progress has been made in the understanding of analytical calculations of dual amplitudes in perturbative Yang-Mills theories. Led by an observation of Witten [1], Britto, Cachazo and Feng (BCFW) have proposed a new recursion relation for tree amplitudes of gluons [2] that naturally arrives at the simplest known expressions for some of those amplitudes in terms of Weyl - Van der Waerden spinor products. Explicit calculations have been performed using this technique [3],[4], extensions to amplitudes involving particles from the electroweak sector [5] have been pursued and a new approach to one loop amplitudes has been proposed [6] employing MHV vertices and unitarity arguments.

¹pdrag@phys.uoa.gr

²R.Kleiss@science.ru.nl

³A.Lazopoulos@science.ru.nl

⁴costas.papadopoulos@cern.ch

The BCFW recursion relation features some remarkable characteristics, among which the on-shell analytic continuation of selected off-shell propagators, the analytic continuation of two selected external momenta in the complex plane and a decomposition of a color helicity amplitude into smaller helicity amplitudes with complex external momenta that doesn't appear to be in direct connection with the decomposition in Feynman diagrams. Moreover, the BCFW relation leads to the use of only three-point (modified) vertices as building blocks of the theory, thus raising questions about the fundamental nature of the Yang-Mills four point vertex.

A proof of the BCFW relation was given by Britto, Chachazo, Feng and Witten [7] which, though it made the connection between the analyticity properties of the color amplitude and the BCFW decomposition obvious, shed little light towards the way the latter relates to the usual Feynman diagram approach.

The simplicity of the final result for tree level gluonic amplitudes implies the existence of major cancellations between Feynman graphs. Since the BCFW formalism avoids those cancellations altogether it is advantageous to find a diagrammatic proof of the BCFW decomposition in order to understand these cancellations in the level of Feynman diagrams. A number of questions immediately arise. Is a diagrammatic proof gauge dependent? If so, are there preferable gauges where simplifications take place? Which gauges are those? Is this decomposition exclusively working for pure YM theories, and if so, what is the key feature of the YM vertices that makes it work? Can we find a BCFW type recursion relation for scalar theories?

We intend to address these questions in the present paper as well as giving a generic diagrammatic proof of the BCFW decomposition. The paper is organized as follows: after clarifying some notation issues in section 2, we present, in section 3 an analysis based on diagram enumeration that hints towards a connection between BCFW decompositions and Feynman graphs. In section 4 we prove some important kinematical identities that will support the main parts of the diagrammatic proof (sections 6 and 7), whereas section 5 describes the gauge in which we are working and the consequences of that choice. We conclude with some remarks on the generality of the proof and the role of gauge invariance in this analysis in section 8.

2 Notation

All amplitudes mentioned in this paper are the color-ordered amplitudes that result after the usual trace-based color decomposition [8]. We will conveniently omit all coupling constants and factors of i as well as the color antenna of all color-ordered amplitudes in what follows.

In non-vanishing tree-level color-ordered amplitudes, one can, without loss of generality, pick out a pair of adjacent legs with opposite helicity and label the positive helicity one by 1 and the negative helicity one by n . Unless otherwise specified we will use this labeling throughout this paper.

The Dirac four-spinors used in this paper are the usual $u_{\pm}(p)$, $\bar{u}_{\pm}(p)$ with the important note that the relation between $u(p)$ and $\bar{u}(p)$ is *not* one of complex conjugation.

The relation with Weyl - Van der Waerden spinors, when using the Weyl representation of γ -matrices, is

$$u_+(p) = (0, \tilde{\lambda}_b(p) \epsilon^{b\dot{a}}) \quad \bar{u}_+(p) = (\lambda_b(p) \epsilon^{b\dot{a}}, 0) \quad (1)$$

$$u_-(p) = (\lambda_a(p), 0) \quad \bar{u}_-(p) = (0, \tilde{\lambda}_{\dot{a}}(p)) \quad (2)$$

Four-momenta can be written in terms of Weyl spinors as

$$p_{\mu} = \frac{1}{2} \bar{u}_-(p) \gamma_{\mu} u_-(p) = \frac{1}{2} \tilde{\lambda}_{\dot{a}}(p) \bar{\sigma}_{\mu}^{\dot{a}a} \lambda_a(p) \quad (3)$$

and products of four-vectors can be performed using the identity

$$\bar{\sigma}^{\mu, \dot{a}a} \bar{\sigma}_{\mu}^{\dot{b}b} = 2 \epsilon^{\dot{a}b} \epsilon^{ab} \quad (4)$$

where $\bar{\sigma}^{\mu} = (\mathcal{I}, -\vec{\sigma})$, and similar identities for the σ 's.

Note that, since

$$\bar{\sigma}^{\mu, \dot{a}a} \sigma_{\mu, b\dot{b}} = 2 \delta_{\dot{b}}^{\dot{a}} \delta_b^a \quad (5)$$

we can write

$$(p \cdot \sigma)_{a\dot{a}} = p^{\mu} \sigma_{\mu, a\dot{a}} = \lambda_a(p) \tilde{\lambda}_{\dot{a}}(p) \quad (6)$$

Moreover, the usual conventions for spinor products⁵ will be adopted:

$$\langle ij \rangle \equiv \bar{u}_+(p_i) u_-(p_j) = \lambda_a(p_i) \lambda_b(p_j) \epsilon^{ab} \quad (7)$$

$$[ij] \equiv -\bar{u}_-(p_i) u_+(p_j) = \tilde{\lambda}_{\dot{a}}(p_i) \tilde{\lambda}_{\dot{b}}(p_j) \epsilon^{\dot{a}\dot{b}} \quad (8)$$

This results to

$$2p \cdot q = \langle pq \rangle [pq] \quad (9)$$

A fairly general definition of the external gluons' polarization vectors⁶

⁵Note the minus sign in the $[ij]$ definition with respect to the definition in [9].

⁶that corresponds to the light-like axial gauge

is (see [9],[10] for similar definitions.)

$$\begin{aligned}\epsilon_+^\mu(\mathbf{p}) &= \frac{\bar{\mathbf{u}}_+(\bar{\mathbf{p}})\gamma^\mu\mathbf{u}_+(\mathbf{p})}{\sqrt{2}\bar{\mathbf{u}}_+(\bar{\mathbf{p}})\mathbf{u}_-(\mathbf{p})} = \frac{\tilde{\lambda}_{\dot{a}}(\mathbf{p})\bar{\sigma}^{\mu,\dot{a}a}\lambda_a(\bar{\mathbf{p}})}{\sqrt{2}\langle\bar{\mathbf{p}}\mathbf{p}\rangle} \\ \epsilon_-^\mu(\mathbf{p}) &= \frac{\bar{\mathbf{u}}_-(\bar{\mathbf{p}})\gamma^\mu\mathbf{u}_-(\mathbf{p})}{\sqrt{2}\bar{\mathbf{u}}_-(\mathbf{p})\mathbf{u}_+(\bar{\mathbf{p}})} = -\frac{\tilde{\lambda}_{\dot{a}}(\bar{\mathbf{p}})\bar{\sigma}^{\mu,\dot{a}a}\lambda_a(\mathbf{p})}{\sqrt{2}[\mathbf{p}\bar{\mathbf{p}}]}\end{aligned}\quad (10)$$

where $\bar{\mathbf{p}}$ is an auxiliary, null four-vector that can be chosen at will for every gluon as long as it is not parallel to \mathbf{p} itself.

Note that one could write

$$\epsilon_{\dot{a}\dot{a}}^+(\mathbf{p}) \equiv \epsilon_\mu^+(\mathbf{p})\sigma_{\dot{a}\dot{a}}^\mu = \sqrt{2}\frac{\lambda_a(\bar{\mathbf{p}})\tilde{\lambda}_{\dot{a}}(\mathbf{p})}{\langle\bar{\mathbf{p}}\mathbf{p}\rangle}\quad (11)$$

$$\epsilon_{\dot{a}\dot{a}}^-(\mathbf{p}) \equiv \epsilon_\mu^-(\mathbf{p})\sigma_{\dot{a}\dot{a}}^\mu = -\sqrt{2}\frac{\lambda_a(\mathbf{p})\tilde{\lambda}_{\dot{a}}(\bar{\mathbf{p}})}{[\mathbf{p}\bar{\mathbf{p}}]}\quad (12)$$

and perform vector products using

$$\mathbf{p}_{\dot{a}\dot{a}}\mathbf{q}_{\dot{b}\dot{b}}\epsilon^{\dot{a}\dot{b}}\epsilon^{\dot{a}\dot{b}} = \langle\mathbf{p}\mathbf{q}\rangle[\mathbf{p}\mathbf{q}] = 2\mathbf{p}\cdot\mathbf{q}\quad (13)$$

It should be noted that nothing of what follows depends on the precise conventions we adopt for the Van der Waerden or Dirac spinors. In fact the present analysis, that deals with pure gluonic amplitudes, could be performed exclusively on the level of four-vectors.

The hat symbol over a function of momenta $\hat{f}(\mathbf{q})$ will denote the function $\hat{f}(\mathbf{q}; z)$ where the argument of f is analytically continued (shifted) by a general four-vector $z\epsilon_\mu$, with

$$\epsilon_\mu \equiv \frac{1}{2}\bar{\mathbf{u}}_-(\mathbf{p}_1)\gamma_\mu\mathbf{u}_-(\mathbf{p}_n) = \frac{1}{2}\tilde{\lambda}_{\dot{a}}(\mathbf{p}_1)\bar{\sigma}_\mu^{\dot{a}a}\lambda_a(\mathbf{p}_n)\quad (14)$$

This has the effect

$$\mathbf{q} \rightarrow \hat{\mathbf{q}} = \mathbf{q} + z\epsilon\quad (15)$$

We will omit the explicit reference to the z -dependence of $\hat{f}(\mathbf{p})$ since this is signified by the hat symbol.

3 Counting contributions

The BCFW recursive relation for pure gluon color ordered amplitudes is given by

$$\mathcal{A}(p_1^{\lambda_1}, \dots, p_n^{\lambda_n}) = \sum_{\lambda_x = \pm 1} \sum_{k=1}^{n-1} \mathcal{A}(\hat{p}_1^{\lambda_1}, \dots, p_k^{\lambda_k}, -\hat{p}_x^{-\lambda_x}) \frac{1}{P_{1\dots k}^2} \mathcal{A}(\hat{p}_x^{\lambda_x}, p_{k+1}^{\lambda_{k+1}}, \dots, \hat{p}_n^{\lambda_n}) \quad (16)$$

where p and λ represent momentum and polarization. In this section we would like to study the diagrammatic content of the BCFW equation. The most appropriate way to do so, is of course to neglect any reference to polarization and also drop the propagator factor, while keeping the ordering of the momenta. In that case of course we count planar graphs (see [12]) and it is obvious that the equation is not self-consistent. In fact the form of the equation is given by

$$\mathcal{A} = \sum \text{---} \bigcirc_{\mathcal{A}_I} \text{---} \bigcirc_{\mathcal{A}_{II}} \text{---} \quad (17)$$

or in a more mathematical form

$$\mathcal{A}^{BCF}{}_{1 \rightarrow n} = \sum_{k=1}^{n-1} \mathcal{A}_{1 \rightarrow k+1} \mathcal{A}_{1 \rightarrow n-k+1} \quad (18)$$

The equation $\mathcal{A}^{BCF}{}_{1 \rightarrow n} = \mathcal{A}_{1 \rightarrow n}$ cannot be true because of the following reasons:


- Firstly, contributions where no propagator line exists between the 1st and the n th particle, are not included in the BCFW equation

$$\mathcal{A}^0 = \text{---} \bigcirc \text{---} + \text{---} \bigcirc \bigcirc \text{---} \quad (19)$$


These contributions are given by

$$\mathcal{A}_{1 \rightarrow n}^0 = \mathcal{A}_{1 \rightarrow n-1} + \sum_{n_1+n_2=n-1} \mathcal{A}_{1 \rightarrow n_1} \mathcal{A}_{1 \rightarrow n_2} \quad (20)$$

- Secondly, BCFW is multiple-counting contributions of the form




or



In fact we suggest that this over counting is exactly equal to the multiplicity of propagator lines connecting particles 1 and n .

To make our arguments more quantitative we start with the Berends-Giele [11] (or Dyson-Schwinger for ordered graphs [13],[14]) recursive equation for a generic theory with 3- and 4-vertices.



$$\mathcal{A}_{1 \rightarrow n} = \sum_{n_1+n_2=n} \mathcal{A}_{1 \rightarrow n_1} \mathcal{A}_{1 \rightarrow n_2} + \sum_{n_1+n_2+n_3=n} \mathcal{A}_{1 \rightarrow n_1} \mathcal{A}_{1 \rightarrow n_2} \mathcal{A}_{1 \rightarrow n_3} \quad (22)$$

and we write the following equation

$$\mathcal{A}_{1 \rightarrow n} = \mathcal{A}^{BCF}_{1 \rightarrow n} + \mathcal{A}_{1 \rightarrow n}^0 - \mathcal{D} \quad (23)$$

where $\mathcal{A}_{1 \rightarrow n}^0$ are the classes of diagrams in eq.19 and the subtracted term \mathcal{D} accounts for the overcounting of diagrams. Since the overcounting relates to diagrams with 2, 3, ... propagators on the line between the first and the last leg, it can be seen that

$$\mathcal{D} = \sum_{M=3}^{n-1} (M-2) \sum_{k=0}^M \binom{M}{k} \mathcal{D}_{M+k}^n \quad (24)$$

where $M-1$ is the number of propagators of the particular overcounted class of diagrams and

$$\mathcal{D}_M^n = \sum_{n_1+\dots+n_M=n} \mathcal{A}_{1 \rightarrow n_1} \dots \mathcal{A}_{1 \rightarrow n_M} \quad (25)$$

counts the number of diagrams within that class. As an example for $n=5$ (6-leg gluon amplitude), the number of Feynman graphs is $\mathcal{A} = 38$, the

number of BCFW graphs is $\mathcal{A}^{BCF} = 29$, the number of graphs without any propagator between the first and the last leg is $\mathcal{A}^0 = 17$ so overcounting should give $\mathcal{D} = 8$. The following graphs are overcounted

(2)

(26)

with the first diagram doubly overcounted (it contains 3 propagators). That diagram corresponds to the $M = 4$ while the other graphs come from the $M = 3$ term of eq.24.

In the table below we give the results for up to $n = 11$ particles.

$1 \rightarrow n$	\mathcal{A}	\mathcal{A}^{BCF}	\mathcal{A}^0	$\mathcal{A}^{BCF} + \mathcal{A}^0 - \mathcal{A}$	\mathcal{D}
3	3	1	2	0	0
4	10	6	5	1	1
5	38	29	17	8	8
6	154	136	64	46	46
7	654	636	259	241	241
8	2871	2992	1098	1219	1219
9	12925	14190	4815	6080	6080
10	59345	67860	21659	30174	30174
11	276835	327080	99385	149630	149630

This analysis hints strongly towards the idea that a connection between Feynman diagrams and the BCFW decomposition might be achieved by grouping together BCFW (hatted) diagrams with the same chain structure along the main line (the same number of propagators hence the same multiplicity) but differently placed cuts: The multiplicity of each group is equal to the number of propagators along the main line which in turns equals the number of possible cuts (one for each propagator along the main line).

4 Kinematical identities

For any function $\hat{f}(\mathbf{p}; z)$ of z that has only simple poles in z and vanishes at $z \rightarrow \infty$, we can perform an expansion over its poles. In particular we have

$$\hat{f}(\mathbf{p}; z) = \sum_j \left[\hat{f}(\mathbf{p}; z)(z - z_j) \right]_{z=z_j} \frac{1}{z - z_j} \quad (27)$$

where the sum is over all the simple poles z_j of $\hat{f}(\mathbf{p}; z)$. This very general identity allows us to analytically continue the function $f(\mathbf{p})$ to the complex plane, make use of the pole expansion and take the limit $z \rightarrow 0$ to return to the real axis, thus obtaining a relation between $f(\mathbf{p})$ and the pole expansion of $\hat{f}(\mathbf{p}; z)$.

Applying this to a momentum antenna we get

$$\frac{1}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_k^2} = \sum_{j=1..k} \left[\frac{1}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_{j-1}^2 \hat{p}_{j+1} \dots \hat{p}_k^2} \right]_{z=z_j} \frac{1}{\hat{p}_j^2} \quad (28)$$

with z_j such that

$$\hat{p}_j(z_j)^2 = (\mathbf{p}_j + z_j \epsilon)^2 = \mathbf{p}_j^2 + 2z_j \mathbf{p}_j \cdot \epsilon = 0 \quad (29)$$

where $\mathbf{p}_1 \dots \mathbf{p}_k$ are arbitrary off-shell four-momenta. In the next sections we will use the above identity with \mathbf{p}_i being sums of on-shell momenta of the form $\mathbf{p}_{1k} = \mathbf{p}_1 + \mathbf{p}_2 + \dots + \mathbf{p}_k$.

Taking the limit $z \rightarrow 0$ at both sides we have

$$\frac{1}{\mathbf{p}_1^2 \mathbf{p}_2^2 \dots \mathbf{p}_k^2} = \sum_{j=1..k} \left[\frac{1}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_{j-1}^2 \hat{p}_{j+1}^2 \dots \hat{p}_k^2} \right]_{z=z_j} \frac{1}{\mathbf{p}_j^2} \quad (30)$$

Further more,

$$\frac{z^\rho}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_k^2} = \sum_{j=1..k} \left[\frac{z^\rho}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_{j-1}^2 \hat{p}_{j+1}^2 \dots \hat{p}_k^2} \right]_{z=z_j} \frac{1}{\hat{p}_j^2} \quad (31)$$

which gives the very useful set of identities, valid for every $\rho < k$

$$\sum_{j=1..k} \left[\frac{z^\rho}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_{j-1}^2 \hat{p}_{j+1} \dots \hat{p}_k^2} \right]_{z=z_j} \frac{1}{\mathbf{p}_j^2} = 0 \quad (32)$$

Finally, if $z = k$, the function in the left hand side of eq.31 is no longer vanishing at $z \rightarrow \infty$. Subtracting its limit at infinity, however, we have a new function that does, so

$$\frac{z^k}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_k^2} - \lim_{z \rightarrow \infty} \frac{z^k}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_k^2} = \sum_{j=1..k} \left[\frac{z^k}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_{j-1}^2 \hat{p}_{j+1} \dots \hat{p}_k^2} \right]_{z=z_j} \frac{1}{\hat{p}_j^2} \quad (33)$$

and taking the limit $z \rightarrow 0$ we get

$$-\lim_{z \rightarrow \infty} \frac{z^k}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_k^2} = \sum_{j=1..k} \left[\frac{z^k}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_{j-1}^2 \hat{p}_{j+1} \dots \hat{p}_k^2} \right]_{z=z_j} \frac{1}{p_j^2} \quad (34)$$

or

$$\frac{-1}{\prod_{j=1}^k 2\epsilon \cdot p_j} = \sum_{j=1..k} \left[\frac{z^k}{\hat{p}_1^2 \hat{p}_2^2 \dots \hat{p}_{j-1}^2 \hat{p}_{j+1} \dots \hat{p}_k^2} \right]_{z=z_j} \frac{1}{p_j^2} \quad (35)$$

5 Choosing a gauge

We have seen in section 3 that classes of Feynman diagrams should somehow correspond to particular BCFW decompositions. Using a particular gauge for the external gluons one can eliminate whole classes of Feynman diagrams.

A consistent definition of the external gluons' polarization vectors was given in eq. 10 where \bar{p} is an auxiliary, null four-vector that can be chosen at will for every gluon as long as it is not parallel to p itself.

We choose to use $\bar{p}_1 = p_n$ and $\bar{p}_n = p_1$

$$\epsilon_{+1}^\mu = \frac{\bar{u}_{+n} \gamma^\mu u_{+1}}{\sqrt{2} \bar{u}_{+n} u_{-1}} = \frac{\tilde{\lambda}_{\dot{a}}(p_1) \bar{\sigma}^{\mu, \dot{a}a} \lambda_a(p_n)}{\sqrt{2} \langle n1 \rangle} \quad (36)$$

$$\epsilon_{-n}^\mu = \frac{\bar{u}_{-1} \gamma^\mu u_{-n}}{\sqrt{2} \bar{u}_{-n} u_{+1}} = -\frac{\tilde{\lambda}_{\dot{a}}(p_1) \bar{\sigma}^{\mu, \dot{a}a} \lambda_a(p_n)}{\sqrt{2} [n1]} \quad (37)$$

A number of simplifications follow immediately. The product of ϵ_1 with ϵ_n or ϵ (defined in eq. 14) vanishes

$$\boxed{\epsilon_1 \cdot \epsilon_n = 0 = \epsilon \cdot \epsilon_1 = \epsilon \cdot \epsilon_n} \quad (38)$$

Moreover

$$\epsilon_{+1} \cdot p_n = 0 = \epsilon_{-n} \cdot p_1. \quad (39)$$

As a consequence, any diagram in which the first and the last leg meet in a three-vertex vanishes.

$$\begin{array}{c} \bullet \\ / \quad \backslash \\ 1 \quad n = 0 \end{array} \quad (40)$$

where the blob denotes any Feynman diagram with the particular off-shell leg.

These simplifications are in the correct direction in view of the fact that there are no BCFW graphs with the first and the last leg in the same vertex. We still have to deal with the case when 1 and n are attached in a four-vertex whose two other lines are off-shell (another situation that doesn't occur in a BCFW decomposition). We will see in the next section how to accommodate these diagrams.

Finally, we should note that the polarization vectors ϵ_{+1}^μ and ϵ_{-n}^μ actually differ by a complex phase. They also differ by a complex factor from the shifting vector ϵ (see 14). Note that in any expression where all these three vectors appear in scalar products, one is allowed to freely interchange them, without altering the result. In a very real sense there is actually only one polarization vector in this gauge.

Finally, the polarization vectors are now invariant under the shifting operation:

$$p_1 \rightarrow p_1 + z\epsilon \quad p_n \rightarrow p_n - z\epsilon \quad (41)$$

If we write

$$p_1^\mu = \frac{1}{2}\bar{u}_{-1}\gamma^\mu u_{-1} \rightarrow \frac{1}{2}\bar{u}_{-1}\gamma^\mu u_{-1} + \frac{1}{2}z\bar{u}_{-1}\gamma^\mu u_{-n} \quad (42)$$

and

$$p_n^\mu = \frac{1}{2}\bar{u}_{-n}\gamma^\mu u_{-n} \rightarrow \frac{1}{2}\bar{u}_{-n}\gamma^\mu u_{-n} + \frac{1}{2}z\bar{u}_{-1}\gamma^\mu u_{-n} \quad (43)$$

we see that the shifting operator effectively sends

$$\begin{aligned} u_{-1} &\rightarrow u_{-1} + zu_{-n} \\ \bar{u}_{-n} &\rightarrow \bar{u}_{-n} - z\bar{u}_{-1} \end{aligned} \quad (44)$$

or, in terms of Weyl - Van der Waerden spinors

$$\begin{aligned} \lambda_a(p_1) &\rightarrow \lambda_a(p_1) + z\lambda_a(p_n) \\ \tilde{\lambda}_{\dot{a}}(p_n) &\rightarrow \tilde{\lambda}_{\dot{a}}(p_n) - z\tilde{\lambda}_{\dot{a}}(p_1) \end{aligned} \quad (45)$$

As a result, the denominators of ϵ_{+1} and ϵ_{-n} become

$$\langle n1 \rangle \rightarrow \langle n1 \rangle + z\langle nn \rangle = \langle n1 \rangle \quad (46)$$

and

$$[n1] \rightarrow [n1] - z[11] = [n1] \quad (47)$$

The only restriction that we impose on the polarization vectors of the other gluons is that they remain invariant under the shifting operator. In case

their assisting vectors \bar{p}_i involve p_1 or p_n this should be arranged in a way that preserves this invariance.

This gauge choice explicitly eliminates any z -dependence from the polarization vectors. The shifting operation affects amplitudes only through the shift at the momenta four-vectors p_1 and p_n , and the induced shift to the momenta of intermediate propagators. Polarization vectors and vertices that do not carry p_1 and p_n are left unchanged. Therefore, a diagram after shifting will be a complex function with poles coming from three-vertices or propagators.

This is in contrast with a general gauge, where the polarization vectors ϵ_{+1} and ϵ_{-n} get an extra factor involving z in their denominator. A pole expansion, in that case, would need to take into account (gauge dependent) poles coming from the gluon polarization vectors.

Counting powers of z in the general gauge shows (see the concluding discussion in [7]) that any diagram vanishes at the limit $z \rightarrow \infty$. This is *not* in general true in the gauge we are working. The $z \rightarrow \infty$ limit of a diagram is used as a guide to group diagrams in classes where these limits cancel.


6 Review of the BCFW decomposition

According to the simplest version of the BCFW decomposition, the color amplitude $A(p_i; h_i)$, where h_i is the helicity of the i 'th gluon, is equal to

$$A(p_1^{h_1}, \dots, p_n^{h_n}) = \sum_{j=2}^{n-1} \sum_h A(\hat{p}_1^{h_1}, \dots, p_j^{h_j}, -\hat{p}_{1\dots j}^h) \frac{1}{p_{1\dots j}^2} A(\hat{p}_{1\dots j}^{-h}, p_{j+1}^{h_{j+1}}, \dots, \hat{p}_n^{h_n}) \quad (48)$$

The sum over j extends over all partitions of the $n - 2$ gluons in two groups. The amplitude is recovered from the sum of $n - 2$ decompositions in lower level amplitudes evaluated at a particular $z = z_j$ such that $\hat{P}_{1\dots j}(z_j)^2 = 0$.

Let us call 'hatted' diagrams graphs of the form



denoting the corresponding Feynman graph where we have multiplied by p_{1A}^2 , we have performed the momenta shift, evaluated it at some $z = z_i$ and

divided back by $p_{1\Lambda}^2$. The value z_i is defined by the demand that the ‘cut’ propagator, carrying momentum $\hat{p}_{1\Lambda}$, vanishes: $\hat{p}_{1\Lambda}^2 = 0$.

In terms of ‘hatted’ graphs, the BCFW decomposition consists in two blobs, one containing the first and one the last leg, the amplitude is hatted and a cut is taken on the propagator that connects the two blobs. A sum over all partitions of n in two integers is employed. Evidently, every particular BCFW decomposition over some propagator P_i is equivalent to the sum of all hatted diagrams with that particular propagator cut.


Thus the sum of all possible decompositions is equivalent to the sum of all hatted diagrams with all possible cuts along the line of propagators that connects the first and the last leg of the original color amplitude.

In what follows we will prove that the sum of all possible hatted diagrams is equivalent to the sum of Feynman diagrams involved in the computation of the particular amplitude.


7 The correspondence of hatted graphs to Feynman diagrams

The special treatment of the two selected external legs 1 and n implies a special classification of the various Feynman diagrams involved in the given amplitude. There are classes of diagrams with zero, one, two, ... propagators on the line between the first and the last leg. Every class consists of subclasses defined by the particular partition of the other $n - 2$ external legs in which they are connected to the main line.

A particular diagram with $n = 20$ would be, for example,



and would correspond to the class



and more precisely to its subclass with 3, 1, 2, 1, 5, 3, 3 on-shell legs in the corresponding blobs⁷. Let m_1, m_2, \dots, m_{k+1} be a particular partition of $n - 2$, such that $\sum m_i = n - 2$. The blobs denote the full set of all possible color-ordered diagrams with one off-shell and m_i on-shell legs.

Let us call this subclass of Feynman diagrams by $F(m_1, \dots, \tilde{m}_j, \tilde{m}_{j+1}, \dots, m_{k+1})$ where the tildes on top of two consecutive m 's mark the presence of a four-vertex attaching the corresponding blobs to the line.

$$F(m_1, \dots, \tilde{m}_j, \tilde{m}_{j+1}, \dots, m_{k+1}) = \frac{1}{\dots} \begin{array}{c} m_1 \quad \tilde{m}_j \quad \tilde{m}_{j+1} \quad m_n \\ \bullet \quad \bullet \quad \bullet \quad \bullet \\ | \quad \diagdown \quad \diagup \quad | \\ \dots \quad \dots \quad \dots \quad \dots \\ 1 \quad \quad \quad \quad n \end{array} \quad (52)$$

The particular subclass of diagrams in eq.51 would then be called $F(3, \tilde{1}, \tilde{2}, \tilde{1}, \tilde{5}, 3, 3)$ and eq.50 would be one of the actual Feynman diagrams contained in that subclass.

In the gauge we are working, the hatting operator would leave the contribution of the blobs, in such a diagram, invariant since the momentum flowing into the blob doesn't contain p_1 or p_n . It would only affect the propagators and vertices along the main line. After the shifting of the momenta a cut can be placed on any of the propagators on the line, thus fixing the value of z at which the residue should be evaluated. We shall denote such a graph with a cut on the j 'th propagator by

$$H(m_1, \dots, m_{k+1}; j) = \left[\hat{F}(m_1, \dots, m_{k+1}) \hat{p}_{1j}^2 \right]_{z=z_i} \frac{1}{p_{1j}^2} \quad (53)$$

Here and in what follows p_{1A_i} denotes the sum of external momenta

$$p_{1A_i} = p_1 + p_{A_1} + p_{A_2} + \dots + p_{A_i} \quad (54)$$

with

$$p_{A_i} = \sum_{k=1}^{m_i} p_k \quad (55)$$

A closer look at the "hatted" diagram will reveal that hatting affects, through the momenta p_1 and p_n , only the three-vertices along the main line. The four-vertices, not carrying any momentum, remain unchanged. The effect on the three-vertices is included in the "hatted" YM three-gluon

⁷A blob with m_i on-shell legs and one off-shell denotes the corresponding sum of color-ordered diagrams with one off-shell leg and m_i on-shell legs.

vertices, which we denote with a crossed white blob in the figures. Each of these can be decomposed in two pieces:

$$\begin{aligned}
\text{---} \circledast \text{---} &= \hat{V}_{\mu\nu\rho} = g_{\mu\nu}(\hat{p}_1 - p_2)_\rho + g_{\nu\rho}(p_2 - \hat{p}_3)_\mu + g_{\rho\mu}(\hat{p}_3 - \hat{p}_1)_\nu \\
&= g_{\mu\nu}(p_1 - p_2)_\rho + g_{\nu\rho}(p_2 - p_3)_\mu + g_{\rho\mu}(p_3 - p_1)_\nu \\
&\quad - z\epsilon^\sigma(2g_{\mu\rho}g_{\nu\sigma} - g_{\mu\sigma}g_{\nu\rho} - g_{\mu\nu}g_{\sigma\rho}) \\
&= V_{\mu\nu\rho} - z\epsilon^\sigma(2g_{\mu\rho}g_{\nu\sigma} - g_{\mu\sigma}g_{\nu\rho} - g_{\mu\nu}g_{\sigma\rho}) \\
&= \text{---} \bullet \text{---} + \text{---} \circ \text{---} \epsilon_\sigma
\end{aligned} \tag{56}$$

where

$$\text{---} \circ \text{---} \epsilon_\sigma = -zV_{\mu\nu\rho\sigma} \tag{57}$$

with $V_{\mu\nu\rho\sigma}$ the QCD four-vertex.

Let us concentrate for the moment on the class of diagrams with the smallest number of propagators along the line between the first and the last leg: diagrams with no propagators along the line can have a three or a four-vertex joining these two legs. The whole class of diagrams with a three-vertex joining the first and the last leg vanishes identically in the gauge we are using. The remaining class with one four-vertex will be dealt with soon.

Next, we have the class with one propagator along the main line.

If we write J_Q^μ for the current coming from the blob Q , we have

$$\begin{aligned}
H(m_1, m_2; 1) &\equiv \text{---} \bullet \text{---} \text{---} \bullet \text{---} \\
&= \left[\epsilon_{1\mu} J_{A\nu} \hat{V}^{\mu\nu\rho} \hat{V}_{\rho\kappa\lambda} J_B^K \epsilon_n^\lambda \right]_{z_0} \frac{1}{p_{1A}^2} \\
&= \left[\epsilon_{1\mu} J_{A\nu} (V^{\mu\nu\rho} - z\epsilon_\sigma V^{\sigma\mu\nu\rho}) (V_{\rho\kappa\lambda} - z\epsilon^\tau V_{\tau\rho\kappa\lambda}) J_B^K \epsilon_n^\lambda \right]_{z_0} \frac{1}{p_{1A}^2} \\
&= \left[\epsilon_{1\mu} J_{A\nu} V^{\mu\nu\rho} V_{\rho\kappa\lambda} J_B^K \epsilon_n^\lambda - z\epsilon_{1\mu} J_{1\nu} \epsilon_\sigma V^{\sigma\mu\nu\rho} V_{\rho\kappa\lambda} J_B^K \epsilon_n^\lambda \right. \\
&\quad \left. - z\epsilon_{1\mu} J_{A\nu} V^{\mu\nu\rho} \epsilon^\tau V_{\tau\rho\kappa\lambda} J_B^K \epsilon_n^\lambda \right. \\
&\quad \left. + z^2 \epsilon_{1\mu} J_{A\nu} \epsilon_\sigma V^{\sigma\mu\nu\rho} \epsilon^\tau V_{\tau\rho\kappa\lambda} J_B^K \epsilon_n^\lambda \right]_{z_0} \frac{1}{p_{1A}^2}
\end{aligned} \tag{58}$$

The first, z -independent term in the bracket equals the Feynman diagram with one propagator. The last term in the bracket vanishes due to eq.38 and we get

$$\begin{aligned}
H(m_1, m_2; 1) &\equiv \text{Diagram with blob A and B connected by a dashed line with a cross} = \\
&= \text{Diagram with blob A and B connected by a solid line} - [2(\epsilon \cdot p_{1A})(\epsilon_1 \cdot J_A)(\epsilon_n \cdot J_B)]_{z_0} \frac{1}{p_{1A}^2}
\end{aligned}$$

with

$$z_0 = -\frac{p_{1A}^2}{2(p_{1A} \epsilon)} \quad (59)$$

where we have used momentum conservation and our freedom to interchange the polarization vectors. So

$$\begin{aligned}
H(m_1, m_2; 1) &\equiv \text{Diagram with blob A and B connected by a dashed line with a cross} = \\
&= \text{Diagram with blob A and B connected by a solid line} + \text{Diagram with two blobs connected by a solid line}
\end{aligned} \quad (60)$$

or

$$H(m_1, m_2; 1) = F(m_1, m_2) + F(\tilde{m}_1, \tilde{m}_2) \quad (61)$$

In the case that the blobs A and B of the first diagram in eq.60 contain exactly one on-shell leg, the BCFW decomposition leads to two hatted QCD 3-vertices. We have verified that the usual QCD 3-vertex evaluated at the hatted kinematics reproduces the formulas

$$A(\hat{1}^+, 2^-, \hat{q}^+) = \sqrt{2} \frac{[\hat{q}\hat{1}]^3}{[\hat{1}2][2\hat{q}]} \quad (62)$$

and

$$A(\hat{1}^+, 2^+, \hat{q}^-) = \sqrt{2} \frac{[\hat{1}\hat{2}]^3}{[2\hat{q}][\hat{q}1]} \quad (63)$$

where $p_q = -p_1 - p_2$ is the off-shell leg and $\hat{p}_q = -p_1 - p_2 - z\epsilon$ is its on-shell continuation⁸. The corresponding formulas involving the last leg (which has negative helicity) work similarly but include angle bracket spinor products.

Proceeding to graphs that contain two propagators between the first and the last leg or graphs that contain one propagator with one four-vertex attached to it, we have

$$\begin{aligned}
 & \begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} + \begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} \quad (64) \\
 & + \begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} + \begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} \\
 = & \begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} + \begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} + \begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} \quad (65)
 \end{aligned}$$

The expressions that correspond to the first two hatted graphs on the left hand side are quadratic in z and are evaluated in different values of z , z_1 and z_2 .

$$\begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \circ \quad \circ \quad \circ \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} = \left[H_0 \frac{1}{\hat{p}_{1B}^2} + H_1 \frac{z}{\hat{p}_{1B}^2} + H_2 \frac{z^2}{\hat{p}_{1B}^2} \right]_{z=z_1} \frac{1}{p_{1A}^2} \quad (66)$$

⁸The other polarizations ($A(\hat{1}^+, 2^+, \hat{q}^+)$, $A(\hat{1}^+, 2^-, \hat{q}^-)$) can be shown to vanish.

with $p_{1B} = p_1 + p_A + p_B$

$$\begin{array}{c} \text{A} \quad \text{B} \quad \text{C} \\ \bullet \quad \bullet \quad \bullet \\ \diagdown \quad | \quad | \quad \diagup \\ \text{---} \quad \text{---} \quad \text{---} \\ \quad \quad \quad \times \end{array} = \left[H_0 \frac{1}{\hat{p}_{1A}^2} + H_1 \frac{z}{\hat{p}_{1A}^2} + H_2 \frac{z^2}{\hat{p}_{1A}^2} \right]_{z=z_2} \frac{1}{p_{1B}^2} \quad (67)$$

The z -independent term of all four graphs gives the Feynman graphs of the right hand side of eq.65, due to the kinematic identity eq.30.

Moreover, the part linear in z of the first two graphs cancels identically due to eq.32 with $\rho = 1$. The quadratic part of the first two graphs cancels exactly the part linear in z of the two last graphs on the left hand side of eq.65.

The generalization of the above mechanism goes as follows: Each hatted diagram with v_3 three-vertices along the main line can be decomposed in a sum of sub-diagrams having $0, 1, \dots, v_3$ white-blob vertices corresponding to $0, 1, \dots, v_3$ powers of z (evaluated at some z_i). If the diagram also has v_4 four-vertices, the number of propagators along the line is $v_3 + v_4 - 1$.

The sum over all possible cuts for the particular diagram can then be written as

$$\begin{aligned} & \sum_{j=1}^{v_3+v_4-1} H(m_1 \dots m_k; j) = \\ & = \sum_{j=1}^{v_3+v_4-1} \left[(H_0 + zH_1 + z^2H_2 + \dots + H_{v_3}z^{v_3}) \prod_{q \neq j} \frac{1}{\hat{p}_{1q}^2} \right]_{z=z_j} \frac{1}{p_{1j}^2} \end{aligned} \quad (68)$$

or

$$\begin{aligned} \sum_{j=1}^{v_3+v_4-1} H(m_1 \dots m_k; j) &= H_0 \sum_{j=1}^{v_3+v_4-1} \left[\frac{1}{\prod_{q \neq j} \hat{p}_{1q}^2} \right]_{z=z_j} \frac{1}{p_{1j}^2} \\ &+ H_1 \sum_{j=1}^{v_3+v_4-1} \left[\frac{z}{\prod_{q \neq j} \hat{p}_{1q}^2} \right]_{z=z_j} \frac{1}{p_{1j}^2} \\ &+ H_2 \sum_{j=1}^{v_3+v_4-1} \left[\frac{z^2}{\prod_{q \neq j} \hat{p}_{1q}^2} \right]_{z=z_j} \frac{1}{p_{1j}^2} \\ &+ \dots \end{aligned}$$

$$+H_{v_3} \sum_{j=1}^{v_3+v_4-1} \left[\frac{z^{v_3}}{\prod_{q \neq j} \hat{p}_{1q}^2} \right]_{z=z_j} \frac{1}{p_{1j}^2} \quad (69)$$

Due to the set of identities eq.32, all terms involving z^λ with $0 < \lambda < v_3 + v_4 - 1$ vanish identically.

Moreover it is easy to see that the z -independent term involving H_0 will give the corresponding Feynman diagram with the help of the identity eq.30.

Let us further distinguish among three cases: diagrams with no four-vertex on the line ($v_3 = k + 1$), with one four-vertex ($v_3 = k$) and with more than one four-vertex ($v_3 < k$). In the third case the right hand side of eq.69 reduces to the z -independent term corresponding to the Feynman diagram. All terms involving z vanish. In other words the sum over all possible cuts of the hatted diagrams with two or more four-vertices on the line, is equal to the corresponding Feynman diagrams.

In the first two cases the sum of all hatted diagrams over all possible cuts gives the corresponding Feynman diagrams plus a number of terms. We will now show that these terms cancel exactly each other due to the structure of the YM vertices.

Diagrams with no four vertex on the line have $v_3 = k + 1$ three-vertices. They are of the form

$$H(m_1, \dots, m_j, m_{j+1}, \dots, m_{k+1}; j) = \frac{\begin{array}{ccccccc} & m_1 & & m_j & m_{j+1} & & m_n \\ & \bullet & & \bullet & \bullet & & \bullet \\ & | & & | & | & & | \\ 1 & \dots & & \dots & \dots & & n \end{array}}{1 \dots n} \quad (70)$$

The highest order term involves H_{k+1} that vanishes identically: it corresponds to diagrams with only crossed white blobs on the line, hence it consists of contractions of ϵ with itself or ϵ_1, ϵ_n . The next to leading order term, H_k is a sum of terms with one QCD vertex and k crossed white-blob vertices.

We have

$$\sum_{j=1}^k H(m_1 \dots m_{k+1}; j) = F(m_1 \dots m_{k+1}) + H_k \sum_{j=1}^k \left[\frac{z^k}{\prod_{q \neq j} \hat{p}_{1q}^2} \right]_{z=z_j} \frac{1}{p_{1j}^2} \quad (71)$$

and using the kinematical identity eq.35 we get

$$\sum_{j=1}^k H(m_1 \dots m_{k+1}; j) = F(m_1 \dots m_{k+1}) - H_k \frac{1}{\prod_{j=1}^k 2\epsilon \cdot p_{1j}} \quad (72)$$

Let us denote $H_{k,r}$ the term where the QCD vertex is coming from the r 'th hatted vertex.

$H_{k,r}$ is the numerator of a diagram containing blobs connected with the main line by white-blob vertices $\epsilon^c M_{\mu\nu\rho c}$ everywhere except the QCD r 'th vertex. A generic piece on the line will contribute by

$$\begin{aligned} \dots \mu \quad \begin{array}{c} \bullet A_i \\ | \\ \circ \\ | \\ \epsilon \end{array} \quad \rho \dots &\equiv M(A_i)_{\mu\rho} = -J_{A_i}^\nu \epsilon^c (2g_{c\nu} g_{\mu\rho} - g_{c\mu} g_{\nu\rho} - g_{c\rho} g_{\mu\nu}) \\ &= -(2(J_{A_i} \cdot \epsilon) g_{\mu\rho} - J_{A_i, \mu} \epsilon_\rho - J_{A_i, \rho} \epsilon_\mu) \end{aligned} \quad (73)$$

where $J_{A_i}^\nu$ is the current (subamplitude) coming from the i 'th blob. The first such white-blob vertex, when contracted with ϵ_1^μ , gives

$$\begin{array}{c} \bullet A_1 \\ | \\ \circ \\ | \\ \epsilon \end{array} \quad \rho \dots = \epsilon_1^\mu M(A_1)_{\mu\rho} = -(\epsilon_1 \cdot J_{A_1}) \epsilon_\rho \quad (74)$$

It is easy to see that a chain of consecutive white-blob vertices gives

$$\begin{array}{c} \bullet A_1 \quad \bullet A_2 \quad \bullet A_s \\ | \quad | \quad | \\ \circ \quad \circ \quad \dots \quad \circ \\ | \quad | \quad | \\ \epsilon \quad \epsilon \quad \dots \quad \epsilon \end{array} \quad \rho = (-1)^\rho (\epsilon_1 \cdot J_{A_1}) (\epsilon \cdot J_{A_2}) \dots (\epsilon \cdot J_{A_s}) \epsilon_\rho \quad (75)$$

Since $H_{k,r}$ has k such vertices we get

$$\begin{aligned} H_{k,r} &= (-1)^k \prod_{j \neq r} (\epsilon \cdot J_{A_j}) \epsilon_1^\mu J_{A_r}^\nu \epsilon_n^\rho \times \\ &\quad \times (g_{\mu\nu} (p_{1,r-1} - p_r)_\rho + g_{\nu\rho} (p_r + p_{1,r})_\mu + g_{\rho\mu} (-p_{1,r} - p_{1,r-1})_\nu) \\ &= (-1)^k \prod_{q \neq r} (\epsilon \cdot J_{A_q}) (\epsilon_n \cdot J_{A_r}) ((p_{1,r-1} + p_{1,r}) \cdot \epsilon_1) \end{aligned} \quad (76)$$

where repeated use was made of our ability to interchange ϵ_1, ϵ_n and ϵ . Note that for the boundary terms, $r = 1$ and $r = k + 1$ the above formula must be understood with the definitions $p_{1,0} = p_{1,k+2} = 0$.

Summing over r we regain H_k , and

$$\begin{aligned}
\sum_{j=1}^k H(\mathbf{m}_1 \dots \mathbf{m}_{k+1}; j) &= F(\mathbf{m}_1 \dots \mathbf{m}_{k+1}) + \\
&+ (-1)^{k+1} \frac{1}{\prod_j 2\epsilon \cdot \mathbf{p}_{1j}} \sum_r (\epsilon_1 \cdot J_{\Lambda_1}) \dots (\epsilon_n \cdot J_{\Lambda_n}) [(\mathbf{p}_{1,r-1} + \mathbf{p}_{1,r}) \cdot \epsilon] \\
&= F(\mathbf{m}_1 \dots \mathbf{m}_k) + \\
&+ (-1)^{k+1} (\epsilon_1 \cdot J_{\Lambda_1}) \dots (\epsilon_n \cdot J_{\Lambda_n}) \sum_r \frac{2\mathbf{p}_{1,r} \cdot \epsilon}{(2\epsilon \cdot \mathbf{p}_{11})(2\epsilon \cdot \mathbf{p}_{12}) \dots (2\epsilon \cdot \mathbf{p}_{1k})} \\
&= F(\mathbf{m}_1 \dots \mathbf{m}_k) + \\
&(-1)^{k+1} (\epsilon_1 \cdot J_{\Lambda_1}) \dots (\epsilon_n \cdot J_{\Lambda_n}) \sum_r \frac{1}{\prod_{q \neq r} (2\epsilon \cdot \mathbf{p}_{1q})} \tag{77}
\end{aligned}$$

Let us consider now a diagram that could occur from the ones above by contracting a propagator, thereby merging two of the three-vertices, say the r 'th and the $r+1$ 'th in one four-vertex. Such a diagram has one four-vertex on the line and one propagator less, that is it has k vertices of which $k-1$ three-vertices and one four-vertex, as well as $k-1$ propagators.

$$H(\mathbf{m}_1, \dots, \tilde{\mathbf{m}}_r, \tilde{\mathbf{m}}_{r+1}, \dots, \mathbf{m}_{k+1}; j) = \frac{1}{\text{Diagram}} \tag{78}$$

The sum over all cuts of such hatted diagrams is

$$\begin{aligned}
\sum_{\substack{j=1 \\ j \neq r}}^k H_r(\mathbf{m}_1, \dots, \tilde{\mathbf{m}}_r, \tilde{\mathbf{m}}_{r+1}, \dots, \mathbf{m}_k; j) &= \\
&= F(\mathbf{m}_1, \dots, \tilde{\mathbf{m}}_r, \tilde{\mathbf{m}}_{r+1}, \dots, \mathbf{m}_k) + \\
&+ H_{k,r} \sum_{\substack{j=1 \\ j \neq r}}^k \left[\frac{z^{k-1}}{\prod_{q \neq j,r} \hat{p}_{1q}^2} \right]_{z=z_j} \frac{1}{p_{1j}^2} \tag{79}
\end{aligned}$$

where r denotes the position of the four-vertex. As before, all contributions involving z^λ for $0 < \lambda < k-1$ vanish due to eq.32. Using the identity eq.35

we can perform the j sum over cuts and get

$$\begin{aligned} \sum_{\substack{j=1 \\ j \neq r}}^k H_r(m_1, \dots, \tilde{m}_r, \tilde{m}_r, \dots, m_k; j) = \\ = F(m_1, \dots, \tilde{m}_r, \tilde{m}_r, \dots, m_k) - \tilde{H}_{k-1,r} \frac{1}{\prod_{q \neq r} 2\epsilon \cdot p_{1q}} \end{aligned} \quad (80)$$

Here $\tilde{H}_{k-1,r}$ stands for the vertex contribution of a diagram with one four vertex and $k - 1$ white-blob three-vertices. It is not difficult to see that this is equal to

$$\tilde{H}_{k-1,r} = (-1)^{k-1} (\epsilon_1 \cdot J_{A_1}) (\epsilon \cdot J_{A_2}) \dots (\epsilon_n \cdot J_{A_k}) \quad (81)$$

Summing up over all possible positions of the four vertex we get

$$\begin{aligned} \sum_{r=1}^k \sum_{\substack{j=1 \\ j \neq r}}^k H_r(m_1 \dots m_k; j) = F(m_1, \dots, \tilde{m}_r, \tilde{m}_r, \dots, m_k) + \\ + (-1)^k (\epsilon_1 \cdot J_{A_1}) \dots (\epsilon_n \cdot J_{A_n}) \sum_r \frac{1}{\prod_{q \neq r} 2\epsilon \cdot p_{1q}} \end{aligned} \quad (82)$$

which exactly cancels the last term of equation eq.77, thus completing the proof that the sum over all cuts of all possible hatted diagrams is equal to the sum of Feynman diagrams.

8 Remarks on generality

We have seen in the previous sections how the BCFW decomposition is related to Feynman diagrams. In fact, when working in the particular gauge that we have chosen, every Feynman diagram is broken in pieces ('hatted' graphs) each of which contains a cut in one of the propagators along the line connecting the two 'special' legs. Then, hatted graphs with the same cut are regrouped together in one decomposition, and the sum of decompositions gives back the whole color amplitude.

In the process of breaking up the Feynman graphs in hatted graphs, some extra terms are produced: those correspond to the $z \rightarrow \infty$ limit of the hatted graph. Thanks to the particular structure of the YM vertices, demonstrated in eq.56, these extra terms cancel exactly among graphs with no four-vertices

and graphs with one four-vertex and one propagator less. In the lowest case of one propagator along the line, the diagram with two three vertices gives an extra term that is exactly equal to the diagram with no propagator and one four-vertex. The latter is a diagram which, having no propagator along the line, didn't fit in an obvious way in any BCFW decomposition.

The absence of a Yang-Mills structure in a scalar theory is what makes a decomposition along the terms of BCFW cumbersome. There one would have to accommodate the diagram in eq.40 by adding an ad hoc term in the recurrence relation which would therefore be less elegant.

The gauge in which we are working explicitly eliminates the z -dependence from the polarization vectors. In other gauges the polarization vectors ϵ_1^+ and ϵ_n^- would be affected by the shift. This would complicate significantly the algebra, as further poles related to vanishing denominators of these polarization vectors will come to play.

If the identity eq.40 doesn't hold, for example, contributions from the class of diagrams in the left hand side of eq.40 would have to be cancelled by contributions from other classes. The appealing cancellations within subclasses of diagrams with a particular partitioning of the external legs would be lost. Keeping track of the terms in fragmenting and regrouping Feynman diagrams into BCFW decompositions would be much harder. Still one could prove the BCFW decomposition by performing the pole expansion on any Feynman diagram minus its $z \rightarrow \infty$ limit but the algebra would be particularly cumbersome.

It is becoming increasingly clear that the BCFW decomposition is a rearrangement of terms of Feynman diagrams within a color ordered gluonic amplitude. A similar impression will presumably occur from an approach of amplitudes with fermionic lines. Under the light of the diagrammatic proof presented here it is seen that the mechanism behind any apparent cancellations in the BCFW recursion formula does not (unfortunately) divulge a deeper principle or even a contingent effect in gluonic amplitudes, but results directly from the structure of the Yang-Mills vertices and the cancellations induced by gauge invariance.

9 Acknowledgments

Author P.D.D. is currently co-funded by the European Social Fund (75%) and National Resources (25%)-EPEAEK B! - PYTHAGORAS. The research of authors A.L. and C.G.P. has been funded by the EU program no.HPMD-

CT-2001-00105. The author C.G.P. also acknowledges HPC-EUROPA Programme for its support.

References

- [1] E.Witten,Commun.Math.Phys.252:189-258,2004 [hep-th/0312171]
- [2] R.Britto, F. Cachazo and B.Feng,Nucl.Phys.B715:499-522,2005 [hep-th/0412308]
- [3] G.Georgiou and V.V.Khoze, JHEP 0405:070 (2004) [hep-th/0404072]
G.Georgiou, E.W.N.Glover and V.V.Khoze, JHEP 0407:048 (2004) [hep-th/0407027]
D.A.Kosower, Phys.Rev.D71:045007 (2005) [hep-th/0406175]
- [4] J.B.Wu and C.J.Zhu,JHEP 0409:063 (2004) [hep-th/0406146]
J.B.Wu and C.J.Zhu,JHEP 0407:032 (2004) [hep-th/0406085]
C.J.Zhu,JHEP 0404:032 (2004)[hep-th/0403115]
- [5] L.J.Dixon, E.W.N.Glover and V.V.Khoze, JHEP 0412:015(2004)[hep-th/0411092]
Z.Bern, D.Forde, D.A.Kosower and P.Mastrogli, [hep-ph/0412167]
S.D.Badger, E.W.N.Glover and V.V.Khoze, JHEP 0503:023 (2005)[hep-th/0412275]
- [6] A.Brandhuber, B.Spence and G.Travaglini, Nucl.Phys.B706:150 (2005)[hep-th/0407214]

Z.bern,L.Dixon and D.A.Kosower, Phys.Rev.D71:105013,2005 [hep-th/0501240]
- [7] R.Britto, F. Cachazo,B.Feng and E.Witten, Phys.Rev.Lett.94:181602,2005 [hep-th/0501052]
- [8] F.A.Berends and W.Giele, Nucl.Phys. B294:700 (1987)
M.L.Mangano, S.J.Parke and Z.Xu, Nucl.Phys.B298:653(1988)
- [9] L.Dixon, Boulder TASI 95:539-584 [hep-ph/9601359]
- [10] R.Kleiss and W.J.Stirling, Nucl.Phys.B262:235-262(1985)
- [11] F.A.Berends and W.Giele, Nucl.Phys.B294:700(1987)
- [12] G. 't Hooft, Nucl. Phys. B **538** (1999) 389 [arXiv:hep-th/9808113].
- [13] E. N. Argyres, R. H. P. Kleiss and C. G. Papadopoulos, Nucl. Phys. B **391** (1993) 42.

- [14] P. Draggiotis, R. H. P. Kleiss and C. G. Papadopoulos, *Phys. Lett. B* **439** (1998) 157 [arXiv:hep-ph/9807207].