

SIMULTANEOUS UNIVERSALITY

L. BERNAL-GONZÁLEZ AND A. JUNG

ABSTRACT. In this paper, the notion of simultaneous universality is introduced, concerning operators having orbits that simultaneously approximate any given vector. This notion is related to the well known concepts of universality and disjoint universality. Several criteria are provided, and several applications to specific operators or sequences of operators are performed, mainly in the setting of sequence spaces or spaces of holomorphic functions.

1. INTRODUCTION

In this paper, we are concerned with the phenomenon of simultaneous approximation by the action of several operators or, more generally, by the action of several sequences of mappings. When the existence of a dense orbit under an operator is proved, we are speaking about universality or hypercyclicity, see below. In many situations, it is possible to show the existence of one vector whose orbits under two or more operators approximate any given vector. Pushing the question quite further, we wonder under what conditions such approximation takes place by using a *common* subsequence. This, together with its connection with other kinds of joint universality, will make up the main aim of the present manuscript.

Next, we fix some related notation and terminology to be used in this work. For a good account of concepts, results and history concerning hypercyclicity, the reader is referred to the books [2, 21].

By \mathbb{N} , \mathbb{N}_0 , \mathbb{R} , \mathbb{C} , \mathbb{D} , $B(a, r)$, $\overline{B}(a, r)$ ($a \in \mathbb{C}$, $r > 0$) we denote, respectively, the set of positive integers, the set $\mathbb{N} \cup \{0\}$, the real line, the complex plane, the open unit disk $\{z \in \mathbb{C} : |z| < 1\}$, the open disk with center a and radius r , and the corresponding closed disk. Let X, Y be two Hausdorff topological spaces, and $T_n : X \rightarrow Y$ ($n = 1, 2, \dots$) be a sequence of continuous mappings. Recall that (T_n) is said to be *universal* whenever there is some (T_n) -orbit which is dense in Y , that is, there exists an element $x_0 \in X$ –called universal for (T_n) – such that

$$\overline{\{T_n x_0 : n \in \mathbb{N}\}} = Y.$$

Note that Y must be separable. We denote by $\mathcal{U}((T_n))$ the set of universal elements for (T_n) . When $X = Y$ and $T : X \rightarrow X$ is a continuous self-mapping,

2010 *Mathematics Subject Classification.* 30E10, 47B33, 47A16, 47B38.

Key words and phrases. hypercyclic operator, composition operator, disjoint universality, simultaneous universality.

then T is called *universal* provided that the sequence (T^n) of iterates of T (i.e., $T^1 = T$, $T^2 = T \circ T$, $T^3 = T \circ T^2$, and so on) is universal, in which case the set $\mathcal{U}((T^n))$ of universal elements will be denoted by $\mathcal{U}(T)$. A sequence $T_n : X \rightarrow Y$ ($n = 1, 2, \dots$) of continuous mappings is said to be *densely universal* if $\mathcal{U}((T_n))$ is dense in X . Birkhoff's transitivity theorem asserts that, if X is a Baire space (in particular, if X is completely metrizable) and Y is second-countable (in particular, if X is metrizable and separable), then (T_n) is densely universal if and only if (T_n) is *transitive* (that is, given nonempty open sets $U \subset X$, $V \subset Y$, there is $N \in \mathbb{N}$ with $T_N(U) \cap V \neq \emptyset$); if this is the case, then $\mathcal{U}((T_n))$ is residual (in fact, a dense G_δ subset) in X . If X lacks isolated points and $T : X \rightarrow X$ is universal, then $\mathcal{U}(T)$ is dense in X (so residual if X is, in addition, completely metrizable).

In the case in which X and Y are topological vector spaces over \mathbb{K} ($= \mathbb{R}$ or \mathbb{C}) and $(T_n) \subset L(X, Y) := \{\text{linear continuous mappings } X \rightarrow Y\}$, the words *hypercyclic* and *universal* are synonymous, although *hypercyclic* is mostly used, as well as the alternative notation $HC((T_n)) := \mathcal{U}((T_n))$ (and $HC(T) := \mathcal{U}(T)$ for $T \in L(X) := L(X, X) = \{\text{operators on } X\}$). In particular, we have if X and Y are F-spaces with Y separable, then $HC((T_n))$ ($HC(T)$, with X separable, resp.) is residual in X as soon as (T_n) is transitive (as soon as T is hypercyclic, resp.). Recall that an F-space is a completely metrizable topological vector space.

Assume now that X, Y are topological spaces, with X a Baire space and Y second-countable, and that $S_n : X \rightarrow Y$ and $T_n : X \rightarrow Y$ ($n \in \mathbb{N}$) are densely universal sequences. Since $\mathcal{U}((S_n)), \mathcal{U}((T_n))$ are dense G_δ subsets of X , we have that $\mathcal{U}((S_n)) \cap \mathcal{U}((T_n))$ is also dense, so non-empty. Hence there is a common hypercyclic element $x \in X$. So, for a given point $y \in Y$, there are sequences $\{n_1 < n_2 < \dots\}$ and $\{m_1 < m_2 < \dots\}$ in \mathbb{N} such that

$$S_{n_j}x \rightarrow y \text{ and } T_{m_j}x \rightarrow y \text{ as } j \rightarrow \infty.$$

Then the following question arises naturally:

Under what conditions on (S_n) and (T_n) one can guarantee the existence of an element $x \in X$ such that, for any given $y \in Y$, there is one sequence $\{n_1 < n_2 < \dots\} \subset \mathbb{N}$ such that

$$S_{n_j}x \longrightarrow y \longleftarrow T_{n_j}x \text{ as } j \rightarrow \infty?$$

Of course, a similar question can be posed for finitely many sequences and for finitely many single operators on X , just by considering the sequences of their iterates in the latter case. With this in mind, the new concept of simultaneous universality will be introduced in the next section, and compared to other related notions existing in the literature, such as those of disjoint hypercyclicity and the weakly mixing property. Several sufficient conditions for simultaneous universality/hypercyclicity will be provided in Section 3. Examples of finite families of simultaneous hypercyclic operators will be furnished in sections 4–6, starting with

multiples of an operator and ending up in the frameworks of sequence spaces and of spaces of analytic functions on complex domains.

2. SIMULTANEOUSLY UNIVERSAL SEQUENCES

Let us define the new concept that is the matter of this paper. If $p \in \mathbb{N}$ and Y is a nonempty set, then by $\Delta(Y^p)$ we denote the diagonal of $Y^p = Y \times \cdots \times Y$ (p times), that is, the subset $\Delta(Y^p) = \{(y, y, \dots, y) : y \in Y\}$. If Y is a topological space, then Y^p is assumed to be endowed with the product topology.

Definition 2.1. Let $p \in \mathbb{N}$ and X, Y be Hausdorff topological spaces. Assume that, for each $j \in \{1, \dots, p\}$, $T_{j,n} : X \rightarrow Y$ ($n \in \mathbb{N}$) is a sequence of continuous mappings. Consider the sequence

$$[T_{1,n}, \dots, T_{p,n}] : x \in X \mapsto (T_{1,n}x, \dots, T_{p,n}x) \in Y^p \quad (n \in \mathbb{N}).$$

Let also $T_1, \dots, T_p : X \rightarrow X$ be continuous mappings.

- (a) We say that the sequences $(T_{1,n}), \dots, (T_{p,n})$ are *simultaneously universal* (or *s-universal*) whenever there exists an element $x_0 \in X$ –called *s-universal* for $(T_{1,n}), \dots, (T_{p,n})$ – satisfying

$$\overline{\{[T_{1,n}, \dots, T_{p,n}]x_0 : n \in \mathbb{N}\}} \supset \Delta(Y^p).$$

The set of such s-universal elements will be denoted by $s\mathcal{U}((T_{1,n}), \dots, (T_{p,n}))$.

- (b) The sequences $(T_{1,n}), \dots, (T_{p,n})$ are said to be *densely simultaneously universal* if the set $s\mathcal{U}((T_{1,n}), \dots, (T_{p,n}))$ is dense in X . And they are called *hereditarily simultaneously universal* (*hereditarily densely simultaneously universal*, resp.) if, for every strictly increasing sequence $(n_k) \subset \mathbb{N}$, the sequences $(T_{1,n_k}), \dots, (T_{p,n_k})$ are s-universal (densely s-universal, resp.).
- (c) The mappings T_1, \dots, T_p are called *s-universal* (*densely s-universal*, *hereditarily s-universal*, *hereditarily densely s-universal*, resp.) if the sequences $(T_1^n), \dots, (T_p^n)$ are s-universal (densely s-universal, hereditarily s-universal, hereditarily densely s-universal, resp.). The set $s\mathcal{U}((T_1^n), \dots, (T_p^n))$ of corresponding s-universal elements will be denoted by $s\mathcal{U}(T_1, \dots, T_p)$.

Remarks 2.2. 1. If Y is first-countable (in particular, if Y is metrizable), then the s-simultaneous universality of $(T_{j,n})_{n \in \mathbb{N}}$ ($1 \leq j \leq p$) means the existence of some $x_0 \in X$ enjoying the property that, for every $y \in Y$, there is a (strictly increasing) sequence $(n_k) \subset \mathbb{N}$ such that $T_{j,n_k}x_0 \rightarrow y$ as $k \rightarrow \infty$ ($j = 1, \dots, p$).

2. In [18, Kapitel 1] the notion of relative universality on a closed subset of the arrival space is introduced under very general assumptions. In the present paper we study a special case of this situation (note that $\Delta(Y^p)$ is closed in Y^p since Y^p is Hausdorff) under more specific hypotheses.

3. According to the introduction, if X, Y are topological vector spaces and $T_{j,n}, T_j \in L(X, Y)$ ($j = 1, \dots, p; n \in \mathbb{N}$), then we use the expressions “s-hypercyclic”,

“densely s-hypercyclic” and “hereditarily densely s-hypercyclic” rather than “s-universal”, “densely s-universal” and “hereditarily densely s-universal”, respectively. In addition, we will denote $s\text{-}HC((T_{1,n}), \dots, (T_{p,n})) := s\mathcal{U}((T_{1,n}), \dots, (T_{p,n}))$ and $s\text{-}HC(T_1, \dots, T_p) := s\mathcal{U}(T_1, \dots, T_p)$ in this case.

4. For a single operator T , hypercyclicity (hereditary hypercyclicity, resp.) is equivalent to dense hypercyclicity (hereditary dense hypercyclicity, resp.).

5. The property of simultaneous universality of $(T_{1,n}), \dots, (T_{p,n})$ is weaker than the property that the sequence $([T_{1,n}, \dots, T_{p,n}])$ is *subspace-universal* for $\Delta(Y^p)$, meaning that the set $\{[T_{1,n}, \dots, T_{p,n}]x_0 : n \in \mathbb{N}\} \cap \Delta(Y^p)$ is dense in $\Delta(Y^p)$ for some $x_0 \in X$ (see e.g. [1, 22, 24] for results on subspace-hypercyclicity/universality).

Before going on, we want to compare s-universality to other related concepts defined in the literature. In 2007, Bès, Peris and the first author ([11], [4]) introduced the notion of disjoint (or d-) universality (sometimes called d-hypercyclicity in the mentioned references). Under the same assumptions and terminology as in Definition 2.1, the sequences $(T_{1,n}), \dots, (T_{p,n})$ are said to be *d-universal* whenever the sequence $[T_{1,n}, \dots, T_{p,n}] : X \rightarrow Y^p$ ($n \in \mathbb{N}$) is universal, that is, whenever there exists some $x_0 \in X$ such that the joint orbit $\{(T_{1,n}x_0, \dots, T_{p,n}x_0) : n \in \mathbb{N}\}$ is dense in Y^p . As a matter of fact, d-universality should not be confused with the universality of the sequence

$$T_{1,n} \oplus \dots \oplus T_{p,n} : (x_1, \dots, x_p) \in X^p \longmapsto (T_{1,n}x_1, \dots, T_{p,n}x_p) \in Y^p.$$

Trivially, disjoint universality of $(T_{1,n}), \dots, (T_{p,n})$ implies universality of the last sequence as well as simultaneous universality of $(T_{1,n}), \dots, (T_{p,n})$. Also, trivially, s-universality implies the universality of each sequence $(T_{j,n})_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) (in particular, Y must be separable). But no other implications among these properties hold, even considering only $p = 2$ and sequences of iterates of single operators. The following examples illustrate this situation:

1. Assume that T is a hypercyclic operator on a topological vector space. Then the operators T, T are s-hypercyclic but not d-hypercyclic.
2. In 1969, S. Rolewicz [26] proved that if $c \in \mathbb{K}$ has modulus > 1 and B is the backward shift $(x_n) \in \ell_2 \mapsto (x_{n+1}) \in \ell_2$, then the operator cB is hypercyclic. In particular, the operators $T = 2B$ and $S = 4B = 2T$ are hypercyclic, but T, S are clearly not s-hypercyclic.
3. Since each of the operators T, S of the latter example is mixing (see the definition at the beginning of the next section, regarding the sequences of iterates; see also [21, p. 46]), the operator $T \oplus S$ is hypercyclic, but T, S are not s-hypercyclic.
4. De la Rosa and Read [15] were able to construct a Banach space X and an operator $T \in L(X)$ such that T is hypercyclic (hence T, T are s-hypercyclic) but T is not weakly mixing on X , meaning that $T \oplus T$ is not hypercyclic on X^2 .

While d-hypercyclic operators must be substantially different, s-hypercyclicity allows more similarity. For instance, an operator can never be d-hypercyclic with a scalar multiple of itself (see [11, p. 299]). Nevertheless, s-hypercyclicity is possible in concrete situations. This will be analyzed in Section 4. Sections 5 and 6 are devoted to more specific operators, namely backward shifts and operators on spaces of analytic functions.

We close this section by establishing, under appropriate assumptions, the existence of large vector subspaces consisting, except for zero, of s-hypercyclic vectors.

Theorem 2.3. (a) *Let X be a topological vector space and $T_j \in L(X)$ ($j = 1, \dots, p$). If T_1, \dots, T_p are s-hypercyclic and at least one of them commutes with the others, then $s\text{-}HC(T_1, \dots, T_p)$ contains, except for 0, a dense linear subspace of X .*

(b) *Let X and Y be two topological vector spaces such that Y is metrizable. Assume that $(T_{j,n}) \subset L(X, Y)$ ($j = 1, \dots, p$) are hereditarily s-hypercyclic sequences. Then $s\text{-}HC((T_{1,n}), \dots, (T_{p,n}))$ contains, except for 0, an infinite dimensional vector subspace of X .*

(c) *Let X and Y be two metrizable separable topological vector spaces. Assume that $(T_{j,n}) \subset L(X, Y)$ ($j = 1, \dots, p$) are hereditarily densely s-hypercyclic sequences. Then $s\text{-}HC((T_{1,n}), \dots, (T_{p,n}))$ contains, except for 0, a dense linear subspace of X .*

Proof. (a) By hypothesis, there is $i \in \{1, \dots, p\}$ such that $T_i T_j = T_j T_i$ ($j = 1, \dots, p$). Therefore $P(T_i)T_j = T_j P(T_i)$ for all j and every polynomial P with coefficients in \mathbb{K} . Let \mathcal{P} denote the set of such polynomials. Of course, the operator T_i is hypercyclic. From a result by Wengenroth [29], the operator $P(T_i)$ has dense range as soon as $P \in \mathcal{P} \setminus \{0\}$. Pick any $x_0 \in s\text{-}HC(T_1, \dots, T_p)$. Let us define $M := \{P(T_i)x_0 : P \in \mathcal{P} \setminus \{0\}\}$. Then M is a linear subspace of X . It is dense because M contains the orbit $\{T_i^n x_0 : n \in \mathbb{N}\}$, that is dense in X as $x_0 \in HC(T_i)$. It remains to show that $M \setminus \{0\} \subset s\text{-}HC(T_1, \dots, T_p)$.

To this end, fix $u \in M \setminus \{0\}$. Then there is $P \in \mathcal{P} \setminus \{0\}$ such that $u = P(T_i)x_0$. It must be proved that

$$Z \supset \Delta(X^p),$$

where $Z := \overline{\{(T_1^n u, \dots, T_p^n u) : n \in \mathbb{N}\}} = \overline{\{(P(T_i)T_1^n x_0, \dots, P(T_i)T_p^n x_0) : n \in \mathbb{N}\}}$, where the last equality follows from commutativity. We know that $\Delta(X^p) \subset \overline{\{(T_1^n x_0, \dots, T_p^n x_0) : n \in \mathbb{N}\}}$. Let $A := \{(T_1^n x_0, \dots, T_p^n x_0) : n \in \mathbb{N}\}$, $\varphi := P(T_i)$ and $\Phi : X^p \rightarrow X^p$ be the mapping defined as $\Phi(x_1, \dots, x_p) := (\varphi(x_1), \dots, \varphi(x_p))$. Then, as φ is continuous, we get

$$Z = \overline{\Phi(A)} \supset \Phi(\overline{A}) \supset \Phi(\Delta(X^p)) = \{(\varphi(x), \dots, \varphi(x)) : x \in X\},$$

so $Z \supset \overline{\{(\varphi(x), \dots, \varphi(x)) : x \in X\}}$. Given $y \in X$ and a neighborhood U of (y, y, \dots, y) , there exists a neighborhood V of y such that $U \supset V^p$. Since φ

has dense range, one can find $x \in X$ with $\varphi(x) \in V$. Then $(\varphi(x), \dots, \varphi(x)) \in U$. In other words, $(y, \dots, y) \in \overline{\{(\varphi(x), \dots, \varphi(x)) : x \in X\}}$, so $(y, \dots, y) \in Z$. Consequently, $Z \supset \Delta(X^p)$, as required.

(b)–(c). By mimicking the proofs of Theorems 1–2 of [3] (in which the results are given for a single sequence (T_n)), we can construct recursively a sequence $(x_N)_{N \in \mathbb{N}} \subset X$ and a family $\{(q(N, k))_{k \in \mathbb{N}} : N \in \mathbb{N}_0\}$ of strictly increasing subsequences of \mathbb{N} satisfying, for all $N \in \mathbb{N}$, the following conditions: $x_N \in G_N \cap s\text{-}HC((T_{1, q(N-1, k)}, \dots, T_{p, q(N-1, k)}))$ and $T_{j, q(l, k)} x_N \rightarrow 0$ as $k \rightarrow \infty$ for all $l \geq N$ and all $j \in \{1, \dots, p\}$, where $G_0 := X$ and $G_N := X \setminus \text{span}\{x_1, \dots, x_{N-1}\}$ ($N \in \mathbb{N}$) if the assumptions of (b) hold, while $\{G_N\}_{N \in \mathbb{N}}$ denotes any fixed open basis of X if the assumptions of (c) hold. Then $M := \text{span}\{x_N : N \in \mathbb{N}\}$ is the sought-after vector subspace. The details are left as an exercise. \square

3. S-UNIVERSALITY CRITERIA

A number of workable sufficient conditions will be useful to detect s-universality. Recall that a sequence of continuous mappings $T_n : X \rightarrow Y$ ($n \in \mathbb{N}$) is called *mixing* provided that, given nonempty open sets $U \subset X$, $V \subset Y$, there is $N \in \mathbb{N}$ such that $T_n(U) \cap V \neq \emptyset$ for all $n \geq N$. The corresponding notion of simultaneous mixing property arises naturally, as well as the one of simultaneous transitivity. Note that $T_n(U) \cap V \neq \emptyset$ is equivalent to $U \cap T_n^{-1}(V) \neq \emptyset$.

Definition 3.1. Let $p \in \mathbb{N}$ and X, Y be Hausdorff topological spaces. Assume that, for each $j \in \{1, \dots, p\}$, $T_{j, n} : X \rightarrow Y$ ($n \in \mathbb{N}$) is a sequence of continuous mappings. Let also $T_1, \dots, T_p : X \rightarrow X$ be continuous mappings. We say that:

- (a) The sequences $(T_{1, n}), \dots, (T_{p, n})$ are *simultaneously transitive* (or *s-transitive*) provided that, for every pair of nonempty open sets $U \subset X$, $V \subset Y$, there is $N \in \mathbb{N}$ such that $U \cap \bigcap_{j=1}^p T_{j, N}^{-1}(V) \neq \emptyset$.
- (b) The sequences $(T_{1, n}), \dots, (T_{p, n})$ are *simultaneously mixing* (or *s-mixing*) provided that, for every pair of nonempty open sets $U \subset X$, $V \subset Y$, there is $N \in \mathbb{N}$ such that $U \cap \bigcap_{j=1}^p T_{j, n}^{-1}(V) \neq \emptyset$ for all $n \geq N$.
- (c) The mappings T_1, \dots, T_p are *simultaneously transitive* (*simultaneously mixing*, resp.) whenever the sequences $(T_1^n), \dots, (T_p^n)$ are s-transitive (s-mixing, resp.).

Remark 3.2. Corresponding concepts of d-transitivity and d-mixing were introduced in [11], where $\bigcap_{j=1}^p T_{j, n}^{-1}(V_j)$ (V_j nonempty open subsets of Y , $j = 1, \dots, p$) appears instead of $\bigcap_{j=1}^p T_{j, n}^{-1}(V)$. Also, most criteria given in this section have their counterparts for the related d-properties as provided in [4] and [11]. A thorough study of d-mixing operators is provided in [8].

Note that, contrary to the one-sequence case, the facts $U \cap \bigcap_{j=1}^p T_{j, N}^{-1}(V) \neq \emptyset$ and $\bigcap_{j=1}^p T_{j, N}(U) \cap V \neq \emptyset$ are not equivalent. Observe also that $\bigcap_{j=1}^p T_{j, n}^{-1}(V) =$

$[T_{1,n}, \dots, T_{p,n}]^{-1}(V^p)$. From the definitions, it is easy to check that the sequences $(T_{1,n}), \dots, (T_{p,n})$ are s-mixing if and only if, for every strictly increasing sequence (n_k) in \mathbb{N} , the sequences $(T_{1,n_k}), \dots, (T_{p,n_k})$ are s-transitive. The following proposition provides what can be called the Birkhoff s-transitivity theorem.

Proposition 3.3. *Under the same assumptions and terminology as in Definition 3.1, let us suppose, in addition, that X is Baire and Y is second-countable. Then we have:*

- (i) *The sequences $(T_{1,n}), \dots, (T_{p,n})$ are s-transitive if and only if they are densely s-universal. If this is the case, then the set $s\mathcal{U}((T_{1,n}), \dots, (T_{p,n}))$ is residual in X .*
- (ii) *The sequences $(T_{1,n}), \dots, (T_{p,n})$ are s-mixing if and only if, for every strictly increasing sequence $(n_k) \subset \mathbb{N}$, the sequences $(T_{1,n_k}), \dots, (T_{p,n_k})$ are densely s-universal.*

Proof. Part (ii) is an immediate consequence of (i). Let us prove (i). Fix a countable open basis (V_m) of Y , as well as a point $x_0 \in X$. Then $x_0 \in s\mathcal{U}((T_{1,n}), \dots, (T_{p,n}))$ if and only if, given a nonempty open set $V \subset Y$, there is $n \in \mathbb{N}$ with $[T_{1,n}, \dots, T_{p,n}]x_0 \in V^p$, that is, $x_0 \in \bigcup_{n \in \mathbb{N}} \bigcap_{j=1}^p T_{j,n}^{-1}(V)$. Since each V contains some V_m and each V_m is a nonempty subset of Y , the last property is the same as $x_0 \in \bigcap_{m \in \mathbb{N}} \bigcup_{n \in \mathbb{N}} \bigcap_{j=1}^p T_{j,n}^{-1}(V_m)$, which shows that

$$s\mathcal{U}((T_{1,n}), \dots, (T_{p,n})) = \bigcap_{m \in \mathbb{N}} \bigcup_{n \in \mathbb{N}} \bigcap_{j=1}^p T_{j,n}^{-1}(V_m). \quad (1)$$

Since the $T_{j,n}$'s are continuous, each set $\bigcap_{j=1}^p T_{j,n}^{-1}(V_m)$ is open. If $(T_{1,n}), \dots, (T_{p,n})$ are s-transitive then every set $\bigcup_{n \in \mathbb{N}} \bigcap_{j=1}^p T_{j,n}^{-1}(V_m)$ ($m \in \mathbb{N}$) is (open and) dense. Hence their (countable) intersection, which equals $s\mathcal{U}((T_{1,n}), \dots, (T_{p,n}))$ by (1), is a dense G_δ subset (so residual) in X because X is Baire. Conversely, assume that the set of s-universal elements is dense in X and fix a nonempty open subset V of Y . Then there is $m \in \mathbb{N}$ with $V \supset V_m$. It follows from (1) that $\bigcup_{n \in \mathbb{N}} \bigcap_{j=1}^p T_{j,n}^{-1}(V_m)$ is dense in X , so the bigger set $\bigcup_{n \in \mathbb{N}} \bigcap_{j=1}^p T_{j,n}^{-1}(V)$ is also dense. But this means that, given a nonempty set $U \subset X$, there is $N \in \mathbb{N}$ such that $U \cap \bigcap_{j=1}^p T_{j,N}^{-1}(V) \neq \emptyset$ or, in other words, the sequences $(T_{1,n}), \dots, (T_{p,n})$ are s-transitive. \square

In the linear case, we state the following set of sufficient conditions, that are inspired by the results contained in [19, Sect. 1c] and the references cited in it.

Theorem 3.4. *Let X and Y be topological vector spaces such that X is Baire and Y is metrizable and separable, and let $(T_{j,n})_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) be sequences in $L(X, Y)$. Assume that there are respective dense subsets X_0 of X and Y_0 of Y satisfying at least one of the following conditions:*

- (A) For every pair of vectors $x \in X_0$, $y \in Y_0$, there exist sequences $(n_k) \subset \mathbb{N}$ and $(x_k) \subset X$ with $x_k \rightarrow 0$, $T_{j,n_k}x \rightarrow 0$ and $T_{j,n_k}x_k \rightarrow y$ ($j = 1, \dots, p$) as $k \rightarrow \infty$.
- (B) For every $x \in X_0$, the sequences $(T_{j,n}x)_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) converge in Y to a common limit and, for every $y \in Y_0$, there exist sequences $(n_k) \subset \mathbb{N}$ and $(x_k) \subset X$ with $x_k \rightarrow 0$ and $T_{j,n_k}x_k \rightarrow y$ ($j = 1, \dots, p$) as $k \rightarrow \infty$.
- (C) For every $x \in X_0$, there exists a sequence $(n_k) \subset \mathbb{N}$ such that the sequences $(T_{j,n_k}x)_{k \in \mathbb{N}}$ ($j = 1, \dots, p$) converge in Y to a common limit and, for every $y \in Y_0$, there exists a sequence $(x_n) \subset X$ such that $x_n \rightarrow 0$ and $T_{j,n}x_n \rightarrow y$ ($j = 1, \dots, p$) as $n \rightarrow \infty$.

Then $(T_{j,n})_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) are densely s -hypercyclic.

Proof. According to Proposition 3.3, we should show that $(T_{j,n})_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) are s -transitive. With this aim, fix a pair of nonempty open sets $U \subset X$, $V \subset Y$. We should exhibit an $N \in \mathbb{N}$ such that $U \cap \bigcap_{j=1}^p T_{j,N}^{-1}(V) \neq \emptyset$.

Assume first that (A) holds. By density, there are $x \in X_0$ and $y \in Y_0$ such that $x \in U$ and $y \in V$. Define $A := U - x$ and $B := V - y$. Then A and B are open neighborhoods of 0 in X and Y respectively. Take a 0-neighborhood $C \subset Y$ satisfying $C + C \subset B$. Consider the sequences (n_k) and (x_k) provided by (A). Then there is $k \in \mathbb{N}$ such that $x_k \in A$, $T_{j,n_k}x \in C$ and $T_{j,n_k}x_k \in y + C$ ($j = 1, \dots, p$). Let $u := x + x_k$ and $N := n_k$. We get $u \in x + A = U$ and $T_{j,N}u = T_{j,N}x + T_{j,N}x_k \in C + y + C \subset y + B = V$ ($j = 1, \dots, p$), so that $u \in U \cap \bigcap_{j=1}^p T_{j,N}^{-1}(V)$.

Suppose now that (B) holds. By density, there is $x \in X_0$ such that $x \in U$. Define $A := U - x$, a neighborhood of 0. By hypothesis, there is $z \in Y$ such that $T_{j,n} \rightarrow z$ as $n \rightarrow \infty$ ($j = 1, \dots, p$). Since Y_0 is dense in Y , there is $y \in Y_0$ with $y \in z + V$. Let $B := V - y + z$, a neighborhood of 0 in Y . Take a 0-neighborhood $C \subset Y$ satisfying $C + C \subset B$. We have that $T_{j,n} \in z + C$ ($j = 1, \dots, p$) for $n \geq n_0$, say. Consider the sequences (n_k) and (x_k) provided by (B) for the vector $y - z$, so that $x_k \rightarrow 0$ and $T_{j,n_k}x_k \rightarrow y - z$ ($j = 1, \dots, p$) as $k \rightarrow \infty$. Choose $k \in \mathbb{N}$ so large that $n_k \geq n_0$, $x_k \in A$ and $T_{j,n_k}x_k \in y - z + C$ ($j = 1, \dots, p$). Let $u := x + x_k$ and $N := n_k$. Then $u \in x + A = U$ and, for every $j = 1, \dots, p$,

$$T_{j,N}u = T_{j,N}x + T_{j,N}x_k \in z + C + y - z + C = y + C + C \subset y + B = V,$$

so that $u \in U \cap \bigcap_{j=1}^p T_{j,N}^{-1}(V)$, as required. Under assumption (C), the proof is similar and left as an exercise. \square

Two of the most popular criteria of hypercyclicity are the so-called blow-up/collapse criterion and the hypercyclicity criterion (see [2, 20, 21]). Now, we can obtain their respective s -versions.

Proposition 3.5. [s -Blow-up/Collapse Criterion] *Let X be a Baire metrizable separable topological vector space, and let $(T_{j,n})_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) be sequences*

in $L(X)$. Suppose that, for every nonempty open subsets U, V of X and every 0-neighborhood $W \subset X$ there is $N \in \mathbb{N}$ such that

$$W \cap \bigcap_{j=1}^p T_{j,N}^{-1}(V) \neq \emptyset \neq U \cap \bigcap_{j=1}^p T_{j,N}^{-1}(W).$$

Then $(T_{j,n})_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) are densely s -hypercyclic.

Proof. Fix a pair of nonempty open sets $U, V \subset X$. Choose vectors $x \in U, y \in V$. It suffices to exhibit sequences $(n_k) \subset \mathbb{N}$ and $(x_k) \subset X$ with $x_k \rightarrow x$ and $T_{j,n_k} x_k \rightarrow y$ ($j = 1, \dots, p$), because this would entail the existence of some $k \in \mathbb{N}$ such that $x_k \in U$ and $T_{j,n_k} x_k \in V$ ($j = 1, \dots, p$), so $x_k \in U \cap \bigcap_{j=1}^p T_{j,n_k}^{-1}(V)$. In other words, the sequences $(T_{j,n})_{n \in \mathbb{N}}$ ($j = 1, \dots, p$) would be s -transitive, hence densely s -hypercyclic by Proposition 3.3.

With this aim, choose a fundamental decreasing sequence (W_k) of 0-neighborhoods. Then $(U_k) := (x + W_k)$ and $(V_k) := (y + W_k)$ are fundamental decreasing sequences of x -neighborhoods and y -neighborhoods, respectively. By hypothesis, for each $k \in \mathbb{N}$, there are $n_k \in \mathbb{N}$ and points x'_k and x''_k such that $x'_k \in W_k \cap \bigcap_{j=1}^p T_{j,n_k}^{-1}(V_k)$ and $x''_k \in U_k \cap \bigcap_{j=1}^p T_{j,n_k}^{-1}(W_k)$. Let $x_k := x'_k + x''_k$. Then $x_k \rightarrow x$ as $k \rightarrow \infty$ because $x'_k \in W_k$ (so $x'_k \rightarrow 0$) and $x''_k \in U_k$ (so $x''_k \rightarrow x$). Finally, $T_{j,n_k} x_k = T_{j,n_k} x'_k + T_{j,n_k} x''_k \rightarrow y + 0 = y$ ($j = 1, \dots, p$) because $T_{j,n_k} x'_k \in V_k$ and $T_{j,n_k} x''_k \in W_k$ for all $k \in \mathbb{N}$. \square

Recall that the convex hull $\text{conv}(A)$ of a subset A of a vector space X is the least convex subset of X containing A .

Definition 3.6. Let X be a Baire metrizable separable locally convex space, $(n_k) \subset \mathbb{N}$ be a strictly increasing sequence and $T_j \in L(X)$ ($j = 1, \dots, p$). We say that T_1, \dots, T_p satisfy the s -hypercyclicity criterion with respect to (n_k) if there are subsets $X_0 \subset X, W_0 \subset X^p$ such that X_0 is dense in X and

$$\overline{W_0} \supset \Delta(X^p)$$

as well as mappings $R_k : W_0 \rightarrow X$ ($k \in \mathbb{N}$) such that

- (i) $T_j^{n_k} \rightarrow 0$ pointwise on X_0 as $k \rightarrow \infty$ ($j = 1, \dots, p$),
- (ii) $R_k \rightarrow 0$ pointwise on W_0 as $k \rightarrow \infty$ and
- (iii) For every $w = (w_1, \dots, w_p) \in W_0$ and every $j \in \{1, \dots, p\}$ there is $y_j \in \text{conv}(\{w_1, \dots, w_p\})$ such that $T_j^{n_k} R_k w \rightarrow y_j$ as $k \rightarrow \infty$.

Theorem 3.7. [s -Hypercyclicity Criterion] *Let X be a Baire metrizable separable locally convex space and $T_j \in L(X)$ ($j = 1, \dots, p$). If T_1, \dots, T_p satisfy the s -hypercyclicity criterion with respect to some $(n_k) \subset \mathbb{N}$, then $(T_1^{n_k}), \dots, (T_p^{n_k})$ are s -mixing. In particular, T_1, \dots, T_p are densely s -hypercyclic.*

Proof. Let $U, V \subset X$ be nonempty open sets. Then there are $x_0 \in U \cap X_0$ and $y_0 \in V$. By local convexity, there is a convex open set \tilde{V} with $y_0 \in \tilde{V} \subset V$. As

$(y_0, \dots, y_0) \in \Delta(X^p) \subset \overline{W_0}$, one can find $w = (w_1, \dots, w_p) \in W_0$ such that $w_j \in \tilde{V}$ for all $j = 1, \dots, p$. Put

$$z_k := x_0 + R_k w \quad (k \in \mathbb{N}).$$

Then, due to (ii), $z_k \rightarrow x_0 + 0 = x_0 \in U$ as $k \rightarrow \infty$. Moreover, for every $j \in \{1, \dots, p\}$ we get thanks to (i) and (iii) that

$$T_j^{n_k} z_k = T_j^{n_k} x_0 + T_j^{n_k} R_k w \longrightarrow 0 + y_j = y_j \quad \text{as } k \rightarrow \infty,$$

where, for each j , $y_j \in \text{conv}(\{w_1, \dots, w_p\}) \subset \text{conv}(\tilde{V}) = \tilde{V} \subset V$. Consequently, there is $k_0 \in \mathbb{N}$ such that, for all $k \geq k_0$, we have $z_k \in U$ and $T_j^{n_k} z_k \in V$ ($j = 1, \dots, p$) or, in other words, $z_k \in U \cap \bigcap_{j=1}^p T_j^{-n_k}(V) \neq \emptyset$, as required. \square

Remarks 3.8. 1. Examples of spaces X satisfying the assumptions of Theorem 3.7 are the Fréchet spaces, that is, the locally convex F-spaces. If local convexity is dropped from the assumptions, then the conclusion still holds if we replace (iii) by the (stronger) condition:

(iii') $T_j^{n_k} R_k w \rightarrow w_j$ as $k \rightarrow \infty$ for every $w = (w_1, \dots, w_p) \in W_0$ and every $j \in \{1, \dots, p\}$.

2. In [11, Proposition 2.6] the following *d-hypercyclicity criterion* was proved, where X is a Fréchet space, $(n_k) \subset \mathbb{N}$ is a strictly increasing sequence and $T_j \in L(X)$ ($j = 1, \dots, p$): Assume that there exist dense subsets $X_0, X_1, \dots, X_p \subset X$ and mappings $S_{k,j} : X_j \rightarrow X$ ($k \in \mathbb{N}; 1 \leq j \leq p$) satisfying $T_j^{n_k} \rightarrow 0$ ($k \rightarrow \infty$) pointwise on X_0 , $S_{k,j} \rightarrow 0$ ($k \rightarrow \infty$) pointwise on X_j , and $T_j^{n_k} S_{k,l} \rightarrow \delta_{j,l} \text{id}_{X_l}$ ($k \rightarrow \infty$) pointwise on X_l ($1 \leq j, l \leq p$). Then $(T_1^{n_k}), \dots, (T_p^{n_k})$ are d-mixing (see [11, Definition 2.1]). In particular, by [11, Proposition 2.3], T_1, \dots, T_p are densely d-hypercyclic.

Now, we can obtain a disjoint hypercyclicity criterion under weaker assumptions. Namely, let us assume that there are dense subsets $X_0 \subset X$, $W_0 \subset X^p$ and mappings $R_k : W_0 \rightarrow X$ ($k \in \mathbb{N}$) satisfying (i)–(ii) of Definition 3.6 together with (iii') of the preceding remark (it is easy to check that these assumptions are weaker than those of the d-hypercyclicity criterion in [11]). Then $(T_1^{n_k}), \dots, (T_p^{n_k})$ are d-mixing. Indeed, let $U, V_1, \dots, V_p \subset X$ be nonempty open sets. By density, there are $x_0 \in U \cap X_0$ and $w = (w_1, \dots, w_p) \in W_0 \cap (V_1 \times \dots \times V_p)$. Let $z_k := x_0 + R_k w$ ($k \in \mathbb{N}$). Then $z_k \rightarrow x_0 + 0 = x_0 \in U$ as $k \rightarrow \infty$. Moreover, for every $j \in \{1, \dots, p\}$ we get $T_j^{n_k} z_k = T_j^{n_k} x_0 + T_j^{n_k} R_k w \longrightarrow 0 + w_j = w_j$ as $k \rightarrow \infty$. Consequently, there is $k_0 \in \mathbb{N}$ such that, for all $k \geq k_0$, we have $z_k \in U$ and $T_j^{n_k} z_k \in V_j$ ($j = 1, \dots, p$), that is, $z_k \in U \cap \bigcap_{j=1}^p T_j^{-n_k}(V_j) \neq \emptyset$, which is the d-mixing property.

3. Several sets of conditions on T_1, \dots, T_p such that these operators satisfy the s-hypercyclicity criterion with respect to a strictly increasing sequence $(n_k) \subset \mathbb{N}$ are –as it is easy to check– the following:

- (a) There are dense subsets $X_0, Y_0 \subset X$ and mappings $S_{k,j} : Y_0 \rightarrow X$ ($k \in \mathbb{N}; 1 \leq j \leq p$) such that (i) holds, $\sum_{j=1}^p S_{k,j} \rightarrow 0$ pointwise on Y_0 and $T_j^{n_k} \sum_{l=1}^p S_{k,l} \rightarrow \text{id}_{Y_0}$ pointwise on Y_0 ($j = 1, \dots, p$).
- (b) There are subsets $X_0, X_1, \dots, X_p \subset X$ in such a way that X_0 is dense in X and $\overline{X_1 \times \dots \times X_p} \supset \Delta(X^p)$ as well as mappings $S_{k,j} : X_j \rightarrow X$ ($k \in \mathbb{N}; 1 \leq j \leq p$) such that (i) holds, $\sum_{j=1}^p S_{k,j} x_j \rightarrow 0$ for all $(x_1, \dots, x_p) \in X_1 \times \dots \times X_p$, and $T_j^{n_k} (\sum_{l=1}^p S_{k,l} x_l) \rightarrow x_j$ for all $(x_1, \dots, x_p) \in X_1 \times \dots \times X_p$ and all $j = 1, \dots, p$.

In view of (b), we see that if T_1, \dots, T_p satisfy the d-hypercyclicity criterion with respect to (n_k) , then they also satisfy the s-hypercyclicity criterion with respect to (n_k) .

Bès and Peris [10] have proved that satisfaction of the hypercyclicity criterion, hereditary hypercyclicity and transitivity of self-sums are equivalent (see also [5]). Moreover, they established a similar result for d-hypercyclicity [11, Theorem 2.7]. Now, we prove that a corresponding statement also holds for s-hypercyclicity, with the d-hypercyclicity criterion replaced by the s-hypercyclicity criterion (Theorem 3.7), so showing that the latter is rather natural.

Proposition 3.9. *Let X be a separable Fréchet space and $T_j \in L(X)$ ($j = 1, \dots, p$). Consider the following statements:*

- (a) T_1, \dots, T_p satisfy the s-hypercyclicity criterion.
 (b) $(T_1^{n_k}), \dots, (T_p^{n_k})$ are hereditarily densely s-hypercyclic for some $(n_k) \subset \mathbb{N}$.
 (c) $\bigoplus_{k=1}^m T_1, \dots, \bigoplus_{k=1}^m T_p$ are s-transitive on X^m for all $m \in \mathbb{N}$.
 (d) $T_1 \oplus T_1, \dots, T_p \oplus T_p$ are s-transitive on X^2 .

Then we have:

- (A) (a), (b) and (c) are equivalent.
 (B) If there exists $i \in \{1, \dots, p\}$ such that $T_i T_j = T_j T_i$ for all $j \in \{1, \dots, p\}$, then (a), (b), (c) are equivalent to (d).

Proof. In the proof of (A), we follow closely the proof of Theorem 2.7 in [11], while the proof of (B) runs similar as the proof of Theorem 2.3, (3) \Rightarrow (1), in [10].

(A) (a) \Rightarrow (b): T_1, \dots, T_p satisfy the s-hypercyclicity criterion with respect to some $(n_k) \subset \mathbb{N}$, so that they also satisfy it for any subsequence (m_k) of (n_k) . By Theorem 3.7, $(T_1^{m_k}), \dots, (T_p^{m_k})$ are s-mixing and therefore densely s-hypercyclic.

(b) \Rightarrow (c): Let $m \in \mathbb{N}$ be fixed and let $\emptyset \neq U_l, V_l \subset X$ open ($l = 1, \dots, m$). It suffices to show that there exists $N \in \mathbb{N}$ such that

$$U_l \cap \bigcap_{j=1}^p T_j^{-N}(V_l) \neq \emptyset \text{ for all } l = 1, \dots, m. \quad (1)$$

Since $(T_1^{m_k}), \dots, (T_p^{m_k})$ are densely s-hypercyclic for each subsequence (m_k) of (n_k) , the sequences $(T_1^{n_k}), \dots, (T_p^{n_k})$ are s-mixing (cf. Proposition 3.3(ii)). Hence, for

each $l \in \{1, \dots, m\}$, there exists $k_0(l) \in \mathbb{N}$ such that $U_l \cap \bigcap_{j=1}^p T_j^{-n_k}(V_l) \neq \emptyset$ for all $k \geq k_0(l)$. Then (1) is satisfied simply by choosing $N := \max\{k_0(1), \dots, k_0(m)\}$.

(c) \Rightarrow (a): Due to the assumption, we have:

(*) For every $m \in \mathbb{N}$ and every $2m$ -tuple $U_1, \dots, U_m, V_1, \dots, V_m$ of nonempty open subsets of X there is $N \in \mathbb{N}$ arbitrarily large such that (1) holds.

Let $(A_n)_{n \in \mathbb{N}}, (B_n)_{n \in \mathbb{N}}$ be bases of nonempty sets of the topology of X . For $n \in \mathbb{N}$, we write $W_n := B(0, 1/n)$ (open d -balls, d being a translation-invariant distance generating the topology of X) and $A_{n,0} := A_n, B_{n,0} := B_n$.

Choose a nonempty open set $A_{1,1}$ with $\text{diam}(A_{1,1}) < 1/2$ and $\overline{A_{1,1}} \subset A_1$. Due to (*) (with $m = 2$), there is $n_1 \in \mathbb{N}$ such that $B_1 \cap \bigcap_{j=1}^p T_j^{-n_1}(W_1) \neq \emptyset$ and $W_1 \cap \bigcap_{j=1}^p T_j^{-n_1}(A_{1,1}) \neq \emptyset$. Thus, there exist a nonempty open set $B_{1,1}$ with $\text{diam}(B_{1,1}) < 1/2$, $\overline{B_{1,1}} \subset B_1$ and $T_j^{n_1}(\overline{B_{1,1}}) \subset W_1$ for all $j = 1, \dots, p$, as well as a point $w_{1,1} \in W_1$ with $T_j^{n_1}w_{1,1} \in A_{1,1}$ for all $j = 1, \dots, p$. Now, for $i = 1, 2$, choose $\overline{A_{i,3-i}}$ open, nonempty, such that $\text{diam}(A_{i,3-i}) < 1/3$, $\overline{A_{i,3-i}} \subset A_{i,2-i}$ and $\overline{A_{1,2}} \cap \overline{A_{2,1}} = \emptyset$. Due to (*) (with $m = 4$), there is $n_2 \in \mathbb{N}$ with $n_2 > n_1$ such that $B_{i,2-i} \cap \bigcap_{j=1}^p T_j^{-n_2}(W_2) \neq \emptyset$ and $W_2 \cap \bigcap_{j=1}^p T_j^{-n_2}(A_{i,3-i}) \neq \emptyset$ for $i = 1, 2$. Thus, there exist nonempty open sets $B_{i,3-i}$ with $\text{diam}(B_{i,3-i}) < 1/3$, $\overline{B_{i,3-i}} \subset B_{i,2-i}$ and $T_j^{n_2}(\overline{B_{i,3-i}}) \subset W_2$ ($i = 1, 2$) for all $j = 1, \dots, p$ as well as points $w_{i,3-i} \in W_2$ ($i = 1, 2$) with $T_j^{n_2}w_{i,3-i} \in A_{i,3-i}$ ($i = 1, 2$) for all $j = 1, \dots, p$.

Continuing this process inductively, by using (*) with $m = 2k$ in step k , we obtain a strictly increasing sequence $(n_k) \subset \mathbb{N}$, nonempty open sets $A_{i,k+1-i}, B_{i,k+1-i}$ with $\text{diam}(A_{i,k+1-i}) < \frac{1}{k+1}$, $\text{diam}(B_{i,k+1-i}) < \frac{1}{k+1}$ and points $w_{i,k+1-i} \in W_k$ ($1 \leq i \leq k; k \in \mathbb{N}$) such that

- (i) $\overline{A_{i,k+1-i}} \subset A_{i,k-i}, \overline{B_{i,k+1-i}} \subset B_{i,k-i}$ for all $1 \leq i \leq k, k \in \mathbb{N}$.
- (ii) For each $k \in \mathbb{N}$, the sets $\overline{A_{i,k+1-i}}, 1 \leq i \leq k$, are pairwise disjoint.
- (iii) $T_j^{n_k}(\overline{B_{i,k+1-i}}) \subset W_k$ ($k \in \mathbb{N}; 1 \leq i \leq k; 1 \leq j \leq p$), and
- (iv) $T_j^{n_k}w_{i,k+1-i} \in A_{i,k+1-i}$ ($k \in \mathbb{N}; 1 \leq i \leq k; 1 \leq j \leq p$).

For each fixed $i \in \mathbb{N}$, the sequences of closed sets $(\overline{A_{i,r}})_{r \in \mathbb{N}}$ and $(\overline{B_{i,r}})_{r \in \mathbb{N}}$ are decreasing (due to (i)) with $\text{diam}(\overline{A_{i,r}}, \overline{B_{i,r}}) < \frac{1}{r+i}$. The completeness of X implies the existence of points $a_i, b_i \in X$ ($i \in \mathbb{N}$) such that $\bigcap_{r \in \mathbb{N}} \overline{A_{i,r}} = \{a_i\}$ and $\bigcap_{r \in \mathbb{N}} \overline{B_{i,r}} = \{b_i\}$.

Put $X_0 := \{b_i : i \in \mathbb{N}\} \subset X$ and $W_0 := \{a_i : i \in \mathbb{N}\}^p \subset X^p$. As $a_i \in \overline{A_{i,1}} \subset A_{i,0} = A_i$ and $b_i \in \overline{B_{i,1}} \subset B_{i,0} = B_i$ for all $i \in \mathbb{N}$, we obtain that X_0 is dense in X and W_0 is dense in X^p . Due to (ii), we have that $a_i \neq a_k$ whenever $i \neq k$ (indeed, if $i < k$, say, then $a_i \in \overline{A_{i,k+1-i}}$ and $a_k \in \overline{A_{k,1}}$, but $\overline{A_{i,k+1-i}} \cap \overline{A_{k,1}} = \emptyset$). Hence,

for each $k \in \mathbb{N}$, the function $R_k : W_0 \rightarrow X$ given by

$$R_k(a_{i_1}, \dots, a_{i_p}) = \begin{cases} \frac{1}{p} \sum_{l=1}^p w_{i_l, k+1-i_l} & \text{if } k \geq \max_{l=1, \dots, p} i_l \\ 0 & \text{otherwise} \end{cases}$$

is well defined. Altogether, we have:

- For all $j = 1, \dots, p$, all $i \in \mathbb{N}$ and all $k \geq i$ one has, due to (iii), that $T_j^{n_k} b_i \in T_j^{n_k} (\overline{B_{i, k+1-i}}) \subset W_k = B(0, 1/k)$, so $T_j^{n_k} \rightarrow 0$ ($k \rightarrow \infty$) pointwise on X_0 for every $j = 1, \dots, p$.
- For every $(a_{i_1}, \dots, a_{i_p}) \in W_0$ and every $k \geq \max_{l=1, \dots, p} i_l$, one has $R_k(a_{i_1}, \dots, a_{i_p}) = \frac{1}{p} \sum_{l=1}^p w_{i_l, k+1-i_l} \rightarrow 0$ ($k \rightarrow \infty$), because $w_{i_l, k+1-i_l} \in W_k = B(0, 1/k)$. Therefore $R_k \rightarrow 0$ ($k \rightarrow \infty$) pointwise on W_0 .
- For all $j = 1, \dots, p$, all $(a_{i_1}, \dots, a_{i_p}) \in W_0$ and all $k \geq \max_{l=1, \dots, p} i_l$, we get $T_j^{n_k} R_k(a_{i_1}, \dots, a_{i_p}) = \frac{1}{p} \sum_{l=1}^p T_j^{n_k} w_{i_l, k+1-i_l}$. Since $T_j^{n_k} w_{i_l, k+1-i_l} \in A_{i_l, k+1-i_l}$ and the sequence of sets $A_{i_l, k+1-i_l}$ ($k \in \mathbb{N}$) collapses to the singleton $\{a_{i_l}\}$ as $k \rightarrow \infty$ for each l , we get $T_j^{n_k} R_k(a_{i_1}, \dots, a_{i_p}) \rightarrow \frac{1}{p} \sum_{l=1}^p a_{i_l} \in \text{conv}(\{a_{i_1}, \dots, a_{i_p}\})$ as $k \rightarrow \infty$.

Thus, T_1, \dots, T_p satisfy the s-hypercyclicity criterion with respect to (n_k) . The proof of (A) is finished.

(B) Obviously, (c) always implies (d). Assume now that (d) holds and that some T_i commutes with all T_j 's. Our goal is to prove that (a) is satisfied.

Let us fix any vector $(x_0, y_0) \in s\text{-}HC(T_1 \oplus T_1, \dots, T_p \oplus T_p)$. We claim that, for each $m \in \mathbb{N}$, the vector $(x_0, T_i^m y_0)$ is also s-hypercyclic for $T_1 \oplus T_1, \dots, T_p \oplus T_p$. Indeed, as T_i is hypercyclic, it has dense range, from which one obtains, inductively, that every set $T_i^m(X)$ is dense in X . Put $A := X \times T_i^m(X)$, so that A is dense in X^2 . Given $(u, v) \in A$ there is $w \in X$ such that $v = T_i^m w$. By s-hypercyclicity, there exists $(n_k) \subset \mathbb{N}$ such that $T_j^{n_k} x_0 \rightarrow u$ and $T_j^{n_k} y_0 \rightarrow w$ ($k \rightarrow \infty$) for all $j = 1, \dots, p$. Hence, for all j , $T_j^{n_k} x_0 \rightarrow u$ and, by commutativity together with continuity of T_i^m , we get $T_j^{n_k} (T_i^m y_0) = T_i^m (T_j^{n_k} y_0) \rightarrow T_i^m w = v$ ($k \rightarrow \infty$). Therefore $\Sigma := \overline{\{(T_1 \oplus T_1)^n, \dots, (T_p \oplus T_p)^n\}(x_0, T_i^m y_0) : n \in \mathbb{N}\}} \supset \Delta(A^p)$. Since $\overline{\Delta(A^p)} \supset \Delta((X^2)^p)$ and Σ is closed, we get $\Sigma \supset \Delta((X^2)^p)$, which proves the claim.

In particular, as y_0 is hypercyclic for T_i , for each nonempty open set $U \subset X$ there exists some $u \in U$ such that (x_0, u) is s-hypercyclic for $T_1 \oplus T_1, \dots, T_p \oplus T_p$. Thus, fixing a decreasing basis (U_k) of neighborhoods of 0 and using induction, we can find for each $k \in \mathbb{N}$ some $u_k \in U_k$ and $n_k \in \mathbb{N}$ with $n_k > n_{k-1}$ (where $n_0 := 0$) such that

- (α) $T_j^{n_k} x_0 \in U_k$ for all $j = 1, \dots, p$ and
- (β) $T_j^{n_k} u_k \in x_0 + U_k$ for all $j = 1, \dots, p$.

We define $X_0 := \{T_i^n x_0 : n \in \mathbb{N}\}$ and $W_0 := X_0^p$. Note that X_0 is dense in X as x_0 is T_i -hypercyclic, so W_0 is dense in X^p (hence $\overline{W_0} \supset \Delta(X^p)$). Now, observe that no orbit of any hypercyclic vector can be finite, that is, $T_i^m x_0 \neq T_i^n x_0$ if $m \neq n$. Thus, for each $k \in \mathbb{N}$, the mapping

$$R_k : (T_i^{m_1} x_0, \dots, T_i^{m_p} x_0) \in W_0 \mapsto \frac{1}{p} \cdot \sum_{l=1}^p T_i^{m_l} u_k \in X \quad (1)$$

is well defined. We have:

- (i) For every $j = 1, \dots, p$ and every $m \in \mathbb{N}$, $T_j^{n_k}(T_i^m x_0) = T_i^m(T_j^{n_k} x_0) \rightarrow T_i^m 0 = 0$ ($k \rightarrow \infty$), where commutativity and continuity of T_i together with property (α) have been used. This shows that $T_j^{n_k} \rightarrow 0$ pointwise on X_0 for all $j = 1, \dots, p$.
- (ii) From the continuity of each $T_i^{m_l}$ and the fact that $u_k \in U_k$ (hence $u_k \rightarrow 0$), it follows that $T_i^{m_l} u_k \rightarrow 0$ ($k \rightarrow \infty$) for every $l = 1, \dots, p$. Then one derives from (1) that $R_k \rightarrow 0$ pointwise on W_0 .
- (iii) For every $j = 1, \dots, p$ and every $(m_1, \dots, m_p) \in \mathbb{N}^p$, it follows from (1) that

$$\begin{aligned} T_j^{n_k} R_k(T_i^{m_1} x_0, \dots, T_i^{m_p} x_0) &= \frac{1}{p} \cdot \sum_{l=1}^p T_j^{n_k} T_i^{m_l} u_k = \frac{1}{p} \cdot \sum_{l=1}^p T_i^{m_l} T_j^{n_k} u_k \\ &\longrightarrow \frac{1}{p} \cdot \sum_{l=1}^p T_i^{m_l} x_0 \in \text{conv}(\{T_i^{m_1} x_0, \dots, T_i^{m_p} x_0\}) \end{aligned}$$

as $k \rightarrow \infty$, because of (β) (which implies $T_j^{n_k} u_k \rightarrow x_0$) together with the commutativity and continuity of each $T_i^{m_l}$.

This tells us that T_1, \dots, T_p satisfy the s-hypercyclicity criterion, as required. \square

We raise here the question whether (d) is equivalent to (a)-(b)-(c) without assuming any commutativity.

4. SCALAR MULTIPLES OF AN OPERATOR

We start by studying s-hypercyclicity of scalar multiples of one operator. We have already pointed out that there is no chance of d-hypercyclicity in this case.

Recall that an operator T on a topological vector space X is called *hereditarily hypercyclic* whenever (T^{n_k}) is universal for every strictly increasing sequence $(n_k) \subset \mathbb{N}$. It is well known –and easy to see– that, if X is an F-space and $T \in L(X)$, then T is hereditarily hypercyclic if and only if T is mixing.

Proposition 4.1. *Let X be a topological vector space, $p \in \mathbb{N}$, $c_1, \dots, c_p \in \mathbb{K}$ and $T, T_1, \dots, T_p \in L(X)$. We have:*

- (a) Assume that X is metrizable and locally convex. If T, c_1T, \dots, c_pT are s -hypercyclic then the c_j 's are unimodular, that is, $|c_1| = \dots = |c_p| = 1$.
- (b) Suppose that X is metrizable. If $T \in L(X)$ is hereditarily hypercyclic and the scalars c_j are unimodular, then T, c_1T, \dots, c_pT are densely s -hypercyclic.

Proof. (a) Assume that T, c_1T, \dots, c_pT are s -hypercyclic, and fix $j \in \{1, \dots, p\}$. Let $c := c_j$. Then T, cT are s -hypercyclic, so there is $x_0 \in s\text{-}HC(T, cT)$. Since X is metrizable, we can find a sequence $(n_k) \subset \mathbb{N}$ such that $T^{n_k}x_0 \rightarrow x_0$ and $c^{n_k}T^{n_k}x_0 \rightarrow x_0$ as $k \rightarrow \infty$. Of course, $x_0 \neq 0$. But X is locally convex, so its topology is defined by a separating family of seminorms. Therefore there is a continuous seminorm q on X such that $q(x_0) > 0$. Consider the sequence of vectors

$$u_k := (c^{n_k} - 1)T^{n_k}x_0 \quad (k \in \mathbb{N}).$$

On the one hand, we have $u_k = c^{n_k}T^{n_k}x_0 - T^{n_k}x_0 \rightarrow x_0 - x_0 = 0$, so $q(u_k) \rightarrow 0$ by the continuity of q . On the other hand, we get $q(u_k) = |c^{n_k} - 1|q(T^{n_k}x_0)$, hence $|c^{n_k} - 1| = \frac{q(u_k)}{q(T^{n_k}x_0)} \rightarrow \frac{0}{q(x_0)} = 0$. Therefore $c^{n_k} \rightarrow 1$ as $k \rightarrow \infty$, which implies $|c| = 1$, that is, $|c_j| = 1$, as required.

(b) The result is trivial if $\mathbb{K} = \mathbb{R}$ (for $c_j = \pm 1$, so $(c_jT)^{2n} = T^{2n}$ for all n and all $j = 1, \dots, p$). The complex case $\mathbb{K} = \mathbb{C}$ is more delicate. Recall that a subset $E \subset \mathbb{T} := \{|z| = 1\}$ is said to be a *Dirichlet set* provided that there is a strictly increasing sequence $(n_k) \subset \mathbb{N}$ such that $\sup_{z \in E} |z^{n_k} - 1| \rightarrow 0$ as $k \rightarrow \infty$. It is well-known that every finite subset of \mathbb{T} is Dirichlet (see [13, Theorem 8.138(a)]). In particular, there exists $(n_k) \subset \mathbb{N}$ strictly increasing such that $c_j^{n_k} \rightarrow 1$ ($j = 1, \dots, p$). According to the hypothesis, we may take $x_0 \in HC((T^{n_k}))$. Given $x \in X$, there is a subsequence $(m_k) \subset (n_k)$ such that $T^{m_k}x_0 \rightarrow x$ and, of course, $c_j^{m_k} \rightarrow 1$ ($j = 1, \dots, p$) as $k \rightarrow \infty$. Therefore, we obtain $(c_jT)^{m_k}x_0 \rightarrow x$ for all $j = 1, \dots, p$ and hence $HC((T^{n_k})) \subset s\text{-}HC(T, c_1T, \dots, c_pT)$. But $HC((T^{n_k}))$ is dense, so $s\text{-}HC(T, c_1T, \dots, c_pT)$ also is. \square

Remarks 4.2. 1. In part (b) of the last proposition, hereditary hypercyclicity is needed in order to obtain common subsequences (n_k) to perform approximations. If this is not claimed, then, by a result due to León and Müller, any unimodular multiple of a hypercyclic operator on any topological vector space is always hypercyclic, even with the same set of hypercyclic vectors (see [23] and [21, pp. 339–340]).

2. It is known that the d -mixing property of T_1, \dots, T_p implies that c_1T_1, \dots, c_pT_p are also d -mixing for all unimodular scalars c_1, \dots, c_p (cf. [7, Remark 24(i)]). However, the corresponding result in case of s -mixing operators does not hold in general. Indeed, for a mixing operator T , the pair T, T is clearly s -mixing, but $T, -T$ are not s -mixing any more. To see this, assume that $T, -T$ are s -mixing. Then Proposition 3.3(ii) would imply that $(T^{n_k}), ((-T)^{n_k})$ are densely s -universal for each strictly increasing sequence (n_k) in \mathbb{N} – but s -universality of the sequences

(T^{2k+1}) and $((-T)^{2k+1}) = (-T^{2k+1})$ is clearly not possible. In connection with this, it is stated in [7, Remark 24(ii)] and actually proved in [28, Proposition 4.9] that in case of unimodular scalars c_1, \dots, c_p every d-hypercyclic vector x_0 for T_1, \dots, T_p is also d-hypercyclic for c_1T_1, \dots, c_pT_p . The proof uses crucially the fact that such a vector x_0 satisfies $(x_0, \dots, x_0) \in HC(T_1 \oplus \dots \oplus T_p)$. Thus, it cannot be adapted for s-hypercyclicity. Hence, we pose the question: Does the equality $s-HC(T_1, \dots, T_p) = s-HC(c_1T_1, \dots, c_pT_p)$ hold?

3. Concerning again part (b) and regarding its proof, we may obtain a much stronger result in the case $\mathbb{K} = \mathbb{C}$ and X a Banach space. Recall that a nonempty subset $E \subset \mathbb{C}$ is said to be *perfect* if it is closed and each point of E is an accumulation point of E . In particular, every perfect set is uncountable. It is well known (see [13, Theorem 8.138(b)]) that there are perfect Dirichlet subsets of \mathbb{T} . We have that if $E \subset \mathbb{T}$ is a perfect Dirichlet set and $T \in L(X)$ is mixing, then the uncountable family of rotations $\{cT : c \in E \cup \{1\}\}$ is *densely uniformly s-hypercyclic*, in the sense that there is a dense set of vectors $x_0 \in X$ satisfying the following: for every $y \in X$ there is $(n_k) \subset \mathbb{N}$ such that $\lim_{k \rightarrow \infty} \sup_{c \in E \cup \{1\}} \|(cT)^{n_k} x_0 - y\| = 0$. Indeed, we can take a sequence $(m_k) \subset \mathbb{N}$ such that $\sup_{c \in E \cup \{1\}} |c^{m_k} - 1| = \sup_{c \in E} |c^{m_k} - 1| \rightarrow 0$ as $k \rightarrow \infty$. As T is mixing, the set $HC((T^{m_k}))$ is dense. If $x_0 \in HC((T^{m_k}))$, then there is a subsequence $(n_k) \subset (m_k)$ with $T^{n_k} x_0 \rightarrow y$. The conclusion follows from the inequality $\|(cT)^{n_k} x_0 - y\| \leq \|c^{n_k}(T^{n_k} x_0 - y)\| + \|(c^{n_k} - 1)y\|$.

4. Proposition 4.1 furnishes examples of pairs of operators –on spaces of sequences or of holomorphic functions (see sections 5–6)– that are s-hypercyclic but not d-hypercyclic: the multiples $2B, -2B$ of the backward shift B on ℓ_q ($1 \leq q < \infty$) or c_0 ; $D, -D$ on $H(\mathbb{C})$ ($Df := f'$); $C_\varphi, -C_\varphi$ on $H(G)$, where $C_\varphi f := f \circ \varphi$, $G \subset \mathbb{C}$ is a simply connected domain and φ is a run-away automorphism of G .

5. BACKWARD SHIFTS AND S-HYPERCYCLICITY

In this section, we consider the sequence spaces c_0 and ℓ_q ($1 \leq q < \infty$) over $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . If $a = (a_n)_{n \in \mathbb{N}}$ is a bounded sequence in $\mathbb{K} \setminus \{0\}$, then B_a will denote the weighted backward shift

$$B_a : (x_0, x_1, x_2, \dots) \in X \mapsto (a_1 x_1, a_2 x_2, \dots) \in X$$

on $X = c_0$ or ℓ_q . The unweighted backward shift B is $B = B_a$, where $a = (1, 1, 1, \dots)$. Salas characterized the hypercyclicity of B_a in terms of the weight sequence a . Bès and Peris [11, Theorem 4.1] did the same for the d-hypercyclicity of different powers of B_a . This characterization happens to hold also for s-hypercyclicity.

Proposition 5.1. *Let $X = c_0$ or ℓ_q ($1 \leq q < \infty$), $p \geq 2$ and let $r_1, \dots, r_p \in \mathbb{N}$ with $r_1 < r_2 < \dots < r_p$ be given. For each $l \in \{1, \dots, p\}$, let $a_l = (a_{l,n})_{n \in \mathbb{N}}$ be a weight sequence. Then the following are equivalent:*

- (i) $B_{a_1}^{r_1}, \dots, B_{a_p}^{r_p}$ are d -hypercyclic.
- (ii) $B_{a_1}^{r_1}, \dots, B_{a_p}^{r_p}$ are s -hypercyclic.
- (iii) For every $M > 0$ and every $k \in \mathbb{N}$ there is $m \in \mathbb{N}$ satisfying, for each $j \in \{0, 1, \dots, k\}$, that $|a_{l,j+1} \cdots a_{l,j+r_l m}| > M$ ($1 \leq l \leq p$) and $\frac{|a_{l,j+1} \cdots a_{l,j+r_l m}|}{|a_{s,j+(r_l-r_s)m+1} \cdots a_{s,j+r_l m}|} > M$ ($1 \leq s < l \leq p$).
- (iv) $B_{a_1}^{r_1}, \dots, B_{a_p}^{r_p}$ satisfy the d -hypercyclicity criterion.
- (v) $B_{a_1}^{r_1}, \dots, B_{a_p}^{r_p}$ satisfy the s -hypercyclicity criterion.

Proof. The equivalence of (i), (iii) and (iv) is proved in [11, Theorem 4.1]. That (i) implies (ii) is trivial. Moreover, (ii) \Rightarrow (iii) is proved in fact in the proof of “(a) \Rightarrow (b)” of the same reference, since only the simultaneous approximation of one vector (namely $e_0 + \cdots + e_q$) is used. Finally, we clearly have (iv) \Rightarrow (v) \Rightarrow (ii). \square

Remarks 5.2. 1. An analogous result about equivalence of d - and s -hypercyclicity also works for powers of weighted bilateral shifts (see Theorem 4.7 of [11] and its proof).

2. Corollary 4.4 in [11] also works with just s -universality, as it is a consequence of Theorem 4.1 there. In particular, we have that B_a, B_a^2, \dots, B_a^p are s -hypercyclic on X if and only if $B_a \oplus B_a^2 \oplus \cdots \oplus B_a^p$ is hypercyclic on X^p . Bès, Martin and Peris [7, p. 855] constructed an operator $T := B_a$ on ℓ_2 such that T is hypercyclic but $T \oplus T^2$ is not hypercyclic on $\ell_2 \oplus \ell_2$, so that T, T^2 is not d -hypercyclic on ℓ_2 . Then we obtain that T, T^2 are even *not s -hypercyclic*. According to [21, Theorem 4.8], the mentioned $T = B_a$ is not mixing. In [8, Sect. 3], a mixing operator $T \in L(\ell_2)$ for which T, T^2 are not d -mixing is exhibited. But the existence of a mixing T on a separable Banach space such that T, T^2 are not d -hypercyclic is unknown so far [8, Question 3.7].

A more delicate question arises when $r_1 \leq r_2 \leq \cdots \leq r_p$. In [11, Corollary 4.2], the following is proved for weighted powers of the unweighted backward shift: if $p \geq 2$ and $r_l \in \mathbb{N}$, $\lambda_l \in \mathbb{K}$ ($1 \leq l \leq p$) with $r_1 \leq r_2 \leq \cdots \leq r_p$, then $\lambda_1 B^{r_1}, \dots, \lambda_p B^{r_p}$ are d -hypercyclic if and only if $r_1 < r_2 < \cdots < r_p$ and $1 < |\lambda_1| < |\lambda_2| < \cdots < |\lambda_p|$. The following result shows that s -hypercyclicity is possible under slightly weaker assumptions.

Proposition 5.3. *Let $p \geq 2$, and let $r_l \in \mathbb{N}$, $\lambda_l \in \mathbb{K}$ ($1 \leq l \leq p$) with $r_1 \leq r_2 \leq \cdots \leq r_p$. Let A denote the set $A := \{j \in \{1, \dots, p-1\} : r_j = r_{j+1}\}$ and consider the conditions*

- (i) $1 < |\lambda_j|$ for all $j \in \{1, \dots, p\}$,
- (ii) $|\lambda_j| < |\lambda_{j+1}|$ for all $j \in \{1, \dots, p-1\} \setminus A$,
- (iii) $|\lambda_j| = |\lambda_{j+1}|$ for all $j \in A$.

Then $\lambda_1 B^{r_1}, \dots, \lambda_p B^{r_p}$ are s -hypercyclic on $X = c_0$ or ℓ_q ($1 \leq q < \infty$) if and only if (i), (ii) and (iii) hold.

Proof. First, suppose that conditions (i),(ii) and (iii) hold. We write $\{1, \dots, p\} \setminus A = \{t_1, \dots, t_d\}$, with $d \in \mathbb{N}$ and $t_1 < \dots < t_d$. As the set $\{\lambda_i/\lambda_j : i, j \in \{1, \dots, p\}\}$ with $|\lambda_i| = |\lambda_j| \subset \mathbb{T}$ is finite, it is a Dirichlet set. Hence there exists a strictly increasing sequence $(n_k) \subset \mathbb{N}$ such that

$$\left(\frac{\lambda_i}{\lambda_j}\right)^{n_k} \rightarrow 1 \quad (k \rightarrow \infty) \quad \text{for all } i, j \in \{1, \dots, p\} \quad \text{with } |\lambda_i| = |\lambda_j|. \quad (1)$$

Consider the set X_0 of finite sequences, that is, $X_0 := c_{00} = \{x = (x_n) \in X : \text{exists } n_0 = n_0(x) \in \mathbb{N} \text{ such that } x_n = 0 \text{ for all } n \geq n_0\}$. Then X_0 is dense in X . If we set $W_0 := \Delta(X_0^p) \subset X^p$, then $\overline{W_0} = \overline{\Delta(X_0^p)} \supset \Delta(X^p)$ because X_0 is dense in X . Now, we set $T_j := \lambda_j B^{r_j}$ ($j = 1, \dots, p$). Define, for each $k \in \mathbb{N}$, the mapping $R_k : W_0 \rightarrow X$ as follows. If $x = (x_1, x_2, \dots, x_N, 0, 0, 0, \dots) \in X_0$ and $w = (x, x, \dots, x)$, then

$$R_k w = \begin{cases} (0_1, u_1, 0_2, u_2, \dots, 0_N, u_N, 0, 0, 0, \dots) & \text{if } n_k \geq N \\ (0, 0, 0, \dots) & \text{if } n_k < N, \end{cases}$$

where $0_1 := (0, 0, \dots, 0)$ [$r_{t_1} n_k$ times], $0_l := (0, 0, \dots, 0)$ [$(r_{t_l} - r_{t_{l-1}})n_k - N$ times] if $l \geq 2$ and $u_l := (\frac{1}{\lambda_{t_l}^{n_k}} x_1, \dots, \frac{1}{\lambda_{t_l}^{n_k}} x_N)$ ($l \geq 1$). We have:

- (a) For each $j \in \{1, \dots, p\}$ and each $x = (x_1, x_2, \dots, x_N, 0, 0, 0, \dots) \in X_0$, $T_j^{n_k} x = 0$ as soon as $r_j n_k > N$, so $T_j^{n_k} \rightarrow 0$ ($k \rightarrow \infty$) pointwise on X_0 .
- (b) For every $w = (x, \dots, x) \in W_0$ as before, the definition of R_k together with (i) yields $R_k w \rightarrow 0$ as $k \rightarrow \infty$.
- (c) Fix $w = (x, \dots, x) \in W_0$, where $x = (x_1, x_2, \dots, x_N, 0, 0, 0, \dots)$. For every $j \in \{1, \dots, p\}$ there is exactly one $l \in \{1, \dots, d\}$ such that $|\lambda_j| = |\lambda_{t_l}|$, due to (ii) and (iii). Finally, if $n_k \geq N$, we have

$$T_j^{n_k} R_k w = \left(\begin{aligned} & \left(\frac{\lambda_j}{\lambda_{t_l}}\right)^{n_k} x_1, \left(\frac{\lambda_j}{\lambda_{t_l}}\right)^{n_k} x_2, \dots, \left(\frac{\lambda_j}{\lambda_{t_l}}\right)^{n_k} x_N, 0, 0, \dots, 0, \\ & \left(\frac{\lambda_j}{\lambda_{t_{l+1}}}\right)^{n_k} x_1, \left(\frac{\lambda_j}{\lambda_{t_{l+1}}}\right)^{n_k} x_2, \dots, \left(\frac{\lambda_j}{\lambda_{t_{l+1}}}\right)^{n_k} x_N, 0, 0, \dots, 0, \dots, \\ & \left(\frac{\lambda_j}{\lambda_{t_d}}\right)^{n_k} x_1, \left(\frac{\lambda_j}{\lambda_{t_d}}\right)^{n_k} x_2, \dots, \left(\frac{\lambda_j}{\lambda_{t_d}}\right)^{n_k} x_N, 0, 0, 0, 0, \dots \end{aligned} \right).$$

It follows from (ii) that $(\frac{\lambda_j}{\lambda_{t_s}})^{n_k} x_\nu \rightarrow 0$ as $k \rightarrow \infty$ for all $s \in \{l+1, \dots, d\}$ and all $\nu \in \{1, \dots, N\}$, while (1) entails that $(\frac{\lambda_j}{\lambda_{t_l}})^{n_k} x_\nu \rightarrow x_\nu$ as $k \rightarrow \infty$ for all $\nu \in \{1, \dots, N\}$. Consequently, $T_j^{n_k} R_k w \rightarrow (x_1, x_2, \dots, x_N, 0, 0, 0, \dots) = x$.

An application of the s-hypercyclicity criterion (see also Remark 3.8.1) concludes the first part of the proof.

Now, suppose that $\lambda_1 B^{r_1}, \dots, \lambda_p B^{r_p}$ are s-hypercyclic. Since hypercyclic operators on normed spaces have norm larger than 1, we obtain

$$1 < \|\lambda_j B^{r_j}\| = |\lambda_j| \|B^{r_j}\| = |\lambda_j|$$

for all $j = 1, \dots, p$ (cf. the proof of Corollary 4.2 in [11]), i.e. condition (i) holds. For each $j \in \{1, \dots, p-1\} \setminus A$, we have $r_j < r_{j+1}$. Hence, as $\lambda_j B^{r_j}, \lambda_{j+1} B^{r_{j+1}}$ are s -hypercyclic, Proposition 5.1, (ii) \Rightarrow (iii), and the same approach as in the proof of Corollary 4.2 in [11] yield $|\lambda_j| < |\lambda_{j+1}|$, i.e. condition (ii) holds. Finally, for each $j \in A$, we have $r_j = r_{j+1}$. Hence, the s -hypercyclicity of

$$\lambda_j B^{r_j}, \lambda_{j+1} B^{r_{j+1}} = \frac{\lambda_{j+1}}{\lambda_j} \cdot \lambda_j B^{r_j}$$

implies $|\lambda_{j+1}/\lambda_j| = 1$ (see Proposition 4.1(a)) and thus $|\lambda_j| = |\lambda_{j+1}|$, i.e. condition (iii) holds. \square

For instance, the operators $2B, 3B^2, -3B^2$, being not d -hypercyclic, are s -hypercyclic. Further study of d -hypercyclicity of weighted unilateral and bilateral backward shifts can be found in [9].

6. S-HYPERCYCLICITY IN SPACES OF HOLOMORPHIC FUNCTIONS

Let $G \subset \mathbb{C}$ be a domain, that is, a nonempty connected open subset of \mathbb{C} . We endow the space $H(G)$ of all holomorphic (or analytic) functions $G \rightarrow \mathbb{C}$ with the topology of uniform convergence on compacta, so that $H(G)$ becomes a separable Fréchet space. In this section we are concerned with s -hypercyclicity of finite sets of operators on $H(G)$ (or on subspaces of it) for certain domains G .

Recall that if X is a topological vector space and $T \in L(X)$, then T is said to be *supercyclic* provided that there exists some $x_0 \in X$ whose projective orbit $\{\lambda T^n x_0 : n \in \mathbb{N}, \lambda \in \mathbb{K}\}$ is dense in X . If $T_1, \dots, T_p \in L(X)$, they are called *d -supercyclic* (see [7]) if there is $x_0 \in X$ such that $\{\lambda [T_1^n, \dots, T_p^n] x_0 : n \in \mathbb{N}, \lambda \in \mathbb{K}\}$ is dense in X^p . Consistently, we say that T_1, \dots, T_p are *s -supercyclic* whenever $\overline{\{\lambda [T_1^n, \dots, T_p^n] x_0 : n \in \mathbb{N}, \lambda \in \mathbb{K}\}} \supset \Delta(X^p)$.

Let $LFT(\mathbb{D})$ denote the family of all linear fractional transformations $\varphi(z) = \frac{az+b}{cz+d}$ of the complex plane such that $\varphi(\mathbb{D}) \subset \mathbb{D}$. The subfamily $\text{Aut}(\mathbb{D})$ of automorphisms of \mathbb{D} consists of all onto members of $LFT(\mathbb{D})$. See e.g. [27, Chapter 1] for terminology related to these families. If $\nu \in \mathbb{R}$, then S_ν denotes the weighted Hardy space $S_\nu = \{f(z) = \sum_{n \geq 0} a_n z^n \in H(\mathbb{D}) : \|f\| := (\sum_{n \geq 0} |a_n|^2 (n+1)^{2\nu})^{1/2} < \infty\}$. Each S_ν is a Hilbert space, and the choices $\nu = -1/2, 0, 1/2$ correspond, respectively, to the classical Bergman, Hardy and Dirichlet spaces. Thanks to the results in [7], we obtain without effort the next two assertions.

Proposition 6.1. *Let $\varphi_1, \dots, \varphi_p \in LFT(\mathbb{D})$ pairwise distinct. Then the following are equivalent:*

- (a) $C_{\varphi_1}, \dots, C_{\varphi_p}$ are s -supercyclic on $H(\mathbb{D})$.
- (b) $\mu_1 C_{\varphi_1}, \dots, \mu_p C_{\varphi_p}$ are s -mixing on $H(\mathbb{D})$ for all nonzero scalars μ_1, \dots, μ_p .
- (c) $C_{\varphi_1}, \dots, C_{\varphi_p}$ are d -supercyclic on $H(\mathbb{D})$.
- (d) $\mu_1 C_{\varphi_1}, \dots, \mu_p C_{\varphi_p}$ are d -mixing on $H(\mathbb{D})$ for all nonzero scalars μ_1, \dots, μ_p .

- (e) $\varphi_1, \dots, \varphi_p$ have no fixed point in \mathbb{D} , and satisfy that if any two φ_l, φ_j have the same attractive fixed point α , then $\varphi'_l(\alpha) = \varphi'_j(\alpha) < 1$ is not possible.

Proof. The equivalence of (c), (d) and (e) is proved in [7, Theorem 4]. The implications (d) \Rightarrow (b) \Rightarrow (a) are trivial. Finally, (a) \Rightarrow (e) is proved in fact in the proof of Theorem 4 in [7]. Indeed, it is used there a result (Lemma 14 in [7]) asserting that if $\varphi_1, \varphi_2 \in LFT(\mathbb{D})$ are hyperbolic and share an attractive fixed point α with $\varphi'_1(\alpha) = \varphi'_2(\alpha)$, then $C_{\varphi_1}, C_{\varphi_2}$ are not d-supercyclic on $H(\mathbb{D})$. But a closer look at its proof shows that $C_{\varphi_1}, C_{\varphi_2}$ are in fact even not s-supercyclic; indeed, via contradiction, only one function g is assumed to be simultaneously approximated by projective orbits. \square

Proposition 6.2. *Let $\varphi_1, \dots, \varphi_p \in LFT(\mathbb{D})$ pairwise distinct and let $\nu < 1/2$. Then the following are equivalent:*

- (a) $C_{\varphi_1}, \dots, C_{\varphi_p}$ are s-supercyclic on S_ν .
- (b) $C_{\varphi_1}, \dots, C_{\varphi_p}$ are s-mixing on S_ν .
- (c) $C_{\varphi_1}, \dots, C_{\varphi_p}$ are d-supercyclic on S_ν .
- (d) $C_{\varphi_1}, \dots, C_{\varphi_p}$ are d-mixing on S_ν .
- (e) Each φ_l is a parabolic automorphism or a hyperbolic map without fixed points in \mathbb{D} , and there are no two φ_l, φ_j having a common fixed point α such that $\varphi'_l(\alpha) = \varphi'_j(\alpha) < 1$.

Proof. The equivalence of (c), (d) and (e) is proved in [7, Theorem 3]. The implications (d) \Rightarrow (b) \Rightarrow (a) are trivial. As for (a) \Rightarrow (e), observe that in the proof of Theorem 3 in [7], only the supercyclicity of each C_{φ_l} is necessary for the first assertion in (e) and that the Comparison Principle [7, Proposition 8] –that also works for s-supercyclicity– implies that $C_{\varphi_1}, \dots, C_{\varphi_p}$ are s-supercyclic on $H(\mathbb{D})$. Now, the second assertion of (e) follows from Proposition 6.1. \square

Remarks 6.3. 1. Recall that if X is an F-space and $T \in L(X)$ is invertible and hypercyclic, then T^{-1} is also hypercyclic. Analogously as in Example 22 in [7], by combining the preceding two propositions, we obtain that there are hyperbolic $\varphi_1, \varphi_2 \in \text{Aut}(\mathbb{D})$ such that $C_{\varphi_1}, C_{\varphi_2}$ are d-hypercyclic (so s-hypercyclic) on $H^2(\mathbb{D})$ (the Hardy space) and on $H(\mathbb{D})$, and $C_{\varphi_1^{-1}} = (C_{\varphi_1})^{-1}, C_{\varphi_2^{-1}} = (C_{\varphi_2})^{-1}$ are even not s-supercyclic on $H^2(\mathbb{D})$ or $H(\mathbb{D})$ (note that φ_1^{-1} and φ_2^{-1} are also hyperbolic). Hence, in general, the d-hypercyclicity of T_1, \dots, T_p does *not* imply the s-hypercyclicity of $T_1^{-1}, \dots, T_p^{-1}$ if T_1, \dots, T_p are invertible. Moreover, finitely many composition operators generated by non-elliptic automorphisms of \mathbb{D} may be *not* s-hypercyclic on $H(\mathbb{D})$ or on $H^2(\mathbb{D})$.

2. Further study of d-hypercyclicity of composition operators, this time on weighted Bergman spaces on \mathbb{D} , is performed in [30].

In 1929 Birkhoff [12] proved that the translation operator τ_a ($a \in \mathbb{C} \setminus \{0\}$) given by $(\tau_a f)(z) = f(z + a)$ is hypercyclic on the space $H(\mathbb{C})$ of entire functions. It is

proved in [4, Prop. 5.5] and [11, Theorem 3.1] that if a_1, \dots, a_p are pairwise distinct nonzero complex numbers, then $\tau_{a_1}, \dots, \tau_{a_p}$ are d-hypercyclic. Trivially, we obtain: if $a_1, \dots, a_p \in \mathbb{C} \setminus \{0\}$, then $\tau_{a_1}, \dots, \tau_{a_p}$ are s-hypercyclic. As the next proposition shows, we may obtain a slight extension to weighted translation operators.

Proposition 6.4. *Let $p \geq 2$, and let $a_1, \dots, a_p, \lambda_1, \dots, \lambda_p \in \mathbb{C} \setminus \{0\}$ such that $|\lambda_j| = |\lambda_l|$ for all $j, l \in \{1, \dots, p\}$ with $a_j = a_l$. Then there is a sequence $(n_k) \subset \mathbb{N}$ such that the sequences $(\lambda_1 \tau_{a_1})^{n_k}, \dots, (\lambda_p \tau_{a_p})^{n_k}$ are s-mixing. In particular, the operators $\lambda_1 \tau_{a_1}, \dots, \lambda_p \tau_{a_p}$ are densely s-hypercyclic on $H(\mathbb{C})$.*

Proof. Select a finite sequence $\{j(1) < j(2) \cdots < j(q)\} \subset \{1, \dots, p\}$ satisfying that, if $b_l := a_{j(l)}$ ($l = 1, \dots, q$), then the b_l 's are pairwise distinct and $\{a_1, \dots, a_p\} = \{b_1, \dots, b_q\}$. Let $\mu_l := \lambda_{j(l)}$. Consider the operators $T_j := \lambda_j \tau_{a_j}$ ($j = 1, \dots, p$) and $S_l := T_{j(l)} = \mu_l \tau_{b_l}$ ($l = 1, \dots, q$).

Let us prove that S_1, \dots, S_q are s-mixing. In fact, by following the approach of the proof of [11, Theorem 3.1], we can prove that they are even d-mixing. To this end, and taking into account that the sets $V(h, r, \varepsilon) := \{f \in H(\mathbb{C}) : |f(z) - h(z)| < \varepsilon \text{ for all } z \in \overline{B}(0, r)\}$ ($h \in H(\mathbb{C}), \varepsilon > 0, r > 0$), form a basis for the topology of $H(\mathbb{C})$, it is enough to prove that, for given $h, g_1, \dots, g_q \in H(\mathbb{C})$ and $\varepsilon, r > 0$, there is $n_0 \in \mathbb{N}$ such that, for every $n \geq n_0$, there exists an entire function f with

$$|f(z) - h(z)| < \varepsilon \text{ and } |(S_l^n f)(z) - g_l(z)| < \varepsilon \quad (z \in \overline{B}(0, r), l = 1, \dots, q). \quad (1)$$

Select $n_0 \in \mathbb{N}$ with $n_0 > \max_{i \neq l} \frac{2r}{|b_i - b_l|} + \max_{1 \leq l \leq q} \frac{2r}{|b_l|}$. Then, for each $n \geq n_0$, the disks $\overline{B}(0, r), \overline{B}(nb_1, r), \dots, \overline{B}(nb_q, r)$ are pairwise disjoint. Pick $s > r$ such that the disks $\overline{B}(0, s), \overline{B}(nb_1, s), \dots, \overline{B}(nb_q, s)$ are still pairwise disjoint. Let $K := \overline{B}(0, r) \cup \overline{B}(nb_1, r) \cup \dots \cup \overline{B}(nb_q, r)$ and $\Omega := B(0, s) \cup B(nb_1, s) \cup \dots \cup B(nb_q, s)$. Note that Ω is an open set, $\Omega \supset K$ and K is a compact subset having connected complement. Consider the function $F : \Omega \rightarrow \mathbb{C}$ defined by

$$F(z) = h(z) \text{ if } z \in B(0, s) \text{ and } F(z) := \mu_l^{-n} g_l(z - nb_l) \text{ if } z \in B(nb_l, s) \text{ (} 1 \leq l \leq q \text{)}.$$

Then $F \in H(\Omega)$. From Runge's approximation theorem (see e.g. [16]), it follows that there exists a polynomial f (so $f \in H(\mathbb{C})$) such that $|f(z) - F(z)| < \varepsilon/(1 + |\mu_l^n|)$ for all $z \in K$. But this implies that $|f(z) - h(z)| < \varepsilon$ on $\overline{B}(0, r)$ and $|\mu_l^n f(z) - g_l(z - nb_l)| < \varepsilon$ on $\overline{B}(nb_l, r)$. Since the last inequality is equivalent to $|\mu_l^n f(z + nb_l) - g_l(z)| < \varepsilon$ on $\overline{B}(0, r)$, (1) is obtained.

As the set $D := \{\lambda_j/\lambda_l : j, l \in \{1, \dots, p\} \text{ with } a_j = a_l\} \subset \mathbb{T}$ is finite, it is a Dirichlet set. Then there is a strictly increasing sequence $(n_k) \subset \mathbb{N}$ such that $\xi^{n_k} \rightarrow 1$ as $k \rightarrow \infty$, for all $\xi \in D$.

Fix a subsequence (m_k) of (n_k) . Since S_1, \dots, S_q are s-mixing, the set $s\text{-}HC((S_1^{m_k}), \dots, (S_q^{m_k}))$ is dense (see Proposition 3.3). Fix f in $s\text{-}HC((S_1^{m_k}), \dots, (S_q^{m_k}))$. For each $\nu \in \{1, \dots, p\}$ there is a unique $l = l(\nu) \in$

$\{1, \dots, q\}$ such that $a_\nu = b_l$, so that $|\lambda_\nu| = |\mu_l|$. Observe that $\xi_\nu := \lambda_\nu / \mu_l \in D$. Then $\xi_\nu^{n_k} \rightarrow 1$, hence $\xi_\nu^{m_k} \rightarrow 1$ ($k \rightarrow \infty$) for all $\nu \in \{1, \dots, p\}$. Given $g \in H(\mathbb{C})$, we can find a subsequence (p_k) of (m_k) with $S_{l(\nu)}^{p_k} f \rightarrow g$ ($k \rightarrow \infty$) uniformly on compacta for every $\nu \in \{1, \dots, p\}$. Since $\xi_\nu^{p_k} \rightarrow 1$ for all ν , we obtain that $T_\nu^{p_k} f = \xi_\nu^{p_k} S_{l(\nu)}^{p_k} f \rightarrow 1 \cdot g = g$ ($k \rightarrow \infty$) uniformly on compacta for every $\nu = 1, \dots, p$. Therefore $f \in s\text{-}HC((T_1^{m_k}), \dots, (T_p^{m_k}))$, which shows that this set is dense. By Proposition 3.3, the sequences $(T_1^{n_k}), \dots, (T_p^{n_k})$ are s-mixing, as required. \square

Another important collection of operators on $H(\mathbb{C})$ is that of differentiation operators. Consider the derivative operator $D : f \in H(\mathbb{C}) \mapsto f' \in H(\mathbb{C})$. Its hypercyclicity on $H(\mathbb{C})$ was proved by MacLane in 1952 [25]. It is shown in [11, Prop. 3.3] that if $p \geq 2$, $r_1, \dots, r_p \in \mathbb{N}$ with $r_1 < \dots < r_p$ and $\lambda_1, \dots, \lambda_p \in \mathbb{C} \setminus \{0\}$, then $\lambda_1 D^{r_1}, \dots, \lambda_p D^{r_p}$ are d-mixing, so densely d-hypercyclic. Concerning s-hypercyclicity, the following proposition shows that somewhat softer assumptions are allowed, although, similarly to the last proposition, we have not been able to obtain the s-mixing property for the whole sequences.

Proposition 6.5. *Let $r_1 \leq \dots \leq r_p$ be positive integers and $\lambda_1, \dots, \lambda_p \in \mathbb{C} \setminus \{0\}$, where $p \geq 2$. Suppose that $|\lambda_j| = |\lambda_l|$ for all $j, l \in \{1, \dots, p\}$ with $r_j = r_l$.*

Then there is a sequence $(n_k) \subset \mathbb{N}$ such that the sequences $(\lambda_1 D^{r_1})^{n_k}, \dots, (\lambda_p D^{r_p})^{n_k}$ are s-mixing. In particular, the operators $\lambda_1 D^{r_1}, \dots, \lambda_p D^{r_p}$ are densely s-hypercyclic on $H(\mathbb{C})$.

Proof. As the set $\{\lambda_j / \lambda_l : j, l \in \{1, \dots, p\} \text{ with } r_j = r_l\} \subset \mathbb{T}$ is finite, it is a Dirichlet set. Then there is a strictly increasing sequence $(n_k) \subset \mathbb{N}$ such that $(\lambda_j / \lambda_l)^{n_k} \rightarrow 1$ as $k \rightarrow \infty$, for all $j, l \in \{1, \dots, p\}$ with $r_j = r_l$. Put $X_0 := \{\text{polynomials}\} = \text{span}\{z^m : m \in \mathbb{N}_0\}$ and $W_0 := \Delta(X_0^p)$. Then X_0 is dense in $X := H(\mathbb{C})$ and $\overline{W_0} = \overline{\Delta(X_0^p)} \supset \Delta(X^p)$. Let $T_j := \lambda_j D^{r_j}$ ($1 \leq j \leq p$). For each $k \in \mathbb{N}$, define the map $R_k : W_0 \rightarrow X$ via

$$R_k(z^m, \dots, z^m) := \sum_{l=1}^p \frac{1}{\tau(l)} \cdot \frac{1}{\lambda_l^{n_k}} \cdot \frac{z^{m+r_l n_k}}{(m+1)(m+2)\cdots(m+r_l n_k)},$$

where $\tau(l) := \text{card}\{i \in \{1, \dots, p\} : r_i = r_l\}$ ($1 \leq l \leq p$). Then R_k is extended to the whole W_0 by linearity. We have:

- (i) $T_j^{n_k} z^m = 0$ as soon as $n_k r_j > m$, so $T_j^{n_k} z^m \rightarrow 0$ as $k \rightarrow \infty$ for all $j \in \{1, \dots, p\}$ and all $m \geq 0$. Therefore, by linearity, $T_j^{n_k} \rightarrow 0$ ($k \rightarrow \infty$) on X_0 for all $j \in \{1, \dots, p\}$.

- (ii) Fix $m \in \mathbb{N}_0$ and a compact set $K \subset \mathbb{C}$. There is $M \in (0, +\infty)$ with $K \subset \overline{B}(0, M)$. Given $k \in \mathbb{N}$, we obtain

$$\begin{aligned} \sup_{z \in K} |R_k(z^m, \dots, z^m)| &\leq \sum_{l=1}^p \frac{1}{\tau(l)} \cdot \frac{1}{\lambda_l^{n_k}} \cdot \frac{M^{m+r_l n_k}}{(m+1)(m+2) \cdots (m+r_l n_k)} \\ &\leq \sum_{l=1}^p \frac{1}{\tau(l)} \frac{M^{m+r_l n_k} / \lambda_l^{n_k}}{(m+1)(m+2) \cdots (m+n_k)} \\ &\leq \sum_{l=1}^p \frac{M^m (M^{r_l} / \lambda_l)^{n_k}}{\tau(l) n_k!} \rightarrow 0 \quad (k \rightarrow \infty) \end{aligned}$$

Hence, by linearity, $R_k \rightarrow 0$ ($k \rightarrow \infty$) pointwise on W_0 .

- (iii) Fix $m \in \mathbb{N}_0$, $j \in \{1, \dots, p\}$ and $k \in \mathbb{N}$ with $n_k > m$. Let us compute the action of $T_j^{n_k} R_k$ on each (z^m, \dots, z^m) . This yields three sums, the first of them corresponding to those $l \in \{1, \dots, p\}$ with $r_l < r_j$, that equals 0. Therefore

$$\begin{aligned} T_j^{n_k} R_k(z^m, \dots, z^m) &= 0 + \sum_{\substack{l=1 \\ r_l=r_j}}^p \frac{1}{\tau(l)} \cdot \left(\frac{\lambda_j}{\lambda_l}\right)^{n_k} \cdot z^m \\ &+ \sum_{\substack{l=1 \\ r_l>r_j}}^p \frac{1}{\tau(l)} \cdot \left(\frac{\lambda_j}{\lambda_l}\right)^{n_k} \cdot \frac{z^{m+(r_l-r_j)n_k}}{(m+1)(m+2) \cdots (m+(r_l-r_j)n_k)} \\ &\rightarrow \frac{1}{\tau(j)} \cdot z^m \cdot \sum_{\substack{l=1 \\ r_l=r_j}}^p 1 + 0 = z^m \quad (k \rightarrow \infty) \end{aligned}$$

uniformly on compacta in \mathbb{C} , because $\tau(j) = \tau(l)$ and $(\frac{\lambda_j}{\lambda_l})^{n_k} \rightarrow 1$ for all (j, l) with $r_j = r_l$. By linearity again, we get $T_j^{n_k} R_k(w, \dots, w) \rightarrow w$ for all $j = 1, \dots, p$ and all $(w, \dots, w) \in W_0$.

The conclusion now follows from Theorem 3.7 (or from Remark 3.8.1). \square

For instance, the operators $5D, D^2, -D^2, e^i D^2, \frac{1}{10} D^3, -3D^4$ are s-hypercyclic, but clearly not d-hypercyclic.

An extension unifying both Birkhoff's and MacLane's theorems takes place by considering convolution operators on $H(\mathbb{C})$, that is, operators commuting with all translations τ_a . Let $\Phi(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{C})$. Then Φ is said to be of *exponential type* provided that there are positive constants A, B such that $|\Phi(z)| \leq A \exp(B|z|)$ for all $z \in \mathbb{C}$. Then its associated differential operator $\Phi(D) = \sum_{n=0}^{\infty} a_n D^n$ given by $\Phi(D)f = \sum_{n=0}^{\infty} a_n f^{(n)}$ ($f \in H(\mathbb{C})$) defines an operator on $H(\mathbb{C})$. Moreover, an operator $T \in L(H(\mathbb{C}))$ is of convolution if and only if $T = \Phi(D)$ for some entire function Φ of exponential type. Note that D and τ_a

are special cases (take $\Phi(z) \equiv z$ and $\Phi(z) \equiv e^{az}$, resp.). Godefroy and Shapiro [17] proved in 1991 that any nonscalar convolution operator is hypercyclic. If G is any domain in \mathbb{C} , then $\Phi(D)$ is also an operator on $H(G)$ whenever Φ is of *subexponential type*, that is, for given $\varepsilon > 0$ there is a constant $A > 0$ such that $|\Phi(z)| \leq A \exp(\varepsilon|z|)$ for all $z \in \mathbb{C}$. We have that also $\Phi(D)$ is hypercyclic on $H(G)$ provided that G is simply connected (i.e. its complement with respect to the one-point compactification \mathbb{C}_∞ of \mathbb{C} is connected) and Φ is not constant. For s-hypercyclicity, we present the following assertion, with which we put an end to this introductory paper on s-universality.

Proposition 6.6. *Assume that $G \subset \mathbb{C}$ is a simply connected domain and that Φ_1, \dots, Φ_p are entire functions of subexponential type (or just of exponential type if $G = \mathbb{C}$). Assume also that the set*

$$U_0 := \left\{ \lambda \in \mathbb{C} : \max_{1 \leq j \leq p} |\Phi_j(\lambda)| < 1 \right\}$$

is nonempty and that each set

$$U_i := \left\{ \lambda \in \mathbb{C} : |\Phi_i(\lambda)| > 1 \text{ and } \max_{1 \leq j \leq p} |\Phi_j(\lambda)| \leq |\Phi_i(\lambda)| \right\} \quad (1 \leq i \leq p)$$

has nonempty interior U_i^0 . Suppose, in addition, that whenever $i, j \in \{1, \dots, p\}$ satisfy $|\Phi_i(\lambda)| = |\Phi_j(\lambda)|$ for some $\lambda \in U_i^0$, there exists $\zeta \in \mathbb{T}$ with $\Phi_j = \zeta \cdot \Phi_i$.

Then there is a sequence $(n_k) \subset \mathbb{N}$ such that the sequences $(\Phi_1(D))^{n_k}, \dots, (\Phi_p(D))^{n_k}$ are s-mixing. In particular, the operators $\Phi_1(D), \dots, \Phi_p(D)$ are densely s-hypercyclic on $H(\mathbb{C})$.

Proof. We write $e_\lambda := \exp(\cdot\lambda)|_G$ for $\lambda \in \mathbb{C}$. It is easy to see that the functions e_λ are linearly independent. Denote $V_i := U_i^0$ ($1 \leq i \leq p$). As U_0, V_1, \dots, V_p are open and nonempty, we obtain that $X_0 := \text{span}\{e_\lambda : \lambda \in U_0\}$ is dense in $X := H(G)$ (because G is simply connected: use Runge's approximation theorem together with the fact that $\text{span}\{\exp(\cdot\lambda) : \lambda \in U_0\}$ is dense in $H(\mathbb{C})$; see e.g. [17, Sect. 5]). Hence $W_0 := \prod_{i=1}^p \text{span}\{e_\lambda : \lambda \in V_i\}$ is dense in X^p .

As $A := \{\zeta \in \mathbb{T} : \text{exist } l, j \in \{1, \dots, p\} \text{ with } \Phi_j = \zeta\Phi_l\} \subset \mathbb{T}$ is finite, it is a Dirichlet set; hence there is a strictly increasing sequence $(n_k) \subset \mathbb{N}$ such that $\zeta^{n_k} \rightarrow 1$ for all $\zeta \in A$.

For each $i \in \{1, \dots, p\}$, we put $T_i := \Phi_i(D)|_{H(G)}$, $E_i := \{j \in \{1, \dots, p\} : \text{exists } \zeta \in \mathbb{T} \text{ with } \Phi_j = \zeta\Phi_i\}$ and $\tau(i) := \text{card}(E_i)$. Notice that if $i \in E_j$, then $E_i = E_j$ (just use that \mathbb{T} is a multiplicative group), hence $\tau(i) = \tau(j)$. Given $i \in \{1, \dots, p\}$ and $v_i \in \text{span}\{e_\lambda : \lambda \in V_i\}$, there are uniquely determined scalars $c_{i,1}, \dots, c_{i,J(i)} \in \mathbb{C}$ and pairwise distinct $\lambda_{i,1}, \dots, \lambda_{i,J(i)} \in V_i$ such that $v_i = \sum_{l=1}^{J(i)} c_{i,l} e_{\lambda_{i,l}}$. For $k \in \mathbb{N}$ we define $R_k : W_0 \rightarrow X$ as

$$R_k w := \sum_{i=1}^p \frac{1}{\tau(i)} \cdot \sum_{l=1}^{J(i)} \frac{c_{i,l}}{\Phi_i(\lambda_{i,l})^{n_k}} \cdot e_{\lambda_{i,l}}, \quad (1)$$

where $w = (v_1, \dots, v_p) \in W_0$ and the v_i 's are as above. We have:

- (i) If $\lambda \in U_0$ and $j \in \{1, \dots, p\}$, then $T_j^{n_k} e_\lambda = \Phi_j(\lambda)^{n_k} e_\lambda \rightarrow 0$ as $k \rightarrow \infty$, because $|\Phi_j(\lambda)| < 1$. By linearity, we get $T_j^{n_k} \rightarrow 0$ on X_0 .
- (ii) Let $w = (v_1, \dots, v_p) \in W_0$, so that $v_i = \sum_{l=1}^{J(i)} c_{i,l} e_{\lambda_{i,l}}$, as above. Since $|\Phi_i(\lambda_{i,l})| > 1$, we get $|\Phi_i(\lambda_{i,l})^{n_k}| \rightarrow +\infty$ as $k \rightarrow \infty$, for each $i \in \{1, \dots, p\}$ and each $l = 1, \dots, J(i)$. From (1) one derives that $R_k w \rightarrow 0$.
- (iii) Again, let $w = (v_1, \dots, v_p) \in W_0$, with $v_i = \sum_{l=1}^{J(i)} c_{i,l} e_{\lambda_{i,l}}$. Fix $j \in \{1, \dots, p\}$ and $k \in \mathbb{N}$. We compute

$$\begin{aligned} T_j^{n_k} R_k w &= \sum_{i=1}^p \frac{1}{\tau(i)} \cdot \sum_{l=1}^{J(i)} \frac{c_{i,l}}{\Phi_i(\lambda_{i,l})^{n_k}} \cdot T_j^{n_k} e_{\lambda_{i,l}} \\ &= \sum_{i=1}^p \frac{1}{\tau(i)} \cdot \sum_{l=1}^{J(i)} c_{i,l} \cdot \left(\frac{\Phi_j(\lambda_{i,l})}{\Phi_i(\lambda_{i,l})} \right)^{n_k} e_{\lambda_{i,l}} = A_k + B_k, \end{aligned}$$

where A_k (B_k , resp.) denotes the part of the preceding sum corresponding to those $i \in E_j$ ($i \notin E_j$, resp.). If $i \in E_j$, there is $\zeta = \zeta_{i,j} \in A$ such that $\Phi_j = \zeta \cdot \Phi_i$, so that $\left(\frac{\Phi_j(\lambda_{i,l})}{\Phi_i(\lambda_{i,l})} \right)^{n_k} = \zeta^{n_k} \rightarrow 1$ as $k \rightarrow \infty$. Note that $\tau(i) = \tau(j)$ if $i \in E_j$. Therefore, on the one hand,

$$A_k \rightarrow \sum_{\substack{i=1 \\ i \in E_j}}^p \frac{1}{\tau(i)} \cdot \sum_{l=1}^{J(i)} c_{i,l} \cdot e_{\lambda_{i,l}} = \frac{1}{\tau(j)} \cdot \sum_{\substack{i=1 \\ i \in E_j}}^p \sum_{l=1}^{J(i)} c_{i,l} \cdot e_{\lambda_{i,l}} = \frac{1}{\tau(j)} \cdot \sum_{\substack{i=1 \\ i \in E_j}}^p v_i.$$

On the other hand, if $i \notin E_j$, we have that $|\Phi_j(\lambda_{i,l})/\Phi_i(\lambda_{i,l})| < 1$ for all $l = 1, \dots, J(i)$ (indeed, as $\lambda_{i,l} \in V_i$, we have $|\Phi_j(\lambda_{i,l})| \leq |\Phi_i(\lambda_{i,l})|$; if we assume $|\Phi_j(\lambda_{i,l})| = |\Phi_i(\lambda_{i,l})|$, then there would exist $\zeta \in \mathbb{T}$ with $\Phi_j = \zeta \cdot \Phi_i$, which would yield $i \in E_j$, a contradiction). Hence $\left(\frac{\Phi_j(\lambda_{i,l})}{\Phi_i(\lambda_{i,l})} \right)^{n_k} \rightarrow 0$, so $B_k \rightarrow 0$. This entails

$$T_j^{n_k} R_k w = A_k + B_k \rightarrow \frac{1}{\tau(j)} \cdot \sum_{\substack{i=1 \\ i \in E_j}}^p v_i \quad (k \rightarrow \infty),$$

and the last vector belongs to $\text{conv}(\{v_1, \dots, v_p\})$ since in the last sum there are exactly $\tau(j)$ summands.

The conclusion follows, once again, from the s-hypercyclicity criterion (Theorem 3.7). \square

Remark 6.7. Proposition 3.4 in [11] (see also [4, Theorem 5.3]) asserts that if U_0 and $W_i := \{\lambda \in \mathbb{C} : |\Phi_i(\lambda)| > 1 \text{ and } \max_{j \neq i} |\Phi_j(\lambda)| < |\Phi_i(\lambda)|\}$ ($1 \leq i \leq p$) are nonempty, then $\Phi_1(D), \dots, \Phi_p(D)$ are d-mixing. If these assumptions are satisfied, then the assumptions of Proposition 6.6 are also satisfied. Note that Proposition 6.6 includes the case $\Phi_1 = \Phi$, $\Phi_j = c_j \Phi$ with $|c_j| = 1$ ($j = 2, \dots, p$).

This paper does not intend to be exhaustive. Of course, many more sets of operators or of sequences of operators may be analyzed under the point of view of s -universality/ s -hypercyclicity. For instance, consider a compact set $K \subset \mathbb{C}$ and the Banach space $(A(K), \|\cdot\|_\infty)$ of continuous functions $K \rightarrow \mathbb{C}$ that are holomorphic on K^0 . Let

$$T_{K,n} : f \in H(\mathbb{D}) \mapsto (S_n f)|_K \in A(K) \quad (n \in \mathbb{N}),$$

where $S_n f$ denotes the n th partial sum of the Taylor series of f around the origin. Assume that $K \subset \mathbb{C} \setminus \mathbb{D}$ and that K has connected complement. Then Costakis and Tsirivas [14, Sect. 3] have recently shown that, given any two strictly increasing sequences $(n_k), (m_k)$ in \mathbb{N} , the sequences (T_{K,n_k}) and (T_{K,m_k}) are –by using our terminology– s -universal. Even more, they have shown that

$$\bigcap \{s\text{-}\mathcal{U}((T_{K,n_k}), (T_{K,m_k})) : K \subset \mathbb{C} \setminus \mathbb{D} \text{ compact, } \mathbb{C} \setminus K \text{ connected}\}$$

is a residual subset of $H(\mathbb{D})$.

Acknowledgements. The first author has been partially supported by Plan Andaluz de Investigación de la Junta de Andalucía FQM-127 Grant P08-FQM-03543 and by MEC Grant MTM2015-65242-C2-1-P. The second author has been supported by DFG-Forschungsstipendium JU 3067/1-1.

REFERENCES

- [1] N. Bamerni, V. Kadets, and A. Kılıçman, *Hypercyclic operators are subspace hypercyclic*, J. Math. Anal. Appl. **435** (2016), 1812–1815.
- [2] F. Bayart and E. Matheron, *Dynamics of Linear Operators*, Cambridge University Press, 2009. Cambridge Tracts in Mathematics.
- [3] L. Bernal-González, *Densely hereditarily hypercyclic sequences and large hypercyclic manifolds*, Proc. Amer. Math. Soc. **127** (1999), 3279–3285.
- [4] L. Bernal-González, *Disjoint hypercyclic operators*, Studia Math. **182** (2007), 113–131.
- [5] L. Bernal-González and K.-G. Grosse-Erdmann, *The Hypercyclicity Criterion for sequences of operators*, Studia Math. **157** (2003), no. 1, 17–32.
- [6] J. Bès and Ö. Martin, *Compositional disjoint hypercyclicity equals disjoint supercyclicity*, Houston J. Math. **38** (2012), 1149–1163.
- [7] J. Bès, Ö. Martin, and A. Peris, *Disjoint hypercyclic linear fractional composition operators*, J. Math. Anal. Appl. **381** (2011), 843–856.
- [8] J. Bès, Ö. Martin, A. Peris, and S. Shkarin, *Disjoint mixing operators*, J. Funct. Anal. **263** (2012), 1283–1322.
- [9] J. Bès, Ö. Martin, and R. Sanders, *Weighted shifts and disjoint hypercyclicity*, J. Operator Theory **72** (2014), 15–40.
- [10] J. Bès and A. Peris, *Hereditarily hypercyclic operators*, J. Funct. Anal. **167** (1999), no. 1, 94–112.
- [11] ———, *Disjointness in Hypercyclicity*, J. Math. Anal. Appl. **336** (2007), 297–315.
- [12] G. D. Birkhoff, *Démonstration d'un théorème élémentaire sur les fonctions entières*, C. R. Acad. Sci. Paris **189** (1929), 473–475.
- [13] L. Bukovský, *The structure of the real line*, Springer Basel, Warsaw, 2011.

- [14] G. Costakis and N. Tsirivas, *Doubly universal Taylor series*, J. Approx. Theory **180** (2014), 21–31.
- [15] M. De La Rosa and C. Read, *A hypercyclic operator whose direct sum $T \oplus T$ is not hypercyclic*, J. Operator Theory **61** (2009), 369–380.
- [16] D. Gaier, *Lectures on complex approximation*, Birkhäuser, Basel-London-Stuttgart, 1987.
- [17] G. Godefroy and J. H. Shapiro, *Operators with dense, invariant, cyclic vectors manifolds*, J. Funct. Anal. **98** (1991), no. 2, 229–269.
- [18] K.-G. Grosse-Erdmann, *Holomorphe Monster und universelle Funktionen*, Vol. 176, Mit. Math. Sem. Giessen, 1987.
- [19] ———, *Universal families and hypercyclic operators*, Bull. Amer. Math. Soc. **36** (1999), no. 3, 345–381.
- [20] ———, *Recent developments in hypercyclicity*, Rev. R. Acad. Cienc. Ser. A Mat. **97** (2003), no. 3, 273–286.
- [21] K.-G. Grosse-Erdmann and A. Peris, *Linear Chaos*, Springer, London, 2011.
- [22] C. M. Le, *On subspace-hypercyclic operators*, Proc. Amer. Math. Soc. **139** (2011), 2847–2852.
- [23] F. León-Saavedra and V. Müller, *Rotations of hypercyclic and supercyclic operators*, Integr. equ. oper. theory **50** (2004), 385–391.
- [24] B. F. Madore and R. A. Martínez-Avendaño, *Subspace hypercyclicity*, J. Math. Anal. Appl. **373** (2011), 502–511.
- [25] G. R. MacLane, *Sequences of derivatives and normal families*, J. Anal. Math. **2** (1952), no. 1, 72–87.
- [26] S. Rolewicz, *On orbits of elements*, Studia Math. **32** (1969), 17–22.
- [27] J. H. Shapiro, *Composition Operators and Classical Function Theory*, Universitext, Springer-Verlag, New York, 1993.
- [28] S. Shkarin, *Universal elements for non-linear operators and their applications*, J. Math. Anal. Appl. **348** (2008), 193–210.
- [29] J. Wengenroth, *Hypercyclic operators on nonlocally convex spaces*, Proc. Amer. Math. Soc. **131** (2003), no. 6, 1759–1761.
- [30] L. Zhang and Z. H. Zhou, *Dynamics of composition operators on weighted Bergman spaces*, Indag. Math. (N.S.) **27** (2016), no. 1, 406–418.

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO
FACULTAD DE MATEMÁTICAS,
UNIVERSIDAD DE SEVILLA
APDO. 1160, AVENIDA REINA MERCEDES
41080 SEVILLA, SPAIN.
E-mail address: lbernal@us.es

FACHBEREICH IV MATHEMATIK
UNIVERSITÄT TRIER
D-54286 TRIER, GERMANY
E-mail address: s4anjung@uni-trier.de