



Pocock, L. V., & Sharp, D. J. (2017). Acute hospital admission of the frail older person: an opportunity to discuss future care. *Age and Ageing*, 46(6), 878–879. [afx052]. <https://doi.org/10.1093/ageing/afx052>

Peer reviewed version

Link to published version (if available):  
[10.1093/ageing/afx052](https://doi.org/10.1093/ageing/afx052)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://academic.oup.com/ageing/article/46/6/878/3748455> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/pure/about/ebr-terms>

## **Acute hospital admission of the frail older person: an opportunity to discuss future care**

More than a third of the UK population now lives to the age of 85<sup>1</sup> and the absolute number of older adults in the population is increasing significantly.<sup>2</sup> Frailty is defined by Clegg et al as “a state of increased vulnerability to poor resolution of homeostasis after a stressor event, which increases the risk of adverse outcomes, including falls, delirium, and disability”.<sup>3</sup> The prevalence increases with age, with up to half of adults over the age of 85 estimated to be frail.<sup>3,4</sup> It is now well reported that increasing frailty is important as a prognostic indicator, with frailty status being strongly associated with both quality of patient outcome and mortality.<sup>5</sup> A recent NHS Benchmarking Network report suggests that 52% of Trusts in the UK now have a specialist frailty unit<sup>6</sup> and, in primary care, the new GP contract will require practices to actively identify their frail patients and review them appropriately.<sup>7</sup>

Measurement of frailty can be performed in several ways and there are a number of tools in use. The Comprehensive Geriatric Assessment (CGA), the gold standard for the management of frailty in older people, is an holistic, multidimensional, interdisciplinary assessment of an individual and has been demonstrated to be associated with improved outcomes in a variety of settings.<sup>8</sup> The CGA can also be used to quantify an individual’s degree of frailty in a Frailty Index (FI-CGA).<sup>9</sup>

In this edition of *Age and Ageing* Hatheway et al present data from secondary analysis of a cohort study, initially reported in 2011, including 409 elderly patients admitted to a tertiary care teaching hospital in Canada.<sup>9</sup> The current study examines the relationship between the recovery of mobility and balance, initial treatment response and underlying frailty. Patients who were more frail at baseline, as reported by the patient or their family, were less likely to recover their balance and mobility (odds of no or incomplete recovery increased by 1.06 with each 0.1 increment in the FI-CGA at baseline) and this was similarly dependent on age (1.010). Improvement in mobility and balance over the first 48 hours was associated with greater improvement overall and with shorter recovery times. Patients with only mild mobility impairment recovered sooner – by day 5 about 50% with mild impairment had recovered, compared to 25% of those with moderate impairment and 10% with severe impairment. Declining mobility in the first 48 hours represented a relative risk of death of 17.1. The relationship between degree of baseline frailty and recovery was independent of the extent of mobility impairment at admission.

This study has immediate clinical relevance and contributes to the growing evidence base on this important topic. Although secondary analysis of existing data can be problematic, as the available data are not collected to address the particular research question, this study makes appropriate use of routinely collected data with a reasonable sample size. To improve generalisability, it would be helpful to see the analysis of data from a number of hospital sites.

Almost two thirds of people in the UK, aged over 85, die during a hospital admission.<sup>10</sup> The likelihood of dying in the 12 months following a hospital admission

increases with age<sup>11</sup>, however it is difficult to predict which older patients are at risk of dying during, or soon after, a hospital admission. This study offers one approach to assessing and managing the frail older patient with acute illness to appropriately tailor their ongoing care. If we are able to identify the patients who are most likely to recover, we can target them for intensive rehabilitation and early discharge planning. Equally, those patients who are at greater risk of dying can be offered the opportunity to have advance care planning discussions and palliative care input, ensuring that unnecessary interventions are minimised.

---

1 National End of Life Care Intelligence Network. *What We Know Now 2014*. London: Public Health England, 2015

2 National End of Life Care Intelligence Network. *Deaths in Older Adults in England*. South West Public Health Observatory, 2010.

3 Clegg A, Young J, Iliffe S, Olde Rikkert M and Rockwood K. Frailty in elderly people, *Lancet* 2013; 381: 752–62.

4 Song X, Mitnitski A and Rockwood K. Prevalence and 10-year outcomes of frailty in older adults in relation to deficit accumulation, *J Am Geriatr Soc*, 58 (2010), pp. 681–687.

5 Singh I, Gallacher J, Davis K, Johansen A, Eeles E and Hubbard RE. Predictors of adverse outcomes on an acute geriatric rehabilitation ward, *Age Ageing* 2012; 41 (2): 242-246.

6 NHS Benchmarking Network, (2016). *Older People's Care in Acute Settings* [online] Available at: <http://www.nhsbenchmarking.nhs.uk/projects/network-projects.php#14> [Accessed 01/03/17]

7 Pulse, (2017). *What is involved in the frailty checks scheme?* [online] Available at: <http://www.pulsetoday.co.uk/hot-topics/gp-contract-2017/18/what-is-involved-in-the-frailty-checks-scheme/20033966.article> [Accessed 08/03/17]

8 British Geriatrics Society. *Fit for Frailty: Consensus best practice guidance for the care of older people living with frailty in community and outpatient settings*. London: British Geriatrics Society, 2014.

9 Hubbard RE, Eeles EMP, Rockwood MRH et al. Assessing balance and mobility to track illness and recovery in older inpatients. *J Gen Intern Med* 2011;26:1471–8.

10 Pocock LV, Ives A, Pring A et al. Factors associated with hospital deaths in the oldest old: a cross-sectional study. *Age Ageing* 2016;45:372–6.

11 Clark D, Armstrong M, Allan A et al. Imminence of death among hospital inpatients: Prevalent cohort study. *Palliative Medicine* 2014; Vol 28, Issue 6, pp. 474 – 479.