



LJMU Research Online

Robertson, CM, Pullinger, SA, Robinson, WR, Smith, ME, Burniston, JG, Waterhouse, JM and Edwards, BJ

Is the diurnal variation in muscle force output detected/detectable when multi-joint movements are analysed using the musclelab force-velocity encoder?

<http://researchonline.ljmu.ac.uk/id/eprint/9311/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)


Robertson, CM, Pullinger, SA, Robinson, WR, Smith, ME, Burniston, JG, Waterhouse, JM and Edwards, BJ (2018) Is the diurnal variation in muscle force output detected/detectable when multi-joint movements are analysed using the musclelab force-velocity encoder? Chronobiology International.

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>



Is the diurnal variation in muscle force output detected/detectable when multi-joint movements are analysed using the MuscleLab force-velocity encoder?

Journal:	<i>Chronobiology International</i>
Manuscript ID	LCBI-2018-0087.R1
Manuscript Type:	Original Reports
Date Submitted by the Author:	n/a
Complete List of Authors:	Robertson, Colin; School for Sport and Exercise Science, University of Bolton Pullinger, Samuel; ASPIRE Academy for Sports Excellence, Sport Science Department Smith, Matt; Ljmu, Research Institute for Sport and Exercise Sciences Robinson, William; Liverpool John Moores University Burniston, Jatin; Liverpool John Moores University, Waterhouse, James; Ljmu, Research Institute for Sport and Exercise Sciences Edwards, Ben; Ljmu, Research Institute for Sport and Exercise Sciences
Keywords:	Strength, upper body, lower body, rectal temperature, time of day

SCHOLARONE™
Manuscripts

1
2
3 **Is the diurnal variation in muscle force output detected/detectable when multi-joint**
4 **movements are analysed using the MuscleLab force-velocity encoder?**
5

6
7 Colin M. Robertson¹, Samuel A. Pullinger^{2,3}, William Robinson², Matt Smith², Jatin G.
8 Burniston², James M. Waterhouse² and Ben J. Edwards²
9

10
11 ¹ *School for Sport and Biological Sciences, University of Bolton, Bolton, Lancashire, UK*

12 ² *Research Institute for Sport and Exercise Sciences, Liverpool John Moores University,*
13 *Liverpool, UK.*

14
15 ³ *Sports Science Department, Aspire Academy, Doha, Qatar.*
16
17

18
19 **Running title:** Time-of-day variations in bench press and back squat performance.
20

21
22 **Key Words:** Strength, upper body, lower body, time of day, rectal temperature.
23
24

25 **Type of submission:** Original investigations
26
27

28 **Address correspondence to:**
29

30 Address correspondence to: -
31
32

33
34 Dr Ben J. Edwards

35
36 Research Institute for Sport and Exercise Sciences,

37
38 Liverpool John Moores University,

39
40 Room 1.25 Tom Reilly Building, Byrom Street Campus,

41
42 Liverpool, L3 23AF. UK.

43
44 Tel: +44 (0) 151 904 6238

45
46 Fax:+44 (0) 151 904 6284

47
48 E-mail: B.J.Edwards@ljmu.ac.uk
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

We have investigated the magnitude of diurnal variation in back squat and bench press performance using the MuscleLab force velocity transducer. Thirty resistance-trained males (mean \pm SD: age 21.7 ± 1.4 years; body mass 80.5 ± 4.5 kg; height 1.79 ± 0.06 m) underwent two sessions at different times of day: morning (M, 07:30 h) and evening (E, 17:30 h). Each session included a period when rectal temperature (T_{rec}) was measured at rest, a 5-min standardized 150 W warm-up on a cycle ergometer, then defined programme of bench press (at 20, 40 and 60 kg) and back squat (at 30, 50 and 70 kg) exercises. A linear encoder was attached to an Olympic bar used for the exercises and average force (AF), peak velocity (PV) and time to peak velocity (tPV) were measured (MuscleLab software; MuscleLab Technology, Langesund, Norway) during the concentric phase of the movements. Values for T_{rec} at rest were higher in the evening compared to morning values (0.48°C , $P < 0.0005$). Daily variations were apparent for both bench press and back squat performance for AF (1.9 and 2.5 %), PV (8.3 and 12.7 %) and tPV (-16.6 and -9.8 %; where a negative number indicates a decrease in the variable from morning to evening). There was a main effect for load where AF and tPV increased and PV decreased from the lightest load to the heaviest for both bench press and back squat (47.1 and 80.2 %; 31.7 and 57.7 %; -42.1 and -73.9 %; $P < 0.0005$ where a negative number indicates a decrease in the variable with increasing load). An interaction was found only for tPV, such that the tPV occurs earlier in the evening than the morning at the highest loads (60 and 70 kg) for both bench press and back squat, respectively (mean difference of 0.32 and 0.62 s). In summary, diurnal variation in back squat and bench press was shown; and the time to peak velocity in complex multi-joint movements occurs earlier during the concentric phase of exercise when back squat or bench press is performed in the evening compared to the morning. This difference can be detected using a

1
2
3 low cost, portable and widely available commercial instrument and enables translation of past
4 laboratory/tightly controlled experimental research in to main-stream coaching practice.
5
6

7 **Introduction**

8
9 A large body of laboratory-based research has shown that in male subjects in a temperate
10 environment (around 17–20°C), muscle power output and force production consistently peak
11 in the mid-afternoon or early evening. This is regardless of the muscle group measured (such
12 as hand, elbow, leg, or back) or the speed of contraction (Drust et al. 2005; Reilly &
13 Waterhouse, 2009). The observation of significant diurnal variation or circadian rhythms in
14 muscle power and force production seems to be somewhat dependent on factors such as the
15 time on task, the motivation of subjects to perform and familiarization of the subjects with the
16 tasks to be performed (Bambaeichi et al. 2003; Giacomoni et al. 2005). Further, it is common
17 in studies investigating time-of-day variation of muscle force output – for single-joint,
18 isokinetic and isometric methods to be employed (Callard et al. 2000, Edwards et al. 2013
19 and Giacomoni et al. 2005). However, the physical demands of sport are such that athletes are
20 required to move and function in multiple directions, at varying rates of speed and for varying
21 periods of time. To train these athletes to meet this challenge is typically achieved *via* multi-
22 directional exercises which are targeted towards enhancing multi-joint and gross muscular
23 activation (Häkkinen & Komi, 1985). Because of this, strength and conditioning coaches
24 routinely use multi-joint exercises (back squat and bench press) to monitor and evaluate
25 muscle force output in a manner which better represents the sporting challenges for which
26 they are preparing the athlete (Baechle & Earle, 2015).
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Whilst there is convincing evidence to support both time-of-day variation of muscle force
41 output (Drust et al. 2005; Giacomoni et al. 2005) and the reliability of isokinetic and
42 isometric maximum voluntary contraction (MVC) methods to measure these differences
43 accurately (Drouin et al. 2004; Morton et al. 2005). It is yet to be established what tool to use
44 and if there is a corresponding diurnal fluctuation in complex multi-joint movements that are
45 more commonly used in applied practice. The ongoing development and validation of a
46 number of accurate and reliable dynamometers which are able to evaluate dynamic muscle
47 work have enabled research to be performed within a standardised laboratory environment
48 with immediately accessible results (Bosco et al. 1995; Jennings et al. 2005; O'Donnell et al.
49 2017). Taken together with dynamometers being practical, cost-friendly and easy to transport,
50
51
52
53
54
55
56
57
58
59
60

1
2
3 athletes, coaches and sport scientists have taken great interest in using these devices as means
4 of determining and monitoring athlete performance (Chiu et al. 2003; García-Ramos et al.
5 2016; Harris et al. 2010). One such device/dynamometer is the MuscleLab linear encoder
6 system, which measures average force (N), along with peak velocity ($\text{m}\cdot\text{s}^{-1}$) and time to peak
7 velocity (s), and can be attached to any moving limb or piece of exercise (resistance)
8 equipment (Bosco et al. 1995). As such, it enables measurement across a broad range of
9 human functions; as opposed to requiring – or stipulating – what movements are needed in
10 order to measure such factors. However, to the knowledge of the authors this device has
11 never been used in a chronobiological research design. Therefore, the aim of this study was to
12 assess the sensitivity of the MuscleLab force-velocity linear encoder (Ergotest version 4010,
13 Norway), to determine whether it was able to detect diurnal variation in multi-joint muscle
14 force output at different masses (low and height) during a bench press (upper body) and back
15 squat (lower body). We hypothesis that the MuscleLab force-velocity linear encoder is
16 sensitive to measure diurnal variation in force output in bench press and back squat at
17 difference masses on the bar.
18
19
20
21
22
23
24
25
26
27
28

29 **Methods**

30 **Subjects**

31 Thirty healthy active males (mean \pm SD: age 21.7 ± 1.4 years; body mass 80.5 ± 4.5 kg;
32 height 1.79 ± 0.06 m) recruited from Liverpool John Moores University Sports Scholarship
33 Scheme participated in this study. Verbal explanation of the experimental procedure was
34 provided to each individual; this included the aims of the study, the possible risks associated
35 with participation and the experimental procedures to be utilized. Any questions were
36 answered. Individuals then provided written, informed consent before participating in the
37 study. The experimental procedures were approved by the Human Ethics Committee at
38 Liverpool John Moores University. All of the participants used in this study had a minimum
39 of two-year's experience of resistance training, and were competent in their use of such
40 equipment and comfortable in this type of training environment. Recruiting participants with
41 this specific type of exercise history meant that the known neuromuscular facilitative
42 responses, which are typically associated with acute increases in muscular strength amongst
43 untrained individuals due to neural adaptations and responses were reduced (Häkkinen,
44 1989).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Participants were asked to avoid any strength training during the entire experimental
4 condition and to avoid taking part in any physical training or hard activity sessions of an
5 aerobic or anaerobic nature. Participants were also required to avoid all alcohol 24 hours (h)
6 prior to each test. None of the participants presented with a history of bone fractures and/or a
7 history of musculoskeletal abnormality; and none of the participants were receiving any
8 pharmacological treatment during the course of this study. The circadian chronotype of the
9 participants was assessed using a composite 'morningness questionnaire' by Smith et al.
10 (1989). The participants' mean 'chronotype' score on a 13 – 52 scale was 33.1 ± 5.5 , hence
11 all of the participants were 'intermediate types'. A circadian type inventory questionnaire of
12 Folkard & Monk (1979) determining languidness/vigorous and flexibility/rigidity of
13 participants was also utilized. The participants' mean scores were 39.8 ± 7.2 and 45.3 ± 6.4
14 for languidness/ vigour and flexibility/rigidity respectively. Only one of the participants was
15 found to be flexible and two were found to be languid. All other participants were deemed
16 rigid and vigorous according to the definitions of the questionnaire. All participants gave
17 their written informed consent. The study was conducted in accordance with the ethical
18 standards of the journal and complied with the Declaration of Helsinki.

29 **Research design**

30 Each participant first completed familiarisation sessions (detailed below), then the 2
31 experimental sessions at 07:30 and 17:30 h. These experimental sessions were
32 counterbalanced in order of administration in order to minimise any potential learning effects
33 (Monk & Leng, 1982), with a minimum of 72 h to ensure recovery between trials.

39 **Protocol: Familiarisation Session**

40 Each participant performed a minimum of four familiarisation sessions under standard
41 laboratory conditions (lighting = 200-250 lux, temperature = 20-22°C, mean humidity = 50±5
42 %), conducted over a two-week period and finishing one week before the study commenced
43 in order to minimize learning effects. The initial familiarisation sessions were used to
44 determine working loads (mass/resistance) that would be appropriate for the cohort. Each
45 participant was asked to perform the back squat with incrementing loads (30, 50, 70 and 90
46 kg) with one repetition at each load, and 5 minute (min) rest was allowed between each
47 effort. For bench press, each participant performed one repetition against each incrementing
48 load (20, 40, 60 and 80 kg) and, again, 5-min rest was given between each load. This was
49 done so that the upper loads required for the experimental trials (back squat, 70 kg; bench
50
51
52
53
54
55
56
57
58
59
60

1
2
3 press, 60 kg) were known to be comfortably within each of the participant's physical
4 capabilities, and so there was minimal likelihood of their failing to perform the
5 required efforts for data collection, especially as this trial was not a measure or test of
6 maximal strength but rather an investigation regarding the components which produce
7 muscle force output. The familiarisation sessions were allocated in a counterbalanced
8 fashion, with half of the participants taking part in the morning (at 07:30 h) and the
9 remaining half during the evening (at 17:30 h), and were separated by at least 3 days,
10 and each participant did this twice.
11
12
13
14
15

16 17 **Testing Procedure**

18 Participants were required to retire at 22:30 h and rise at 06:30 h; and be in the laboratory 30
19 min prior to measurements being taken so 07:00 and 17:00 h, respectively. Subjects came
20 into the morning session having fasted and were not allowed to consume food 4 h before the
21 evening session. On arrival compliance to the protocols sleeping, food intake and exercise
22 restrictions were assessed verbally and this was high. Thereafter, rectal temperature (T_{rec})
23 was measured using a soft flexible rectal probe (Mini-thermistor, Grant Instruments Ltd,
24 Shepreth, UK) inserted ~10 cm beyond the external anal sphincter. While subjects remained
25 in a standardised position (semi-supine and awake), T_{rec} was recorded continuously for a 30
26 min period by means of a Squirrel 1000 data logger (Grant Instruments Ltd, Shepreth, UK),
27 with the average value recorded over the last 5-min being retained for analysis. This standard
28 body position was used since core body temperature is subject to other influences than that
29 produced by the endogenous 24 h period oscillator, such as muscle activity, feeding and sleep
30 (Bougard et al. 2009, Edwards et al. 2002). As with the familiarisation period, all procedures
31 were conducted under standard laboratory conditions for lighting, humidity and temperature
32 as previously described. The daily energy intake on test days was controlled and limited to
33 2500 Kcal, which complies with the British Nutrition Foundation's recommended daily
34 calorific intake for a mature man (www.nutrition.org.uk/home.asp, accessed on 17.04.2018).
35 Water and non-caffeinated/non-alcoholic, calorie-free beverages were allowed ad libitum up
36 to a total volume of 3 L per day.
37
38
39
40
41
42
43
44
45
46
47
48
49

50
51 Participants were required to perform a standardized 5-min warm-up on a cycle ergometer
52 (Monark 817E, Stockholm, Sweden) at 150 W. Following the warm-up procedure, the
53 methods for assessing muscle force output were performed (See Figure 1).
54
55
56
57
58
59
60

1
2
3 ****Insert Figure 1 near here****
4
5

6 **Back Squat and Bench Press** (*MuscleLab, Ergotest version 4010, Norway*)

7
8 The decision not to employ one-repetition-maximum (1RM) testing in this present study was
9 largely due to the varying nature of the elite athletic disciplines the participants came from,
10 and that as Buckner et al. (2017) report the 1RM is a specific skill which warrants specific
11 training, and therefore favours those with experience of using it in training. Whilst muscle
12 strength is commonly measured *via* the performance of a one-repetition maximum (1RM)
13 protocol, Buckner et al. (2017) state that a true measurement of strength remains tenuous with
14 this approach. Further, that low-load alternatives to traditional resistance training result in
15 muscle hypertrophic changes similar to those resulting from traditional high-load resistance
16 training, but with less robust changes observed with maximal strength as measured by the
17 1RM. For this present study, each of the participants performed the two exercises in a
18 randomised order, performing either back squat or bench press (or vice versa), performing
19 three lifts at each exercise against progressively increasing loads (bench press: 20, 40 and 60
20 kg; back squat 30, 50 and 70 kg), with 5-min rest allowed between each lift (See Figure 1).
21 There is evidence to suggest that the optimal training load for performance enhancement is
22 one that emphasizes the athlete's maximum power output (Ammar et al. 2018, Comfort et al.
23 2011), in bench press and squat exercise, with moderate loads (from >30 to <70 % of 1RM)
24 appearing to provide the ideal load for power production. However, in line with
25 recommendations provided by Buckner et al. (2017), and the findings of Pareja-Blanco et al.
26 (2016), the present study utilised loads above 80 % of those measured as a better indication
27 of strength ability, and therefore prompting a maximal effort. The upper values here (bench
28 press: 60 kg, and back squat: 70 kg) equate to 85 % (bench press) and 80% (back squat) of
29 the upper load achieved during the familiarisation phase. As such, the participants were
30 performing outside of the loading realms of a routine training stimulus, and confidently
31 within a maximal loading/effort range. The individual's own body mass was factored into the
32 back squat exercise, as this is a whole-body movement, but not into the bench press. The
33 MuscleLab force-velocity linear encoder was attached to an Eleiko Olympic bar (20 kg)
34 which was set upon rests within a standard squat rack so that the participant had a 90° knee
35 flexion position (settings measured and recorded during the familiarisation process). From
36 this position, the participant was instructed to drive the bar upwards as forcefully as possible;
37 the value recorded during the test was for the concentric phase of the action only. The
38 MuscleLab system measured both the average force produced, the peak velocity and time-to-
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 peak velocity for each individual lift. This process was repeated three times, with 5-min rest
4 allowed between each individual lift, and against the progressive workloads as described
5 above. For the bench press, the bar was set so that it rested just above (~2.5 cm) the
6 participant's chest and, again, the instruction given was to push against the bar as forcefully
7 as possible. Again, this was repeated three times against the workloads described above, with
8 5-min rest between each push. The highest of the three average force outputs (and associated
9 peak velocity and time to peak velocity values) were used for analysis for each mass on the
10 bar for both bench press and back squat respectively. Only the lower and highest loads of the
11 back squat and bench tests were analysed. This was to allow the participant to perform at
12 greater velocities (lowest load) and challenging overall muscle force output when the ability
13 to move quickly has been reduced (at the highest loads). The middle load was used purely for
14 the purpose of preparing the participant within the trial for the incremental increase in
15 demand, to both reduce the potential for injury and to allow them to adjust more
16 progressively to an increase external load. In order to reduce the likelihood of injury, two
17 people were positioned either side of the participants as they performed their lifts, to
18 intervene if there was a problem. In addition, there were safety supports in place on every
19 occasion (so that, if any participant had to release the weight for any reason, it would not fall
20 upon them).
21
22
23
24
25
26
27
28
29
30
31
32

33 **Statistical analysis**

34
35 The Statistical Package for the Social Sciences (SPSS), version 22, for Windows were used.
36 All data were checked for normality using the Shapiro-Wilk test. Differences between
37 conditions were evaluated using a general linear model with repeated measures, within
38 subject factor 'time of day' (2 levels) and within subject factor 'load on bar' (2 levels) and
39 interaction. To correct violations of sphericity, the degrees of freedom were corrected in a
40 normal way, using Huynh-Feldt ($\epsilon > 0.75$) or Greenhouse-Geisser ($\epsilon < 0.75$) values for ϵ , as
41 appropriate. Graphical comparisons between means and Bonferroni pairwise comparisons
42 were made where main effects were present. Only the data obtained from the first and final
43 lifts of the back squat and bench tests were analysed, as they represent the two extremities of
44 the force-velocity relationship; lesser resistances typically allowing for greater peak velocities
45 and greater resistances usually leading to larger average force values. The α -level of
46 statistical significance was set at $P < 0.05$, and values of '0.000' given by the statistics
47 package are shown here as $P < 0.0005$ (Kinear & Gray, 1995). Effect sizes (ES) were
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 calculated from the ratio of the mean difference to the pooled standard deviation. The
4 magnitude of the ES was classified as trivial (≤ 0.2), small ($> 0.2-0.6$), moderate ($> 0.6-1.2$),
5 large ($> 1.2-2.0$) and very large (> 2.0) based on guidelines from Batterham & Hopkins
6 (2006). The results are presented as the mean \pm the standard deviation throughout the text
7 unless otherwise stated. Ninety-five percent confidence intervals are presented where
8 appropriate.

11 **Results**

12 *Resting rectal temperature*

13
14 There was a significant time-of-day difference with higher resting T_{rec} values in the E (mean
15 difference = 0.49°C , $P < 0.0005$, 95 % CI: $0.34-0.64^{\circ}\text{C}$; ES=1.36) compared to the M
16 condition (See Table 1).
17
18

19
20
21 ****Insert Table 1 and Figure 2 near here****

22 *Muscle Strength Measurements*

23 **Bench press**

24
25 Daily variations were apparent for the bench press for AF, PV and tPV (See Table 1). Where
26 AF (mean difference of two loads = 10.8 N , $P < 0.0005$, 95 % CI: $6.4-15.3\text{ N}$; ES=0.21), and
27 PV values were higher in the evening than the morning (mean difference of two loads = 0.13
28 ms^{-1} , $P < 0.0005$, 95 % CI: $0.08-0.19\text{ ms}^{-1}$; ES=0.52). And tPV values occurred earlier in the
29 evening than the morning (mean difference of two loads = 0.12 s , $P = 0.001$, 95 % CI: $0.07-$
30 0.18 s ; ES=0.45). There was a main effect for load ($P < 0.05$) such that for AF (mean
31 difference = 363 N , $P = 0.001$, 95 % CI: $357-369\text{ N}$), and tPV (mean difference = 0.62 s , $P <$
32 0.0005 , 95 % CI: $0.45-0.78\text{ s}$) values were higher at greater masses on the bar than lower
33 ones. And PV (mean difference = 1.09 ms^{-1} , $P = 0.001$, 95 % CI: $0.95-1.22\text{ ms}^{-1}$; See Table 1)
34 values were lower with higher masses. There was a significant interaction for tPV such that
35 the time to peak velocity occurred earlier in the evening than the morning for the highest load
36 (mean difference of 0.62 s , $P = 0.006$; See Figure 2 and Table 1).
37
38
39
40
41
42
43
44
45

46 **Back squat**

47
48 Daily variations were apparent for the back squat for AF, PV and tPV (See Table 1). Where
49 AF (mean difference of two loads = 27.6 N , $P < 0.0005$, 95 % CI: $17.7-37.5\text{ N}$; ES=0.18),
50 and PV values were higher in the evening than the morning (mean difference of two loads =
51 0.13 ms^{-1} , $P < 0.0005$, 95 % CI: $0.08-0.17\text{ ms}^{-1}$; ES=0.51). And tPV values occurred earlier
52 in the evening than the morning (mean difference of two loads = 0.07 s , $P = 0.001$, 95 % CI:
53
54
55
56
57

0.04-0.10 s; ES=0.31). There was a main effect for load ($P < 0.05$) such that for AF (mean difference = 330 N, $P = 0.001$, 95 % CI: 320-340 N), and tPV (mean difference = 0.32 s, $P < 0.0005$, 95 % CI: 0.25-0.39 s) values were higher at greater masses on the bar than lower ones. And PV (mean difference = 0.48 ms^{-1} , $P < 0.0005$, 95 % CI: 0.37-0.59 ms^{-1}) See Table 1) values were lower with heavier masses. There was a significant interaction for tPV such that the time to peak velocity occurred earlier in the evening than the morning for the highest load (mean difference of 0.32 s, $P = 0.001$ See Figure 2 and Table 1).

Discussion

We report the MuscleLab linear encoder (Ergotest, version 4010, Norway) is capable of detecting a diurnal variation in muscle force output when used with back squat and bench press (multi-joint) exercises. More importantly, this piece of apparatus is able to detect this fluctuation for the three dimensions of muscle force output: average force, peak velocity and time to peak velocity. Where average force and peak velocity increased from morning to evening by 2.5, 12.7 % and time to peak velocity decreased by 16.6 %, for bench press (for 20 and 60 kg). And, similarly average force and peak velocity increased from morning to evening by 1.9, 8.3 % and time to peak velocity decreased by 9.8 %, for back squat (for 30 and 70 kg). Further, a diurnal variation in resting rectal temperature of 0.49°C was found, a difference previously reported. Taken together these findings are in agreement with existing literature regarding time-of-day variation in rectal temperature and muscle force output (See Edwards et al. 2013), which has consistently demonstrated that the highest values of muscle force output coincide with the daily peak of the rhythm for core temperature (15:00–18:00 h; Reilly & Waterhouse, 2009) and the least values with the lowest core temperatures (03:00–06:00 h).

A main effect for 'load' was also found as expected, for both bench press and back squat, where average force and time to peak velocity increased and peak velocity decreased, from lower to higher loads on the bar (47.1 and 80.2 %; 31.7 and 57.7 %; -42.1 and -73.9 %; $P < 0.0005$ where a negative number indicates a decrease in the variable with increasing load). That force production increases with increasing load, agrees with other authors who have used different methodology's such as equipment (force platforms and motion capture systems) and strength techniques (power clean, hang clean and hang power clean; Ammar et al. 2018). This is a typical relationship in terms of the production of overall power; as the external load increases, the rate (ms^{-1}) at which the movement is achieved is inhibited by the

1
2
3 greater mass. As such, the strength (force output/power) component of the power equation
4 (power = force x speed) is the primary influence of the increase 'load' we have observed
5 during the evening. In the present study, only the data obtained from the first and final lifts of
6 the back squat and bench tests were analysed. The lower resistance (mass on bar) creates an
7 opportunity for the participant to perform at greater velocities, whilst the greater resistance
8 will ascertain a difference in overall muscle force output when the ability to move quickly
9 has been reduced. Essentially, they represent the two extremities of the force-velocity
10 relationship, lesser resistances typically allowing for greater peak velocities and greater
11 resistances commonly leading to larger average force values.
12
13
14
15
16
17

18
19 A main finding of the current study was that an interaction was found for time of day and
20 load on the bar. Such that participants could achieve maximal rates of velocity in less time
21 during the heavier lifts, in both lower and upper body exercise in the evening. Although, the
22 mechanisms regarding development of RFD remain largely unknown (Rodriguez-Rossell et
23 al. 2017), this evening superiority in rate of force development (RFD) provides an insight as
24 to how greater peak velocities and power outputs in the evening are achieved. This time of
25 day effect on RFD may be linked with a selective effect on fast twitch protein isoforms (e.g.
26 IIA or x myosin heavy chain profile) or on the processes of excitation contraction coupling.
27
28
29
30
31
32

33 At present, the exact mechanisms for the observed diurnal variation in muscle performance
34 are unknown. We have covered this previously (Edwards et al. 2013) but in brief it has been
35 attributed to **i)** input from the body clock and proteins and peripheral clocks - that is, an
36 endogenous component to the daily variation in muscle force production - has been suggested
37 to be important (Zhang et al. 2009). The evidence relating to this is limited. **ii)** The causal
38 link between core temperature and performance has been investigated, although this has been
39 shown to be complex, due to a multiplicity of components and mechanisms which require
40 further research (Edwards et al. 2013). Further, **iii)** peripheral or muscle-related variables
41 (contractility – time to peak velocity in agreement with the current study's
42 findings, metabolism, morphology of muscle fibres, accrued damage during exercise and
43 local muscle temperature), which can be influenced by hormonal and ionic muscle
44 process variations (Reilly & Waterhouse, 2009; Tamm et al. 2009), and/or **iv)** central/
45 neurological factors (central nervous system command, alertness, motivation, and mood;
46 Castaingts et al. 2004; Giacomoni et al. 2005; Racinais et al. 2005a, 2010) have been
47 suggested.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 To the best of our knowledge, this is the first study using linear encoder technology in a
4 chronobiological research design. We have duplicated time of day and muscle force and
5 power findings previously reported in only laboratory based strength assessment (such as
6 isokinetic and isometric methods); but used complex exercises that better replicate the
7 movements associated with athletic performance (Chapman et al. 1998, LeSuer et al. 1997).
8 In the case of the lower limbs the back squat and in the case of the upper limbs/torso the
9 bench press. Power output is arguably the primary determinant of sport success; however
10 RFD is also important on its own and can be considered an underpinning mechanism of
11 power output (Bevan et al. 2010; Hawley et al. 1992; Stone et al. 2002, Taber et al. 2016).
12 The findings presented in this study are, therefore, of far greater relevance to athletic
13 performance that previously research in this area and provide an ecologically meaningful
14 assessment of the vital physical qualities (particularly rate of force development) that
15 contribute towards sport performance allowing previously published laboratory based and
16 tightly controlled experimental research to be readily translated into main-stream coaching
17 practice. In the case of the current study's findings have implications for training RFD in
18 evening and impact on sporting performance.
19
20
21
22
23
24
25
26
27
28
29

30 Lastly, effective managing, planning, periodization and phase potentiation of the specific
31 component parts of muscle force output is essential for optimal training (DeWeese et al.
32 2015). The linear encoder equipment lends itself for this use, as well as measurement of
33 adaption during micro- and mesocycles and providing real time information to submaximal
34 and one-repetition maximum tests (1 RM). The Linear encoder system can fit to any
35 traditional gym based equipment for specificity assessment, hence giving the researcher or
36 practitioner freedom of choice to choose exercises that employ the required sports-specific
37 technical abilities and important skills being developed an investigated; whilst also providing
38 insights and relationships of average force, peak velocity and time to peak velocity of specific
39 muscle force output. Unlike the traditional approaches of muscle assessment such as Wingate
40 anaerobic power test, force plate (for RFD) and/ or isokinetic or isometric dynamometry which
41 limit the range and type of movements being assessed and measurement of single muscles
42 might (as in the instance of isokinetic dynamometry) mask the force output of a group of
43 muscles (Chtourou et al. 2011, Souissi et al. 2010).
44
45
46
47
48
49
50
51
52
53

54 **Limitations**

1
2
3 We have shown clear diurnal variation for muscle performance using the linear encoder but
4 further work should extend this to a 24 h period allowing circadian characteristics to be
5 determined (such as mesor, acrophase and amplitude), but also using protocols where in the
6 endogenous component to performance can be elicited (either using a constant routine or
7 ultra-short wake-cycle experimental design). In the current study we measured only
8 resting rectal temperature, future work should measure pre and post exercise rectal and
9 muscle temperature to better explore the core and muscle temperature causal link
10 with gross muscular performance (See Edwards et al. 2013 and Pullinger et al. 2018). In
11 our study, we recruited intermediate chronotypes, future work could investigate the
12 sensitivity of linear encoders to find a diurnal variation for muscle performance using
13 outright morning or evening types.
14
15
16
17
18
19
20
21

22 **Summary**

23 The MuscleLab linear encoder provides a suitable tool for measuring diurnal variations in
24 muscle force output on multi-joint movements which more closely resemble athletic
25 performance. The advantages of this approach allow the time-of-day variation in muscle
26 force output to be measured in complex multi-joint human movements, over an array of
27 submaximal to maximal loads rather than isolated single joint maximal efforts.
28
29
30
31
32
33
34

35 **Acknowledgements**

36 We would like to thank the participants who participated in the study. We would also like to
37 acknowledge the intellectual input of Professor James M. Waterhouse who sadly passed away
38 in October 2016.
39
40
41
42
43
44

45 **Declaration of interest**

46 The authors report no conflicts of interest, the linear encoder was bought from internal funds
47 and there is no link between our laboratory and chronobiology group and the MuscleLab
48 Company. The authors are responsible for the content and writing of this article.
49
50
51
52
53

54 **Practical implications**

1
2
3 The present study supports the established notion of an evening preference (or optimal time
4 of day) for strength training, but due to the diurnal nature of the current investigation it is not
5 possible to quantify how much of this is due to the endogenous component of circadian
6 rhythmicity. That said, as the majority of athletes will live and train in typical diurnal
7 conditions such that they sleep at night and wake and are active in the day time, the findings
8 of this study support the idea strength training should be undertaken in the early to late
9 evening (17:00 to 19:00 h). Specifically where maximal or near-maximal external loads are
10 involved or a requirement for producing peak velocity outputs (explosive power work,
11 sprinting, or plyometric training for example).
12
13

14
15
16
17
18
19 Participants could achieve maximal rates of velocity in less time during the heavier lifts in
20 both lower and upper body exercise in the evening. Although the mechanisms regarding
21 development of RFD remain largely unknown (Rodriguez-Rossell et al. 2017), this evening
22 superiority in rate of force development (RFD) provides an insight as to how greater peak
23 velocities and power outputs in the evening are achieved. Research investigating the potential
24 time of day specificity to training as previously attempted by other authors (such as Sedliak et
25 al. 2008), but now employing linear encoders (measuring muscle force output, RFD and peak
26 velocity) and conducting upper and lower body tasks such as in this study are warranted.
27
28
29
30
31

32
33
34 Past laboratory based and tightly controlled experimental research can now be translated into
35 main-stream coaching practice. New and emerging intervention and methods that look to use
36 either chronobiotics such as bright light or exercise, to directly shift the body clock or dawn
37 simulators to improve mood (Thompson et al. 2014) or investigate the causal relationship
38 between performance and Trec or muscle temperature – to ultimately increase morning gross
39 muscular performance and aid athletes that have to compete or undergo training in the
40 morning could be tested in a real world context using MuscleLab (linear encoder).
41
42
43
44
45

46 **ORCID**

47
48
49 Jatin G. Burniston <http://orcid.org/0000-0001-7303-9318>

50 Ben J. Edwards <https://orcid.org/0000-0001-8913-0941>

51 Samuel A. Pullinger <http://orcid.org/0000-0001-7680-3991>

52 Colin M. Robertson <https://orcid.org/0000-0002-6186-8466>
53
54
55
56
57
58
59

List of Figures and Tables

Figure 1: Schematic of the protocol for the morning (M 07:30 h) and evening (E, 17:30 h) conditions. Rectal (T_{rec}) temperature was measured after the subjects had reclined for 30 min at the start of the protocol (solid black box). Average force, peak velocity and time to peak velocity were measured three times for each weight (black lines), for bench press and back squat respectively. There was 5-min recovery time for each lift.

Figure 2: Mean and 95 % confidence intervals (corrected for between-subject variability) for average force, peak velocity and time to peak velocity for Bench Press (for 20 and 60 kg) and Back Squat (for 30 and 70 kg) respectively at 07:30 and 17:30 h. # indicates a time of day effect; + indicates a load effect; * indicates interaction between 'load' and 'time of day'.

Table 1: Mean (\pm SD) values for rectal temperature, bench press and back squat for morning (07:30 h) and evening (17:30 h) conditions. Where only a *P*-value for time of day is given this indicates a paired *t*-test has been undertaken to remove load and interaction effects. Statistical significance ($P < 0.05$) is indicated in bold.

References

Ammar, A., Riemann, B.L., Masmoudi, L., Blaumann, M., Abdelkarim, O. & Hökelmann, A. 2018. Kinetic and kinematic patterns during high intensity clean movement: searching for optimal load. *Journal of Sport Sciences*. 36(12):1319-1330. doi: 10.1080/02640414.2017.1376521. PMID: 28895467.

Baechle, TR, Earle, RW. *Essentials of strength training and conditioning* (4th edition), 2015; Champaign, IL: Human Kinetics.

Batterham AM, Hopkins WG. 2006. Making meaningful inferences about magnitudes. *Int J Sports Physiol Perform* 1(1):50–57. doi:10.1123/ijsp.1.1.50. PMID: 19114737.

Bambaeichi E, Reilly T, Cable N, Giacomoni M. 2005. The influence of time of day and partial sleep loss on muscle strength in eumenorrheic females. *Ergonomics*. 48(11-14):1499–511. doi:10.1080/00140130500101437. PMID: 16338716.

Bevan HR, Bunce PJ, Owen NJ, Bennett MA, Cook CJ, Cunningham DJ, Kilduff LP. 2010. Optimal loading for the development of peak power output in professional rugby players. *Journal of strength and conditioning research / National Strength & Conditioning Association*, 24(1), 43-47. doi:10.1519/JSC.0b013e3181c63c64.

Bosco C, Belli A, Astrua M, Tihanyi J, Pozzo R, Kellis S, Tsarpela O, Foti C, Manno R, Tranquilli C. 1995. A dynamometer for evaluation of dynamic muscle work. *Eur J Appl Physiol Occup Physiol*. 70(5):379-86. PMID: 7671871.

Bougard C, Bessot N, Moussay S, Sesboüé B, Gauthier A. 2009. Effects of waking time and breakfast intake prior to evaluation of physical performance in the early morning. *Chronobiol Int*. 26(2):307-323. doi:10.1080/07420520902774532. PMID: 19212843.

Buckner SL, Jessee MB, Mattocks KT, Mouser JG, Counts BR, Dankel SJ, Loenneke JP. 2017. Determining Strength: A Case for Multiple Methods of Measurement. *Sports Medicine*. 47(2):193-195. doi: 10.1007/s40279-016-0580-3. PMID: 27380100.

1
2
3 Callard D, Davenne D, Gauthier A, Lagarde D, Van Hoecke J. 2000. Circadian Rhythms in
4 Human Muscular Efficiency: Continuous Physical Exercise Versus Continuous Rest. A
5 Crossover Study. *Chronobiol. Int.* 17(5):693-704. doi:10.1081/CBI-100101075.
6 PMID:11023216.
7
8

9
10 Castaingts V, Martin A, VanHoecke J, Perot C. 2004. Neuromuscular efficiency of the triceps
11 surae in induced and voluntary contractions: morning and evening evaluations. *Chronobiol.*
12 *Int.* 21(4-5):631–643. doi: 10.1081/CBI-120039207. PMID:15470959.
13
14

15
16 Chapman PP, Whitehead JR, Binkert RH. 1998. The 225-lb Reps-to-Fatigue Test as a
17 Submaximal Estimate of 1-RM Bench Press Performance in College Football Players. *J*
18 *Strength Cond Res.* 12(4):258-261. doi:10.1519/00124278-199811000-00010
19
20

21
22 Chiu LZ, Fry AC, Weiss LW, Schilling BK, Brown LE, Smith SL. 2003. Postactivation
23 potentiation response in athletic and recreationally trained individuals. *Strength Cond Res.*
24 17(4):671-677. PMID:14636093.
25
26

27
28 Chtourou H, Zarrouk N, Chaouachi A, Dogui M, Behm DG, Chamari K, Hug F, Souissi, N.
29 2011. Diurnal variation in wingate-test performance and associated electromyographic
30 parameters. *Chronobiol Int.* 28(8):706–713. doi:10.3109/07420528.2011.596295.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

61
62 Comfort P, Allen M, & Graham-Smith P. (2011). Comparisons of Peak Ground
63 Reaction Force and Rate of Force Development During Variations of the Power Clean. *The*
64 *Journal of Strength and Conditioning Research.* 25(5):1235-1239. PMID:
65 21522071 doi: 10.1519/JSC.0b013e3181d6dc0d.
66
67
68
69
70

71
72 DeWeese BH, Sams ML, Williams JH, Bellon CR. 2015. The nature of speed: Enhancing
73 sprint abilities through a short to long training approach, in: *Techniques in Track & Field and*
74 *Cross Country.* pp 8-24.
75
76
77
78
79
80

81
82 Drust B, Waterhouse J, Atkinson G, Edwards B, Reilly T. 2005. Circadian rhythms in sports
83 performance: an update. *Chronobiol Int.* 22(1):21-40. doi:10.1081/CBI-200041039.
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

1
2
3 Edwards BJ, Pullinger SA, Kerry JW, Robinson WR, Reilly TP, Robertson CM, Waterhouse
4 JM. 2013. Does raising morning rectal temperature to evening levels offset the diurnal
5 variation in muscle force production? *Chronobiol Int.* 30(4):486-501.
6
7 doi:10.3109/07420528.2012.741174. PMID: 23281719.
8
9

10
11
12 Edwards B, Waterhouse J, Reilly T, Atkinson G. 2002. A comparison of the suitabilities of
13 rectal, gut, and insulated axilla temperatures for measurement of the circadian rhythm of core
14 temperature in field studies. *Chronobiol Int.* 19(3):579-597. doi:10.1081/CBI-120004227.
15
16 PMID:12069039.
17
18

19
20
21 Folkard S, Monk TH. 1979. Towards a predictive test of adjustment to shiftwork.
22
23 *Ergonomics*, 22(1):79-91. doi:10.1080/00140137908924591. PMID: 436816.
24
25

26
27 García-Ramos A, Stirn I, Strojnik V, Padial P, De la Fuente B, Argüelles-Cienfuegos J,
28 Feriche B. 2016. Comparison of the force-, velocity-, and power-time curves recorded with a
29 force plate and a linear velocity transducer. *Sports Biomech.* 15(3):329-41.
30
31 doi:10.1080/14763141.2016.1161821. PMID: 27239893.
32
33

34
35 Giacomoni M, Edwards B, Bambaiechi E, Waterhouse J, Reilly T. 2005. Gender differences
36 in the circadian variations in muscle strength assessed with and without superimposed
37 electrical twitches. *Ergonomics.* 48(11-14):1473-1487. doi:10.1080/00140130500101452.
38
39 PMID:16338714.
40
41

42
43 Häkkinen K, Komi. PV. 1985. Changes in electrical and mechanical behaviour of leg
44 extensor muscles during heavy resistance strength training. *Scand J Sports Sci.* 7(2):55-64.
45
46

47
48 Hakkinen K. 1989. Neuromuscular and hormonal adaptations during strength and power
49 training. *J Sports Med Phys Fitness.* 29(1):9-26. PMID: 2671501.
50
51

52
53 Harris NK, Cronin J, Taylor K-L, Jidovsteff B, Sheppard, JM. 2010. Understanding Position
54 Transducer Technology for Strength and Conditioning Practitioners. *Strength Cond J.*
55
56 32(4):66-79. doi:10.1519/SSC.0b013e3181eb341b.
57
58

1
2
3
4 Hawley JA, Williams, MM, Vickovic MM, Handcock, P. J. 1992. Muscle power predicts
5 freestyle swimming performance. *Br J Sports Med.* 26(3):151-155.
6 doi:10.1136/bjism.26.3.151. PMID:1422650.
7
8

9
10 Jennings CL, Viljoen W, Durandt J, Lambert MI. 2005; The reliability of the FitroDyne as a
11 measure of muscle power. *J Strength Cond Res.* 19(4):859-63. doi:10.1519/R-15984.1.
12 PMID:16287353.
13
14

15
16
17 Kinear PR, Gray LD. (1995). *SPSS for windows made simple.* London:Lawrence Ehbaum.
18 pp. 228.
19
20

21
22 LeSuer DA, McCormick JH, Mayhew JL, Wasserstein RL, Arnold MD. 1997. The Accuracy
23 of Prediction Equations for Estimating 1-RM Performance in the Bench Press, Squat and
24 Deadlift. *J Strength Cond Res,* 11(4):211-213. doi:10.1519/00124278-199711000-00001.
25
26

27
28 Monk TH, Leng VC. 1982. Time of day effects in simple repetitive tasks: some possible
29 mechanisms. *Acta Psychol.* 51:207-221. doi:10.1016/0001-6918(82)90035-X
30
31

32
33 Morton JP, Atkinson G, MacLaren DPM, Cable NT, Gilbert G, Broome C, McArdle A, Drust
34 B. 2005. Reliability of maximal muscle force and voluntary activation as markers of exercise-
35 induced muscle damage. *Eur. J. Appl. Physiol.* 94:541-548. doi 10.1007/s00421-005-1373-9.
36
37

38
39 O'Donnell S, Tavares F, McMaster D, Chambers S, Driller M. 2017. The validity and
40 reliability of the GymAware linear position transducer for measuring counter-movement
41 jump performance in female athletes. *Measurement in Physical Education and Exercise*
42 *Science.* 22:101-107. doi:10.1080/1091367X.2017.1399892.
43
44
45

46
47 Pareja-Blanco F, Rodríguez-Rosell D, Sánchez-Medina L, Sanchis-Moysi J, Dorado C,
48 Mora-Custodio R, Yáñez-García JM, Morales-Alamo D, Pérez-Suárez I, Calbet, JAL,
49 González-Badillo JJ. 2016. Effects of velocity loss during resistance training on athletic
50 performance, strength gains and muscle adaptations. *Scandinavian Journal of Medicine and*
51 *Science in Sports.* 27(7):724-735. PMID: 27038416. doi: 10.1111/sms.12678.
52
53
54
55

1
2
3
4 Pullinger SA, Oksa J, Brocklehurst EL, Iveson RP, Newlove A, Burniston JG, Doran DA,
5 Waterhouse JM, Edwards BJ. 2018. Controlling rectal and muscle temperatures: Can we
6 offset diurnal variation in repeated sprint performance? *Chronobiol Int.* 1-10. doi:
7 10.1080/07420528.2018.1444626. PMID: 29557676. [Epub ahead of print].
8
9

10
11
12 Racinais S, Connes P, Bishop D, Blonc S, Hue O. 2005. Morning versus evening power
13 output and repeated-sprint ability. *Chronobiol Int* 22(6):1029–39.
14 doi:10.1080/07420520500397918. PMID: 16393706.
15
16

17
18
19 Racinais S, Oksa J. 2010. Temperature and neuromuscular function. *Scan J Med Sci Sports*
20 20(3):1–18. doi:10.1111/j.1600-0838.2010.01204.x. PMID: 21029186.
21
22

23
24 Reilly T, Waterhouse J. 2009. Sports performance: is there evidence that the body clock plays
25 a role? *Eur J Appl Physiol.* 106(3):321-332. doi:10.1007/s00421-009-1066-x. PMID:
26 19418063.
27
28

29
30 Rodríguez-Rosell D, Pareja-Blanco F, Aagaard P, González-Badillo JJ. 2017. Physiological
31 and methodological aspects of rate of force development assessment in human skeletal
32 muscle. *Clin Physiol Funct Imaging.* doi: 10.1111/cpf.12495. [Epub ahead of print]
33
34

35
36
37 Sedliak M, Finni T, Peltonen J, Häkkinen, JK. 2008. Effect of time-of-day-specific strength
38 training on maximum strength and EMG activity of the leg extensors in men. *J Sports Sci.*
39 26(10):1005-1014. doi:10.1080/02640410801930150. PMID: 18608836.
40
41

42
43 Smith C, Reilly C, Midkiff K. 1989. Evaluation of three circadian rhythm questionnaires with
44 suggestions for an improved measure of morningness. *J Appl Psych.* 74(5):728-738.
45 doi:10.1037/0021-9010.74.5.728. PMID: 2793773.
46
47
48

49
50
51 Souissi N, Driss T, Chamari K, Vandewalle H, Davenne D, Gam A, Fillard J-R, Joussetin E.
52 2010. Diurnal variation in wingate test performances: influences of active warm-up.
53 *Chronobiol Int.* 27(3):640-652. doi:10.3109/07420528.2010.483157. PMID: 20524806.
54
55
56

1
2
3
4 Stone MH, Moir G., Glaister, M, Sanders R. 2002. How much strength is necessary? *Phys*
5 *Ther Sport.* 3(2): 88-96. doi:10.1054/ptsp.2001.0102.
6
7

8
9 Taber C, Bellon CR, Abbott H, Bingham GE. 2016. Roles of maximal strength and rate of
10 force development in maximizing muscular power. *Strength & conditioning Journal*
11 38(1):71-78. doi:10.1519/SSC.000000000000193.
12
13


14
15 Tamm AS, Largerquist O, Ley AL, Collins DF. 2009. Chronotype influences diurnal
16 variations in the excitability of the human motor cortex and the ability to generate torque
17 during a maximum voluntary contraction. *J. Biol. Rhythms.* 24(3):211–224.
18 doi:10.1177/0748730409334135. PMID:19465698 .
19
20
21
22

23
24 Thompson A, Jones H, Gregson W, Atkinson G. 2014. Effects of dawn simulation on
25 markers of sleep inertia and postwaking performance in humans. *Eur J Appl Physiol.*
26 114(5):1049–56. doi: 10.1007/s00421-014-2831-z. PMID: 2450 9892.
27
28
29

30
31 Drouin JM, Valovich TC, Shultz SJ, Gansneder BM, Perrin DH. 2004. Reliability and
32 validity of the Biodex System 3 Pro Isokinetic Dynamometer velocity, torque and position
33 measurements. *European Journal of Applied Physiology.* 91(1):22-29. doi: 10.1007/s00421-
34 003-0933-0.
35
36
37

38
39 Zhang X, Dube TJ, Esser KA. 2009. The role of clock genes in 685 cardiometabolic disease:
40 Working around the clock: Circadian rhythms and skeletal muscle. *J Appl Physiol*
41 107(5):1647–54. doi:10.1152/jappphysiol.00725.2009. PMID: 19696362.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60







1 *Insert probe*
2 *and lay down*
3



32 **Figure 1.**

Bench press

Back squat



20 60
Load (kg)

30 70
Load (kg)

Table 1

Measure	Variable	Morning (M)	Evening (E)	<i>P</i> -value Time of day
Resting Rectal temperature	T _{rec} (°C)	36.5 ± 0.3	37.0 ± 0.4	<0.0005
Bench Press				
20 kg	AF (N)	317.3 ± 46.5	328.3 ± 48.6	= 0.0004
	PV (ms ⁻¹)	1.81 ± 0.34	1.94 ± 0.35	= 0.002
	TpV (s)	0.31 ± 0.07	0.27 ± 0.07	< 0.0005
60 kg	AF (N)	680.5 ± 51.1	691.1 ± 56.5	< 0.0005
	PV (ms ⁻¹)	0.73 ± 0.26	0.86 ± 0.34	< 0.0005
	TpV (s)	1.01 ± 0.58	0.80 ± 0.43	< 0.0005
Back Squat				
30 kg	AF (N)	1326.3 ± 150.4	1352.4 ± 162.0	= 0.0002
	PV (ms ⁻¹)	1.76 ± 0.28	1.91 ± 0.32	= 0.0002
	TpV (s)	0.45 ± 0.15	0.42 ± 0.13	= 0.011
70 kg	AF (N)	1654.9 ± 156.2	1683.9 ± 155.7	< 0.0005
	PV (ms ⁻¹)	1.30 ± 0.29	1.41 ± 0.30	< 0.0005
	TpV (s)	0.81 ± 0.32	0.71 ± 0.28	< 0.0005