



Journal of Aggression, Conflict and Peace Research

**Attachment Style, Psychotic Phenomena and the  
Relationship with Aggression: An Investigation in a General  
Population Sample**

Journal:	<i>Journal of Aggression, Conflict and Peace Research</i>
Manuscript ID	JACPR-04-2018-0356.R2
Manuscript Type:	Empirical Paper
Keywords:	psychosis, attachment, aggression, risk, schizophrenia, trauma

SCHOLARONE™  
Manuscripts

1  
2  
3 **Attachment Style, Psychotic Phenomena and the Relationship with Aggression: An**  
4 **Investigation in a General Population Sample**  
5  
6  
7  
8  
9

10 **Abstract**

11  
12 Purpose: This study aimed to explore the relationship between attachment style, sub-clinical  
13 symptoms of psychosis and aggression in a general population sample.  
14  
15

16  
17 Design: Using both convenience and snowball sampling, participants in the community  
18 (n=213) completed an online questionnaire including previously validated measures of adult  
19 attachment, aggression and psychotic like events.  
20  
21

22  
23 Findings: Results suggested that there were statistically significant correlations between all  
24 study variables. Multiple linear regression demonstrated that total psychotic-like experiences and  
25 attachment scores significantly predicted variance in total aggression. Moderation approaches  
26 revealed that the relationship between psychotic-like events and aggression was stronger in  
27 individuals with more insecure attachment styles.  
28  
29  
30

31  
32  
33 Research limitations/implications: This generalisability of the results is compromised by the  
34 sampling methodology and the use of self-report tools. However, the significant results would support  
35 larger scale replications investigating similar variables.  
36  
37  
38

39  
40 Originality/value: This study suggests there is a relationship between psychotic like  
41 experiences and facets of aggression in the general population. The findings suggest that attachment  
42 is a contributing factor to aggression associated with psychotic like experiences, and highlight the  
43 need for similar investigations within clinical samples. The results imply that attachment may be a  
44 useful construct for explanatory models of the relationship between adverse childhood experiences,  
45 psychotic experiences and aggression.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Introduction

This study aimed to explore the relationships between attachment, sub-clinical symptoms of psychosis and aggression in a community sample.

### Attachment

Attachment theory suggests that human beings are predisposed to form bonds with primary care-givers. A child's experiences with attachment figures will influence his or her internal working models of self and others (Bowlby, 1982), forming a prototype for later relationships (Bartholomew and Horowitz, 1991). The quality of early attachments is related to the child's reliance on attachment figures as a source of security (Ainsworth et al., 1978). A child's internal model of others is influenced by the degree to which the attachment figure is seen as someone who is responsive to the need for support and protection; their internal model of self is influenced by whether or not the self is judged to be someone to whom the attachment figure responds to in a helpful way (Bowlby, 1973).

Adult models of attachment develop Bowlby's (1982) theory of the internal working model, by conceptualising a person's model of self and others as anxiety (sometimes termed dependence) and avoidance in attachment (Crowell, Fraley and Shaver, 1999). Bartholomew and Horowitz (1991) proposed four adult attachment styles on the basis of these dimensions:

- Secure: Individuals who have a positive model of self (low anxiety), and a positive model of others (low avoidance), will feel comfortable with intimacy and autonomy.
- Preoccupied: Individuals who have a negative model of self (high anxiety), but a positive model of others, (low avoidance) may be preoccupied with retaining relationships
- Dismissing: Individuals who have a positive model of self (low anxiety), but a negative view of others, (high avoidance) may be dismissing of intimate relationships
- Fearful: Individuals who have a negative model of self (high anxiety), and a negative model of others, (high avoidance), may be fearful of intimacy and close relationships.

### Psychosis

1  
2  
3 Psychotic experiences have historically been conceptualised as symptoms of mental disorder  
4 which include impairments in an individual's perception of reality, such as hallucinations and  
5 delusions (e.g. Parker, 2014). While these experiences have traditionally been dichotomised as  
6 present or absent for diagnostic purposes (van Os et al., 2000) a body of evidence has demonstrated  
7 that these symptoms occur in the general population as well as clinical samples (e.g. van Os et al.,  
8 2000; Verdoux and van Os, 2002; Stefanis et al., 2002 Loch et al., 2011). The presence of a range of  
9 psychotic and psychotic-like experiences (PLEs) in individuals who do not meet the diagnostic  
10 threshold of a case in need of treatment has informed the concept of a continuum of psychosis  
11 (Esterberg and Compton, 2009). Investigating these experiences in non-clinical samples allows for  
12 exploration of a range of psychotic experiences and correlates, rather than including only those at the  
13 endpoint of the distribution of the psychotic dimension (Verdoux and van Os, 2002).  
14  
15  
16  
17  
18  
19  
20  
21  
22

### 23 *Psychosis and Attachment*

24  
25  
26 Across research designs, evidence suggests that adverse events in childhood significantly  
27 increase the likelihood of an individual experiencing symptoms of psychosis in adulthood. A review  
28 (Read et al., 2005) found that of individuals with psychosis (from both inpatient and community  
29 samples) 69% of females and 59% of males had been subject to abuse in childhood. This research  
30 compliments earlier findings that individuals with psychosis are more likely to have experienced  
31 parental loss (Granville-Grossman, 1966). Similarly, Morgan et al. (2007) found that, controlling for  
32 parental history of mental illness, individuals with first-episode psychosis were more likely to have  
33 been separated from their parents for at least a year during childhood and to have experienced  
34 maternal death. A more recent meta-analysis identified a medium-large effect size of increased rates  
35 of childhood adversity in individuals with schizophrenia in comparison to control samples (Matheson  
36 et al., 2013).  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 These adverse childhood events may have a negative impact on attachment style. Models of  
47 psychosis highlight that negative beliefs about the self and the world can create vulnerability and  
48 maintenance (Penn et al., 1997). This suggests that previous experiences of relationships and  
49 traumas contribute to negative beliefs and symptoms of psychosis (Garety et al., 2001). Current  
50 experiences of relationships and interpersonal functioning have been associated with relapse and  
51 recovery in individuals with psychosis. Warner and Atkinson (1988) found that patients who perceived  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 their parents negatively and had less contact with them had more relapses and a more severe course  
4 of schizophrenia.

5  
6  
7 Berry et al. (2006) found significant associations between insecure attachment and non-  
8 clinical psychotic phenomena in a community sample. Positive psychotic phenomena were associated  
9 with anxiety in attachment, and social anhedonia with avoidance in attachment. This suggests that  
10 attachment may be a useful construct in understanding PLEs in the general population.  
11  
12  
13

#### 14 15 Attachment and Aggression

16  
17  
18 There is an established relationship between insecure attachment, aggression and offending  
19 behaviours. A literature review of offenders and attachment style found that insecure attachment was  
20 over-represented in the general offending population, with prevalence ranging from 64-97% (Ratnip,  
21 2013). Some results suggested no differences in attachment style across offenders, with others  
22 suggesting this relationship was stronger in sexual offenders.  
23  
24  
25

26  
27  
28 Other evidence (Bartholomew and Horowitz, 1991; Kobak and Sceery, 1988; Mallinckrodt,  
29 2000) suggests that attachment style is related to different expressions of anger and aggression. For  
30 example, dismissing attachment has been associated with hostility, interpersonal coldness, and  
31 emotional detachment; pre-occupied attachment has been associated with over-intrusive behaviour,  
32 dominance and aggression and fearful attachment have been associated with difficulties in  
33 assertiveness, social inhibition, anger and aggression.  
34  
35  
36  
37  
38

39  
40  
41 Mikulincer and Shaver (2007) found that individuals with a dismissive attachment style were  
42 more likely to reactively express anger, as well as be more critical and devaluing of other's emotional  
43 needs and controlling through their anger. Individuals with a pre-occupied attachment style were more  
44 likely to express irritation and anxiety and externalise blame, as well as find it difficult to be soothed by  
45 others. Individuals with a fearful attachment style were more likely to show extreme expressions of  
46 anger and be violent towards themselves and others. Ratip (2013) also found that of secure  
47 psychiatric patients, those with a secure attachment style had lower anger temperament scores  
48 compared to those with insecure attachment styles.  
49  
50  
51  
52  
53

#### 54 Violence and Psychosis

1  
2  
3 A significant body of research has investigated the relationship between violence and  
4 psychosis. A meta-analysis of 204 studies, suggested psychosis was associated with a 49-68%  
5 increase in the odds of violence (Douglas, Guy and Hart, 2009). However, this review highlighted that  
6 there was substantial dispersion amongst effect sizes as a function of moderator variables and  
7 challenges synthesising data due to methodological differences. This review identified positive  
8 symptoms of psychosis (among others) as associated with violence, however the authors highlight the  
9 need for further investigation of the symptom-violence association. The authors conclude that  
10 although individuals with psychosis were at an increased risk of violence in comparison to individuals  
11 with no mental disorder, there was no elevated risk when compared to individuals with externalising  
12 psychopathologies.  
13  
14  
15  
16  
17  
18  
19  
20

21 A comprehensive review of violence in first episode psychosis suggested that while rates of  
22 violence were relatively high in this population (in comparison to individuals in a later stage of illness),  
23 the frequency of violence decreased with severity. Severity of psychotic symptoms was associated  
24 with serious violence, but not with less severe forms of aggression (Large and Niessen, 2011).  
25 Similar relationships have been explored in community samples, suggesting that PLEs are associated  
26 with aggressive behaviour, in particular hallucinatory behaviour (Nederlof, Muris and Hovens, 2012).  
27 Mojtabai (2006) found that unusual perceptual experiences and paranoid ideation were associated  
28 with violence. Kinoshita et al. (2011) suggested that most forms of PLE were associated with violence  
29 towards objects, and feelings of persecution and hearing voices were associated with violence  
30 towards people.  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 Research exploring this link suggests that variables such as personality (Eysenck and  
41 Eysenck, 1970), dispositional anxiety, anger (Posner, Russell and Peterson, 2005) and drug use  
42 (Allen et al., 1997) may be relevant. A number of studies have suggested that substance use  
43 accounts for a large proportion of violence in individuals with psychosis (Soyka, 2000; Fazel et al.,  
44 2009). However, while substance use may increase the risk of criminality in individuals with  
45 psychosis, it does not fully explain the risk of violence (Short et al., 2013). A recent meta-analysis  
46 suggested that experiences of childhood maltreatment are a significant risk factor for violence to  
47 others in individuals with psychosis (Green, Browne and Chou, 2017), emphasising the need to  
48 explore the mechanisms of this relationship.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 In a sample of individuals with schizophrenia, attachment style, mentalisation (the ability to  
4 understand and attribute cognitive and affective mental states to oneself and others) and personality  
5 variables have been found to discriminate between impulsive and premeditated aggression (Bo et al.,  
6 2013). In particular, pre-meditated aggression was associated with: positive representations of self  
7 and negative representations of other; diminished mentalisation and more severe personality  
8 pathology. However, given evidence that the association between aggression towards others and  
9 psychotic experiences is present in non-clinical samples, there is a need to investigate the role of  
10 attachment in this relationship in the general population.  
11  
12  
13  
14  
15  
16  
17

18 Given the established links between attachment, violence and psychosis, the primary aim of  
19 this study is to investigate whether attachment style explains some of the variance in aggression  
20 towards others associated with PLEs in a general population sample. The secondary aim is to  
21 establish whether this relationship changes depending on either the type of attachment, type of PLE  
22 or type of aggression.  
23  
24  
25  
26  
27

## 28 **Method and Materials**

### 29 Participants and Procedure

30  
31  
32 An a-priori power calculation using g\*power (Faul et al., 2009) suggested a minimum sample  
33 size of 138 participants to achieve 95% power for fixed model, linear multiple regression with 5  
34 variables. Recruitment was online via convenience sampling, participants were approached directly  
35 via email and indirectly via social media (facebook and a 'call for participants' website). Participants  
36 were encouraged to share the link to the study (snowball sampling). In total, 213 respondents  
37 completed the survey. Respondents were taken to an information page, advising them they would be  
38 participating in a study exploring 'attachment, experiences and emotions', and explaining their right to  
39 withdraw and anonymity. If participants consented, they were asked to complete demographic  
40 information before psychometric assessments. At the end of the survey, debrief information and  
41 details of supportive organisations were provided.  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 The research was given a favourable opinion by the University of Nottingham's Faculty of  
52 Medicine and Health Sciences Research Ethics Committee.  
53  
54  
55

### 56 Measures

1  
2  
3 *Psychosis: Community Assessment Psychic Experience (CAPE)-42; Stefanis et al.*  
4  
5 (2002)

6  
7 The CAPE-42 is a 42-item scale that has proved to be a stable, reliable and valid assessment  
8 of self-reported PLE (Mossaheb et al. 2012; Fonseca-Pedrero et al., 2012). The instrument has 18  
9 positive-symptom items, 14 negative-symptom items and 8 depressive-symptom items. Stefanis et al.  
10 (2002) identified that the scale is valid, and that dimensions of psychosis are approximately normally  
11 distributed in the general population. The tool has separate symptom and distress scales, only the  
12 symptom scale was used in the current study. Total scale internal consistency (Cronbach's alpha)  
13 was excellent in this study ( $\alpha=.925$ ).  
14  
15  
16  
17  
18  
19

20 *Adult Attachment: Psychosis Attachment Measure (PAM); Berry et al. (2006).*

21  
22  
23 The PAM was developed for use in both clinical and non-clinical populations. The scale has  
24 demonstrated good internal reliability and good-to-acceptable test-retest reliability (Berry et al., 2006).  
25 The scale has 16 items, each on a four point likert scale. Scores translate into two subscales: anxiety  
26 and avoidance. Eight items assess attachment avoidance and eight items assess attachment anxiety.  
27  
28  
29 Total scale internal consistency was good in this study ( $\alpha=.814$ ).  
30  
31  
32

33 *Aggression: The Aggression Questionnaire, Buss and Perry (1992)*

34  
35 The aggression questionnaire is a self-report tool designed to measure facets of aggression  
36 towards others, and has demonstrated strong internal consistency and stability over time (Buss and  
37 Perry, 1992). More recent studies have identified good test-retest reliability and internal consistency,  
38 for both subscales and general questionnaire (Harris, 1995; Gerevich, Bacskai and Czobor, 2007).  
39  
40  
41

42 The scale provides four sub-scales:

- 43  
44
- 45 • Physical aggression – This scale relates to behavioural aspects of physical  
46 aggression including actual violence.
  - 47 • Verbal aggression – This scale relates to behavioural aspects of verbal aggression,  
48 including arguments and disagreements.
  - 49 • Anger – This scale relates to the affective experience of anger and difficulties  
50 controlling temper.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- Hostility – This scale relates to cognitive aspects of hostility, including suspicion and resentment.

The scale is comprised of 29 items, each answered on a five-point likert scale. Total scale internal consistency was excellent in this study ( $\alpha=.915$ ).

#### *Substance Use*

Questions regarding substance use (including cannabis, legal high and alcohol use) were included as potential confounding variables. This included frequency of intake of alcohol, cannabis and other illicit substances each week, and how often an individual would drink more than three drinks on one occasion.

#### *Data Analysis*

Missing data were replaced by mean imputation to maintain sample size, and power of statistical analysis. No data were excluded from analyses, as no individual missed more than 1 question per scale.

Data were positively skewed but unimodal. A 1-sample K-test was conducted on each scale, all of which were significant ( $p<.01$ ), suggesting distributions were non- normal. Correlational data was analysed using non-parametric analyses. Data were reviewed for assumptions of linear multiple regression, and outliers via distance (more than 3 standard deviations from the norm) and influence were removed from analysis. All analyses were conducted using IBM SPSS Statistics 22.

### **Results**

Of 213 participants, the median age was 32.64 years (range 18-64), 42 were male and 169 were female. 82.9% of the sample identified themselves as British, and the majority of participants had an Undergraduate level of education (1 PhD, 9 Doctorate, 47 postgraduate, 102 undergraduate, 34 A- Levels, 18 GCSEs).

#### *Spearman's Correlation Coefficients*

A Spearman's non-parametric correlation was used to assess bivariate relationships between variables. Positive, statistically significant, relationships were found between all total measure scores.

1  
2  
3 A strong positive correlation was found between total attachment and total PLE scores ( $r = .643$ ,  $p$   
4  $<.001$ ); a weaker positive correlation was seen between total attachment and total aggression scores  
5 ( $r = .346$ ,  $p <.001$ ) a moderate positive correlation was seen between total PLE scores and total  
6 aggression ( $r = .546$ ,  $p <.001$ ).  
7  
8  
9

10 Relationships differed between sub-scales. All PLE domains showed positive, statistically  
11 significant ( $p \leq .001$ ) correlations with aggression scores. PLE domains (positive, depressive and  
12 negative) were most closely associated with hostility scores ( $r(211) = .441$ ,  $p <.001$ .;  $r(211) = .576$ ,  
13  $p <.001$ .;  $r(211) = .504$ ,  $p <.001$ .) respectively. Physical aggression was most strongly correlated with  
14 positive symptoms;  $r(211) = .319$ ,  $p <.001$ . Verbal aggression was most strongly correlated with negative  
15 symptoms;  $r(211) = .206$ ,  $p <.002$ . Anger scores were most strongly correlated with depressive  
16 symptoms;  $r(211) = .429$ ,  $p <.001$ . Hostility scores were also most strongly correlated with depressive  
17 symptom scores;  $r(211) = .576$ ,  $p <.001$ .  
18  
19  
20  
21  
22  
23  
24  
25

26 Anxious and Avoidant attachment were both positively correlated with hostility;  $r(211) = .486$ ,  
27  $p <.001$ ;  $r(211) = .417$ ,  $p <.001$  respectively. Physical aggression was most strongly correlated with  
28 avoidant attachment scores, with a weak, positive relationship;  $r(211) = .184$ ,  $p <.01$ . Verbal Aggression  
29 was not significantly related to Avoidant or Anxious attachment. Anger and Hostility Scores were both  
30 positively related to Anxious Attachment ( $r(211) = .265$ ,  $p <.001$ ;  $r(211) = .486$ ,  $p <.001$ . respectively).  
31  
32  
33  
34  
35

36 There were positive, statistically significant, relationships between all the sub-domains of PLE  
37 and Attachment. Positive symptoms were most strongly correlated with avoidant attachment  
38 ( $r(211) = .451$ ,  $p <.001$ ), depressive symptoms were most strongly correlated with anxious attachment  
39 ( $r(211) = .554$ ,  $p <.001$ ), and negative symptoms are most strongly correlated with avoidant attachment  
40 ( $r(211) = .480$ ,  $p <.001$ ).  
41  
42  
43  
44

#### 45 Multiple Linear Regression

##### 46 *Influence of total PLE scores and total attachment scores on total aggression scores*

47  
48  
49  
50 A multiple-linear regression analysis was completed, including total PLE and attachment  
51 scores as predictors, and total aggression score as the criterion variable. Assumptions for multiple-  
52 linear regression were met, and data were assessed for multi-collinearity and removed if applicable.  
53  
54  
55  
56 Factors were entered using the 'enter' method.  
57  
58  
59  
60

1  
2  
3 The results indicated that the two factors explained 32.7% of the variance in total aggression  
4 (adjusted  $R^2=32.7$ ,  $F(2,211)=46.864$ ,  $p<0.001$ ). Total PLE score was the most influential, and the only  
5 significant predictor in the model ( $\beta=.573$ ,  $t=7.579$ ,  $p<0.0001$ ), total attachment score did not  
6 significantly predict additional variance in total aggression ( $\beta=-.002$ ,  $t=-.010$ ,  $p>0.05$ ).  
7  
8  
9

10 The model became slightly stronger when measures of legal high and cannabis use were  
11 entered prior to attachment and PLE scores (adjusted  $R^2=41.2$ ,  $F(4,209)=35.87$ ,  $p<0.001$ ). However,  
12 neither cannabis ( $\beta=.007$ ,  $t=.126$ ,  $p>0.05$ ) nor legal high use ( $\beta=.070$ ,  $t=.126$ ,  $p>0.05$ ) were significant  
13 predictors in the model.  
14  
15  
16  
17

#### 18 *Influence of Subscales of PLE and Attachment on Aggression Scores*

19 Analysis was then conducted to investigate the impact of sub-scales of attachment (anxiety  
20 and avoidance) and PLEs (positive, negative and depressive symptoms) on total aggression.  
21  
22

23 The results indicated that the five factors explained 41.1% of the variance in total aggression  
24 scores (adjusted  $R^2=41.1$ ,  $F(5,208)=28.55$ ,  $p<0.001$ ). Positive symptoms ( $\beta=.223$ ,  $t=3.070$ ,  $p<0.01$ ),  
25 negative symptoms ( $\beta=.226$ ,  $t=2.805$ ,  $p>0.01$ ) and depressive symptoms ( $\beta=.253$ ,  $t=2.996$ ,  $p>0.01$ )  
26 were significant predictors in the model. Neither anxious ( $\beta=.029$ ,  $t=.425$ ,  $p>0.05$ ) nor avoidant  
27 ( $\beta=.021$ ,  $t=.302$ ,  $p>0.05$ ) attachment contributed significantly to the prediction of total aggression.  
28  
29  
30  
31  
32  
33  
34

35 The five-factor regression analysis was also completed for aggression sub-domains. These  
36 results are summarised in Table 1.  
37  
38  
39  
40  
41

42 *Table 1. About here*  
43  
44  
45  
46  
47

48 The five-factor analysis provides the strongest explanatory model for hostility, and the  
49 weakest for verbal aggression. Across all models, PLE sub-domains were significant predictors,  
50 attachment sub-scales were not.  
51  
52  
53

#### 54 *Interaction Between Total PLE and Attachment Scores in Predicting Total Aggression*

1  
2  
3 An interaction term was created for total PLE x total Attachment scores, to see if this  
4 interaction predicted variance in total aggression scores. Data were centred to account for collinearity  
5 in this analysis. While the total model was significant (adjusted  $R^2=37.2$ ,  $F(3,209)=42.80$ ,  $p<0.001$ ),  
6 the interaction term did not significantly predict variance in aggression ( $\beta=.033$ ,  $t=.0483$ ,  $p>0.05$ ). In  
7 keeping with the initial two-factor model, the only significant predictor of variance in aggression was  
8 total PLE ( $\beta=.562$ ,  $t=6.709$ ,  $p<0.0001$ ), total attachment scores were not significant predictors  
9 ( $\beta=.052$ ,  $t=.724$ ,  $p>0.05$ ).  
10  
11  
12  
13  
14  
15

16 However, using the Aiken and West (1991) moderation approach it can be seen that those in  
17 higher attachment score groups (i.e. more insecure attachments) demonstrated a stronger  
18 relationship between PLE and aggression. Bivariate correlation between Total PLE and Total  
19 Aggression was  $r=0.62$  for individuals with high attachment scores;  $r=0.45$  for individuals with  
20 moderate attachment scores, and  $r=0.43$  for those with low attachment scores. See figure 1. for a  
21 graph of linear regression lines for the relationship between total PLEs and total aggression in the  
22 three attachment groups.  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 Figure 1. About here  
33  
34  
35  
36

### 37 Discussion

38  
39  
40 This study found that, as predicted, there were significant relationships between measures of  
41 attachment, PLEs and aggression.  
42  
43

44 These findings support a link between Attachment and PLE, highlighting that attachment is an  
45 important factor in understanding psychotic experiences (Penn et al., 1997; Mickelson, Kessler and  
46 Shaver, 1997). Previous research using community samples has suggested that positive symptoms  
47 were closely related to anxious attachment and anhedonia related to avoidant attachment (Berry et  
48 al., 2006). This study found that positive symptoms were more strongly correlated with avoidant  
49 attachment, and depressive symptoms most closely related to anxious. This discrepancy may be due  
50 to differing measurement tools or samples, however further research is necessary.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 An avoidant attachment style suggests a negative internal model of others, in the current  
4 study this was associated with increased positive PLEs. The traumagenic neurodevelopmental model  
5 of psychosis (Read et al., 2014) highlights a heightened physiological responsivity to stressors (as a  
6 consequence of childhood adversity) as a potential factor in the aetiology of psychosis. It is plausible  
7 that a negative internal working model of others is associated with an increased perception of others  
8 as unsafe or threatening, resulting in an increase in the threat response associated with PLEs  
9 (including paranoia) in the general population. This finding would warrant replication and further  
10 exploration.  
11  
12  
13  
14  
15  
16  
17

18 These analyses suggested that participants reporting greater anxiety or avoidance in  
19 attachments also reported increased aggression, in keeping with previous findings (Ratip, 2013;  
20 Bartholemew and Horowitz, 1991; Kobak and Sreery, 1988; Mallinckrodt, 2000). When considering  
21 subscales of aggression, this was true for physical aggression, anger and hostility, but not for verbal  
22 aggression. This is plausibly due to verbal aggression being influenced by other factors, but would  
23 warrant further exploration. Hostility showed the strongest relationship with attachment styles, and  
24 was most closely related to anxious attachment. Physical aggression was only significantly related to  
25 avoidant attachment.  
26  
27  
28  
29  
30  
31  
32

33 Buss and Perry (1992) consider anger expression to fall into either cognitive, affective or  
34 instrumental (behavioural) domains. Verbal and physical aggression are both behavioural aspects of  
35 aggression, therefore the current results suggest that the behavioural manifestation of aggression has  
36 weaker relationships with attachment (and PLEs) than cognitive (hostility) or affective (anger)  
37 aggression. Future research is necessary to clarify this relationship. Total attachment scores were a  
38 stronger predictor of aggression than either anxious or avoidant attachment problems alone.  
39  
40  
41  
42  
43  
44

45 PLE scores had a stronger relationship with aggression than attachment scores. Similarly to  
46 attachment, all PLE sub-scales showed the strongest relationship with hostility and weakest with  
47 physical and particularly verbal aggression, although this relationship remained significant. Each  
48 aggression domain was linked most significantly with different PLE symptoms; physical aggression  
49 with positive symptoms, verbal aggression with negative symptoms and anger and hostility with  
50 depressive symptoms.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3           These results support evidence of a link between specific symptoms of psychosis and  
4 aggression (Douglas, Guy and Hart, 2009; Jones et al., 2010) particularly in community samples,  
5 (Nederlof, Muris and Hovens, 2012; Mojtabai, 2006; Kinoshita et al., 2011), although direct  
6 comparison is limited due to differing methodology. Findings suggest that particular symptoms of  
7 psychosis may relate to different facets of aggression. The finding that positive PLEs were most  
8 closely related to physical aggression is in keeping with previous evidence that hallucinations are  
9 associated with aggression in community samples (Nederlof, Muris and Hovens, 2012). However,  
10 while the symptom-violence link may be present in community samples as well as clinical samples,  
11 there remains a need to understand the mechanisms of this relationship.  
12  
13  
14  
15  
16  
17  
18

19           Total PLE and attachment scores predicted variance in aggression, although PLE was the  
20 only significant predictor in the model. Attachment scores did strengthen the models of verbal  
21 aggression and hostility, suggesting that attachment plays a more influential role in these dimensions  
22 of aggression.  
23  
24  
25  
26

27           The five factor model of aggression (Attachment domains and PLE domains) was statistically  
28 significant. These factors accounted for over half of variance in hostility scores. Future research may  
29 benefit from considering whether hostile aggression is seen more frequently in individuals with  
30 psychosis and attachment problems. Formulation and management plans relating to hostility may  
31 benefit from considering both symptom and attachment variables.  
32  
33  
34  
35  
36

37           Although including the interaction term did not improve models of aggression, the relationship  
38 between PLE and aggression was stronger in individuals with more insecure attachment styles. This  
39 suggests that PLEs were more closely associated with aggression in individuals reporting the highest  
40 levels of attachment insecurity. This finding requires replication, but suggests that attachment may be  
41 a relevant factor in the assessment and management of individuals experiencing PLE.  
42  
43  
44  
45  
46

47           A more insecure attachment style is associated with childhood adversity and early trauma,  
48 and the current findings support the importance of considering early trauma in the assessment and  
49 formulation of risk in individuals experiencing symptoms of psychosis. The findings suggest that there  
50 was a closer relationship between PLEs and aggression in individuals with greater anxiety and  
51 avoidance in attachment. It is plausible that individuals with a more negative internal working model of  
52 self and others are more prone to experiencing PLEs associated with threat (e.g. paranoia) or a  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 derogatory view of the self. While this hypothesis remains tentative, there is some evidence that  
4 trauma affects the nature of PLEs in the community (Freeman and Fowler, 2009), and that negative  
5 emotions in hallucinatory experiences and persecutory delusions are associated with risk in  
6 individuals with schizophrenia (Cheung et al., 1997). The current research suggests that attachment  
7 may be a useful framework for understanding the relationship between early adversity, symptoms of  
8 psychosis and risk of aggression towards others.  
9  
10  
11  
12

13  
14 The current findings also support the hypothesis that these relationships are present in the  
15 general population as well in samples of individuals with schizophrenia (or other psychotic disorders).  
16 Longitudinal research suggests that clinical psychosis may be a rare outcome of progressively more  
17 persistent PLEs (Dominguez et al., 2009). The current study suggests that individuals with more  
18 insecure attachment styles were more prone to PLEs, which were more closely associated with  
19 aggression than in individuals with more secure attachment styles. This would tentatively suggest that  
20 attachment may be a useful area of formulation and intervention for at-risk populations and early  
21 intervention services, although further research is necessary to establish this. Investigating the  
22 potential protective role of systemic interventions which incorporate attachment perspectives in the  
23 treatment of children with complex trauma (e.g. Kinniburgh et al., 2017) may be a useful area of future  
24 research into the prevention of aggression associated with symptoms of psychosis.  
25  
26  
27  
28  
29  
30  
31  
32  
33

### 34 35 Limitations

36  
37 This generalisability of these results is limited by the sampling methodology, there is an  
38 inherent bias in a convenience sample. All participants self-selected for the study, and there is some  
39 evidence that those who self-select are more likely to be experiencing psychological difficulties  
40 (Freeman et al. 2005). The study was also biased towards female respondents, given that there is  
41 evidence that male and female psychiatric inpatients vary in their expression of aggression (e.g  
42 Picken et al., 2010) this again limits the generalisability of the results, particularly to clinical samples.  
43 Similarly, there is a reliance on self-reporting, leaving the results subject to responder bias and social  
44 desirability. Given the self-report nature of the psychometric tools, it is plausible that the results are  
45 affected by self-appraisal, for example individuals with less secure attachment styles may perceive  
46 themselves as more aggressive. Future replications may benefit from having behavioural correlates or  
47 informant information to support self-report measures.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



This study employed a cross-sectional design, and therefore conclusions about inference and causality are theoretical. The significant results would support larger scale, possibly longitudinal, replications.

Relatively few participants reported recent substance misuse in the current study, future replications may benefit from larger samples to more fully investigate the potential effect of substance use on the identified relationships.

### Conclusions

This research identifies relationships between attachment, PLE and aggression in individuals in the community. The results suggest that attachment is a contributing factor to aggression associated with PLE and that this relationship exists in community samples. It tentatively supports the hypothesis that addressing risk in individuals with PLE should go beyond symptom-focussed work, and consider associated attachment problems. The findings suggest that risk assessments and formulations should consider specific symptoms and problems with attachment as contributing to aggression, in particular hostility (or cognitive aggression). These results highlight the need to consider attachment problems as part of the aetiology of PLE and associated behaviours. There is substantial evidence that insecure attachment leads to poor engagement and poorer outcomes for individuals with psychosis (Gumley et al., 2014), these findings suggest that attachment is central to the formulation and treatment of individuals experiencing PLE.

### **References**

- Aiken, L. and West, S. (1991), *Multiple regression: Testing and interpreting interactions*, Sage, Newbury Park, London.
- Ainsworth, M.D.S., Blehar, M.C., Waters, E., and Wall, S. (1978), *Patterns of attachment: A psychological study of the strange situation*, Erlbaum, Hillsdale, N J.
- Allen, T.J., Moeller, F.G., Rhoades, H.M. and Cherek, D.R. (1998), "Impulsivity and history of drug dependence", *Drug & Alcohol Dependence*, Vol. 50 No. 2, pp.137-145.
- Bartholomew, K. and Horowitz, L.M. (1991), "Attachment styles among young adults: a test of a four-category model", *Journal of personality and social psychology*, Vol. 61 No.2, pp.226-244.



1  
2  
3 Berry, K., Wearden, A., Barrowclough, C. and Liversidge, T. (2006), "Attachment styles, interpersonal  
4 relationships and psychotic phenomena in a non-clinical student sample", *Personality and Individual*  
5 *Differences*, Vol. 41 No. 4, pp.707-718.

6  
7  
8  
9 Bo, S., Abu-Akel, A., Bertelsen, P., Kongerslev, M. and Haahr, U.H. (2013), "Attachment, mentalizing  
10 and personality pathology severity in premeditated and impulsive aggression in schizophrenia",  
11 *International Journal of Forensic Mental Health*, Vol. 12 No.2, pp.126-138.

12  
13  
14  
15 Bowlby, J. (1973), *Attachment and loss: Volume II: Separation, anxiety and anger*, Basic Books, New  
16 York.

17  
18  
19 Bowlby, J. (1982), "Attachment and loss: Retrospect and prospect", *American journal of*  
20 *Orthopsychiatry*, Vol. 52 No. 4, pp.664-678.

21  
22  
23  
24 Buss, H. and Perry, M. (1992), "The aggression questionnaire", *Journal of personality and social*  
25 *psychology*, Vol. 63 No. 3, pp. 452-459.

26  
27  
28  
29 Cheung, P., Schweitzer, I., Crowley, K. and Tuckwell, V. (1997), "Violence in schizophrenia: role of  
30 hallucinations and delusions", *Schizophrenia research*, Vol. 26 No. 2-3, pp.181-190.

31  
32  
33  
34  
35  
36  
37  
38 Crowell, J.A., Fraley, R.C. and Shaver, P.R. (1999), "Measurement of individual differences in  
39 adolescent and adult attachment", in Cassidy, J. and Shaver, P. (Eds.), *Handbook of Attachment:*  
40 *Theory, research, and clinical applications*, Guildford Press, New York, pp. 434-465.

41  
42  
43  
44 Dominguez, M.D.G., Wichers, M., Lieb, R., Wittchen, H.U. and van Os, J. (2009), "Evidence that  
45 onset of clinical psychosis is an outcome of progressively more persistent subclinical psychotic  
46 experiences: an 8-year cohort study", *Schizophrenia bulletin*, Vol. 37 No. 1, pp.84-93.

47  
48  
49  
50  
51  
52  
53 Douglas, K.S., Guy, L.S. and Hart, S.D. (2009), "Psychosis as a risk factor for violence to others: a  
54 meta-analysis", *Psychological bulletin*, Vol. 135 No. 5, pp. 679-706.

55  
56  
57  
58  
59  
60 Esterberg, M.L. and Compton, M.T. (2009), "The psychosis continuum and categorical versus  
61 dimensional diagnostic approaches", *Current psychiatry reports*, Vol. 11 No. 3, pp.179-184.

62  
63  
64  
65  
66  
67  
68  
69  
70 Eysenck, S.B. and Eysenck, H.J. (1970), "Crime and personality: an empirical study of the three-  
71 factor theory", *The British Journal of Criminology*, Vol. 10 No. 3, pp.225-239.

1  
2  
3 Faul, F., Erdfelder, E., Buchner, A. and Lang, A.G. (2009), "Statistical power analyses using G\* Power  
4 3.1: Tests for correlation and regression analyses", *Behavior research methods*, Vol. 41 No. 4,  
5 pp.1149-1160.  
6

7  
8  
9 Fazel, S., Långström, N., Hjern, A., Grann, M. and Lichtenstein, P. (2009), "Schizophrenia, substance  
10 abuse, and violent crime", *JAMA*, Vol. 301 No. 19, pp.2016-2023.  
11

12  
13 Freeman, D. and Fowler, D. (2009), "Routes to psychotic symptoms: trauma, anxiety and psychosis-  
14 like experiences", *Psychiatry research*, Vol. 169 No. 2, pp.107-112.  
15

16  
17 Freeman, D., Garety, P.A., Bebbington, P.E., Smith, B., Rollinson, R., Fowler, D., Kuipers, E., Ray, K.  
18 and Dunn, G. (2005), "Psychological investigation of the structure of paranoia in a non-clinical  
19 population", *The British Journal of Psychiatry*, Vol. 186 No. 5, pp.427-435.  
20  
21

22  
23 Fonseca-Pedrero, E., Paino, M. D. L. M., Lemos-Giraldez, S. and Muiz, J. (2012), "Validation of the  
24 Community Assessment Psychic Experiences-42 (CAPE-42) in Spanish college students and patients  
25 with psychosis", *Actas espanolas de psiquiatria*, Vol. 40 No. 4, pp. 169-176.  
26  
27

28  
29 Garety, P.A., Kuipers, E., Fowler, D., Freeman, D. and Bebbington, P.E. (2001), "A cognitive model of  
30 the positive symptoms of psychosis", *Psychological medicine*, Vol. 31 No. 2, pp.189-195.  
31  
32

33  
34 Gerevich, J., Bácskai, E. and Czobor, P. (2007), "The generalizability of the Buss–Perry Aggression  
35 Questionnaire", *International Journal of Methods in Psychiatric Research*, Vol. 16 No.3, pp.124-136.  
36  
37

38  
39 Granville-Grossman, K.L. (1966), "Early bereavement and schizophrenia", *The British Journal of*  
40 *Psychiatry*, Vol. 112 No. 491, pp.1027-1034.  
41  
42

43  
44 Green, K., Browne, K. and Chou, S. (2017), "The relationship between childhood maltreatment and  
45 violence to others in individuals with psychosis: a systematic review and meta-analysis", *Trauma,*  
46 *Violence, & Abuse*, doi: 1524838017708786.  
47  
48

49  
50 Gumley, A.I., Taylor, H.E.F., Schwannauer, M. and MacBeth, A. (2014), "A systematic review of  
51 attachment and psychosis: measurement, construct validity and outcomes", *Acta Psychiatrica*  
52 *Scandinavica*, Vol. 129 No. 4, pp.257-274.  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Harris, J.A. (1995), "Confirmatory factor analysis of the Aggression Questionnaire", *Behaviour*  
4 *research and therapy*, No. 33 Vol. 8, pp.991-993.  
5  
6  
7 Jones, R.M., Van den Bree, M., Ferriter, M. and Taylor, P.J. (2010), "Childhood risk factors for  
8 offending before first psychiatric admission for people with schizophrenia: a case-control study of  
9 high security hospital admissions", *Behavioral sciences & the law*, Vol. 28 No. 3, pp.351-365.  
10  
11  
12  
13 Kinniburgh, K.J., Blaustein, M., Spinazzola, J. and Van der Kolk, B.A. (2017), "Attachment, Self-  
14 Regulation, and Competency: A comprehensive intervention framework for children with complex  
15 trauma", *Psychiatric annals*, Vol. 35 No. 5, pp.424-430.  
16  
17  
18  
19 Kinoshita, Y., Shimodera, S., Nishida, A., Kinoshita, K., Watanabe, N., Oshima, N., Akechi, T.,  
20 Sasaki, T., Inoue, S., Furukawa, T.A. and Okazaki, Y. (2011), "Psychotic-like experiences are  
21 associated with violent behavior in adolescents", *Schizophrenia Research*, Vol. 126 No. 1, pp.245-  
22 251.  
23  
24  
25  
26  
27 Kobak, R.R. and Sceery, A. (1988), "Attachment in late adolescence: Working models, affect  
28 regulation, and representations of self and others", *Child development*, Vol. 59 No. 1, pp.135-146.  
29  
30  
31  
32 Large, M.M. and Nielssen, O. (2011), "Violence in first-episode psychosis: a systematic review and  
33 meta-analysis", *Schizophrenia research*, Vol. 125 No. 2-3, pp.209-220.  
34  
35  
36  
37 Loch, A.A., Wang, Y.P., Rössler, W., Tófoli, L.F., Silveira, C.M. and Andrade, L.H. (2011), "The  
38 psychosis continuum in the general population: findings from the Sao Paulo Epidemiologic Catchment  
39 Area Study", *European archives of psychiatry and clinical neuroscience*, Vol. 261 No. 7, pp.519-527.  
40  
41  
42  
43 Matheson, S.L., Shepherd, A.M., Pinchbeck, R.M., Laurens, K.R. and Carr, V.J. (2013), "Childhood  
44 adversity in schizophrenia: a systematic meta-analysis", *Psychological medicine*, Vol. 43 No. 2,  
45 pp.225-238.  
46  
47  
48  
49 Mallinckrodt, B. (2000), "Attachment, social competencies, social support, and interpersonal process  
50 in psychotherapy", *Psychotherapy research*, Vol. 10 No. 3, pp.239-266.  
51  
52  
53  
54 Mickelson, K.D., Kessler, R.C. and Shaver, P.R. (1997), "Adult attachment in a nationally  
55 representative sample", *Journal of personality and social psychology*, Vol. 73 No. 5, pp. 1092-1106.  
56  
57  
58  
59  
60

- 1  
2  
3 Mikulincer, M. and Shaver, P.R. (2007), "Boosting attachment security to promote mental health,  
4 prosocial values, and inter-group tolerance", *Psychological Inquiry*, Vol. 18 No. 3, pp.139-156.  
5  
6  
7 Mojtabai, R. (2006), "Psychotic-like experiences and interpersonal violence in the general  
8 population", *Social Psychiatry and Psychiatric Epidemiology*, Vol. 41 No. 3, pp.183-190.  
9  
10  
11 Morgan, C., Kirkbride, J., Leff, J., Craig, T., Hutchinson, G., McKenzie, K., Morgan, K., Dazzan, P.,  
12 Doody, G.A., Jones, P. and Murray, R. (2007), "Parental separation, loss and psychosis in different  
13 ethnic groups: a case-control study", *Psychological medicine*, Vol. 37 No. 4, pp.495-503.  
14  
15  
16  
17 Mossaheb, N., Becker, J., Schaefer, M.R., Klier, C.M., Schloegelhofer, M., Papageorgiou, K. and  
18 Amming, G.P. (2012), "The Community Assessment of Psychic Experience (CAPE) questionnaire  
19 as a screening-instrument in the detection of individuals at ultra-high risk for  
20 psychosis", *Schizophrenia research*, Vol. 141 No. 2, pp.210-214.  
21  
22  
23  
24  
25  
26 Nederlof, A.F., Muris, P. and Hovens, J.E. (2012), "Psychotic-like experiences and aggressive  
27 behavior in a non-clinical sample", *Personality and Individual Differences*, Vol. 53 No. 1, pp.33-37.  
28  
29  
30  
31 Parker, G.F. (2014), "DSM-5 and psychotic and mood disorders", *Journal of the American Academy  
32 of Psychiatry and the Law Online*, Vol. 42 No.2, pp.182-190.  
33  
34  
35  
36 Penn, D.L., Corrigan, P.W., Bentall, R.P., Racenstein, J. and Newman, L. (1997), "Social cognition in  
37 schizophrenia", *Psychological bulletin*, Vol. 121 No. 1, pp. 114-132.  
38  
39  
40  
41 Picken, A.L., Berry, K., TARRIER, N. and Barrowclough, C. (2010), "Traumatic events, posttraumatic  
42 stress disorder, attachment style, and working alliance in a sample of people with psychosis", *The  
43 Journal of nervous and mental disease*, Vol. 198 No. 10, pp.775-778.  
44  
45  
46  
47 Posner, J., Russell, J.A. and Peterson, B.S. (2005), "The circumplex model of affect: An integrative  
48 approach to affective neuroscience, cognitive development, and psychopathology", *Development and  
49 psychopathology*, Vol. 17 No. 3, pp.715-734.  
50  
51  
52  
53 Ratip, D.E. (2013), "The role of attachment in violent offending and violence in institutions", *Doctoral  
54 dissertation, University of Birmingham*. Available at:  
55 <http://etheses.bham.ac.uk/4157/1/Ratip13ForenPsyD.pdf> (accessed 19 July 2018).  
56  
57  
58  
59  
60

1  
2  
3 Read, J., Fosse, R., Moskowitz, A. and Perry, B. (2014), "The traumagenic neurodevelopmental  
4 model of psychosis revisited", *Neuropsychiatry*, Vol. 4 No. 1, pp.65-79.

6  
7 Read, J., Os, J.V., Morrison, A.P. and Ross, C.A. (2005), "Childhood trauma, psychosis and  
8 schizophrenia: a literature review with theoretical and clinical implications", *Acta Psychiatrica*  
9  
10 *Scandinavica*, Vol. 112 No. 5, pp.330-350.

12  
13 Short, T., Thomas, S., Mullen, P. and Ogloff, J.R. (2013), "Comparing violence in schizophrenia  
14 patients with and without comorbid substance-use disorders to community controls", *Acta Psychiatrica*  
15  
16 *Scandinavica*, Vol. 128 No. 4, pp.306-313.

18  
19 Soyka, M., (2000), "Substance misuse, psychiatric disorder and violent and disturbed behaviour", *The*  
20  
21 *British Journal of Psychiatry*, Vol. 176 No. 4, pp.345-350.

23  
24 Stefanis, N.C., Hanssen, M., Smiris, N.K., Avramopoulos, D.A., Evdokimidis, I.K., Stefanis, C.N.,  
25  
26 Verdoux, H. and Van Os, J. (2002), "Evidence that three dimensions of psychosis have a distribution  
27 in the general population", *Psychological medicine*, Vol. 32 No. 2, pp.347-358.

29  
30 Van Os, J., Hanssen, M., Bijl, R.V. and Ravelli, A. (2000), "Strauss (1969) revisited: a psychosis  
31  
32 continuum in the general population?", *Schizophrenia research*, Vol. 45 No. 1-2, pp.11-20.

34  
35 Verdoux, H. and van Os, J. (2002), "Psychotic symptoms in non-clinical populations and the  
36  
37 continuum of psychosis", *Schizophrenia research*, Vol. 54 No. 1, pp.59-65.

38  
39 Warner, R. and Atkinson, M. (1988), "The relationship between schizophrenic patients' perceptions of  
40  
41 their parents and the course of their illness", *The British Journal of Psychiatry*, Vol. 153 No. 3, pp.344-  
42  
43 353.

Table 1. Five factor Regression models with Physical Aggression, Verbal Aggression, Anger and Hostility Scores as the Criterion variables (N=213)

Criterion Variable	F Value	Sig.	R <sup>2</sup> Variance
Physical Aggression	F(5,209)= 9.25	.000	18.4%
Verbal Aggression	F(5,210)= 2.94	.014	6.7%
Anger	F(5,210)= 16.40	.000	28.5%
Hostility	F(5,210)= 44.71	.000	52.0%




Figure 1. linear regression lines for the relationship between total PLEs and total aggression by attachment group