



Age, diet, growth and reproduction of a Population of *Cobitis paludica* (de Buen, 1930) in the Palancar Stream (southwest of Europe, Spain) (Pisces: Cobitidae)

M. C. Soriguer, C. Vallespín, C. Gomez-Cama & J. A. Hernando

Departamento de Biología Animal, Biología Vegetal y Ecología, Facultad de Ciencias del Mar, Universidad de Cádiz, Campus Río San Pedro, Puerto Real, Cádiz, Spain
E-mail: mila.soriguer@uca.es

Received 2 June 1998; in revised form 4 May 2000; accepted 13 June 2000

Key words: Cobitidae, length–weight, fecundity, feeding habits

Abstract

A total sample of 99 specimens of the vulnerable fish species *Cobitis paludica* (de Buen, 1930), was collected on a monthly basis from a tributary of the River Guadalquivir between September 1990 and September 1991. Female longevity was 1 year longer than that of males, the former reaching age-class 5+. The diet of *C. paludica* consisted mainly of Ostracoda and chironomid larvae, benthic prey of small size and occasionally other resources, such as Macrobiotid eggs spawned during winter. Detritus in gastrointestinal content was scarce, which differs from other populations. Back-calculations of length were performed from scale annuli measurements. The relationship between standard length and scale radius was different for males and females. Females were larger than males of the same age. Both back-calculated length and theoretical length, calculated by applying von Bertalanffy's model, were larger for females. The relationship between female standard length (SL) and fecundity (F) was $\log F = -7.465 + 5.5753 \log SL$.

Introduction

Cobitis paludica (de Buen, 1930) is an endemic Iberian species, catalogued as vulnerable (IUCN Conservation Categories, Blanco & Gonzalez, 1992). Knowledge of its biology is restricted (Lobón-Cerviá & Zabala, 1984; Soriguer 1996), in particular its feeding habits and diet (Hernando, 1978). However, there is information on the diet of the related European *C. taenia* (Maksunov, 1969; Spataru, 1970; Robotham, 1977). *C. paludica* is a bottom-dweller found in slow-flowing streams. The secretive nature of this fish is probably the main reason for this paucity of information.

In the Palancar Stream, *C. paludica* coexist with the cyprinids *Barbus sclateri* and *Leuciscus pyrenaicus*, which are characterised by their adaptability. A diet analysis is the starting point for obtaining knowledge of their trophic requirements and their possible interactions with other species (competition, predation, etc.).

This study of *C. paludica* was aimed primarily at determining the biology, diet variation with season and age, as well as growth rate, fecundity and insight into their reproductive cycle, in a small stream subject to an extreme low-water period.

Materials and methods

Study area

The study area was a 200 m section of the Palancar Stream, a small, second-order tributary on the left bank of the central watercourse of the Guadalquivir River. It is located in the Natural Park of 'Sierras Subbéticas', lies within co-ordinates 37° 27' 46" N; 4° 16' 40" – 37° 27' 59" N; 4° 15' 40" W.

Field and laboratory methods

Monthly samples, from September 1990 to September 1991, were by electro-fishing, applying the same

effort on each occasion. Due to the vulnerable status of this species, collections were kept to the minimum. Captured samples were frozen, and for each specimen, the following variables were determined: total length (TL), standard length (SL), total weight (TW), somatic weight (SW: corresponding to fresh weight after removal of the gastrointestinal tract and gonads), and gastrointestinal contents weight (GCW: determined as the difference between full and empty gastrointestinal weights).

The specimens were sexed on capture into three categories: male, female and juvenile. The wet weight of gonads (testis or ovary weight) was determined to the nearest 0.0001 g. Ovaries were preserved whole in Gilson's fluid after weighting (Bagenal & Braun, 1978). Scales below the first radius of the dorsal fin and above the lateral line were removed from each sample. These were cleaned using 7% NaOH, and placed on two slides, for age determination. The number of oocytes present in each gonad were counted using a system of subsampling by area. Their diameters were measured in mm using a profile projector. Oocytes were then separated into vitelline and transparent, and the mean diameter was calculated, together with the proportion of the different sizes present (Bagenal & Braun, 1978; Crim & Glebe, 1990). Absolute fecundity was estimated as the number of mature eggs present in the ovary immediately before spawning.

Gastrointestinal contents were viewed whole with a WILDTM M10 binocular microscope. The different taxa found were identified and the number of prey for each taxon was calculated. To carry out the analysis, the specimens were grouped by age and month of sampling. Assessment of the diet was based on frequency of occurrence of the different diet components (Guziur, 1976). The proportion of each type of prey was determined and expressed as π = number of prey in each category (NP)/total number of prey (TNP). The ratios of standard length–total weight (SL–TW) and standard length–somatic weight (SL–SW) have been calculated for male, female and juvenile groups separately, using the expression:

$$W = a SL^b.$$

The study of annual growth was based on back-calculations of length (Lea, 1910; Lagler, 1956; Bagenal & Tesch, 1978; Carlander, 1981). For this purpose, using the scales removed from each individual fish, the distance from the scale centre to each of the rings **n** (**Rn**) and to the edge (**R**) was meas-

ured to the nearest 0.001 mm using a profile projector. Growth in length was described using von Bertalanffy's equation of theoretical growth in length (Von Bertalanffy, 1938, 1957):

$$L_t = L_\infty * (1 - e^{-k*(t-t_0)}).$$

L_∞ and k were estimated from Ford's equation (1993) and Walford (1946), as well as

$$t_0 = t + (1/k) * \ln[(L_\infty - L_t)/L_\infty].$$

To make a comparison of growth we used the parameter

$$\omega = k * L_\infty$$

(Gallucci & Quinn, 1979), and the index of overall fish growth performance (Munro & Pauli, 1983)

$$\phi' = \log k + 2 \log L_\infty.$$

We separately calculated the instantaneous rates of growth in length L_i from the mean back-calculated lengths, and the instantaneous rates of growth in weight G_i from mean weights computed from the relationship TL–TW for males and females, using the expressions:

$$L_i = \ln L_{t+1} - \ln L_t,$$

$$G_i = \ln W_{t+1} - \ln W_t,$$

where L_{t+1} and W_{t+1} are mean length and mean weight, respectively, of the age-class $t+1$, and L_t and W_t , mean length and weight of the previous age-class.

Results

The relationship between total and standard length (TL–SL) was: $TL = 0.113531 + 1.14441 SL$ ($r = 0.99$; $p < 0.000$).

Females reached age 5+ and attained 90 mm SL. The oldest males captured belonged to age-class 4+ and ranged from 55 to 85 mm SL. Females were not mature until they reached 65 mm SL.

The overall male–female ratio was 1:2, but this proportion varied with age-class (Figure 1). At age 1+, mature specimens were 23.8% of the total catch; of these, 80% were males and 20% females. At age 2+, mature specimens increased to 65.2%, of which 40% were males and 60% females. At age 3+, 100% of

Table 1. Coefficients, sample number and determining coefficient R^2 , in SL-TW and SL-SW ratios, of *Cobitis paludica* from the Palancar Stream, Spain

	SL-TW			SL-EW			
	N	$\log a$	B	R^2	$\log a$	b	R^2
Juveniles	36	-6.4763	3.8493	0.9779	-6.2155	3.6288	0.9622
Males	17	-4.619	2.7906	0.8004	-4.9824	2.947	0.8354
Females	44	-5.3484	3.1644	0.9586	-5.4405	3.1664	0.9584
Total population	97	-6.0438	3.5415	0.9793	-5.734	3.3301	0.9691




Figure 1. Sex ratio at each age-class of *C. paludica* in the Palancar Stream, Spain (I: Juvenile, M: Males, F: Females).




Figure 2. Percentage occurrence of each item in the diet of *C. paludica* from the Palancar Stream, Spain.

the captured specimens were mature, 38% males and 62% females. Males comprised only 8% of age-class 4+, and the remaining 92% were females. At age 5+, 100% are females. Thus, females became numerically more predominant with increasing age.

In respect of occurrence rates in diet, 14 taxa of invertebrates were found in the 99 gastrointestinal tracts analysed, as well as a constant low rate of diatoms and detritus. The main foods, expressed as percentage occurrence, were Diptera, chironomid and ceratopogonid larvae, Ostracoda and Cladocera, macrobiotid spawns and Hydracarina (Figure 2). A comparison of the percentage occurrence of different resources in the diet indicates that there were no significant differences (U of Mann-Whitney $p > 0.05$) between months. A seasonal analysis of the numerical percentage of each type of prey ($\pi * 100$) (Figure 3) showed the amounts of Chironomidae and Ostracoda were significant, the latter predominating in autumn, while macrobiotid spawns were more frequent in winter. Figure 4 shows

the percentage occurrence at different ages: Ceratopogonidae diminished with age, whereas Chironomidae and Ostracoda remained similar for all ages, except 2+; Cladocera, Copepod and Macrobiotidae became more abundant with increasing age. We frequently found chironomid nymphs in the stomachs of specimens aged 5+. Occurrence of diatoms and detritus was isolated. Vegetation remains were scarce. The U of Mann-Whitney's test showed that there were significant differences ($p < 0.05$) in the diet of age-classes 0+, 1+, 2+ and 5+, whereas diet is homogeneous for age classes 3+ and 4+. In terms of the numerical proportions of each type of prey (Figure 4), those of age 0+ consumed mainly Copepods and Ceratopogonidae; those of age 1+ consumed mainly Cladocera and Ostracoda, with some chironomid larvae; those of age 2+ consumed less Cladocera and more macrobiotid spawns; those of ages 3+ and 4+ showed very similar diets, consisting mainly of Ostracoda and macrobiotid spawns; age-class 5+ was not well represented; only

Table 2. Relationship between SL and scale radius *R*, for males and females, of *Cobitis paludica* from Palancar Stream, Spain

Parameter	Estimate	Standard error	T Statistic	p-value	
Males					
Intercept	4.14678	0.027185	190.933	0.0000	
Slope	0.297356	0.0355547	8.36842	0.0000	
Analysis of Variance					
Source	Sum of Squares	Df	Mean Squared	F-ratio	p-value
Model	0.117656	1	0.117656	70.03	0.0000
Residual	0.0268812	16	0.0016807		
Total (Corr.)	0.144537	17			
Correlation Coefficient= 0.90223					
R-squared= 81.4019 percent					
Females					
Intercept	4.34353	0.0172471	251.841	0.0000	
Slope	0.485973	0.0266336	18.2467	0.0000	
Analysis of Variance					
Source	Sum of Squares	Df	Mean Squared	F-ratio	p-value
Model	1.28301	1	1.28301	332.94	0.0000
Residual	0.146436	38	0.00385357		
Total (Corr.)	1.42944	39			
Correlation Coefficient= 0.947395					
R-squared= 89.7558 percent					

Table 3. Comparison of regression lines of *Cobitis paludica* from Palancar Stream, Spain. Dependent variable: lnSL. Independent variable: ln *R*. Level codes: sex. Number of complete cases: 58 and number of regression lines: 2

Parameter	Estimate	Standard Error	T Statistic	p-value
Constant	4.34535	0.0157401	257.952	0.0000
ln <i>R</i>	0.485973	0.0243064	19.9936	0.0000
sex= males	-0.196749	0.0338949	-5.80468	0.0000
ln <i>R</i> *sex=males	-0.188437	0.054825	-3.43706	0.0011

two specimens were caught, and analysis showed a fairly even mix of prey found in the other age classes.

Table 1 shows the growth rate variables in SL-TW and SL-SW ratios for the total population of *C. paludica*, and separately for juveniles, males and females. The **b** and **b'** values in SL-TW and SL-SW ratios, respectively, were larger for juveniles than for mature samples, and larger for females than males.




Figure 3. Seasonal variation in the numerical proportion of each type of prey in the diet of *C. paludica* from the Palancar Stream, Spain.




Figure 4. Variation with age of the percentage occurrence and numerical proportion of each item in the diet of *C. paludica* from the Palancar Stream, Spain.

The relationship between standard length and scale radius in males and females was fitted to the equations:

$$\ln SL = 4.14678 + 0.297536 * RT \text{ for males,}$$

and

$$\ln SL = 4.34353 + 0.485973 * RT \text{ for females}$$

The relationship between these variables, in both sexes, is statistically significant at the 99% confidence level (Table 2). There were significant differences in this relationship between the two sexes (Table 3), and the expressions for back-calculating the lengths are:

$$\text{For females: } Ln = Ls * (Rn/R)^{0.48597}$$

$$\text{For males: } Ln = Ls * (Rn/R)^{0.29754}$$

Table 4 show the mean standard lengths obtained for each group and age, as well as length increments between successive age-classes.




Figure 5. Monthly distribution of oocyte diameters of *C. paludica* from the Palancar Stream, Spain.

Back-calculated lengths from male and female scale measurements were similar only for ages 1+ and 2+ (ANOVA $p > 0.05$). However, for ages 3+ and 4+ they were significantly different (ANOVA $p < 0.05$). When the possible differences of size are considered as a function of sex and age (by MULTIFACTOR ANOVA), in age classes 1+ and 3+, significant differences of size in function both of age and sex were seen;

in age class 4+, differences were only significant in respect of sex, while in age class 2+ there were no significant differences. The length increment at intervals I, II, III and IV was greater in females than in males.

Constants in Von Bertalanffy's (1938) equation of theoretical growth in length, k , L_{∞} , and t_0 , as well as variables ω , ϕ' for each sex, are shown in Table 6. Lobón-Cerviá (1982) finds that computed infinite lengths are smaller than sizes attained by the population. In our case, this is not so. None of the captured specimens reach the calculated L_{∞} .

The equations of theoretical growth in length for this species in the Palancar Stream are as follows:

$$\text{FEMALES: } L_t = 110.094 * (1 - e^{-0.2088665(t+0.896726)})$$

$$\text{MALES: } L_t = 66.83 * (1 - e^{-0.439856(t+0.867196)})$$

Theoretical lengths of male and female (Table 6) show that, from the third year of life (2+), females are larger than males.

The instantaneous growth rates for length (L_1) and weight (G_1), of males and females separately, shown in Table 7, indicate a decrease in both growth rates with age. Also, these values are much smaller for males, especially younger males, but tend to become more similar with age.

A monthly GSI analysis could not be carried out because no females were captured in May and June. Monthly analysis of the frequency distribution of oocyte diameters (Figure 5) showed a mode in November, corresponding to translucent oocytes of approximately 0.1 mm in diameter. In winter, oocytes increased in diameter, with the proportion of vitelline eggs increasing progressively. In April, oocytes attained a suitable size for spawning and these became predominant in July, with no translucent oocytes occurring. In September, translucent oocytes accounted for the dominant mode again. There were also small numbers of intra-ovarian oocytes of size suitable for spawning, which were re-absorbed.

Absolute fecundity of females captured in April was calculated from the expression: $\log F = -7.465 + 5.5753 \log SL$; $r = 0.9998$ ($p < 0.01$).

Discussion

Given the vulnerable status of the species, the number of specimens captured is less than that desirable for a definitive study. The ratio of sexes in the population

of *C. paludica* in the Palancar Stream is very different from that found in this species in the Jarama River, where it is close to parity (Lobón-Cerviá & Zabala, 1984), and in *C. taenia*, in Italy (Gambetta, 1934; Lodi, 1967; Marconato & Rasotto, 1989) and England (Robotham, 1981). As with females of the populations of *C. taenia*, the longevity of female *C. paludica* in the Palancar Stream is a year longer than that of males. By age 4+, females were clearly predominant. The reduced proportion of males as they age appears to be a result of earlier mortality, although hermaphroditism has been reported in other populations (Lodi, 1967).

Regarding their diet, *C. paludica* in the Palancar Stream consumed only a limited variety of resources. The most frequently found prey in their diet were chironomid larvae and Ostracoda, similar to the population of *C. taenia* studied by Moretti (1948) in Italy. Detritus and diatoms in gastrointestinal contents of this population was limited. However, these components predominate in the diet of this species in the marshlands of the Guadalquivir River (Hernando, 1974), and are also usually predominant in the diet of *C. taenia* (Skora, 1966; Maksunov, 1969) or only when other resources are scarce (Robotham, 1977). Prey of relatively large size, such as Ephemeroptera (Caenidae and Baetidae) and Trichoptera (Ecnomidae and Leptoceridae), were limited. Only at age 0+ are there significant differences in the diet in respect of age, although these were only differences in the consumption of isolated resources, such as macrobiotid spawns, and not in monthly variations in the diet.

Male and female growth rates were different. Parameters \mathbf{b} and \mathbf{b}' in SL-TW and SL-SW ratios were much higher for juveniles, and males showed lower values than females. The back-calculated length of females was greater than that of males. The population growth rate in the Palancar Stream was similar to that of the population in the Jarama River (Lobón-Cerviá, 1982), being somewhat lower for females and higher for males in the Palancar population. L_{∞} and k values in Von Bertalanffy's growth equation can be used as indicators for making comparisons of growth (Misra, 1980). The value of k was greater for females than for males in this population, which was not the case with the population in the Jarama River (Lobón-Cerviá & Zabala, 1984), where values for both sexes were also greater than those estimated for this population. Considering the index of overall fish growth performance, which is recommended by Moreau et al. (1986) because it has a smaller variation and can be used to compare similar taxa, once again, the growth of the

Table 4. Standard back-calculated lengths obtained for *Cobitis paludica* from Palancar Stream, Spain

Age	N	L I	L II	L III	L IV	L V
Males						
I+	3	36.75 +/-4.9				
II+	6	37.28 +/- 5.6	48.54+/-3.9			
III+	7	37.49 +/-3.4	48.32+/-2.7	54.85 +/-2.4		
IV+	1	36.77	43.97	53.94	58.94	
Mean		37.38+/-3.22	47.83+/-2.9	54.75+/-2.3	58.94	
Δ			10.5	6.92	4.19	
Females						
I+	2	39.40+/-6.0				
II+	6	31.38+/-5.8	46.48+/-4.7			
III+	6	30.40+/-3.7	46.62+/-6.2	60.79+/-4.2		
IV+	6	36.36+/-4.4	51.31+/-6.3	62.40+/-5.3	70.32+/-3.1	
V+	3	35.90+/-6.0	49.45+/-5.8	61.56+/-5.17	70.89+/-5.9	77.87+/-5
Mean		35.90+/-6	49.45+/-5.8	61.50+/-5.2	70.87+/-5.2	77.87+/-5
Δ			13.6	12.1	9.4	7.0

Table 5. Values of constant terms in Von Bertalanffy's equation of theoretical growth in length, k , L_{∞} , and t_0 , together with parameters ω and ϕ' for each sex, of *Cobitis paludica* from Palancar Stream, Spain

	K	L_{∞}	t_0	ω	ϕ'
Females	0.208866	110.094	-0.896726	22.99	3.403395
Males	0.439856	66.83	-0.867196	29.52	3.2932535

population studied was lower than the Jarama River population, for both males and females, and higher than that obtained by Slavik (1995) for *C. taenia*. As is the case of other populations studied (Robotham, 1981; Lobón-Cerviá & Zabala, 1984; Slavik, 1995), growth rates reached high values during the first years of life and low values in the later years, the decrease in growth being due to sexual maturity.

The vitelline oocyte mode of a size suitable for spawning in the month of July indicates a long reproductive period, already noted by Lobón-Cerviá & Zabala (1984) in this species, and by Marconato & Rasotto (1989) in *C. taenia*. These results cannot be taken to indicate the existence of two spawning periods, but they may indicate that spawning is sequential, starting in late April. Fecundity was similar to that described for this species in the Iberian Peninsula, with values ranging from 281–1397 oocytes for standard length interval 60–80 mm. Females of a smaller size

Table 6. Theoretical standard lengths obtained for *C. paludica* from von Bertalanffy's model (Palancar Stream, Spain)

	Males	Females
L_0	21.19	18.80
L_1	37.43	36.00
L_2	47.90	49.98
L_3	54.63	61.61
L_4	58.97	70.50
L_5		77.97

in the Palancar Stream were not as fecund as those of the same size in the Jarama River (Lobón-Cerviá &

Table 7. Instantaneous rates of growth in length (L_i) and weight (G_i), for *C. paludica* males and females from Palancar Stream, Spain

Age interval	L_i		G_i	
	Females	Males	Females	Males
I–II	0.320	0.238	0.987	0.481
II–III	0.218	0.135	0.814	0.307
III–IV	0.142	0.074	0.287	0.197
IV–V	0.094		0.147	

Zabala, 1984), whereas larger females had a higher fecundity than in the Jarama River.

Acknowledgements

Assistance with the English language presentation was provided by Royston Snart, CPI, Cádiz.

References

- Bagenal, T. B. & E. Braum, 1978. Eggs and early life history. In Bagenal, T. B. (ed.), *Methods for Assessment of Fish Production in Freshwaters*. IBP Handbook n° 3. Blackwell Scientific Publications, Oxford: 165–201.
- Bagenal, T. B. & F. W. Tesch, 1978. Age and growth. In: Bagenal, T. B. (ed.), *Methods for Assessment of Fish Production in Freshwaters*. IBP Handbook n° 3. Oxford: Blackwell Scientific Publications: 101–136.
- Bertalanffy, L. Von., 1938. A quantitative theory of organic growth. *Hum Biol.* 4: 229–236.
- Bertalanffy, L. Von. 1957. Quantitative laws in metabolism and growth. *B. Rev. Biol.*, 32: 217–231.
- Blanco, J. C. & J. L. Gonzalez, 1992. Red book of the Spanish vertebrates. Ministerio de Agricultura Pesca y Alimentación. Madrid: 714 pp. (in Spanish).
- Carlander, K. D., 1981. Caution on the use of the regression method of back-calculating length from scale measurements. *Fisheries* 6 (1): 2–4.
- Crim L. W. & B. D. Glebe, 1990. Reproduction. In Schreck, C. B. & P. B. Moyle (eds), *Methods for Fish Biology*. American Fisheries Society, Bethesda, Maryland: 529–554.
- Ford, E., 1933. An account of the herring investigation conducted at Plymouth during the years 1924–1933. *J. Mar. Biol.* 19: 305–384.
- Gallucci, V.F. & T.J. Quinn. 1979. Reparametrizing fitting and testing a simple growth model. *Trans. am. Fish. Soc.* 108: 14–25.
- Gambetta, L., 1934. Sulla variabil Cobite fluvial *C. taenia* e sul rapagsorto numerico dei sessi. *Boll. Mus. Zool. Anat. com. Torino Ser.* 3, 44: 193–194.
- Guziur, J. 1976. The feeding of two year old carp (*Cyprinus carpio* L.) in a vendance lake Klawoj. *Ekol. pol.* 24: 211–235.
- Hernando, J.A. 1978. Fish community structure of the Guadalquivir river marshlands. Doctoral Thesis. University of Sevilla. 365 pp. (in Spanish).
- Keast, A. & D. Webb, 1966. Mouth and body form relative to feeding ecology in the fish fauna of a small lake, Lake Opinicon, Ontario. *J. Fish. Res. Bd Can.* 23: 1845–1873.
- Lagler, K. F., 1956. *Freshwater Fishery Biology*. W.C. Brown Company Publishers Dubuque, Iowa: 421 pp.
- Lea, E., 1910. On the method used in the herring investigations. *Cons. Permanent International pour l'Exploitation de la Mer, Publications de Circonstance* 53: 7–174.
- Lobón-Cerviá, J. & A. Zabala, 1984. Observations on the reproduction of *Cobitis paludica*, de Buen, 1930, in the Jarama river. *Cybiurn* 8: 63–68.
- Lodi, E., 1967. Sex reversal of *Cobitis taenia* (L.). *Experientia* 23: 446–447.
- Maksunov, J. A. 1969. The Aral loach, *Urata aralensis*. *Problemy Ichthyol* 9: 320–325.
- Marconato, A. & M. B. Rasotto, 1989. The biology of a population of spined loach, *Cobitis taenia* L. *Boll. Zool.* 56: 73–80.
- Misra, R. K. 1980. Statistical comparison of several growth curves of the von Bertalanffy type. *Can. J. Fish. aquat. Sci.* 37: 920–926.
- Moreau, J., C., Bambino & D. Pauly. 1986. Indices of overall fish growth performance of 100 Tilapia (Cichlidae) populations: In MacLean, J. L., L. B. Dizon & L.V. Hosillos (eds), *The First Asian Fisheries Forum*. Manila, Philippines. Asian Fisheries Society. Pags: 201–206.
- Munro, J.L. & D. Pauli, 1983. A simple method for comparing the growth of fishes and invertebrates. *Fishbyte* 1: 5–6.
- Robothan, P. W. J., 1977. Feeding habits and diet in two populations of Spined loach, *Cobitis taenia* (L.). *Freshwat. Biol.* 7: 469–477.
- Skora, S., 1966. *Koza (C. taenia) Srzeki Wschidnies der Steinbisser aus dem Wscodnia – Bach*. *Acta Hydrob. Krakow* 8: 425–435.
- Slavik, O. & P. Rab, 1995. Effect of the microhabitat on the age and growth of two steam-dwelling populations of Spinet loach, *Cobitis taenia*. *Folia Zool.* 44: 167–174.
- Soriguer, M. C., 1996. Life cycle and trophic ecology of fishes from Palancar river (Sierras Subbéticas, Cordoba). Doctoral Thesis. University of Cordoba: 336 pp. (in Spanish).
- Spataru, P., 1979. Biologia nutritiei la zvirloga, *C. taenia taenia* din complexul de balti crapinajijila. *Ann. Univ. Bucurest Ser. Stiint. Natur.* 16: 163–168.
- Walford, L. A., 1946. A new graphic method for describing the growth of animals. *Bull. Mar. Biol. lab. Woods Hole* 90: 141–147.