



Short communication: a survey of grass-clover ley management and creation of a near infra-red reflectance spectroscopy equation to predict clover concentration

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Thomson, A. L., Humphries, D. J., Archer, J. E., Grant, N. W. and Reynolds, C. K. (2018) Short communication: a survey of grass-clover ley management and creation of a near infra-red reflectance spectroscopy equation to predict clover concentration. *Animal Feed Science and Technology*, 245. pp. 48-53. ISSN 0377-8401 doi: <https://doi.org/10.1016/j.anifeedsci.2018.09.003> Available at <http://centaur.reading.ac.uk/79139/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.anifeedsci.2018.09.003>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Short communication: A survey of grass-clover ley management and creation**
2 **of a Near Infra-Red Reflectance Spectroscopy equation to predict clover**
3 **concentration**

4 A. L. Thomson¹, D. J. Humphries¹, J. Archer², N. Grant², and C. K. Reynolds¹

5

6 ¹ *Centre for Dairy Research, School of Agriculture, Policy and Development,*

7 *University of Reading, Whiteknights Road, Reading, RG6 6AR, UK*

8 ² *Agri-Food and Biosciences Institute, Large Park, Hillsborough, Co Down, BT26*

9 *6DR, Northern Ireland*

10

11 Corresponding author: C. K. Reynolds. Email: c.k.reynolds@reading.ac.uk

12

13 **Keywords:** grass-clover, agronomy, silages, clover concentration, management,

14 NIRS,

15

16 **Summary**

17 The purpose of the present study was, firstly, to examine current practice for the
18 agronomy of grass-clover mixed swards used for silage-making in the UK, and
19 secondly, to develop and validate a Near Infra-Red Reflectance Spectroscopy (NIRS)
20 equation capable of predicting clover concentration (CC) in undried and unmilled
21 grass-clover silage samples. A calibration set of 94 grass-clover (white, *trifolium*
22 *repens*, and red, *trifolium pratense*) mixture silage samples were sourced from UK
23 farms and an accompanying questionnaire was used to obtain information on the
24 sward agronomy used to produce each sample. Questionnaire data highlighted that
25 (i) reducing the use of fertiliser inputs (ii) increasing uptake of new varieties, and (iii)
26 increasing the farmer's ability to measure botanical composition as potential strategies
27 for improving the utilisation of clover in grass swards. Botanical composition was
28 measured by hand separation for each sample and a new NIRS equation was created
29 and assessed using blind validation with an independent set of 30 grass-clover
30 samples. The relative standard error of cross validation (SECV, as a percentage of the
31 measured mean) of the optimised equation produced was 36.8%, and, in an
32 independent validation test, the ratio of standard error of prediction to the standard
33 deviation of the reference data set (RPD) was 1.56. The equation could be improved
34 by increasing accuracy at high CCs but showed promise as a simple tool to assist
35 growers in sward management decisions.

36

37 **Introduction**

38 The use of mixed grass-clover swards for both grazing and silage production is now
39 relatively wide-spread across temperate European agricultural systems and
40 particularly in the UK where 70% of grass swards on dairy farms are thought to contain

41 clover (DEFRA, 2015). Grass-legume swards offer a sustainable approach to reduce
42 fertiliser input into grasslands, as the atmospheric nitrogen (**N**) fixed by clover can be
43 utilised by grass, an example of niche complementarity between two species (Nyfeler
44 et al., 2011, Phelan et al., 2015). Utilising niche complementarity in this way is
45 becoming an area of increasing interest for both binary and complex (3+ species)
46 sward mixtures. Combining species of different functional groups, can not only
47 increase productivity and minimise the need for inputs, but may also supply different
48 beneficial nutrients, minerals, and secondary plant compounds to livestock (Provenza
49 et al., 2007). A key determinant of the success of mixed swards is determining and
50 maintaining the correct concentration ratio of species or functional groups so that they
51 work in harmony. Previous research has shown that, in general, the best results can
52 be achieved by an even distribution of species within a sward, with no single species
53 becoming dominant (Finn et al., 2013). Where one species is over-dominant the others
54 may not reach their production potential or fulfil their niche functionalities, and the
55 productivity of the whole sward could be reduced (Kirwan et al., 2007, Lüscher et al.,
56 2014).

57 To date, we are not aware of any published surveys that document the
58 management strategies farmers utilise for grass-clover leys, and to what extent these
59 conform to best practice guidelines for maintaining species evenness. Additionally,
60 there is a need for increased development of practical methods by which growers can
61 manage species evenness within a sward, beginning with simple binary mixtures. The
62 first step to improved management is the ability to measure the botanical composition
63 of a sward with ease. Near Infra-Red Spectroscopy (**NIRS**) analysis offers a quick and
64 inexpensive method, already routinely used for silage analysis, by which the
65 composition of a mixed sample might be determined. Prediction equations for NIRS

66 analysis of clover in a mixed grass-clover silage sample have been successfully
67 reported previously using dried samples for calibration (Wachendorf et al., 1999,
68 Cougnon et al., 2014) however no prediction equations currently exist which are
69 appropriate for the UK where silage analysis is performed on undried (fresh) and
70 unmilled samples. Once the botanical composition of a sward is known, management
71 may be adjusted to suit one species or another by varying cutting height, cutting
72 frequency, fertilisation or grazing intensity (Yarrow and Penning, 1994, Phelan et al.,
73 2014).

74 The objectives of the present study were therefore to develop an NIRS equation
75 to measure the botanical composition of fresh grass-clover silages appropriate for
76 uptake by laboratories in the UK, and secondly, to assess current management
77 practices of grass-clover swards to better understand where further research into
78 management of botanical composition is required.

79

80 **Material and methods**

81

82 *Experimental design*

83 Ninety-four grass-clover silages (58 baled and 36 clamped) were sourced from 50
84 commercial farms spread throughout England, Scotland, and Wales, and brought to
85 the University of Reading's Centre for Dairy Research (**CEDAR**; Arborfield, UK) for
86 processing. A further 95th sample was created by combining one of the original 94
87 samples with additional grass silage to create a new sample, this was done to create
88 a greater quantity of material for other *in vivo* analyses. The samples were obtained
89 to evaluate the use of NIRS analysis for nutrient concentrations as described
90 previously (Thomson et al., 2018) over three consecutive years (2012/13, 2013/14,

91 and 2014/15). The quantity of each silage collected was approximately 500 kg. Where
92 the clover species was known ($n = 65$) 66% of samples were red clover (*trifolium*
93 *repens*), 20% were white clover (*trifolium pratense*) and 14% were a mixture of both.
94 The number of samples that were first, second, third and fourth cuts were 36, 20, 16,
95 and 4 respectively (22 samples unknown). Sample processing is described in detail
96 by Thomson *et al.* (2018), however, in brief, samples were mixed either in a feeder
97 wagon containing knives (Hi-Spec Mix Max, Hi Spec Engineering, Co. Carlow, Ireland
98 for 45 min), or in a DataRanger feed mixer without knives (American Calan,
99 Northwood, NH, USA) for unchopped and pre-chopped samples respectively. After
100 mixing, representative subsamples of each silage sample were stored separately at -
101 20°C for future analysis by manual separation and NIRS.

102

103 *Silage Questionnaire*

104 A questionnaire was given to each farmer who donated a silage sample to the study.
105 The questionnaire comprised 17 questions (Supplementary Table 1) relating to the
106 timing of establishment, fertiliser applications, and harvesting; the composition of seed
107 mixtures used; and ensiling practices. For the botanical composition of the seed
108 mixture, the variety sown was recorded for ryegrass and clover whereas any other
109 components were simply recorded at the species level as variety was rarely provided.
110 In addition farmers were asked to retrospectively estimate the percentage of clover in
111 the sward at the time of harvest (Question 9, Supplementary Table 1). Farms were
112 permitted to contribute more than one silage sample to the study provided the samples
113 originated from differing cuts, years, or swards. Separate questionnaires were
114 completed for each of the samples. Questionnaire forms were returned for 64 of the
115 94 samples however not all questions were answered on all returned questionnaires

116 and in some instances answers were insufficiently detailed to be included. These 64
117 completed questionnaires originated from 36 individual farms, reflecting that a number
118 of farms returned more than one questionnaire, each relating to a different crop of
119 silage.

120

121 *Sample analysis and NIRS scanning*

122 Approximately 200 g of silage was manually aspeciated into clover, grass and other
123 species as a means of determining the clover concentration (**CC**) in the silage.
124 Resulting fractions were then oven dried at 60°C for 72 h to determine CC on a DM
125 basis. A second 2 kg subsample of frozen material was sent to the Agri-Food and
126 Biosciences Institute (**AFBI**; Hillsborough, Northern Ireland) where all samples were
127 hand-chopped to 2.5 cm length. Two separate packages were created from each
128 sample, each containing 100 g of undried and unmilled silage wrapped in non-PVC
129 cling film which were placed in coarse transport cells for scanning (Park et al., 1999).
130 NIRS spectra for each scan, recorded as Log 1/Reflectance over a 400-2498 nm range
131 (2 nm gaps), were obtained using a Foss NIRSystems 6500 machine (Foss, Hillerød,
132 Denmark) and ISI v.3.10 software (Infrasoft International, Port Matilda, PA, USA).

133

134 *Data Analysis*

135 *Statistical analysis.* Survey data are summarised and reported as percentages of
136 recieved responses for each question. A one-way ANOVA followed by a post-hoc
137 Tukey test was used to determine the effect of cut number in the year (1st, 2nd, 3rd or
138 4th) on CC, where CC was the independent variable, in Genstat 16th Edition (VSN
139 International, Hemel Hempsted, UK). *P* values < 0.05 were considered statistically

140 significant. As the survey only concerns one group of farmers (rather than contrasting
141 multiple groups) no further statistical analysis was deemed necessary.

142

143 *Data pre-treatment and production of new NIRS equations.* Creation of a new
144 prediction equation for CC was performed as described by Thomson *et al.* (2018)
145 using WinISI III v1.50 (Infrasoft International, Port Matilda, PA, USA). Calibration was
146 carried out as Modified Partial Least Squares regressions over the range 1100-2498
147 nm using a 2 nm gap. To prevent any sub-sampling error the root mean square
148 difference of each sub-sample was calculated (an upper limit of 5000 was used to
149 judge poor replication however none of the samples in the calibration set were above
150 this limit). Raw data and two derivatives were tested (Raw (0,0,1,1), 1st Derivative
151 (1,4,4,1) and 2nd Derivative (2,10,5,1)) and three scatter corrections (Standard
152 Normal Variate Detrending (**SNVD**), Normal Multiplicative Scatter Correction (**NMSC**)
153 and Weighted Multiplicative Scatter Correction (**WMSC**)) for each of the derivatives.
154 The maximum number of terms set for each equation was 11. For cross validation,
155 three elimination passes were carried out and the cross validation value was set at 6
156 (i.e. the calibration set was divided into six groups with one group removed
157 sequentially and predicted using a calibration formed using the remaining samples)
158 and the combined standard error of cross validation (**SECV**) was obtained in addition
159 to the standard error of calibration (**SEC**). The optimal equation (lowest SECV) was
160 compared against an equation produced using the UK industry standard data-
161 pretreatment method: first derivative (1,4,4,1) SNVD scatter correction and a
162 repeatability file (a file containing multiple spectra from the same sample measured
163 under different conditions) (Park *et al.*, 1997). For the purposes of a blind validation
164 test, 30 independant grass-clover silage samples of known CC (for which NIR spectra

165 had been previously obtained at AFBI in a prior study) were used to assess prediction
166 accuracy of the new equation using root mean square error of prediction (**RMSEP**)
167 and the ratio of the standard error of prediction to the standard deviation of the
168 measured sample set (**RPD**) (Williams, 2014).

169

170 **Results and Discussion**

171 *Grass-clover management questionnaire*

172 The final set of 95 samples had a mean concentration of clover, grass and other
173 species of 310, 640 and 50 g/kg DM, respectively (median CC = 280 g/kg DM). The
174 distribution of CC, shown in Figure 1a, indicated that low CC silages were more
175 common than high CC silages within the sample set. Ploughing or subsoiling were the
176 most common forms of cultivation prior to establishment with 48% of those providing
177 cultivation information employing these methods. A further 34% of respondents used
178 light cultivation such as discs, tines or a power harrow and 18% reported using a
179 minimum or no tillage approach. In 18 instances (31% of responses) the sward was
180 established by under-sowing. The timing of crop establishment (where known) was
181 roughly evenly split between spring and autumn with 20 sown in March, April or May,
182 4 sown in June or July, and 22 sown in August or September.

183 All farms providing seed information reported sowing more than one variety of
184 grass and 22 (52% of responses) sowed more than one variety of clover within a single
185 ley. Sowing for increased varietal richness helps mitigate the risk of any one variety
186 performing poorly (Surault et al., 2010). In total 38 different grass varieties including
187 varieties of italian ryegrass, ryegrass hybrids, timothy, cocksfoot, fescues, and
188 festuloliums were sown, but the predominant species was perennial ryegrass (23 of
189 the 38 varieties). In comparison, only 19 different varieties of clover were represented

190 within the sample set which was likely reflective of the wider range of grass varieties
191 available on the market. The two most frequent grass varieties sown were 'Solid' (34%
192 of responses) and 'Tetragraze' (26% of responses) which are both hybrids of Italian
193 and perennial ryegrass. 'High sugar' grass varieties 'AberDart' and 'AberStar' were the
194 joint 5th most frequently sown grasses (each sown in 12% of reported swards)
195 indicating good uptake, possibly as a result of research showing the use of high-sugar
196 grasses in combination with red clover produces a favourable balance of
197 metabolisable protein and fermentable energy in the ruminant diet (Merry et al., 2006).
198 Of the four most frequently sown clover varieties, all were red clovers with the varieties
199 'Merviot' and 'Milvus' being the most popular (sown in 33% and 21% of reported
200 swards). These are both older varieties (for example, Merviot was first introduced to
201 the UK recommended list of varieties in 1980) perhaps indicating a need for greater
202 adoption of newer clover varieties to take advantage of genetic gains (Frick et al.,
203 2008, Capstaff and Miller, 2018). A recent study showed that, out of 12 clover varieties
204 sown, 10 new varieties showed increased persistency within a 3-year ley when
205 compared to Merviot or Milvus (Marshall et al., 2012).

206 Information relating to applications that occurred in the year prior to harvesting
207 was given for 46 of the silages. Of the 46 silages, 23 silages (50%) had slurry applied
208 and 12 (26%) had been fertilised using an inorganic fertiliser containing nitrogen (N).
209 A further 5 (11%) had an application of farm yard manure. Only 5 silages (11%) had
210 been reported as having no fertiliser applied. Where excess applications of N are
211 applied, clover adapts by reducing the rate of fixation, and is more likely to be
212 outcompeted by grass (Nyfeler et al., 2011). Farmers were not required to state the
213 timing of the application; therefore, the applications may have been early in the year
214 or during establishment for first year silages, however, even out-of-season N

215 applications have been shown to reduce CC over the summer months (Laidlaw et al.,
216 1992). Those applying slurry may benefit from the application of potassium and
217 phosphorous for soil fertility, however, even slurry may contain enough N to adversely
218 affect N fixation efficiency by clover and sward CC if applied in excess of 50 T/ha
219 (Nesheim et al., 1990). This evidence suggests that few farmers are maximising the
220 N-fixation potential of clover and possibly seeing reduced economic performance as
221 a result, particularly where expensive inorganic N fertiliser is applied to clover-
222 containing swards.

223 The most common months for taking silage cuts were May, July and September
224 for 1st, 2nd, and 3rd cuts respectively. Stepwise increases in CC were seen in 1st to 3rd
225 cuts ($P < 0.001$) as shown in Figure 1b. The use of silage additives was relatively
226 common with 13 different brands of additive reported. Of these 13 additives used, 10
227 were bacterial inoculants and the remainder contained either enzymes or salt as the
228 active ingredient. Of 29 total responses to the question on the use of additives, 5
229 specified that no additive had been used.

230 Farmers were retrospectively asked to estimate the CC of the sample based on
231 their memory of the sward at harvest. Fifty-five estimations were received and
232 compared against reference values obtained by hand separation (Figure 2). The
233 majority of farmers estimated a value between 300 to 700 g/kg DM clover. As a result
234 samples containing a measured CC of less than 400 g/kg DM were generally over-
235 estimated while samples containing greater than 400 g/kg DM CC were often under-
236 estimated. The number of farmers that successfully predicted CC to within ± 100 and
237 ± 200 g/kg DM was low at 15 and 31 out of 55 respectively. One possible explanation
238 for this would be poor uniformity within the sward meaning that the sample taken was
239 not representative of the general crop (Marriott et al., 1997), although this explanation

240 is more valid for baled samples as opposed to those ensiled in a clamp where mixing
241 is performed in the forage harvester. The inability of many farmers to recall an accurate
242 estimate of the CC of their forage highlights the need for tools to be developed which
243 automate this process and provide lasting records; one option being the use of NIRS
244 on resulting silage which has been explored in the present study. Another option which
245 has been investigated previously is to determine n-alkane concentration which differs
246 distinctively between species (Jurado et al., 2015), however, the use of a laboratory
247 assay of this kind is expensive and time consuming.

248

249 *Creation and validation of an NIRS equation to predict clover concentration*

250 The best-performing NIRS prediction equation for CC was produced using 182
251 spectra, and NMSC (2,10, 5,1) data pre-treatment which gave an SEC of 8.99 and an
252 SECV of 36.8% of the measured mean. Using the industry standard data pre-
253 treatment of SNVD (1,4,4,1) slightly reduced calibration performance by a small
254 margin with an SEC of 10.0 and an SECV of 37.5% of the measured mean.

255 Spectra from 30 independent grass-clover silage samples of known CC were
256 used in a blind validation test of the industry standard CC equation. The measured
257 mean CC within this independent set of samples was 440 g/kg DM, which was greater
258 than that of the calibration set. Using this data set the equation gave an RMSEP of
259 52.2% of the measured mean and an RPD value of 1.56. Plotting measured against
260 predicted CC in this test indicated over-estimation at low CC and under-estimation at
261 high CC with the average bias being towards under-estimation (Figure 3). For samples
262 that contained 1000 g/kg clover by DM, the equation underpredicted CC by 300-400
263 g/kg DM. A group of samples with 150 g/kg DM CC showed the best prediction
264 accuracy. Prediction for samples with the same botanical composition often differed

265 by 150-400 g/kg DM CC indicating low repeatability. Other studies in which prediction
266 equations for CC have been produced have shown more robust prediction accuracy
267 than the present study, for example Cougnon *et al.* (2014) produced an equation with
268 an RPD of 3.8 using just 42 silage samples as a calibration set. In another study
269 Wachendorf *et al.* (1999) produced separate equations for freshly cut grass-red clover
270 and grass-white clover mixtures with large calibration sets ($n = 282$ and 183
271 respectively) where the relative SECV were 29.9 and 23.5% respectively, which again
272 shows an improvement over equations in the present study. A major difference
273 between the calibration samples used in the present study and those used by
274 Cougnon *et al.* (2014) and Wachendorf *et al.* (1999) is that no sample preparation
275 such as drying or milling was used in line with UK recommendations whereas, in the
276 previous studies samples were dried and milled. The drying and milling process
277 reduces heterogeneity within the sample (which is particularly important to
278 representatively subsample mixtures for analysis) and also removes peaks produced
279 by water molecules from the resulting NIRS spectra, reducing noise, and improving
280 interpretation and repeatability (Sorensen, 2004). The reference technique used to
281 measure CC (manual separation prior to drying and milling) was the same in the
282 present study and in the studies of Cougnon *et al.* (2014) and Wachendorf *et al.*
283 (1999), however, the drawback of this technique is that results can be subjective,
284 particularly on chopped samples as were used in the present study, due to the lack of
285 species-defining characteristics on some particles. Using a reference technique prone
286 to human error such as this reduces the likelihood of producing a robust prediction
287 equation. We conclude that, even though the equation produced in the present study
288 was less robust than in previous examples, the equation may still be of use to guide
289 on-farm decisions, particularly for swards containing low CCs.

290

291 **Acknowledgements**

292 Funding for the project was provided by AHDB Dairy as part of the Soils, Forage and
293 Grassland Partnership. Collaborators within the partnership included the Forage
294 Analytical Assurance (FAA) Group who offered in-kind contributions to the project and
295 particular thanks are given to group chairs Dr Dave Davies and Dr Jonathan Blake for
296 their invaluable advice throughout the study. We would like to acknowledge all the
297 farmers who contributed silage samples to the project. Further thanks go to Mr C.
298 Green, Dr. M. Dittmann, Mr. P. Kirton, Mr. C. Fawdry, Mr. R. Szoka and Ms. D.
299 Cockman for technical inputs.

300

301 **References**

302 Capstaff, N. M. and A. J. Miller. 2018. Improving the Yield and Nutritional Quality of
303 Forage Crops. *Frontiers in Plant Science* 9.
304 Cougnon, M., C. Van Waes, P. Dardenne, J. Baert, and D. Reheul. 2014.
305 Comparison of near infrared reflectance spectroscopy calibration strategies
306 for the botanical composition of grass-clover mixtures. *Grass and Forage*
307 *Science* 69:167-175.
308 DEFRA. 2015. Farm Practices Survey: Basic Payment Scheme and greening -
309 England. *In National Statistics*.
310 Finn, J. A., L. Kirwan, J. Connolly, M. Teresa Sebastia, A. Helgadottir, O. H.
311 Baadshaug, G. Belanger, A. Black, C. Brophy, R. P. Collins, J. Cop, S.
312 Dalmannsdottir, I. Delgado, A. Elgersma, M. Fothergill, B. E. Frankow-
313 Lindberg, A. Ghesquiere, B. Golinska, P. Golinski, P. Grieu, A.-M.
314 Gustavsson, M. Hoglind, O. Huguenin-Elie, M. Jorgensen, Z. Kadziuliene, P.
315 Kurki, R. Llorba, T. Lunnan, C. Porqueddu, M. Suter, U. Thumm, and A.
316 Lüscher. 2013. Ecosystem function enhanced by combining four functional
317 types of plant species in intensively managed grassland mixtures: a 3-year
318 continental-scale field experiment. *Journal of Applied Ecology* 50:365-375.

- 319 Frick, R., B. Jeangros, M. Demenga, D. Suter, and H. U. Hirschi. 2008. Red clover
320 variety tests. *Revue Suisse D Agriculture* 40:245-248.
- 321 Jurado, N. V., A. E. Tanner, S. R. Blevins, H. M. McNair, R. W. Mayes, and R. M.
322 Lewis. 2015. Long-chain alcohols did not improve predictions of the
323 composition of tall fescue and red clover mixtures over n-alkanes alone.
324 *Grass and Forage Science* 70:499-506.
- 325 Kirwan, L., A. Lüscher, M. T. Sebastia, J. A. Finn, R. P. Collins, C. Porqueddu, A.
326 Helgadottir, O. H. Baadshaug, C. Brophy, C. Coran, S. Dalmannsdottir, I.
327 Delgado, A. Elgersma, M. Fothergill, B. E. Frankow-Lindberg, P. Golinski, P.
328 Grieu, A. M. Gustavsson, M. Hoglind, O. Huguenin-Elie, C. Iliadis, M.
329 Jorgensen, Z. Kadziulienė, T. Karyotis, T. Lunnan, M. Malengier, S. Maltoni,
330 V. Meyer, D. Nyfeler, P. Nykanen-Kurki, J. Parente, H. J. Smit, U. Thumm,
331 and J. Connolly. 2007. Evenness drives consistent diversity effects in
332 intensive grassland systems across 28 European sites. *Journal of Ecology*
333 95:530-539.
- 334 Laidlaw, A. S., N. G. Teuber, and J. A. Withers. 1992. Out-of-season management of
335 grass clover swards to manipulate clover content. *Grass and Forage Science*
336 47:220-229.
- 337 Lüscher, A., I. Mueller-Harvey, J. F. Soussana, R. M. Rees, and J. L. Peyraud. 2014.
338 Potential of legume-based grassland-livestock systems in Europe: a review.
339 *Grass and Forage Science* 69:206-228.
- 340 Marriott, C. A., G. R. Bolton, and E. I. Duff. 1997. Factors affecting the stolon growth
341 of white clover in ryegrass/clover patches. *Grass and Forage Science* 52:147-
342 155.
- 343 Marshall, A. H., M. Lowe, and J. Vale. 2012. Improved persistence of red clover
344 (*Trifolium pratense* L.) varieties in mixed swards. *In Grassland - a European*
345 *Resource?*. 24th General Meeting of the European-Grassland-Federation.
346 Lublin, Poland, 3rd – 7th June. pp 73-75.
- 347 Merry, R. J., M. R. F. Lee, D. R. Davies, R. J. Dewhurst, J. M. Moorby, N. D. Scollan,
348 and M. K. Theodorou. 2006. Effects of high-sugar ryegrass silage and
349 mixtures with red clover silage on ruminant digestion. 1. In vitro and in vivo
350 studies of nitrogen utilization. *Journal of Animal Science* 84:3049-3060.
- 351 Nesheim, L., B. C. Boller, J. Lehmann, and U. Walther. 1990. The effect of nitrogen
352 in cattle slurry and mineral fertilizers on nitrogen-fixation by white clover the

353 effect of nitrogen in cattle slurry and mineral fertilizers on nitrogen-fixation by
354 white clover. *Grass and Forage Science* 45:91-97.

355 Nyfeler, D., O. Huguenin-Elie, M. Suter, E. Frossard, and A. Lüscher. 2011. Grass-
356 legume mixtures can yield more nitrogen than legume pure stands due to
357 mutual stimulation of nitrogen uptake from symbiotic and non-symbiotic
358 sources. *Agriculture Ecosystems & Environment* 140:155-163.

359 Park, R. S., R. E. Agnew, F. J. Gordon, and R. J. Barnes. 1999. The development
360 and transfer of undried grass silage calibrations between near infrared
361 reflectance spectroscopy instruments. *Animal Feed Science and Technology*
362 78:325-340.

363 Park, R. S., F. J. Gordon, R. E. Agnew, R. J. Barnes, and R. W. J. Steen. 1997. The
364 use of Near Infrared Reflectance Spectroscopy on dried samples to predict
365 biological parameters of grass silage. *Animal Feed Science and Technology*
366 68:235-246.

367 Phelan, P., I. A. Casey, and J. Humphreys. 2014. The effects of simulated summer-
368 to-winter grazing management on herbage production in a grass-clover
369 sward. *Grass and Forage Science* 69:251-265.

370 Phelan, P., A. P. Moloney, E. J. McGeough, J. Humphreys, J. Bertilsson, E. G.
371 O'Riordan, and P. O'Kiely. 2015. Forage Legumes for Grazing and
372 Conserving in Ruminant Production Systems. *Critical Reviews in Plant*
373 *Sciences* 34:281-326.

374 Provenza, F. D., J. J. Villalba, J. Haskell, J. W. MacAdam, T. C. Griggs, and R. D.
375 Wiedmeier. 2007. The value to herbivores of plant physical and chemical
376 diversity in time and space. *Crop Science* 47:382-398.

377 Sorensen, L. K. 2004. Prediction of fermentation parameters in grass and corn silage
378 by near infrared spectroscopy. *Journal of Dairy Science* 87:3826-3835.

379 Surault, F., B. Julier, and C. Huyghe. 2010. Forage Production of Grasslands
380 Composed by One, Two or Three Varieties of Perennial Ryegrass. *In*
381 *Sustainable Use of Genetic Diversity in Forage and Turf Breeding*. pp 239-
382 244.

383 Thomson, A. L., D. J. Humphries, C. Rymer, J. Archer, N. Grant, and C. K.
384 Reynolds. 2018. Assessing the accuracy of current near infra-red reflectance
385 spectroscopy analysis for fresh grass-clover mixture silages and development

386 of new equations for this purpose. *Animal Feed Science and Technology*
387 239:94-106.

388 Wachendorf, M., B. Ingwersen, and F. Taube. 1999. Prediction of the clover content
389 of red clover- and white clover-grass mixtures by near-infrared reflectance
390 spectroscopy. *Grass and Forage Science* 54:87-90.

391 Williams, P. 2014. Tutorial: The RPD statistic: a tutorial note. *NIR news* 25:22-26.

392 Yarrow, N. H. and P. D. Penning. 1994. Managing grass clover swards to produce
393 differing clover proportions. *Grass and Forage Science* 49:496-501.

394

395 **Figure captions**

396

397 **Figure 1** The distribution of clover concentration within a set of 94 grass-clover
398 silages sourced from working farms across the UK over various cuts (1st - 4th) and
399 years (2012-2015; Figure 1a) and the effect of cut number on clover concentration
400 (Figure 1b).

401 **Figure 2** The relationship between actual clover concentration (●) and the grower's
402 prediction of clover concentration (○) in a range of 54 grass-clover silage samples
403 sourced from working UK farms over several years (2012-2015). White drop lines
404 indicate over-prediction and dark drop lines indicate under-prediction of clover
405 concentration.

406 **Figure 3** The results of a blind validation of a Near Infra-Red Spectroscopy
407 prediction equation for clover concentration calibrated using a set of 94 diverse fresh
408 grass-clover silage samples which were manually aspeciated to produce reference
409 values and validated using an independent set of 30 grass-red clover silage samples
410 of 6 known clover concentrations (0, 150, 450, 600, and 1000 g/kg dry matter)

411