

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/323936624>

Effects of Medicinal Plants on Nocturia: A Systematic Review

Article · March 2018

DOI: 10.9734/JPRJ/2018/40251

CITATIONS

0

READS

19

3 authors, including:



Saeid Heidari Soureshjani

Shahrekord University of Medical Sciences

31 PUBLICATIONS 37 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Anti-cancer activity of medicinal plants [View project](#)



Effects of Medicinal Plants on Nocturia: A Systematic Review

**Mahnaz Yavangi¹, Ramazan Khodadadi-Arpanahi²
and Saeid Heidari-Soureshjani^{3*}**

¹*Endometrium and Endometriosis Research Center, Hamedan University of Medical Sciences, Hamadan, Iran.*

²*Deputy of Health, Shahrekord University of Medical Sciences, Shahrekord, Iran.*

³*Deputy of Research and Technology, Shahrekord University of Medical Sciences, Shahrekord, Iran.*

Authors' contributions

This work was carried out in collaboration between all authors. Author MY designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors MY and SHS managed the analyses of the study. Authors SHS and RKA managed the literature Searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2018/40251

Editor(s):

(1) Wenbin Zeng, School of Pharmaceutical Sciences, Central South University, Hunan, China.

Reviewers:

(1) M. Vani, Vijaya Institute of Pharmaceutical Sciences for Women, India.

(2) Emeritus Dr. Mohd Ismail Bin Mohd Tambi, Malaysia.

Complete Peer review History: <http://www.sciencedomain.org/review-history/23725>

Review Article

Received 27th December 2017

Accepted 4th March 2018

Published 20th March 2018

ABSTRACT

Nocturia is one of the common disorders that are associated with certain complications such as fracture and declined quality of life in addition to disrupting sleep. We, therefore, sought to review the evidence on the effects and action mechanisms of medicinal plants on nocturia.

Methods: The search terms of interest were used to retrieve relevant articles published between 2000 and 2018 and indexed in the databases *Institute for Scientific Information*, *PubMed*, and *Scopus* using *EndNote* software.

Peganum harmala and *Cannabis sativa* are two of the plants that have been reported to be effective on nocturia. In addition, certain plant-based derivatives such as SagaPro and certain combinations such as Gosha-jinki-gan, paladin kashaya, and PRO 160/120 from *Viola odorata* L. (Violaceae), C.A.Mey. From *Echium amoenum* Fisch. (Boraginaceae), and *Physalis alkekengi* L. (Solanaceae) help improve nocturia through different mechanisms.

*Corresponding author: E-mail: heidari_62@yahoo.com, heidari_1983@yahoo.com;

As with other types of treatments, plants and plant-based compounds cannot be used as monotherapy in treating nocturia, and therefore should be used as a complementary therapy alongside available treatments for this disorder to induce the best therapeutic effects.

Keywords: Medicinal plant; nocturia; lower urinary tract; symptoms.

1. INTRODUCTION

Nocturia is one of the most common disorders of lower urinary tract symptoms (LUTS) [1] and refers to waking up to urinate more than once each night [2]. This disorder is associated with certain complications such as benign prostatic obstruction and overactive bladder as well as serious adverse effects on quality of life in the affected people [1,3]. The people with nocturia are faced with disruption in night's sleep [4,5] that affects different aspects of health adversely. Inadequate night's sleep leads ultimately to increased mortality rate [6]. In addition, these people are likely to fall down and to develop fracture especially femur fracture [7]. Because this disease is multifactorial, single drug therapies are not sufficiently efficacious [8]. Therefore, nocturia is treated by multiple treatments including improvement of lifestyle, chemical drugs, alpha-blockers, antimuscarinic agents, antidiuretic therapy, and surgery [9]. The treatments for night time urination mainly include urotherapy and different drugs such as desmopressin, imipramine, and anticholinergic drugs as well as behavioral therapies [10,11]. However, these treatments should be promoted [10]. Medicinal plants can represent effective treatments for different diseases including urinary tract disorders. The popularity of these plants has increased considerably because of comparatively lower cost and fewer side effects [12-19]. Regarding the complications of nocturia, the present review was conducted to investigate the effects and action mechanisms of the medicinal plants on this night time disorder.

2. MATERIALS AND METHODS

To conduct this systematic review, the search terms *Nocturia* or *Nocturnal voids* in combination with *Herb*, *Medicinal plant*, and *Phyto* were used to retrieve relevant articles published between 2000 and 2018 and indexed in the databases *Institute for Scientific Information*, *PubMed*, and *Scopus* using *EndNote* software. Then, the plants and products that were reported to be effective on and therefore control nocturia were identified and, according to the research team's comments, included in the study. Studies that

directly examined their effect on nocturia and showed positive effects were investigated. The studies that Fig. 1 illustrates how the articles were examined for inclusion in the analysis and explains the inclusion criteria.

3. RESULTS

The medicinal plants such as *Serenoa repens*, *Pygeum africanum*, *Urtica dioica*, and *Cucurbita pepo* were reported to be the most important medicinal plants effective on nocturia and certain compounds such as isoflavones, lycopene, selenium, and beta-sitosterol are being addressed for treating the symptoms associated with benign prostate hyperplasia (BPH) such as nocturia [20]. Because BPH causes certain symptoms such as nocturia, decreased voided urine volume, the sensory urgency to urinate, the frequency of urination during the day or night, inability to start and stop a urine stream [21], most studies have reported nocturia to be one of the symptoms to measure the severity of BPH. Table 1 summarizes the evidence on the plants and the phytochemical compounds that are effective on nocturia.

Several compounds and formulations exist that are used in traditional medicine to treat nocturia. Goshajinki-gan (GJG) is a combination made up of 10 plants that are used in Japanese traditional medicine. This formula includes *Rehmannia Radix*, *Achyranthis Radix*, *Copri Fructus*, *Dioscoreae Rhizoma*, *Plantaginis Semen*, *Alismatis Rhizoma*, *Hoclen*, *Moutan Cortex*, *Cinnamomi Cortex* and *Aconiti Tuber*.

The six-week treatment with this combination (2.5 g GJG mixture x 3/day) can help improve disorders of urination such as nocturia [25]. Another study showed that this combination, without interaction with alpha1-blockers or antimuscarinic agents, contributed to treating nocturia [26]. However, Yoshimura et al. reported that furosemide was more efficient in decreasing nocturia than GJG [27]. Pataladi kashaya that compose of Patola – *Trichosanthes dioica*, *Triphala*, *Amla* – *Emblica officinalis*, *Haritaki* – *Terminalia chebula*, *Vibhitaki* – *Terminalia bellirica*, *Arishta* – *Nimba* – *Neem* – *Azadirachta*

indica, Guduchi – *Tinospora cordifolia*, Dhavani – *Solanum indicum*, Vrisha – Vasa – *Adhatoda vasica*, Karanja – *Pongamia pinnata*, is a drug used in Ayurvedic medicine that is useful to improve BPH symptoms including nocturia in addition to decreasing such symptoms [28].

Oelke et al. studied a plant-based combination called PRO 160/120 (extracts from saw palmetto fruits and stinging nettle roots), and reported that

nocturia score, according to the *International Prostate Symptom Score* (IPSS), decreased [29]. A study on the effects of hydroalcoholic extracts of *Viola odorata* L. (Violaceae), C.A.Mey. (Boraginaceae), *Echium amoenum* Fisch., and *Physalis alkekengi* L. (Solanaceae) showed that this combination caused IPSS including nocturia to decline [30]. It has been even reported that herbal capsules work better in treating nocturia than allopathic medicine [31].




Fig. 1. The flowchart of examining articles according to exclusion and inclusion criteria

Table 1. Medicinal plants and their phytochemical compounds effective on nocturia

Plants/ phytochemicals name	Part of plant/ Isolated from	Main effects and mechanisms	References
<i>Cannabis sativa</i>	Extracts containing delta-9-tetrahydrocannabinol and cannabidiol	Effect on cannabinoid receptors in the bladder and nervous system	[22]
<i>Peganum harmala</i>	Extract	Improvement in BPH symptoms including nocturia	[23]
SagaPro	<i>Angelica rchangelica</i>	Reduce the score of nocturnal bladder capacity index (NBC index)	[24]

A number of non-clinical and *in vitro* studies have reported that some plants are effective on nocturia. For example, *Pygeum africanum* [32-34], *rye pollen* [35], and *Serenoa repens* [36-41], and also plant compound such as Cernilton [32,42,43] and beta-sitosterols (B-sitosterol) [44] have been found to decrease one of the complications associated with nocturia.

Antiandrogenic action, the antiproliferative effect exerted through the inhibition of growth factors, anti-inflammatory/anti-oedematous effect and prolactin signal modulation are certain mechanisms that have been proposed for treating BPH and associated complications [40]. However, the action mechanisms of the medicinal plants in treating nocturia remain ambiguous. Certain plants and their derivatives have been reported to control nocturia through decreasing irritable bladder via affecting spinal kappa-opioid receptors [45] distributing water among different organs of the body and decreasing urination [45] and decreasing oxidative stress. Oxidative stress through damage to the urothelium imposes an adverse effect on sensitizing bladder afferent [46]. Besides that, oxidative stress leads to the overactive bladder through mediating capsaicin-sensitive C-fibers [47].

Plants and their compounds exert good antioxidant effects due to phenolic compounds and vitamins E and C; therefore, they can control oxidative stress and help treat nocturia. Although plants and their derivatives can effectively decrease LUTS, some studies have reported that monotherapy is not effective in decreasing the symptoms especially nocturia [48-50]. In addition, it is necessary to use a combination of the plants or their compounds to treat the disease so that more optimal results can be achieved [51].

Accordingly, as with routine treatments for these two disorders, phytotherapies should be used as complementary therapies to achieve better outcomes. In addition, some plants can exert diuretic effects. Prescribing these plants for the people with this disorder should be taken into account to avoid an increase in the associated complications [52].

Regarding nocturnal enuresis in children, there is convincing evidence on the effects of herbal drugs in treating this disorder [53]. Therefore, finely-designed clinical trials on their active compounds and with large sample size should be conducted to supplement available evidence on this subject.

Nocturia is multifactorial disease and more common among women [54,55] but In the elderly nocturia is more frequent among men [54] and it should be noted that in the treatment with medicinal plants, the causes of nocturia and gender differences should be considered. For example, nocturnal polyuria and decreased nocturnal bladder capacity are the major causes of nocturia in women [56]. Testosterone deficiency and prostatic hyperplasia (BPH) are the leading cause of nocturia in men [57,58]. Hence, in the use of medicinal plants, must considering the gender factor and underlying disorders that cause nocturia,

4. CONCLUSION

Plants and plant-based compounds cannot be used alone to treat nocturia and should be used as a complementary therapy with routine treatments to produce best therapeutic effects. Many studies have been conducted on nocturia as a symptom of BPH but very few studies have investigated it independently. Further clinical

trials should be therefore conducted to obtain more reliable evidence.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Weiss JP. Nocturia: Focus on etiology and consequences. *Reviews in Urology*. 2012;14(3-4):48-55.
2. van Kerrebroeck P, Abrams P, Chaikin D, Donovan J, Fonda D, Jackson S, et al. The standardisation of terminology in nocturia: Report from the Standardisation Subcommittee of the International Continence Society. *Neurourology and Urodynamics*. 2002;21(2):179-83.
3. Tikkinen KA, Johnson TM, 2nd, Tammela TL, Sintonen H, Haukka J, Huhtala H, et al. Nocturia frequency, bother, and quality of life: how often is too often? A population-based study in Finland. *European Urology*. 2010;57(3):488-96.
4. Nakagawa H, Niu K, Hozawa A, Ikeda Y, Kaiho Y, Ohmori-Matsuda K, et al. Impact of nocturia on bone fracture and mortality in older individuals: A Japanese longitudinal cohort study. *The Journal of Urology*. 2010;184(4):1413-8.
5. Ancoli-Israel S, Bliwise DL, Nørgaard JP. The effect of nocturia on sleep. *Sleep Medicine Reviews*. 2011;15(2):91-7.
6. Ikehara S, Iso H, Date C, Kikuchi S, Watanabe Y, Wada Y, et al. Association of sleep duration with mortality from cardiovascular disease and other causes for Japanese men and women: The JACC study. *Sleep*. 2009;32(3):295-301.
7. Reeves P, Irwin D, Kelleher C, Milsom I, Kopp Z, Calvert N, et al. The current and future burden and cost of overactive bladder in five European countries. *European Urology*. 2006;50(5):1050-7.
8. Yazici CM, Kurt O. Combination therapies for the management of nocturia and its comorbidities. *Research and Reports in Urology*. 2015;7:57-63.
9. Park HK, Kim HG. Current evaluation and treatment of nocturia. *Korean Journal of Urology*. 2013;54(8):492-8.
10. DiBianco JM, Morley C, Al-Omar O. Nocturnal enuresis: A topic review and institution experience. *Avicenna Journal of Medicine*. 2014;4(4):77-86.
11. Sinha R, Raut S. Management of nocturnal enuresis - myths and facts. *World Journal of Nephrology*. 2016;5(4):328-38.
12. Shabanian S, Khalili S, Lorigooini Z, Malekpour A, Heidari-Soureshjani S. The effect of vaginal cream containing ginger in users of clotrimazole vaginal cream on vaginal candidiasis. *Journal of Advanced Pharmaceutical Technology & Research*. 2017;8(2):80-4.
13. Yavangi M. Use of Iranian medicinal plants effective on male fertility indices. *Journal of Global Pharma Technology*. 2017;8(10):36-43.
14. Yavangi M. A systematic review of Iranian medicinal plants effective on female infertility. *Journal of Global Pharma Technology*. 2017;8(10):44-9.
15. Nikfarjam M, Bahmani M, Heidari-Soureshjani S. Phytotherapy for anxiety in Iran: A review of the most important Anti-anxiety medicinal plants.
16. Nikfarjam M, Bahmani M, Heidari-Soureshjani S. Phytotherapy for depression: A review of the most important medicinal plants of flora of Iran effective on depression.
17. Heidari-Soureshjani S, Asadi-Samani M, Yang Q, Saeedi-Boroujeni A. Phytotherapy of nephrotoxicity-induced by cancer drugs: an updated review. *J Nephropathol*. 2017;6(3):254-63.
18. Asadi-Samani M, Raeisi R, Heidari-Soureshjani S, Asadi-Samani M. A review for discovering hepatoprotective herbal drugs with least side effects on kidney. *Journal of Nephrotherapeutics*. 2017;6(2):38-48.
19. Shirani-Boroujeni M, Heidari-Soureshjani S, Keivani Hafshejani Z. Impact of oral capsule of *Peganum harmala* on alleviating urinary symptoms in men with benign prostatic hyperplasia; a randomized clinical trial. *Journal of Renal Injury Prevention*. 2017;6(2):127-31.

20. Pagano E, Laudato M, Griffo M, Capasso R. Phytotherapy of benign prostatic hyperplasia. A Minireview. *Phytother Res.* 2014;28(7):949-55.
21. Lund TD, Blake C, Bu L, Hamaker AN, Lephart ED. Equol an isoflavonoid: Potential for improved prostate health, in vitro and in vivo evidence. *Reproductive Biology and Endocrinology: RB&E.* 2011; 9-4.
22. Brady CM, DasGupta R, Dalton C, Wiseman OJ, Berkley KJ, Fowler CJ. An open-label pilot study of cannabis-based extracts for bladder dysfunction in advanced multiple sclerosis. *Mult Scler J.* 2004;10(4):425-33.
23. Shirani-Boroujeni M, Heidari-Soureshjani S, Keivani Hafshejani Z. Impact of oral capsule of Peganum harmala on alleviating urinary symptoms in men with benign prostatic hyperplasia; a randomized clinical trial. *Journal of Renal Injury Prevention.* 2017;6(2):127-31.
24. Sigurdsson S, Geirsson G, Gudmundsdottir H, Egilsdottir PB, Gudbjarnason S. A parallel, randomized, double-blind, placebo-controlled study to investigate the effect of SagaPro on nocturia in men. *Scand J Urol.* 2013; 47(1):26-32.
25. Ogushi T, Takahashi S. Effect of Chinese herbal medicine on overactive bladder. *Hinyokika Kiyō Acta Urologica Japonica.* 2007;53(12):857-62.
26. Yagi H, Nishio K, Sato R, Arai G, Soh S, Okada H. Clinical efficacy and tolerability of Gosha-jinki-gan, a Japanese traditional herbal medicine, for nocturia. *Journal of Traditional and Complementary Medicine.* 2016;6(1):126-9.
27. Yoshimura K, Shimizu Y, Masui K, Okuno H, Ueda T, Soda T, et al. Furosemide versus Gosha-Jinki-Gan, a Blended Herbal Medicine, for Nocturnal Polyuria: A Randomized Crossover Trial. *Luts.* 2012; 4(2):77-81.
28. Pradeep EK, Umashankar KS, Hegde G. A clinical study on the efficacy of pataladi kashaya in the management of 'vatashteela' vis a vis 'bph' (benign prostatic hypertrophy). *International Journal of Research in Ayurveda and Pharmacy.* 2013;4(6):814-9.
29. Oelke M, Berges R, Schlafke S, Burkart M. Fixed-dose combination PRO 160/120 of sabal and urtica extracts improves nocturia in men with LUTS suggestive of BPH: Re-evaluation of four controlled clinical studies. *World J Urol.* 2014;32(5):1149-54.
30. Beiraghdar F, Einollahi B, Ghadyani A, Panahi Y, Hadjiakhoondi A, Vazirian M, et al. A two-week, double-blind, placebo-controlled trial of Viola odorata, Echimium amoenum and Physalis alkekengi mixture in symptomatic benign prostate hyperplasia (BPH) men. *Pharm Biol.* 2017;55(1):1800-5.
31. Nadeem HMR, Mohiuddin E, Shahab-ud-Din, Daniyal M, Usmanghani K. Comparison of Anti BPH capsule (herbal) and Terazosin HCl in the treatment of benign prostate hyperplasia. *Pak J Pharm Sci.* 2017;30(1):289-93.
32. MacDonald R, Ishani A, Rutks I, Wilt TJ. A systematic review of Cernilton for the treatment of benign prostatic hyperplasia. *Bju Int.* 2000;85(7):836-41.
33. Wilt T, Ishani A, Mac Donald R, Rutks I, Stark G. Pygeum africanum for benign prostatic hyperplasia. *The Cochrane Database of Systematic Reviews.* 2002(1):CD001044.
34. Ishani A, MacDonald R, Nelson D, Rutks I, Wilt TJ. Pygeum africanum for the treatment of patients with benign prostatic hyperplasia: A systematic review and quantitative meta-analysis. *Am J Med.* 2000;109(8):654-64.
35. MacDonald R, Ishani A, Rutks I, Wilt TJ. A systematic review of Cernilton for the treatment of benign prostatic hyperplasia. *Bju Int.* 2000;85(7):836-41.
36. Wilt T, Ishani A, Stark G, MacDonald R, Mulrow C, Lau J. Serenoa repens for benign prostatic hyperplasia. *The Cochrane Database of Systematic Reviews.* 2000(2):CD001423.
37. Wilt TJ, Ishani A, Rutks I, MacDonald R. Phytotherapy for benign prostatic hyperplasia. *Public Health Nutrition.* 2000;3(4A):459-72.
38. Wilt T, Ishani A, Mac Donald R. Serenoa repens for benign prostatic hyperplasia. *The Cochrane Database of Systematic Reviews.* 2002;3:CD001423.
39. Lowe FC. The Role of Serenoa repens in the Clinical Management of Lower Urinary Tract Symptoms Due to Benign Prostatic Hyperplasia. *European Urology, Supplements.* 2009;8(13):894-7.
40. Geavlete P, Multescu R, Geavlete B. Serenoa repens extract in the treatment of

- benign prostatic hyperplasia. *Therapeutic Advances in Urology*. 2011;3(4):193-8.
41. Boyle P, Robertson C, Lowe F, Roehrborn C. Updated meta-analysis of clinical trials of *Serenoa repens* extract in the treatment of symptomatic benign prostatic hyperplasia. *Bju Int*. 2004;93(6):751-6.
 42. Wilt T, Mac Donald R, Ishani A, Rutks I, Stark G. Cernilton for benign prostatic hyperplasia. *The Cochrane Database of Systematic Reviews*. 2000;2:CD001042.
 43. Wilt TJ, Macdonald R, Ishani A, Rutks I, Stark G. WITHDRAWN: Cernilton for benign prostatic hyperplasia. *The Cochrane Database of Systematic Reviews*. 2011;5:CD001042.
 44. Wilt T, Ishani A, MacDonald R, Stark G, Mulrow C, Lau J. Beta-sitosterols for benign prostatic hyperplasia. *The Cochrane Database of Systematic Reviews*. 2000;2:CD001043.
 45. Yamada K, Suzuki E, Nakaki T, Watanabe S, Kanba S. *Aconiti tuber* increases plasma nitrite and nitrate levels in humans. *Journal of Ethnopharmacology*. 2005;96(1-2):165-9.
 46. Aikawa K, Leggett R, Levin RM. Effect of age on hydrogen peroxide mediated contraction damage in the male rat bladder. *The Journal of Urology*. 2003;170(5):2082-5.
 47. Masuda H, Kihara K, Saito K, Matsuoka Y, Yoshida S, Chancellor MB, et al. Reactive oxygen species mediate detrusor overactivity via sensitization of afferent pathway in the bladder of anaesthetized rats. *Bju Int*. 2008;101(6):775-80.
 48. MacDonald R, Tacklind JW, Rutks I, Wilt TJ. *Serenoa repens* monotherapy for benign prostatic hyperplasia (BPH): An updated Cochrane systematic review. *Bju Int*. 2012;109(12):1756-61.
 49. Tacklind J, MacDonald R, Rutks I, Stanke JU, Wilt TJ. *Serenoa repens* for benign prostatic hyperplasia. *Cochrane Db Syst Rev*. 2012;12.
 50. Barry MJ, Meleth S, Lee JY, Kreder KJ, Avins AL, Nickel JC, et al. Effect of increasing doses of saw palmetto extract on lower urinary tract symptoms a randomized trial. *Jama-J Am Med Assoc*. 2011;306(12):1344-51.
 51. Russo A, Capogrosso P, La Croce G, Ventimiglia E, Boeri L, Briganti A, et al. *Serenoa repens*, selenium and lycopene to manage lower urinary tract symptoms suggestive for benign prostatic hyperplasia. *Expert Opin Drug Saf*. 2016;15(12):1661-70.
 52. Clare BA, Conroy RS, Spelman K. The diuretic effect in human subjects of an extract of *taraxacum officinale folium* over a single day. *J Altern Complem Med*. 2009;15(8):929-34.
 53. Huang T, Shu X, Huang YS, Cheuk DK. Complementary and miscellaneous interventions for nocturnal enuresis in children. *The Cochrane Database of Systematic Reviews*. 2011;12:CD005230.
 54. Tikkinen KA, Tammela TL, Huhtala H, Auvinen A. Is nocturia equally common among men and women? A population based study in Finland. *J Urol*. 2006;175(2):596-600.
 55. Wein A, Lose GR, Fonda D. Nocturia in men, women and the elderly: A practical approach. *BJU Int*. 2002;90(Suppl3):28-31.
 56. Kim SO, Kim JS, Kim HS, Hwang EC, Oh KJ, Kwon D, et al. Age related change of nocturia in women. *International Neurourology Journal*. 2010;14(4):245-249.
 57. Jin MH, Moon DG. Practical management of nocturia in urology. *Indian Journal of Urology: IJU: Journal of the Urological Society of India*. 2008;24(3):289-294.
 58. Shigehara K, Izumi K, Mizokami A, Namiki M. Testosterone deficiency and nocturia: A review. *The World Journal of Men's Health*. 2017;35(1):14-21.

© 2018 Yavangi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/23725>