



Title	Bioaccessibility and bioavailability of a marine-derived multimineral, Aquamin-Magnesium
Author(s)	Felice, Valeria D.; O'Gorman, Denise M.; O'Brien, Nora M.; Hyland, Niall P.
Publication date	2018
Original citation	Felice, V., O'Gorman, D., O'Brien, N. and Hyland, N. (2018) 'Bioaccessibility and Bioavailability of a Marine-Derived Multimineral, Aquamin-Magnesium', <i>Nutrients</i> , 10(7), 912 (8pp). doi: 10.3390/nu10070912
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.mdpi.com/2072-6643/10/7/912 http://dx.doi.org/10.3390/nu10070912 Access to the full text of the published version may require a subscription.
Rights	© 2018, the Authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/). http://creativecommons.org/licenses/by/4.0/
Item downloaded from	http://hdl.handle.net/10468/6935

Downloaded on 2018-09-30T19:29:52Z



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Communication

Bioaccessibility and Bioavailability of a Marine-Derived Multimineral, Aquamin-Magnesium

Valeria D. Felice ^{1,2,3,*}, Denise M. O’Gorman ⁴, Nora M. O’Brien ² and Niall P. Hyland ^{1,3}

¹ Department of Pharmacology and Therapeutics and Department of Physiology, University College Cork, T12 K8AF Cork, Ireland; n.hyland@ucc.ie

² School of Food and Nutritional Sciences, University College Cork, T12 K8AF Cork, Ireland; nob@ucc.ie

³ APC Microbiome Ireland, University College Cork, T12 K8AF Cork, Ireland

⁴ Marigot Ltd., P43 NN62 Cork, Ireland; denise.ogorman@marigot.ie

* Correspondence: valeria.felice@marigot.ie; Tel.: +353-(0)21-420-5876

Received: 31 May 2018; Accepted: 11 July 2018; Published: 17 July 2018



Abstract: Introduction: Magnesium is an essential mineral involved in a range of key biochemical pathways. Several magnesium supplements are present on the market and their degree of bioavailability differs depending on the form of magnesium salt used. Aquamin-Mg is a natural source of magnesium, containing 72 additional trace minerals derived from the clean waters off the Irish coast. However, the *in vitro* bioaccessibility and bioavailability of Aquamin-Mg in comparison with other supplement sources of magnesium has yet to be tested. **Method:** Aquamin-Mg, magnesium chloride (MgCl₂) and magnesium oxide (MgO) were subjected to gastrointestinal digestion according to the harmonized INFOGEST *in vitro* digestion method and *in vitro* bioavailability tested using the Caco-2 cell model. Magnesium concentration was measured by atomic absorption spectrophotometry (AAS). **Results:** Magnesium recovery from both Aquamin-Mg and MgCl₂ was greater than for MgO. Magnesium from all three sources was transported across the epithelial monolayer with Aquamin-Mg displaying a comparable profile to the more bioavailable MgCl₂. **Conclusions:** Our data support that magnesium derived from a marine-derived multimineral product is bioavailable to a significantly greater degree than MgO and displays a similar profile to the more bioavailable MgCl₂ and may offer additional health benefits given its multimineral profile.

Keywords: Aquamin; multimineral supplement; magnesium bioavailability

1. Introduction

Magnesium is an essential mineral for the human body and is involved in a wide range of crucial physiological processes [1]. Magnesium can be obtained from the diet, being naturally present in foods such as green leafy vegetables, seeds, beans, whole grains, fish and nuts, amongst others. However, dietary magnesium intake has been shown to be insufficient in the Western population due to industrial food processing that reduces the nutrient contents including magnesium, as well as changes in dietary habits [2]. Deficiency in magnesium dietary intake may lead to hypomagnesemia which has been associated with several disorders including diabetes, osteoporosis and cardiovascular disease [3–5]. Early symptoms of hypomagnesemia are non-specific and include loss of appetite, nausea, vomiting, lethargy, fatigue and weakness with more pronounced hypomagnesemia characterised by increased neuromuscular excitability including muscle cramps, tremor, tetany and generalized seizures [6].

The market currently offers various supplement preparations to reach the recommended magnesium daily intake. These supplements differ in the type of magnesium salt used which can be either organic (i.e., magnesium citrate and magnesium aspartate) or inorganic (i.e., MgO and MgCl₂), their dosage and bioavailability. For example, magnesium from MgCl₂ has high bioavailability equivalent to organic

magnesium supplements such as magnesium lactate and aspartate [7]. Moreover, these three sources of magnesium have significantly greater bioavailability than MgO [7]. Magnesium derived from magnesium hydroxide (Mg(OH)₂) (Mablet) has been shown to be absorbed into the circulation and, hence, bioavailable in healthy male adults [8]. In a previous study, the bioavailability of magnesium from formulations containing different combinations of magnesium salts displayed similar bioavailability, however the daily dose of magnesium differed [9].

Aquamin-Mg is a natural source of magnesium in the form of Mg(OH)₂ derived from the clean waters off the Irish coast. In addition to magnesium, Aquamin-Mg also contains 72 additional trace minerals (Marigot Ltd., Cork, Ireland, Table 1) with the same profile of Lithothamnion Aquamin which has previously been shown to be a highly bioavailable source of calcium [10].

Here we describe for the first time the *in vitro* bioaccessibility and bioavailability of Aquamin-Mg in comparison with two commercially available sources of magnesium, MgCl₂ and MgO.

Table 1. Mineral composition of Aquamin-Mg. (ppm, parts per million).

Mineral	ppm	Mineral	ppm
Aluminum	461	Molybdenum	1.78
Antimony	<0.5	Neodymium	2.810
Arsenic	1.811	Nickel	<0.5
Barium	1.83	Niobium	<0.5
Beryllium	<0.5	Osmium	0.002
Bismuth	<0.5	Palladium	0.301
Boron	186	Phosphorus	117
Cadmium	0.394	Platinum	0.002
Calcium	23,000	Potassium	88.21
Carbon	10,100	Praseodymium	0.697
Cerium	3.411	Rhenium	0.001
Cesium	0.008	Rhodium	0.010
Chloride	613.4	Rubidium	0.063
Chromium	5.83	Ruthenium	0.135
Cobalt	<0.5	Samarium	0.576
Copper	5.09	Scandium	1.050
Dysprosium	0.829	Selenium	<0.5
Erbium	0.613	Silicon	657
Europium	0.197	Silver	<0.5
Fluoride	1.1	Sodium	1467
Gadolinium	0.770	Strontium	84.7
Gallium	0.163	Sulfur	3335
Germanium	0.020	Tantalum	0.016
Gold	<0.5	Tellurium	<0.5
Hafnium	0.046	Terbium	0.140
Holmium	0.194	Thallium	<0.5
Indium	<0.001	Thorium	0.860
Iodine	9.1	Thulium	0.081
Iridium	0.002	Tin	0.179
Iron	1213	Titanium	18.5
Lanthanum	1.01	Tungsten	2.08
Lead	0.604	Vanadium	16.0
Lithium	<0.5	Ytterbium	0.498
Lutetium	0.116	Yttrium	7.38
Magnesium	404,400	Zinc	2.37
Manganese	486	Zirconium	<0.5
Mercury	0.009		

2. Methods

2.1. Harmonized INFOGEST *In Vitro* Digestion Protocol

To determine bioaccessibility, the amount of magnesium subjected to digestion for each compound was calculated according to the recommended dietary allowance for men (RDA, 420 mg/day). 5.6 mg of magnesium from Aquamin-Mg, MgCl₂ and MgO were digested according to the harmonized INFOGEST *in vitro* digestion method published by Minekus and colleagues [11]. Four to five independent digestions were carried out for each compound (Aquamin-Mg, MgCl₂ and MgO). Aquamin-Mg, MgCl₂ and MgO were exposed to simulated gastric fluid (composition: 6.9 mM KCl, 0.9 mM KH₂PO₄, 25 mM NaHCO₃, 47.2 mM NaCl, 0.1 mM MgCl₂(H₂O)₆, 0.5 mM (NH₄)₂CO₃). Pepsin and calcium chloride were added to the mixture to achieve a final concentration of 2000 U/mL and 0.075 mM respectively. Hydrochloric acid (HCl, 6 M) was then used to acidify the mixture to pH 3 and water was added to reach a final volume of 20 mL. Samples were then incubated in a stirring water bath at 37 °C and 95 rpm for 2 h. The pH was checked after 1 hour and adjusted if necessary. The simulated intestinal fluid (composition: 6.8 mM KCl, 0.8 mM KH₂PO₄, 85 mM NaHCO₃, 38.4 mM NaCl, 0.33 mM MgCl₂(H₂O)₆) was then added together with pancreatin (the concentration was based on trypsin activity, 100 U/mL) and bile salts for a final concentration of 10 mM. Calcium chloride was also added to achieve a final concentration of 0.3 mM. Sodium hydroxide (NaOH, 1 M) was used to bring the pH to 7 and the necessary amount of water added to reach a final volume of 20 mL. Samples were incubated in a stirring water bath at 37 °C and 95 rpm for 2 h. The pH was checked after one hour and adjusted if necessary. A control sample containing all reagents included in the digestion protocol except the experimental powders was also subjected to the procedure. Upon completion of the incubation period, aliquots (1 mL) of each sample were frozen in liquid nitrogen. Prior to the analysis, one sample from each treatment was filtered using 0.2 µm cell culture sterile filters. The amount of magnesium recovered from these samples was then compared to non-filtered samples.

2.2. Caco-2 Cell Bioavailability Assay

Caco-2 cells are human epithelial colorectal adenocarcinoma cells that, upon differentiation, express numerous morphological and biochemical characteristics of small intestinal enterocytes. This *in vitro* model is widely used to study mineral bioavailability from different sources and their transport mechanisms [12].

For Caco-2 bioavailability experiments 60 mg of magnesium derived from Aquamin-Mg, MgCl₂ and MgO were subjected to the harmonized INFOGEST *in vitro* digestion protocol described above (data not shown), and unfiltered samples were used. 60 mg of magnesium was chosen in order to ensure that sufficient concentrations of magnesium could be achieved to perform the Caco-2 experiments. Three independent digestions were carried out for each compound (Aquamin-Mg, MgCl₂ and MgO) and these were subsequently used in the Caco-2 bioavailability assay. Caco-2 cells (supplied by the European Collection of Authenticated Cell Cultures (ECACC)) were cultured in Dulbecco's modified eagle's medium (DMEM) supplemented with 1% non-essential amino acids and 10% foetal bovine serum (FBS) and were stored in a humidified incubator at 37 °C and 5% CO₂. For all experiments, cells were seeded at a density of 1 × 10⁵ cells/mL on 6-well Transwell plates with inserts of 24 mm diameter and differentiated for 21 days. Media was changed every other day.

2.3. Bioavailability of Magnesium from Aquamin-Mg, MgCl₂ and MgO Using Caco-2 Cells

On the day of the experiment, media was removed from all wells and 1 mL of fresh media was added to the luminal side and 2 mL to the basolateral side of each well. Transepithelial electrical resistance (TEER) was measured to confirm the integrity of the epithelial monolayer. Cells were then incubated with either Aquamin-Mg, MgCl₂- or MgO-derived magnesium in the luminal side (concentrations of 25, 50, 100 and 150 µg/mL) for 2 h at 37 °C. Samples from each independent digestion were used in a corresponding independent bioavailability study which was conducted in

triplicate. Two controls were included in the assay, a blank sample, only containing media, and a digest sample, containing all reagents included in the digestion protocol except the experimental powders. At the end of the incubation time, TEER values were recorded again to ensure that the treatments did not have any effect on the integrity of the monolayer. Luminal and basolateral samples were collected and stored at 4 °C. Magnesium concentration of luminal and basolateral samples was measured by atomic absorption spectrophotometry (AAS). Three independent experiments were carried out. Each treatment was randomly assigned and was performed in duplicate.

2.4. Atomic Absorption Spectrophotometry (AAS)

The magnesium content of the digested samples, as well as luminal and basolateral samples was determined by AAS. Samples were diluted in Milli-Q water prior to analysis. Lanthanum chloride (0.1%) was also added to eliminate any phosphate interferences. A commercially available magnesium standard (Spectrosol from BDH Chemicals Ltd., Dublin, Ireland) was used. Standard solutions were prepared using Milli-Q water containing lanthanum chloride (0.1%) and ranged from 0 to 1 mg/L.

2.5. Statistical Analyses

Data are expressed as mean \pm standard error of the mean (SEM). Statistical analysis was carried out using the Kruskal–Wallis test, followed by Dunn’s multiple comparison test for the digestion study. For the bioavailability study, we calculated the residuals of the data to determine whether there were outliers and statistical analysis was performed using one-way analysis of variance (ANOVA) followed by Bonferroni post-hoc test. Values of $p < 0.05$ were considered statistically significant.

3. Results

3.1. Magnesium Recovery from Aquamin-Mg, MgCl₂ and MgO Following In Vitro Digestion

Aquamin-Mg-derived magnesium, showed an in vitro bioaccessibility more similar to highly soluble alternative such as MgCl₂ than to MgO (Aquamin-Mg, 122.6 ± 4.1 $\mu\text{g}/\text{mL}$, $n = 5$; MgCl₂, 115.4 ± 6.0 $\mu\text{g}/\text{mL}$, $n = 4$; MgO, 73.39 ± 10.20 $\mu\text{g}/\text{mL}$, $n = 4$; in unfiltered samples), which is characterised by low solubility (Figure 1, $p < 0.05$ Aquamin-Mg vs. MgO). To determine whether magnesium was lost during filtration for the Caco-2 cell culture experiments, magnesium concentration in filtered samples was also determined and these were lower for all the treatments relative to unfiltered samples (Aquamin-Mg, 99.3 ± 6.2 $\mu\text{g}/\text{mL}$, $n = 5$; MgCl₂, 92.2 ± 5.6 $\mu\text{g}/\text{mL}$, $n = 4$; MgO, 64.5 ± 9.3 $\mu\text{g}/\text{mL}$ in filtered samples).




Figure 1. Magnesium recovery from Aquamin-Mg, MgCl₂ and MgO following in vitro digestion. The percentage of magnesium recovery from Aquamin-Mg was significantly higher than MgO (* $p < 0.5$, $n = 4-5$) in unfiltered samples.

3.2. Transepithelial Electrical Resistance (TEER)

As an indicator of cell viability we measured TEER at the start of each experiment and after treatment. The results from the transepithelial resistance measurements confirmed that after 21 days of culture, Caco-2 cells formed an integral monolayer. Moreover, none of the treatments, at any concentration tested, affected epithelial integrity following 2 h incubation at 37 °C (Figure 2).




Figure 2. Transepithelial electrical resistance in Caco-2 cells following treatment with Aquamin-Mg, MgCl₂ and MgO. At time zero (T0) cells were differentiated in an integral monolayer. Transepithelial electrical resistance (TEER) values confirmed that the treatments did not compromise the integrity of the monolayer at any of the concentrations tested following 2 h incubation (T2h) (*n* = 3).

3.3. Bioavailability of Magnesium from Aquamin-Mg, MgCl₂ and MgO Using the Caco-2 Cell Model

Digestates were added to the luminal compartment. The reduced amount of magnesium from MgO is reflective of the reduced bioaccessibility (Figure 3a). These results are in accordance with the in vitro digestion data (60 mg; data not shown). When the concentration of magnesium was measured in the basolateral chamber, magnesium derived from Aquamin-Mg and MgCl₂ showed the same degree of bioavailability at all the concentrations tested, while only a small amount of magnesium from MgO was transported across the epithelium and thus bioavailable. Both Aquamin-Mg and MgCl₂ were significantly more bioavailable than MgO at the highest concentration tested (150 μg/mL) (Figure 3b, Table 2).




Figure 3. Bioavailability of magnesium from Aquamin-Mg, MgCl₂ and MgO using Caco-2 cells. (a) Magnesium derived from Aquamin-Mg, MgCl₂ and MgO applied into the luminal side. Following in vitro digestion, the amount of bioaccessible magnesium was significantly higher for Aquamin-Mg and MgCl₂ compared to MgO (***p* < 0.001 for Aquamin-Mg vs. MgO at 25 $\mu\text{g/ml}$, *****p* < 0.0001 for MgCl₂ vs. MgO at 25 $\mu\text{g/ml}$ and *****p* < 0.0001 for Aquamin-Mg and MgCl₂ vs. MgO at 50, 100 and 150 $\mu\text{g/ml}$, *n* = 3); (b) Magnesium derived from Aquamin-Mg, MgCl₂ and MgO measured in the basolateral side after 2 h incubation at 37 °C. At the highest concentration tested, magnesium from Aquamin-Mg and MgCl₂ was significantly more bioavailable than MgO (***p* < 0.01 for Aquamin-Mg and MgCl₂ vs. MgO at 150 $\mu\text{g/ml}$, *n* = 3).

Table 2. Magnesium derived from Aquamin-Mg, MgCl₂ and MgO measured in the basolateral side after 2 h incubation at 37 °C expressed as $\mu\text{g/ml}$ (*n* = 3).

Magnesium Source	Sample Concentration ($\mu\text{g/ml}$)	Basolateral Side $\mu\text{g/ml}$ Mean \pm Standard Error of the Mean (SEM)
Aquamin-Mg	25	0.37 \pm 0.17
	50	0.44 \pm 0.16
	100	0.69 \pm 0.16
	150	0.89 \pm 0.17
MgCl ₂	25	0.14 \pm 0.06
	50	0.33 \pm 0.13
	100	0.51 \pm 0.09
	150	0.86 \pm 0.13
MgO	25	0.05 \pm 0.03
	50	0.08 \pm 0.02
	100	0.25 \pm 0.01
	150	0.10 \pm 0.05

4. Discussion

The aim of these studies was to examine the bioaccessibility and bioavailability of magnesium from Aquamin-Mg compared to $MgCl_2$ and MgO using the Caco-2 cell model. In this model both active saturated and passive non-saturated pathways have been previously identified for magnesium transport [13]. The study from Thongon and Krishnamrain has indeed shown that in Caco-2 monolayers, magnesium transported from the apical to the basolateral side (representing magnesium absorption) against magnesium in the apical solution (representing magnesium concentration) was curvilinear as previously shown in humans [13,14]. Furthermore, the same study has shown that treatment with omeprazole selectively inhibited the non-saturable passive component, without affecting the saturable active component of magnesium transport which was abolished using the Transient Receptor Potential Cation Channel Subfamily M Member 6 (TRPM6) inhibitor Ruthenium Red (RR) [13]. This evidence shows that magnesium can be transported through both a paracellular and a transcellular pathway and that the Caco-2 monolayer is a suitable *in vitro* model of intestinal magnesium absorption. In the context of our findings, however, we cannot comment on which pathway was responsible for the apical to basolateral transport of magnesium and further research is warranted in order to elucidate these mechanisms. Our results show for the first-time, however, direct evidence that Aquamin-Mg-derived magnesium is highly bioaccessible following *in vitro* digestion and magnesium is transported across the intestinal epithelium in this well-established *in vitro* model. Moreover, the degree of bioaccessibility and bioavailability of Aquamin-Mg was comparable to $MgCl_2$ while being superior to MgO .

$MgCl_2$ and MgO represent a high bioavailable and low bioavailable source of magnesium respectively, and our *in vitro* data are in keeping with *in vivo* data demonstrating that the mean urinary excretion of magnesium in healthy volunteers was significantly higher for $MgCl_2$ than MgO [7]. Interestingly, in this study, $MgCl_2$ bioavailability was comparable to that of organic magnesium forms such as magnesium aspartate and magnesium lactate [7].

Magnesium in Aquamin-Mg is in the form of $Mg(OH)_2$. However, as well as magnesium, Aquamin-Mg also provides 72 additional trace minerals all derived from sea water (Marigot Ltd., Cork, Ireland, Table 1). In support of $Mg(OH)_2$ as a magnesium supplement, the pharmacokinetic profile of a single oral dose of $Mg(OH)_2$ in healthy male adults showed that the bioavailability of magnesium from $Mg(OH)_2$ was 15% [8]. Moreover, none of the participants recruited reported any side effect following supplementation suggesting that $Mg(OH)_2$ may be a clinically relevant option for oral magnesium supplementation [8]. In a second human study the degree of bioavailability of $Mg(OH)_2$ was compared to other sources of magnesium, including $MgCl_2$ measured as urinary elimination of magnesium [9]. In this study it was found that $Mg(OH)_2$ was required at a higher dose to reach the same level of bioavailability [9].

The solubility of magnesium in the gastrointestinal tract plays a key role in magnesium absorption. Our bioaccessibility results demonstrate that magnesium from Aquamin-Mg is soluble as $MgCl_2$ and hence potentially available for absorption. Our bioavailability data support that $Mg(OH)_2$, derived from Aquamin-Mg, displays a similar profile and transport characteristics as magnesium derived from $MgCl_2$ at the same concentrations, suggesting that Aquamin-Mg represents a source of magnesium coupled with potential health benefits of a multimineral supplement. Currently, however, as Aquamin-Mg is not formulated as an oral supplement (tablets and capsules), further comparisons with other formulated magnesium supplements were not possible.

Moreover, Aquamin-Mg is composed of multiple minerals and whether these affect its bioaccessibility or bioavailability is difficult to determine.

5. Conclusions

In conclusion, our data suggests that Aquamin-Mg-derived magnesium is bioaccessible and bioavailable to a significantly greater degree than magnesium oxide while displaying a comparable profile to magnesium chloride. Nonetheless, our *in vitro* results are qualitatively consistent with the clinical study from Firoz and Graber showing that magnesium from $MgCl_2$ has significantly

greater bioavailability than MgO. Further research is warranted to investigate the bioaccessibility and bioavailability of Aquamin-Mg in clinical studies.

Author Contributions: V.D.F., N.P.H., N.M.O. and D.M.O. conceived and designed the experiments. D.M.O. provided Aquamin-Mg. V.D.F. performed the experiments and V.D.F. and N.P.H. analysed the data. V.D.F. and N.P.H. wrote the manuscript; N.M.O., and D.M.O., edited the manuscript.

Funding: This research was funded by the Irish Research Council Enterprise Partnership Scheme (EPS) Postdoctoral Fellowship, grant number EPSPD/2015/52. Marigot Ltd. provided the mineral-rich algae extract Aquamin and their financial support. APC Microbiome Ireland is funded by SFI, through the Irish Government's National Development Plan (Grant 12/RC/2273).

Acknowledgments: Felice has received an Irish Research Council Enterprise Postdoctoral Fellowship. We would like also to acknowledge Marigot Ltd. who provided the mineral-rich algae extract Aquamin and their financial support. Thank you also to O'Callaghan, University College Cork, and Shane O'Connell (Marigot Ltd.), for their help in the preparation of this manuscript.

Conflicts of Interest: Felice and O'Gorman are employees of Marigot and Hyland and O'Brien have received research support from Marigot Ltd.

References

1. Saris, N.E.; Mervaala, E.; Karppanen, H.; Khawaja, J.A.; Lewenstam, A. Magnesium. An update on physiological, clinical and analytical aspects. *Clin. Chim. Acta* **2000**, *294*, 1–26. [[CrossRef](#)]
2. Marier, J.R. Magnesium content of the food supply in the modern-day world. *Magnesium* **1986**, *5*, 1–8. [[PubMed](#)]
3. Rosique-Esteban, N.; Guasch-Ferré, M.; Hernández-Alonso, P.; Salas-Salvadó, J. Dietary magnesium and cardiovascular disease: A review with emphasis in epidemiological studies. *Nutrients* **2018**, *10*, 168. [[CrossRef](#)] [[PubMed](#)]
4. Guerrero-Romero, F.; Jaquez-Chairez, F.O.; Rodríguez-Morán, M. Magnesium in metabolic syndrome: A review based on randomized, double-blind clinical trials. *Magnes. Res.* **2016**, *29*, 146–153. [[PubMed](#)]
5. Castiglioni, S.; Cazzaniga, A.; Albisetti, W.; Maier, J.A.M. Magnesium and osteoporosis: Current state of knowledge and future research directions. *Nutrients* **2013**, *5*, 3022–3033.
6. Grober, U.; Schmidt, J.; Kisters, K. Magnesium in prevention and therapy. *Nutrients* **2015**, *7*, 8199–8226. [[CrossRef](#)] [[PubMed](#)]
7. Firoz, M.; Graber, M. Bioavailability of US commercial magnesium preparations. *Magnes. Res.* **2001**, *14*, 257–262. [[PubMed](#)]
8. Dolberg, M.K.B.; Nielsen, L.P.; Dahl, R. Pharmacokinetic profile of oral magnesium hydroxide. *Basic Clin. Pharmacol. Toxicol.* **2017**, *120*, 264–269. [[CrossRef](#)] [[PubMed](#)]
9. Bøhmer, T.; Røseth, A.; Holm, H.; Weberg-Teigen, S.; Wahl, L. Bioavailability of oral magnesium supplementation in female students evaluated from elimination of magnesium in 24-hour urine. *Magnes. Trace Elem.* **1990**, *9*, 272–278.
10. Zenk, J.L.; Frestedt, J.L.; Kuskowski, M.A. Effect of Calcium Derived from Lithothamnion sp. on markers of calcium metabolism in premenopausal women. *J. Med. Food* **2018**, *21*, 154–158. [[CrossRef](#)] [[PubMed](#)]
11. Minekus, M.; Alvinger, M.; Alvito, P.; Balance, S.; Bohn, T.; Bourlieu, C.; Carrière, F.; Boutrou, R.; Corredig, M.; Dupont, D.; et al. A standardised static in vitro digestion method suitable for food—An international consensus. *Food Funct.* **2014**, *5*, 1113–1124. [[CrossRef](#)]
12. Glahn, R. The use of Caco-2 cells in defining nutrient bioavailability: Application to iron bioavailability of foods. In *Designing Functional Foods*; McClements, D.J., Decker, E.A., Eds.; Woodhead Publishing: Cambridge, UK, 2009; pp. 340–361.
13. Thongon, N.; Krishnamra, N. Omeprazole decreases magnesium transport across Caco-2 monolayers. *World J. Gastroenterol.* **2011**, *17*, 1574–1583. [[CrossRef](#)] [[PubMed](#)]
14. Fine, K.D.; Ana, C.A.S.; Porter, J.L.; Fordtran, J.S. Intestinal absorption of magnesium from food and supplements. *J. Clin. Investig.* **1991**, *88*, 396–402. [[CrossRef](#)] [[PubMed](#)]

