



This article is published as part of a themed issue of ***Photochemical & Photobiological Sciences*** in honour of

[Esther Oliveros](#)

Guest edited by **Marie-Thérèse Maurette** and **Guillermo Orellana**

Published in [issue 7, 2009](#)

Other articles in this issue include:

---

[Sensitized formation of oxidatively generated damage to cellular DNA by UVA radiation](#)

J. Cadet, T. Douki, J-L. Ravanat and P. Di Mascio, *Photochem. Photobiol. Sci.*, 2009, **8**, 903

[Physical and chemical quenching rates and their influence on stereoselective photooxygenation of oxazolidinone-functionalized enecarbamates](#)

M. R. Solomon, J. Sivaguru, S. Jockusch, W. Adam and N. J. Turro, *Photochem. Photobiol. Sci.*, 2009, **8**, 912

[Water disinfection with Ru\(II\) photosensitisers supported on ionic porous silicones](#)

F. Manjón, D. García-Fresnadillo and G. Orellana, *Photochem. Photobiol. Sci.*, 2009, **8**, 926

[Photophysics and photochemistry of rose bengal bound to human serum albumin](#)

E. Alarcón, A. M. Edwards, A. Aspée, C. D. Borsarelli and E. A. Lissi, *Photochem. Photobiol. Sci.*, 2009, **8**, 933

[Photolysis of an asymmetrically substituted diazene in solution and in the crystalline state](#)

P. A. Hoijemberg, S. D. Karlen, C. N. Sanramé, P. F. Aramendía and M. A. García-Garibay, *Photochem. Photobiol. Sci.*, 2009, **8**, 961

[Heterogeneous photocatalytic degradation of gallic acid under different experimental conditions](#)

N. Quici and M. I. Litter, *Photochem. Photobiol. Sci.*, 2009, **8**, 975

[Photolysis of ferric ions in the presence of sulfate or chloride ions: implications for the photo-Fenton process](#)

A. Machulek Jr., J. Ermírio F. Moraes, L. T. Okano, C. A. Silvério and F. H. Quina, *Photochem. Photobiol. Sci.*, 2009, **8**, 985

[Light and dark biocidal activity of cationic poly\(arylene ethynylene\) conjugated polyelectrolytes](#)

T. S. Corbitt, L. Ding, E. Ji, L. K. Ista, K. Ogawa, G. P. Lopez, K. S. Schanze and D. G. Whitten, *Photochem. Photobiol. Sci.*, 2009, **8**, 998

[Optimisation of the chemical generation of singlet oxygen \( \$^1\text{O}\_2\$ ,  \$^1\Delta\_g\$ \) from the hydrogen peroxide-lanthanum\(III\) catalytic system using an improved NIR spectrometer](#)

C. Pierlot, J. Barbillat, V. Nardello-Rataj, D. Mathieu, M. Sergent, J. Marko and J-M. Aubry, *Photochem. Photobiol. Sci.*, 2009, **8**, 1024

[Photocatalytic efficiencies of self-cleaning glasses. Influence of physical factors](#)

L. Peruchon, E. Puzenat, J. M. Herrmann and C. Guillard, *Photochem. Photobiol. Sci.*, 2009, **8**, 1040

# Physical and chemical quenching rates and their influence on stereoselective photooxygenation of oxazolidinone-functionalized enecarbamates†‡

Marissa R. Solomon,<sup>a</sup> J. Sivaguru,<sup>\*b</sup> Steffen Jockusch,<sup>a</sup> Waldemar Adam<sup>\*c,d</sup> and Nicholas J. Turro<sup>\*\*a,e</sup>

Received 24th February 2009, Accepted 12th March 2009

First published as an Advance Article on the web 24th March 2009


DOI: 10.1039/b903848e

Physical and chemical quenching rate constants were measured for the reaction of singlet oxygen with oxazolidinone-functionalized enecarbamates to investigate the role of vibrational deactivation in product stereoselectivity.

Stereocontrol in photoreactions is problematic for photochemists, chiefly owing to the short lifetimes of reactive species and intermediates.<sup>1,2,3</sup> In an attempt to influence stereoselection, photochemists have employed a variety of confined media with varying degree of success.<sup>3–7</sup> In spite of some elegant methodologies that are employed to carry out asymmetric phototransformations,<sup>3–7</sup> excited-state stereocontrol in solution presents new opportunities and challenges to chemists.

Previous investigations in our laboratories have focused on the reaction of the enecarbamates **1–2** with singlet oxygen (<sup>1</sup>O<sub>2</sub>) to form the dioxetane **3**. The dioxetane subsequently decomposes to the ketone **4** (methyldeoxybenzoin, MDB) and the chiral aldehyde **5** (Scheme 1). In studying the stereoselectivity in the dioxetane product **3** we employed optically pure enecarbamates. The enantioselectivity in the MDB product **4** was obtained after utilizing an epimeric mixture (50/50 *R/S*-isomers at the C-3' position) of the starting enecarbamate.

In exploring this system, several remarkable features came to light with respect to the approach of <sup>1</sup>O<sub>2</sub> on to the enecarbamates **1**: (a) <sup>1</sup>O<sub>2</sub> reacts with the *Z*-enecarbamates (*Z*-**1**) to produce the dioxetane product **3** with diastereoselectivity values >95% (Scheme 1, top).<sup>8–10</sup> That is, >98% of the approaching <sup>1</sup>O<sub>2</sub> reacted with one face of the π bond to form one dioxetane product and <2% reacted with the other face to form the other dioxetane product; (b) selectivity in both dioxetane product **3**, and in the MDB product **4** is independent of the size of the C-4 alkyl group (Me, <sup>i</sup>Pr or <sup>t</sup>Bu) of the enecarbamates;<sup>11,12</sup> (c) the *R/S* configuration of the MDB product **4** is dependent on the C-4 *R/S* configuration of the enecarbamates;<sup>11</sup> (d) for the *Z*-enecarbamates, the selectivity in the dioxetane product **3**, is independent of the C-3'



**Scheme 1** Photooxidative cleavage of *Z/E* enecarbamates **1–2** with singlet oxygen (<sup>1</sup>O<sub>2</sub>).

*R/S* configuration of the enecarbamates;<sup>10,11,13</sup> (e) the *E*-isomer shows a solvent and temperature dependence of stereoselectivity in the MDB product **4**, in contrast to the corresponding *Z*-isomer.<sup>11,14,15</sup>

These results raised key questions about the role of the C-4 position and its ability to direct the approach of <sup>1</sup>O<sub>2</sub>—a small enophile that is blocked to the same extent by substituents of varying size.<sup>11,12</sup> Consequently, in view of these results, we hypothesized that factors apart from steric ones were playing a crucial role in the observed selectivity.<sup>11,13,14</sup>

To rationalize our observations we envisioned two distinct quenching processes during the photooxidation of enecarbamates **1–2** by <sup>1</sup>O<sub>2</sub>: the formation of dioxetane **3** (chemical quenching), and/or deactivation of <sup>1</sup>O<sub>2</sub> to <sup>3</sup>O<sub>2</sub>, the triplet ground state (physical quenching). The rate constant for the reaction leading to dioxetane **3** is called the chemical quenching rate constant (*k*<sub>cq</sub>) and the rate constant for deactivation (leading to <sup>3</sup>O<sub>2</sub>) is called physical quenching rate constant (*k*<sub>pq</sub>). The sum of both rate constants is *k*<sub>q</sub>, the rate constant for total quenching (eqn (1)).

$$k_q = k_{cq} + k_{pq} \quad (1)$$

<sup>1</sup>O<sub>2</sub> is known to be deactivated from its singlet excited state to its triplet ground state when it encounters C–H bonds.<sup>16</sup> For

<sup>a</sup>The Department of Chemistry, Columbia University, New York, NY 10027, USA

<sup>b</sup>Department of Chemistry and Molecular Biology, North Dakota State University, Fargo, ND 58105, USA. E-mail: sivaguru.jayaraman@ndsu.edu

<sup>c</sup>Institut für Organische Chemie, Universität Würzburg, D-97074, Würzburg, Germany. E-mail: wadam@chemie.uni-wuerzburg.de

<sup>d</sup>The Department of Chemistry, University of Puerto Rico, Facundo Bueso 110, Rio Piedras, PR 00931, USA

<sup>e</sup>The Department of Chemical Engineering, Columbia University, New York, NY 10027, USA. E-mail: njt3@columbia.edu

† This article was published as part of the themed issue in honour of Esther Oliveros.

‡ Electronic supplementary information (ESI) available: General procedures and details of rate constants determination. See DOI: 10.1039/b903848e

this reason, the lifetime of  $^1\text{O}_2$  is longer in deuterated solvents than in proteated solvents. This physical quenching phenomenon is referred to as vibrational deactivation of  $^1\text{O}_2$  since the coupling of the C–H vibration with the electronic excited state of molecular oxygen leads to relaxation of the singlet excited state. To better understand the role of vibrational deactivation in the observed high stereoselectivities, we performed a systematic study of the rate constants for both chemical ( $k_{\text{cq}}$ ) and physical ( $k_{\text{pq}}$ ) quenching. Because **1** and **2** are electron deficient alkenes, these quenching rate constants are expected to be relatively low and therefore challenging to measure accurately. In addition, sorting out the role of physical and chemical quenching adds another level of experimental difficulty. In spite of the difficulty of extracting accurate relative rate constants for the chemical portion of the quenching, we felt that a careful study of the total quenching would be important for exploring structure–reactivity relationships.

The chemical quenching rate constants ( $k_{\text{cq}}$ ) were measured using competitive kinetics. *trans*-4-Octene was utilized as a standard ( $k_{\text{cq}} = 4 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ ).<sup>17,18</sup> The relative ratios of the standard and enecarbamate peaks were monitored by  $^1\text{H-NMR}$  spectroscopy upon reaction with  $^1\text{O}_2$ .<sup>12</sup> After determining  $k_{\text{cq}}$  for **1a** (3'*R* and 3'*S*), these values were used as additional standards for subsequent measurements.<sup>12</sup>

Total quenching rate constants ( $k_{\text{q}}$ ) of  $^1\text{O}_2$  by enecarbamates were determined by using a chemiluminescence method, in which  $^1\text{O}_2$  was generated by thermal decomposition of 1,4-dimethylnaphthalene-1,4-endoperoxide.<sup>19</sup> Chemiluminescence (phosphorescence of  $^1\text{O}_2$ ) at 1270 nm was monitored to determine Stern–Volmer quenching constants by varying the quencher (enecarbamates **1** or **2**) concentration. To convert the Stern–Volmer quenching rate constants ( $K_{\text{SV}}$ ) into the total quenching rate constants ( $k_{\text{q}}$ ), the known total rate constant of *trans*-4-octene ( $k_{\text{q}} = 1.8 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ )<sup>17,18</sup> was used as standard to calculate  $\tau_0$ , the  $^1\text{O}_2$  lifetime under our experimental conditions (eqn (2)).<sup>12</sup> After determining  $k_{\text{q}}$  and  $k_{\text{cq}}$  for the enecarbamates experimentally,  $k_{\text{pq}}$  was derived from these two values using eqn (1).

$$K_{\text{SV}} = k_{\text{q}} \tau_0 \quad (2)$$

As we had employed a 50/50 epimeric mixture of enecarbamates at C-3' position, the relative rates of photooxygenation, known as

**Table 1** Conversion and stereoselectivity in the MDB product **4** upon photooxidative cleavage of enecarbamates with  $^1\text{O}_2$  in  $\text{CDCl}_3$  at  $15^\circ\text{C}^a$

Substrate	Configuration		$s^a$ [enhanced MDB enantiomer]
	C-4	C-3'	
<i>Z</i> ( <i>Me</i> )- <b>1a</b>	<i>R</i>	<i>R/S</i>	1.2 [ <i>R</i> ]
<i>Z</i> ( <i>Me</i> )- <b>1b</b>	<i>S</i>	<i>R/S</i>	1.5 [ <i>S</i> ]
<i>E</i> ( <i>Me</i> )- <b>1c</b>	<i>R</i>	<i>R/S</i>	9.2 [ <i>R</i> ]
<i>E</i> ( <i>Me</i> )- <b>1d</b>	<i>S</i>	<i>R/S</i>	10 [ <i>S</i> ]
<i>Z</i> ( <i>iPr</i> )- <b>2a</b>	<i>R</i>	<i>R/S</i>	2.2 [ <i>R</i> ]
<i>Z</i> ( <i>iPr</i> )- <b>2b</b>	<i>S</i>	<i>R/S</i>	1.4 [ <i>S</i> ]
<i>E</i> ( <i>iPr</i> )- <b>2c</b>	<i>R</i>	<i>R/S</i>	5.0 [ <i>R</i> ]
<i>E</i> ( <i>iPr</i> )- <b>2d</b>	<i>S</i>	<i>R/S</i>	3.4 [ <i>S</i> ]

<sup>a</sup> Data taken from previously published reports: ref. 11, 15, 20.

the stereoselectivity factor ( $s$  factor),<sup>21</sup> could be derived from the observed enantiomeric excess ( $ee$ ) values in **4** based on eqn (3). The stereoselectivity factor ( $s$ ), is a quantitative measure (corrected for the extent of conversion) of the relative reaction rates for the two stereoisomers in question (eqn (3)). In the present case, the  $s$  factor may be computed for the two-epimeric enecarbamates [C-3'(*R*) and C-3'(*S*) epimers] from the substrate conversion and the MDB enantiomeric excess by using eqn (3).<sup>22</sup> In the present case,  $k_1$  is the reactive rate constant leading to formation of the predominant MDB isomer and  $k_2$  the reactive rate constant leading to formation of the subordinate isomer. A large  $s$  value translates to high enantiomeric excess in the MDB product but corrected for conversion. From Table 1, it is evident that the  $s$  factor and consequently  $ee$  of the MDB products is higher in the *E*-enecarbamates than in the *Z*-enecarbamates. By exploring the chemical and total quenching rates independently, it would be of relevance to the relative rates of the individual epimers, as it would shed light into the factors that control the photooxygenation process.

$$s = \frac{k_1}{k_2} = \frac{\ln[1 - C(1 + ee)]}{\ln[1 - C(1 - ee)]} \quad (3)$$

The experimentally measured rate constants are presented in Table 2. The  $k_{\text{pq}}/k_{\text{cq}}$  values are experimentally the same for all the entries in the table regardless of alkene geometry

**Table 2** Chemical quenching rate constant ( $k_{\text{cq}}$ ), physical quenching rate ( $k_{\text{pq}}$ ) and total quenching rate constant ( $k_{\text{q}}$ ) for photooxygenation of enecarbamates

Substrate <sup>a</sup>	Configuration		Quenching rate constants <sup>b</sup> /M <sup>-1</sup> s <sup>-1</sup>				$s_{\text{cq}}^f$
	C-4	C-3'	$k_{\text{cq}} (\times 10^{-3})^c$	$k_{\text{q}} (\times 10^{-3})^d$	$k_{\text{pq}} (\times 10^{-3})$	$k_{\text{pq}}/k_{\text{cq}}$	
<i>Z</i> ( <i>Me</i> )- <b>1a</b>	<i>R</i>	<i>R</i>	2.4	43 <sup>e</sup>	41	17	1.3
<i>Z</i> ( <i>Me</i> )- <b>1a</b>	<i>R</i>	<i>S</i>	1.8	44 <sup>e</sup>	42	23	
<i>Z</i> ( <i>iPr</i> )- <b>2a</b>	<i>R</i>	<i>R</i>	1.7	26	24	16	1.2
<i>Z</i> ( <i>iPr</i> )- <b>2a</b>	<i>R</i>	<i>S</i>	1.3	24	22	17	
<i>Z</i> ( <i>iPr</i> )- <b>2b</b>	<i>S</i>	<i>S</i>	1.6	27	25	16	1.8
<i>E</i> ( <i>iPr</i> )- <b>2c</b>	<i>R</i>	<i>R</i>	1.6	23 <sup>e</sup>	21	13	
<i>E</i> ( <i>iPr</i> )- <b>2c</b>	<i>R</i>	<i>S</i>	0.9	20	19	21	

<sup>a</sup> Concentration of the substrates is given in ESI. <sup>†</sup> <sup>b</sup> Experimental error within 25–35%; average of 2 or more runs. <sup>c</sup> Data from ref. 11, determined by competitive kinetics using *trans*-4-octene as standard; in  $\text{CDCl}_3$  at  $20 \pm 1^\circ\text{C}$ . <sup>d</sup>  $\text{CDCl}_3$ ,  $22 \pm 1^\circ\text{C}$ ; for experimental details see ESI. <sup>†</sup> <sup>e</sup> Values obtained from optical antipode. <sup>f</sup>  $s_{\text{cq}}$  is the ratio of the chemical quenching rates of the C-3' epimers.

**Table 3** Stereoselectivity factors obtained upon photo-oxygenation of proteated and C-4 deuterated *E(Me)-1d* in CDCl<sub>3</sub>

Substrate	Temperature/°C	<i>s</i> [enhanced MDB enantiomer]
<i>E(Me)-1d</i>	15	10 [S] <sup>a</sup>
	-15	39 [S] <sup>a</sup>
	-40	71 [S] <sup>a</sup>
<i>E(Me)-1d(D<sub>3</sub>)</i>	15	11 [S]
	-15	25 [S]
	-40	41 [S]

<sup>a</sup> Data taken from previously published report: ref. 11.

(*E* vs. *Z*), the C-4 substituent or the C-4 and C-3' *R/S* configuration. The expected differences in physical quenching observed for the C-3' diastereomers are not evident. This, however, does not suggest the non-existence of vibrational quenching as one of the stereoselective mechanism in operation in the photooxygenation of the enecarbamates. The reactive rate constants are low, as expected, due to the electron deficient alkene double bonds<sup>23</sup> in enecarbamates. It must be noted that when calculations of rate constants are performed with  $k_q$  that are one order of magnitude larger than  $k_{cq}$ , then associated inaccuracy undermines possible conclusions.

As computed in Table 2, the ratio of chemical quenching rate constants  $s_{cq}$  (Table 2) is significantly higher for the *E* enecarbamates ( $s_{cq}$  of 1.8) compared to the corresponding *Z* enecarbamates ( $s_{cq}$  of 1.2). As the ratio of chemical quenching rate constants are not corrected for conversion they seem not to correlate with the *s* factor computed from *ee* values (Table 1). Nevertheless, we were encouraged to see the chemical quenching trend in line with the *s* factors computed from the *ee* values. We could not correlate physical quenching rates as it is not corrected for conversion. To overcome this, we took an indirect approach to verify the physical quenching process and to quantify the vibrational quenching contribution to stereoselective photooxidation of enecarbamates (in terms of the *ee* values in the product) by studying deuterated and the corresponding H enecarbamates (Table 3).

Deuteration of the vibrationally distinguishing groups allowed us to investigate vibrational quenching as a stereodifferentiation mechanism. The C-4 substituents of the *E(Me)-1d* enecarbamate (optically pure C-4 substituents and *R/S* epimer C-3' position) were deuterated and the *s* values at various temperatures were compared to the results for its proteated counterpart (Table 3). Upon inspection of Table 3, it is clear that deuterated and proteated enecarbamates gave different stereoselectivities under identical conditions. For example, at -40 °C, an *s* value of 41 favoring the *S*-MDB product was observed with the deuterated derivative, whereas an *s* value of 71 was observed with the corresponding H-derivative. The differences in stereodifferentiation decreased as the temperature was raised from -40 °C to 15 °C. At 15 °C, almost identical values were obtained for both the proteated (*s* = 10) and deuterated (*s* = 11) derivatives (Table 3). A rationale for the observation of a nearly doubled *s* values at -40 °C for proteated enecarbamates compared to deuterated enecarbamates is needed to quantify the selectivity. We believe that the difference in the selectivity between the two derivatives is a reflection of the ability of the proteated derivative to deactivate <sup>1</sup>O<sub>2</sub> selectively from one of the stereotopic faces of the epimeric enecarbamate (C-3' epimers) leading to noticeably

different stereoselectivities at a given temperature. As the rate constants (physical and chemical quenching rate constants) are dependent on the temperature, one would expect variation with the reaction temperature. Based on the *s* values, the difference in rate constants (physical and chemical) for a given pair of C-3' epimers is pronounced at lower temperatures leading to higher selectivities. At the present time, with our limited experimental data, we could only speculate about the origin of this difference. One possibility is the dynamic movement/restrictions involved leading to conformational changes in enecarbamates would be temperature dependent. If there is conformational bias towards a particular conformation of C-3' epimers, lowering the temperature could likely result in a conformation that interacts (chemically and/or physically) with <sup>1</sup>O<sub>2</sub> effectively from one C-3' epimer than the other. If sterics alone is involved in determining the extent of stereoselectivity, identical *s* values would be expected for both the proteated and the corresponding deuterated derivatives even at -40 °C. To rationalize the difference, factors apart from sterics would have to be involved in the stereodifferentiation process leading to the different stereoselectivities for the proteated and deuterated derivatives of *E(Me)-1d* enecarbamates. We attribute the additional stereoselectivity (nearly doubling of the *s* value in the case of *E(Me)-1d* at -40 °C; Table 3) obtained in the proteated enecarbamate to vibrational quenching of singlet oxygen by C-H bonds. This quenching is more pronounced at lower temperatures.

Thus our investigation has opened an opportunity to quantify the extent of physical quenching during the photooxygenation processes in terms of the enantiomeric excess in the photoproduct. Deuteration of enecarbamates offers a more accessible means of exploring this stereodifferentiating mechanism in the photooxygenation process. Efforts are underway in our laboratories to quantify the rate constants for the individual epimers of both the proteated and deuterated enecarbamates to obtain a more direct evidence of vibrational quenching.

## Acknowledgements

The authors at Columbia University thank the NSF for its generous support of this research through grants CHE 04-15516 and CHE 07-17518. J.S. thanks NDSU for a Faculty Start-up award and the National Science Foundation for the CAREER award (CHE-0748525). W.A. is grateful for previous funding from the Deutsche Forschungsgemeinschaft, the Fonds der Chemischen Industrie and the Alexander von Humboldt-Stiftung.

## References

- 1 Y. Inoue, Asymmetric photochemical reactions in solution, *Chem. Rev.*, 1992, **92**, 741–770.
- 2 H. Rau, Asymmetric photochemistry in solution, *Chem. Rev.*, 1983, **83**, 535–547.
- 3 Y. Inoue and V. Ramamurthy, eds., *Chiral Photochemistry*, Marcel Dekker, 2004.
- 4 M. A. Garcia-Garibay, Engineering Carbene Rearrangements in Crystals: From Molecular Information to Solid-State Reactivity, *Acc. Chem. Res.*, 2003, **36**, 491–498.
- 5 J. N. Gamlin, R. Jones, M. Leibovitch, B. Patrick, J. R. Scheffer and J. Trotter, The Ionic Auxiliary Concept in Solid State Organic Photochemistry, *Acc. Chem. Res.*, 1996, **29**, 203–209.
- 6 J. Sivaguru, A. Natarajan, L. S. Kaanumalle, J. Shailaja, S. Uppili, A. Joy and V. Ramamurthy, Asymmetric Photoreactions within Zeolites:

- Role of Confinement and Alkali Metal Ions, *Acc. Chem. Res.*, 2003, **36**, 509–521.
- 7 J. Sivaguru, R. B. Sunoj, T. Wada, Y. Origane, Y. Inoue and V. Ramamurthy, Enhanced Diastereoselectivity *via* Confinement: Diastereoselective Photoisomerization of 2,3-Diphenyl-1-benzoyl-cyclopropane Derivatives within Zeolites, *J. Org. Chem.*, 2004, **69**, 5528–5536.
- 8 W. Adam, S. G. Bosio and N. J. Turro, Control of the Mode Selectivity (Ene Reaction *versus* [2 + 2] Cycloaddition) in the Photooxygenation of Ene Carbamates: Directing Effect of an Alkenylic Nitrogen Functionality, *J. Am. Chem. Soc.*, 2002, **124**, 14004–14005.
- 9 W. Adam, S. G. Bosio and N. J. Turro, Highly Diastereoselective Dioxetane Formation in the Photooxygenation of Enecarbamates with an Oxazolidinone Chiral Auxiliary: Steric Control in the [2 + 2] Cycloaddition of Singlet Oxygen through Conformational Alignment, *J. Am. Chem. Soc.*, 2002, **124**, 8814–8815.
- 10 W. Adam, S. G. Bosio, N. J. Turro and B. T. Wolff, Enecarbamates as Selective Substrates in Oxidations: Chiral-Auxiliary-Controlled Mode Selectivity and Diastereoselectivity in the [2 + 2] Cycloaddition and Ene Reaction of Singlet Oxygen and in the Epoxidation by DMD and mCPBA, *J. Org. Chem.*, 2004, **69**, 1704–1715.
- 11 J. Sivaguru, M. R. Solomon, H. Saito, T. Poon, S. Jockusch, W. Adam, Y. Inoue and N. J. Turro, Conformationally controlled (entropy effects), stereoselective vibrational quenching of singlet oxygen in the oxidative cleavage of oxazolidinone-functionalized enecarbamates through solvent and temperature variations, *Tetrahedron*, 2006, **62**, 6707–6717.
- 12 Further details are included in the ESI†.
- 13 M. Solomon, J. Sivaguru, S. Jockusch, W. Adam and N. J. Turro, Vibrational deactivation of singlet oxygen: does it play a role in stereoselectivity during photooxygenation?, *Photochem. Photobiol. Sci.*, 2008, **7**, 531–533.
- 14 J. Sivaguru, M. R. Solomon, T. Poon, S. Jockusch, S. G. Bosio, W. Adam and N. J. Turro, The Reaction of Singlet Oxygen with Enecarbamates: A Mechanistic Playground for Investigating Chemoselectivity, Stereoselectivity, and Vibratoselectivity of Photooxidations, *Acc. Chem. Res.*, 2008, **41**, 387–400.
- 15 T. Poon, J. Sivaguru, R. Franz, S. Jockusch, C. Martinez, I. Washington, W. Adam, Y. Inoue and N. J. Turro, Temperature and Solvent Control of the Stereoselectivity in the Reactions of Singlet Oxygen with Oxazolidinone-Substituted Enecarbamates, *J. Am. Chem. Soc.*, 2004, **126**, 10498–10499.
- 16 C. Schweitzer and R. Schmidt, Physical Mechanisms of Generation and Deactivation of Singlet Oxygen, *Chem. Rev.*, 2003, **103**, 1685–1757.
- 17 F. Wilkinson, W. P. Helman and A. B. Ross, Rate constants for the decay and reactions of the lowest electronically excited singlet state of molecular oxygen in solution. An expanded and revised compilation, *J. Phys. Chem. Ref. Data*, 1995, **24**, 663–1021.
- 18 C. Tanielian and R. Mechin, Interaction of singlet molecular oxygen with disubstituted olefins. Evidence for a physical quenching induced by the hydrocarbon chain, *J. Phys. Chem.*, 1988, **92**, 265–267.
- 19 S. Ben-Shabat, Y. Itagaki, S. Jockusch, J. R. Sparrow, N. J. Turro and K. Nakanishi, Formation of a nonaoxirane from A2E, a lipofuscin fluorophore related to macular degeneration, and evidence of singlet oxygen involvement, *Angew. Chem., Int. Ed.*, 2002, **41**, 814–817.
- 20 T. Poon, N. J. Turro, J. Chapman, P. Lakshminarasimhan, X. Lei, S. Jockusch, R. Franz, I. Washington, W. Adam and S. G. Bosio, Stereochemical Features of the Physical and Chemical Interactions of Singlet Oxygen with Enecarbamates, *Org. Lett.*, 2003, **5**, 4951–4953.
- 21 H. B. Kagan and J. C. Fiaud, Kinetic resolution, *Top. Stereochem.*, 1988, **18**, 249–330.
- 22 J. Sivaguru, T. Poon, C. Hooper, H. Saito, M. R. Solomon, S. Jockusch, W. Adam, Y. Inoue and N. J. Turro, A comparative mechanistic analysis of the stereoselectivity trends observed in the oxidation of chiral oxazolidinone-functionalized enecarbamates by singlet oxygen, ozone, and triazolinedione, *Tetrahedron*, 2006, **62**, 10647–10659.
- 23 C. Vever-Bizet, M. Dellinger, D. Brault, M. Rougee and R. V. Bensasson, Singlet molecular oxygen quenching by saturated and unsaturated fatty-acids and by cholesterol, *Photochem. Photobiol.*, 1989, **50**, 321–325.