

References

1. Zell R, Krumbholz A, Wulzler P. Influenza A virus PB1–F2 gene [letter]. *Emerg Infect Dis*. 2006;12:1607–8.
2. Chen GW, Yang CC, Tsao KC, Huang CG, Lee LA, Yang WZ, et al. Influenza A virus PB1–F2 gene in recent Taiwanese isolates. *Emerg Infect Dis*. 2004;10:630–6.
3. Zamarin D, Ortigoza MB, Palese P. Influenza A virus PB1–F2 protein contributes to viral pathogenesis in mice. *J Virol*. 2006;80:7976–83.
4. Chen GW, Chang SC, Mok CK, Lo YL, Kung YN, Huang JH, et al. Genomic signatures of human versus avian influenza A viruses. *Emerg Infect Dis* 2006;9:1353–60.

Address for correspondence: Guang-Wu Chen, Chang Gung University, Department of Computer Science and Information Engineering, 259 Wen-Hua 1st Rd, Kwei-shan, Taoyuan, Taiwan 333; email: gwchen@mail.cgu.edu.tw

Enterovirus 75 and Aseptic Meningitis, Spain, 2005

To the Editor: Although most human enterovirus (EV) (genus *Enterovirus*, family *Picornaviridae*) infections are asymptomatic, they can cause upper respiratory illness, febrile rash, aseptic meningitis, pleurodynia, encephalitis, acute flaccid paralysis, and neonatal sepsislike disease (1). Most EVs have been implicated in aseptic meningitis, most notably echovirus (E) 30, 9, 6, and 11 and coxsackie B virus (CBV) type 5 (2); other serotypes are less frequently associated with neurologic disease.

New EV serotypes have come to light, chiefly as a result of molecular typing methods (3–6). EV75 was proposed as a new serotype of the EV genus in 2004 (5). Retrospective analysis showed that it had circulated sporadically in Asia, the United

States, and Africa since at least 1974. Only 8 isolates of this serotype have been reported worldwide, in 1974, 1985, 1986, 1987 (n = 2), 1998, and 2000 (n = 2). Infection in those cases was associated with respiratory disease, acute flaccid paralysis, neonatal jaundice, failure to thrive, or unspecified neurologic disease or was asymptomatic. At the time of writing this manuscript, EV75 had not been linked to aseptic meningitis.

From May 2005 through January 2006, 106 EVs were received for typing from Spanish hospital laboratories; 46 of them were from patients with aseptic meningitis, 10 from patients or contacts of patients with acute flaccid paralysis, 27 from patients with fever, 7 from patients with respiratory diseases, and 16 from other patients. Twenty EVs could not be typed by serum neutralization (7); however, 3' terminus VP1 gene sequence analysis (8) showed that they were E18 (n = 7), CBV3 (n = 1), and E16 (n = 2); 2 could not be typed with serologic or molecular methods because the 3' terminus of VP1 gene amplification was negative. The analysis of the 3' terminus of VP1 gene of the remaining 5 cerebrospinal fluid (CSF) and 3 nasopharyngeal isolates showed that they were similar to the recently proposed EV75 serotype (5). These 8 isolates were obtained from samples from children in Bilbao (n = 3), Granada (n = 3), Barcelona (n = 1), and the Canary Islands (n = 1). In 4 patients with aseptic meningitis, EV75 was isolated from CSF. EV75 was isolated from CSF of a fifth patient who had symptoms of fever and irritability. The remaining 3 EV75 isolates were from nasopharyngeal swabs of children who had fever, respiratory disease, or gastroenteritis. All isolates were grown in cell lines (rhabdomyosarcoma, lung adenocarcinoma, and human fetal lung fibroblast) and identified as EV by immunofluorescence with pan-EV antibody assays (Pan Entero Blend Chemicon,

Temecula, CA, USA, and Monoclonal Mouse Anti-Enterovirus, Dako, Glostrup, Denmark).

Phylogenetic analysis of the isolates from 2005 was performed on the basis of complete VP1 gene sequence (GenBank accession nos. DQ468137–DQ468142). The 5' terminal domain was obtained by reverse transcriptase–PCR with specific primers EV75_sense: 5'-GAAAGCTTYTTC-CAAGGGGA-3' and EV75_anti: 5'-GAGAAGTGKGCACCAWCCATC-3'. Phylogenetic analysis of the Spanish isolates and representatives of all other species B EVs showed that the Spanish isolates clustered (bootstrap value 100, Figure) with strains USA/OK85-10362, ETH74-1341, USA/VA86-10363, USA/CT87-10364-5, OMA98-10366, and BAN00-10367-8 (accession nos. AY556063–AY556070), corresponding to the proposed EV75. The Spanish isolates constitute a subgroup (bootstrap value 100, Figure). The similarity between the Spanish cluster and other EV75 isolates was 82.8%–85.4% at the nucleic acid level. Although the entire VP1 sequence was not available for the isolates from 2006, the VP1 3' terminal analysis showed the strains belonged to the same cluster.

To our knowledge, this is the first isolation of EV75 in Spain. Indeed, isolation of EV75 has not been reported in Europe. Given that the European EV75 isolate grows easily in a variety of cell lines, is detected by common EV genus-specific antibodies, and that EV surveillance and typing were performed in Spain since 1988 (2), EV75 might have begun to circulate in Spain recently. However, because isolates are not obtained from all aseptic meningitis patients and many EVs are detected by PCR but never typed, we cannot rule out the possibility of previous asymptomatic circulation.

The European strains of EV75 appear to represent a different evolutionary lineage than those previously




Figure. Phylogenetic analysis of complete VP1 sequences of Spanish enterovirus (EV) isolates (GenBank accession nos. DQ468137–DQ468142), the new proposed EV75 sequences (AY556063–AY556070 and AY919545), and prototype EV sequences (echovirus [E] 5, AJ241425; E31, AJ241435; E2, AF081315; E15, AJ241429; E14, AJ241428; E17, AF081330; coxsackie B virus [CBV] 2, AF081312; E26, AJ241433; E27, AF081338; E1, AJ241422; E8, AF081325; E4, AF081319; E21, AF081334; E30, AF081340; E25, AF081336; E29, AJ241434; CBV5, AF114383; CBV6, AF081313; E13, AF081327; EV69, AF081349; E24, AJ241432; E33, AF081346; E3, AF081316; E12, X77708; CBV3, M16572; CBV1, M16560; E6, AF081322; coxsackie A virus [CAV] 9, D00627; E16, AY302542; E9, AF524866; E7, AJ241426; E32, AF081345; E19, AJ241430; E11, AF081326; CBV4, X05690; E18, AF081331; E20, AJ241431; EV70, D17602; CAV6, AF081297; CAV13, AF081303; EV74, AY208118). Phylogenetic trees were constructed with the neighbor-joining method (MEGA version 3.0, available from <http://www.megasoftware.net>) with Kimura 2-parameter substitution model. Significance of phylogenies was estimated by bootstrap analysis with 1,000 pseudoreplicate datasets. Closed and open circles show Spanish and previously reported EV75 isolates, respectively.

described in the United States, Asia, and Africa (9). Only 1 of those EV75s was obtained from CSF (a nonspecific neurologic syndrome). Thus, EV75 has not been associated with aseptic meningitis, despite the fact that EV infections are a common cause of aseptic meningitis. Most of the Spanish isolates (5 of 8) were associated with aseptic meningitis in children. Although the number of EV75-associated cases was not high (as a

percentage of the number of EVs isolated from aseptic meningitis patients, 10.8%), the wide distribution of the cases may indicate wide circulation. To avoid outbreaks of aseptic meningitis caused by previously noncirculating EVs (EV13, 2001 [10]) and to help define the extent of circulation of newly identified EV types, careful surveillance of aseptic meningitis should be undertaken.

Ana Avellón,* Gurutze Rubio,†
Gustavo Palacios,‡
Inmaculada Casas,*
Nuria Rabella,§ Gabriel Reina,¶
Cármén Pérez,# W. Ian Lipkin,‡
and Gloria Trallero*

*Carlos III Institute of Health, Majadahonda, Spain; †Cruces Hospital, Bilbao, Spain; ‡Columbia University, New York, New York, USA; §Sant Pau y Santa Creu Hospital, Barcelona, Spain; ¶Virgen de las Nieves Hospital, Granada, Spain; and #Doctor Negrín Hospital, Las Palmas de Gran Canaria, Spain

References

- Melnick JL. Enteroviruses: polioviruses, coxsackieviruses, echoviruses, and newer enteroviruses. In: Fields BN, Knipe DM, Howley PM, editors. *Virology*. 3rd ed. Philadelphia: Lippincott-Raven; 1996. p. 655–712.
- Trallero G, Casas I, Tenorio A, Echevarria JE, Castellanos A, Lozano A, et al. Enteroviruses in Spain: virological and epidemiological studies over 10 years (1988–97). *Epidemiol Infect*. 2000;124:497–506.
- Oberste M, Schnurr D, Maher K, al-Busaidy S, Pallansch M. Molecular identification of new picornaviruses and characterization of a proposed enterovirus 73 serotype. *J Gen Virol*. 2001;82:409–16.
- Norder H, Bjerregaard L, Magnus L, Lina B, Aymard M, Chomel JJ. Sequencing of “untypable” enteroviruses reveals two new types, EV-77 and EV-78, within human enterovirus type B and substitutions in the BC loop of the VP1 protein for known types. *J Gen Virol*. 2003;84:827–36.
- Oberste MS, Michele SM, Maher K, Schnurr D, Cisterna D, Junttila N, et al. Molecular identification and characterization of two proposed new enterovirus serotypes, EV74 and EV75. *J Gen Virol*. 2004;85:3205–12.
- Oberste MS, Maher K, Michele SM, Belliot G, Uddin M, Pallansch MA. Enteroviruses 76, 89, 90 and 91 represent a novel group within the species human enterovirus A. *J Gen Virol*. 2005;86:445–51.
- Melnick JL, Wimberly IL. Lyophilized combination pools of enterovirus equine antisera: new LBM pools prepared from reserves of antisera stored frozen for two decades. *Bull World Health Organ*. 1985;63:543–50.
- Casas I, Palacios GF, Trallero G, Cisterna D, Freire MC, Tenorio A. Molecular characterization of human enteroviruses in clinical samples: comparison between VP2, VP1, and RNA polymerase regions using RT nested PCR assays and direct sequen-

- cing of products. *J Med Virol.* 2001; 65:138–48.
9. Palacios G, Casas I, Cisterna D, Trallero G, Tenorio A, Freire C. Molecular epidemiology of echovirus 30: temporal circulation and substitution of single lineages. *J Virol.* 2002;76:4940–9.
10. Avellon A, Casas I, Trallero G, Perez C, Tenorio A, Palacios G. Molecular analysis of echovirus 13 isolates and aseptic meningitis, Spain. *Emerg Infect Dis.* 2003;9:934–41.

Address for correspondence: Ana Avellón, Spanish Enterovirus Reference Laboratory, National Center of Microbiology, Carlos III Institute of Health, Carretera de Majadahonda a Pozuelo, Km 2, Majadahonda 28220, Madrid, Spain; email: aavellon@isci.iii.es

Ciprofloxacin-resistant *Salmonella* Kentucky in Travelers

To the Editor: Ciprofloxacin is the treatment of choice of severe nontyphoidal *Salmonella* infections in adults. Resistance to ciprofloxacin has been found exceptionally in nontyphoidal *Salmonella enterica* isolates and only in serotypes Typhimurium, Choleraesuis, and Schwarzengrund (1–8). Such isolates have been collected from humans and animals in Europe, Asia, and North America.

We report the emergence of ciprofloxacin-resistant isolates of *S. Kentucky* since 2002 in French travelers returning from northeast and eastern Africa. From 2000 through 2005, 197 *S. Kentucky* isolates from humans (1 per patient) were serotyped, from 69,759 total *S. enterica* isolates serotyped at the French National Reference Centre for *Salmonella*. Antimicrobial drug sus-

ceptibility was determined for 186 isolates by the disk-diffusion method with 32 antimicrobial drugs, as previously described (9). Resistance to several drugs, amoxicillin (18%), gentamicin (16%), nalidixic acid (21%), sulfonamides (24%), and tetracycline (24%), has been observed from 2000 through 2005.

A total of 17 (9%) ciprofloxacin-resistant *S. Kentucky* strains were isolated. A resistant isolate that was untypable by conventional serotyping (rough) but that had a pulsed-field gel electrophoresis (PFGE) profile associated with serotype Kentucky, was included in this study. Ciprofloxacin MIC levels in these isolates, determined by standard agar doubling dilution as previously described (2), were 4–16 mg/L. The first ciprofloxacin-resistant strain was isolated in December 2002 from a French tourist who had gastroenteritis during a Nile cruise in Egypt. In 2004 and 2005, 17 ciprofloxacin-resistant isolates were identified in unrelated adults who lived in different cities of France at different times of the year. The 16 patients we contacted acquired the infection during or immediately after travel to Egypt (10 patients), Kenya and Tanzania (3), or Sudan (1). In 2 cases, gastroenteritis occurred 2 months after travel to Egypt. None of the investigated cases were fatal or life-threatening.

The 18 ciprofloxacin-resistant isolates (17 serotype Kentucky and 1 rough) displayed various susceptibility patterns, from single resistance to quinolones to multiple resistance (up to 9 antimicrobial agents). To identify mutations responsible for ciprofloxacin resistance, the quinolone resistance-determining regions (QRDRs) of *gyrA*, *gyrB*, *parC*, and *parE* were amplified by PCR and sequenced as described previously (3,9), except that different forward primers for *gyrB* (5'-TTATCGACGC-CGCGCGTGCGC-3') and *parE* (5'-CGCGTAACTGCATCG-GGTTC-3')

were used. The 18 ciprofloxacin-resistant isolates had different double mutations in *gyrA* leading to amino acid substitutions, Ser83Phe and Asp87Gly (8 isolates), Ser83Phe and Asp87Asn (5), and Ser83Phe and Asp87Tyr (5), but had identical mutations in *parC* (resulting in Ser80Ile). An additional substitution was observed in ParC, Thr57Ser. This substitution, however, did not appear to be associated with quinolone resistance because it was also identified in nalidixic acid-susceptible isolates. No isolates had substitutions in the QRDRs of GyrB and ParE. All isolates tested by PCR for the plasmid-mediated quinolone resistance-conferring gene *qnrA* (9) were negative. In the presence of the efflux pump inhibitor Phe-Arg- β -naphthylamide, the MICs of ciprofloxacin were reduced from 4-fold to 16-fold, which suggests that an active efflux mechanism was present (2). The involvement of the AcrAB-TolC efflux system was determined by measuring AcrA expression with a method previously described (5). A moderate production of AcrA (3- to 4-fold increase when compared with the baseline production of AcrA in reference strain 98K) was observed in all but 1 ciprofloxacin-resistant isolate. This isolate overproduced (6-fold) AcrA, which correlated with a higher ciprofloxacin MIC (16 mg/L).

The 18 ciprofloxacin-resistant isolates and 14 ciprofloxacin-susceptible *S. Kentucky* isolates used for comparison were genotyped by PFGE with *Xba*I restriction and PulseNet's running conditions, as described previously (9). Each profile that differed by ≥ 1 clear band > 50 kb was considered a distinct profile. The 18 resistant isolates displayed 9 profiles that differed by 1 to 3 bands (Dice correlation coefficient 55%) (Figure). Profile X1c was predominant (7 [39%] of 18). The 6 pansusceptible isolates tested displayed 5 different patterns unrelated to those of resistant isolates. Use of a