

Field, Experimental, and Modeling Study of Arsenic Partitioning across a Redox Transition in a Bangladesh Aquifer


Hun Bok Jung,^{†,§} Benjamin C. Bostick,[‡] and Yan Zheng^{*,†,‡}

[†]School of Earth and Environmental Sciences, Queens College, and Graduate School and University Center, City University of New York, Flushing, New York 11367, United States

[‡]Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York 10964, United States

S Supporting Information

ABSTRACT: To understand redox-dependent arsenic partitioning, we performed batch sorption and desorption experiments using aquifer sands subjected to chemical and mineralogical characterization. Sands collected from the redox transition zone between reducing groundwater and oxic river water at the Meghna riverbank with HCl extractable Fe(III)/Fe ratio ranging from 0.32 to 0.74 are representative of the redox conditions of aquifers common in nature. One brown suboxic sediment displayed a partitioning coefficient (K_d) of 7–8 L kg⁻¹ at equilibrium with 100 μg L⁻¹ As(III), while two gray reducing sediments showed K_d of 1–2 L kg⁻¹. Lactate amendment to aquifer sands containing 91 mg kg⁻¹ P-extractable As resulted in the reduction of As and Fe with sediment Fe(III)/Fe decreasing from 0.54 to 0.44, and mobilized an equivalent of 64 mg kg⁻¹ As over a month. Desorption of As from nonlactate-amended sediment was negligible with little change in sediment Fe(III)/Fe. This release of As is consistent with microbial reduction of Fe(III) oxyhydroxides and the resulting decrease in the number of surface sites on Fe(III) oxyhydroxides. Arsenic partitioning (K_d) in iron-rich, sulfur-poor aquifers with circumneutral pH is redox-dependent and can be estimated by HCl leachable sediment Fe(III)/Fe ratio with typical Fe concentrations.



INTRODUCTION

Quantification of arsenic partitioning between solute and solid phases is important to assess transport of this toxic element in aquifers. The sorption reactions of As have been extensively investigated for synthetic Fe, Mn, and Al oxyhydroxides and for clay minerals.^{1–4} A handful of studies examined natural soil or sediments under oxic conditions and found that sorption behavior was mostly similar to that of pure Fe(III) oxyhydroxides.^{5–7} A knowledge gap exists for As partitioning in reducing aquifers conducive to As mobilization due to difficulties in sample preservation for sorption experiments and mineralogical characterization.⁸

Sorption isotherms of As(III) onto sands from a coastal aquifer of Waquoit Bay, MA⁹ suggest a redox-dependent As partitioning,^{10,11} showing increasing As(III) sorption with higher Fe(III) oxides content. In the Ganges-Brahmaputra-Meghna Delta (GBMD, South Asia), the As(III) sorption coefficient (K_d) at equilibrium with a typical groundwater As concentration of ~100 μg L⁻¹ established by batch experiment or field investigation ranged from 2 to 6 L kg⁻¹ for Holocene reducing aquifer sediments,^{8,12,13} and were approximately 20–30 L kg⁻¹ for brown sediments from deeper depths.^{14–16} To the best of our knowledge, batch sorption studies have not evaluated As partitioning in sediments of Bangladesh Holocene shallow aquifer, frequently enriched in groundwater arsenic

under reducing conditions that are typical for many aquifers with elevated As. Because transport in aquifers is thought to influence the distribution of As in aquifers,¹⁷ improved understanding of As partitioning has implications on interpretation of the source and fate of As in the GBMD⁸ and elsewhere.

The objectives of this study are to quantify As partitioning by batch sorption–desorption experiments using natural sediment and porewater samples with minimal disturbance of redox condition, and to understand parameters and mechanisms controlling the mobility of As in iron-rich and sulfur-poor aquifer sediment through mineralogical characterization and geochemical modeling. Along the Meghna River bank (Bangladesh) where iron- and arsenic-rich anoxic reducing groundwater interacts with oxic river water, a redox transition zone spanning a wide range of redox conditions within small spatial distances and with enrichment of sediment As¹⁸ up to hundreds or thousands of mg kg⁻¹ provides an ideal setting. The study is especially relevant to As transport in floodplain aquifers of South Asia where elevated groundwater As is

Received: September 19, 2011

Revised: November 17, 2011

Accepted: December 22, 2011

Published: December 22, 2011




Figure 1. Top: Schematic diagram showing the location of porewater and sediment core sampling collected in Jan 2006 (black font) and Nov 2007 (gray font) with the distance from the river shore in Jan 2006. Open bars and solid bars indicate porewater piezometers and sediment cores, respectively. Middle: Depth profiles of porewater dissolved Fe, As, and As(III) concentrations. Bottom: Depth profiles of sediment HCl-leachable Fe(III) and P-extractable total As and As(III) concentrations.

widespread¹⁹ with similar hydrogeological settings,^{20–22} and has implications for As transport in aquifers.

MATERIALS AND METHODS

Sediment and Porewater Samples. In addition to shallow well groundwater and river water (Table S1), sediment and porewater depth profiles (Figure 1) were collected from surface to 6–7 m depth near the Meghna River shore (23.6° N,

90.6° E) in Gazaria, Bangladesh (Figure S1) using a soil probe (AMS Inc., USA) and a stainless steel drive point piezometer system (Retract-A-Tip, AMS Inc.) at corresponding depths in Jan 2006 and Nov 2007. Sampling and analysis procedures are similar to those of Jung et al.⁹ and are described in the Supporting Information in detail. Sorbed sediment As was determined by N₂-purged 1 M sodium phosphate (Fisher) solution containing 0.1 M ascorbic acid for 36 h in a crimp-

sealed amber vial to characterize sediment samples, including the sorption–desorption experiments (Tables S2, S6, and S7). A hot 1.2 N HCl extraction was conducted at 80 °C for 1 h for the same samples to determine As, Fe, Mn, S, and P associated with Fe oxides. For the sorption experiment, three sediment samples at depths of 1.5, 3.6, and 5.4 m from core RS39, paired with porewater samples (PZ10) from the same depth interval, were collected on Nov 4, 2007. For the desorption experiment, a section of RS19-4 (depth 1.5–1.8 m) collected on Jan 25, 2006 was used. Sediment cores were double-sealed in nitrogen-filled Mylar bags with oxygen absorbent (SorbentSystems) during transportation, and refrigerated at 4 °C prior to use.

X-ray Absorption Spectroscopy. The arsenic speciation and Fe mineralogy were analyzed by X-ray absorption spectroscopy (XAS) at the Stanford Synchrotron Radiation Laboratory. Iron mineralogy was obtained using extended X-ray absorption fine structure (EXAFS) spectroscopy. Arsenic speciation was determined using X-ray absorption near edge spectroscopy (XANES). Due to the very similar spectra between less crystalline goethite and ferrihydrite, which may be hard to differentiate based on linear combination fitting approaches,²³ the concentrations of ferrihydrite estimated represent the sum of ferrihydrite and freshly precipitated (poorly crystalline) goethite (Supporting Information).

Batch Sorption Experiment. One suboxic brown sediment with Fe(III)/Fe of 0.58 and two reducing gray sediments with Fe(III)/Fe ratio of ~0.37 from RS39, containing HCl-leachable or P-extractable As of ~1 mg kg⁻¹ (Table S2; HCl- and P-extraction were conducted using different aliquots of the same sample), were subjected to batch sorption experiment in the field in a N₂ filled glovebox immediately after sample collection. About 10 mL of porewater (pH 6.0–6.5) from the same depth (PZ10, Table S3) was added to 15-mL amber serum vials containing 5–6 g of homogenized sediment spiked with As(III) and As(V) stock solutions freshly prepared using reagent grade NaAsO₂ and NaHAsO₄·7H₂O to yield concentrations from 0 to ~7000 µg L⁻¹ As (Table S4). The vials were crimp-sealed without headspace and intermittently shaken, kept in a N₂-filled chamber except during transportation in N₂-filled Mylar bags, and sampled after 7 days. The supernatant was analyzed for As(III) immediately by voltammetry after filtration through a 0.45-µm membrane filter and acidification to 1% HCl and for total As by ICP-MS (Supporting Information). The pH of the supernatant likely remained circumneutral during the experiment because sediment and porewater samples were from the same depth, and thus were already at equilibrium.

Batch Desorption Experiment. To ascertain the mobility of As found at high concentrations (46–600 mg kg⁻¹ P-extractable As), a sediment core section (1.5–1.8 m depth) from RS19-4 was homogenized inside a N₂-filled chamber, and reacted with suboxic nanopure water (>18 MΩ) or artificial groundwater for 1 month. Artificial groundwater and nanopure water were purged with pure nitrogen gas to reach a dissolved oxygen value of ~0.5 mg L⁻¹, determined by a CHEMets dissolved oxygen kit. The homogenized sample contained a P-extractable As concentration of 91 mg kg⁻¹ with 57 mg kg⁻¹ as As(III), HCl-leachable Fe and As concentration of 6933 and 86 mg kg⁻¹, respectively. Artificial groundwater composition was based on that of the typical groundwater composition from adjacent tube wells at similar depths: 1 mM NaHCO₃, 0.02 mM KH₂PO₄, 2.5 mM CaCO₃, 0.16 mM MgSO₄, 0.50 mM MgCl₂,

and 0.06 mM KCl, and was buffered to a pH of 6.8 using 20 mM PIPES (piperazine-*N,N'*-bis(2-ethanesulfonic acid)).

There were four types of amendment. Type I used nanopure water (pH = 6.0) with 50 mg L⁻¹ kanamycin, an antibiotic. Type II used artificial groundwater (pH = 6.8) with 50 mg L⁻¹ kanamycin. Type III is the same as Type II but without kanamycin, serving as a biotic control. Type IV is Type III plus 1 mM lactate, serving as an enhanced biotic control. Kanamycin was intended to limit indigenous microbial activities. Each series started in 5 replicates of 15-mL serum bottles containing 5 mL of nanopure water or artificial groundwater and ~1 g of sediment. All glassware was autoclaved. After flushing the headspace with N₂ gas and crimp sealing, bottles were shaken at 140 rpm over 1 month in a N₂ filled anaerobic chamber (Coy Laboratory Products) without H₂ but with oxygen absorbent (SorbentSystems). The supernatant and sediment for 4 types of incubation were sampled sacrificially from a replicate of each type on the 2nd, 5th, 10th, 17th, and 30th day. Total Fe, Mn, S, P, and As in filtered supernatant and sediment extracts by hot HCl and anoxic sodium phosphate were analyzed, as well as Fe(II) and As(III) (Supporting Information).

Langmuir Isotherm and Surface Complexation Modeling (SCM). Batch sorption equilibrium data were fitted to Langmuir isotherms to determine sorption capacity (S_{max}) and K_{La} , a constant representing the binding strength, and the effective K_d values at equilibrium with 100 µg L⁻¹ As, typical of Bangladesh groundwater.⁸ A semimechanistic SCM of sorption experimental data was conducted in PHREEQC version 2.15 with MINTEQA2 version 4.0 database, using the acidity constants and equilibrium constants (K ; Table S5) for a diffuse double-layer surface complexation model of hydrous ferric oxide (HFO),²⁴ given that Fe(III) oxyhydroxides are the dominant reactive mineral phase in suboxic sediment (Table 1). Other Fe phases and primary minerals such as quartz, feldspar, and mica vary little along the depth profiles,¹⁸ and are not included. For both amorphous and crystalline Fe(III) oxide minerals, the same sorption equilibrium constants were applied because intrinsic surface complexation constants of metals and anions for both HFO and goethite have been found to be similar.²⁵ Surface complexation of Si was not considered because the K value is not available from MINTEQA2 database. Surface site density was adjusted based on HCl leachable Fe(III) concentrations, assuming that Fe minerals are goethite-like phases with ~0.01 mol surface sites per mol Fe(III)²⁶ to fit As data in sorption SCM modeling. Based on the composition of porewater (Table S3), pH, Ca, and alkalinity were fixed to 6.0, 0.3 mM, and 0.6 mM for brown sediment, respectively, and were 6.3, 1 mM, and 2 mM for gray sediment, respectively. P concentrations of 0.005 and 0.04 mM were used for brown and gray sediments, respectively, according to aqueous composition at the end of the experiment. In desorption SCM modeling for type III without lactate and type IV with lactate, all P-extractable As of 91 mg kg⁻¹ was subjected to adsorption reactions. For each sampling time point, solid-phase arsenic redox speciation was considered for modeling. Based on P-extractable sediment As data, the percentage of As(III) was initially 50%, decreasing to 40% on day 17, and 30% on day 30 in type III. In type IV, the percentage of As(III) was initially 50%, increasing to 60% on day 2, 70% on day 5, 80% on day 10, and 100% on days 17 and 30. The surface site density was estimated using HCl leachable Fe(III) concentration for each sampling point. For type III, Fe minerals were presumed to be like ferrihydrite with 0.20 mol surface sites per mol Fe(III)²⁴

Table 1. Iron Mineralogy of Meghna Riverbank Sediment Collected from Gazaria, Bangladesh in January 2006

sample no	latitude	longitude	depth (m)	HCl leach Fe(III)/Fe	HCl leach Fe(III) mg/kg	Fe mineralogy									
						ferrihydrate	hematite	goethite	magnetite	biotite	siderite	FeS	Chi2	Red_Chi2	
RS20-1	23° 35.006'	90° 35.160'	0.3	0.74	7715	45 ± 13%	0 ± 3%	0 ± 6%	0 ± 4%	21 ± 5%	34 ± 6%	0 ± 3%	145	0.85	
RS20-4	25 m from the shore		1.8	0.54	2783	52 ± 8%	0 ± 2%	0 ± 4%	0 ± 3%	21 ± 3%	26 ± 4%	0 ± 2%	74	0.43	
RS20-7			6.1	0.49	5029	49 ± 10%	0 ± 2%	0 ± 5%	0 ± 3%	20 ± 4%	30 ± 5%	0 ± 2%	107	0.63	
RS21-1	23° 35.008'	90° 35.153'	0.3	0.64	5109	62 ± 6%	0 ± 1%	0 ± 3%	0 ± 2%	20 ± 2%	19 ± 3%	0 ± 1%	44	0.26	
RS21-4	14 m from the shore		1.4	0.50	2529	38 ± 19%	0 ± 4%	7 ± 9%	4 ± 6%	16 ± 8%	26 ± 9%	9 ± 4%	315	1.84	
RS21-7			6.7	0.46	2935	48 ± 18%	5 ± 4%	0 ± 9%	0 ± 6%	28 ± 7%	19 ± 9%	0 ± 4%	314	1.84	
RS19-1	23° 35.011'	90° 35.149'	0.3	0.65	5748	59 ± 8%	0 ± 2%	4 ± 4%	0 ± 3%	14 ± 3%	21 ± 4%	1 ± 2%	57	0.34	
RS19-3	7 m from the shore		0.9	0.52	2741	45 ± 19%	0 ± 4%	0 ± 9%	0 ± 6%	23 ± 7%	30 ± 9%	1 ± 4%	316	1.85	
RS19-4			1.8	0.32	2246	39 ± 16%	0 ± 4%	0 ± 8%	0 ± 6%	33 ± 6%	27 ± 8%	1 ± 3%	209	1.22	

during a period of 1 month, whereas for type IV, it was assumed that Fe minerals were initially ferrihydrite-like Fe(III) mineral with 0.20 mol surface sites per mol Fe(III), and finally transformed to magnetite-like Fe(II, III) minerals with ~0.01 mol surface sites per mol Fe(III). Based on aqueous data, pH, Ca, alkalinity, and P were fixed to 6.8, 2.5, 5, and 0.01 mM, respectively.

RESULTS AND DISCUSSION

Redox Transition Zone. A redox transition zone between approximately 0 and 2 m depth is evident from depth profiles of porewater and sediment Fe and As (Figure 1). Porewater Fe and As reach peak concentrations of 20–30 mg L⁻¹ and 200–300 μg L⁻¹ between 3 and 4 m although PZ7 and 10 depth profiles show a less pronounced peak most likely due to deeper infiltration of oxic river water at the shore (Figure 1). Dissolved As speciation is dominated by As(III) (90 ± 5%; *n* = 7) at depth ≥ 2 m but has less As(III) (57 ± 7%; *n* = 3) at depth < 2 m. HCl-leachable sediment Fe(III) was highest at depths < 1 m (6.3 ± 2.7 g kg⁻¹, *n* = 7) and decreased rapidly with depth (3.3 ± 1.7 g kg⁻¹ at depth of 1–2 m, *n* = 6; 3.1 ± 1.5 g kg⁻¹ at depth > 2 m, *n* = 10). This decrease correlates to decreases in the Fe(III)/Fe ratio (0.65 ± 0.07 at depth < 1 m, 0.53 ± 0.12 at depth of 1–2 m, and 0.40 ± 0.11 at depth > 2 m). This redox transition zone develops because the river water level declines faster than the groundwater table after the monsoon season (April–October) and causes a hydraulic gradient from the aquifer toward the river,⁸ driving discharge of anoxic groundwater (DO < ~0.1 mg L⁻¹) to oxic river water (DO > 5 mg L⁻¹). A summary of the average chemical composition of groundwater, porewater, and river water is available in Supporting Information (Table S1).

Iron Mineralogy and Sediment Arsenic. In most cases, the dominant Fe mineral in 9 samples analyzed was ferrihydrite-like, accounting for 49 ± 8% of Fe minerals at depths < 2 m (Table 1, Figure S2). Small concentrations of crystalline goethite and hematite were also detected in 2 samples (Table 1), and these concentrations probably represent minimum values given that microcrystalline Fe(III) oxides have spectra that are similar to ferrihydrite and thus may in some cases not be quantified separately. The preponderance of ferrihydrite is somewhat surprising because it commonly converts to more stable oxides.^{27,28} We explain this apparent discrepancy as follows. First, ferrihydrite commonly forms through the rapid oxidation of Fe(II) at neutral pH,²⁷ and the study site is characterized by the active discharge of Fe(II)-rich groundwater. Second, coprecipitated silica, organic matter, and trace metals from groundwater may stabilize the ferrihydrite.²⁹ Kinetics of seasonal transformation of ferrihydrite to stable Fe phases is likely affected by the seasonal transition of hydrological condition between dry season and rainy season. The prevalence of Fe(III) minerals at shallow depths and then decreasing with depth is consistent with the chemical extraction data (Figure 1, Table 1). A high quantity of Fe(II) minerals (26 ± 5%; fit as siderite), as well as iron silicates (22 ± 6%; fit as biotite), are also present in the samples. The presence of biotite or other Fe silicates indicates that sediments are young because it weathers rapidly to secondary Fe(III) oxides. Although green rust may possibly be present in Bangladesh aquifer,⁸ green rust was not considered due to its metastable nature.²⁷

P-extractable sediment As concentration is elevated between 0 and 2 m depth and within ~15 m distance from the river, showing 46 mg kg⁻¹ As at RS21, 284, and 600 mg kg⁻¹ As at

RS19 (Figure 1). XRF analyses of another sediment core (RS34) collected 2 m away from the shore found that bulk sediment As concentration is 100 mg kg^{-1} at 1.4 m depth. Sediment As speciation determined by P-extraction and XANES (Figure 1, Figure S2, Table S6) suggests mixed As(III) and As(V) (Supporting Information).

Langmuir Sorption Isotherms. As(III) sorption to sediment core RS39 was equilibrated within 10 h. Adsorption data fit well to Langmuir isotherms (Figure 2; $\text{RMSE} \leq 0.6$)




Figure 2. Equilibrium isotherms of As sorption to one brown (diamond) and two gray sediment (triangle and square) samples (RS39) collected in Nov 2007. (A) As(III) spiking experiment; (B) As(V) spiking experiment. Dashed lines represent Langmuir isotherm fit. Solid lines represent surface complexation model fit and sorbed As species is 100% As(III) in SCM.

that differ considerably depending on the color (redox state) of the sediment. Based on these isotherms, the calculated K_d values at equilibrium with $100 \mu\text{g L}^{-1}$ solute As are 7–8 and 1–2 L kg^{-1} for brown and gray sediments, respectively (Table S2). These values are comparable regardless of whether sediments were spiked with As(III) or As(V). For gray sediment, as solute As increases from 10 to $1000 \mu\text{g L}^{-1}$, a typical groundwater As concentration range in the GBMD aquifers, the K_d value of the gray sediment is nearly constant at 1–2 L kg^{-1} , similar to a relatively constant K_d of $\sim 4 \text{ L kg}^{-1}$ that was proposed for reducing shallow aquifer of GBMD across a range of As concentrations of nearly 3 orders of magnitude.¹³

The sorption capacity (S_{max}) for brown sediment was also about 2–3 times higher than that for gray sediment (Table S2). Regardless of whether As(III) or As(V) was spiked, the dissolved As was primarily As(III) ($94 \pm 11\%$ and $89 \pm 14\%$ for As(III) and As(V) additions, respectively, Table S4) possibly

due to redox processes readily reducing As(V) to As(III). This may in part explain the similar sorption behaviors in As(III) and As(V) spiked experiments because it is essentially As(III) equilibrium (Figure 2).

Surface Complexation Model. For As(III) and As(V) spikes, the best-fitted surface site density was 0.6 and $0.2 \mu\text{M g}^{-1}$ for brown and gray sediments, respectively ($\text{RMSE} < 0.8$), assuming that all sorbed As is As(III) in SCM (Figure 2). The 3-fold higher surface site density in brown sediment than gray sediment indicates that they have different concentrations of similar assemblage of minerals controlling adsorption. This is supported by differences in the concentration of HCl-leachable Fe(III) (Table S2), with $\sim 0.01 \text{ mol}$ surface sites per mol Fe(III), assuming that Fe(III) phases are principally responsible for As retention. Using a surface site density of 3.5 sites nm^{-2} ,³⁰ an Fe mineral surface area of $\sim 20 \text{ m}^2 \text{ g}^{-1}$ yields Fe concentrations that are consistent with observed HCl-leachable Fe(III) concentrations of ~ 2700 and $\sim 1200 \text{ mg kg}^{-1}$ for brown and gray sediments, respectively (Table S2). This surface area is consistent with a range of microcrystalline Fe oxides such as goethite but is much lower than what is common for ferrihydrite, which was identified by XAS spectroscopy. This difference can be explained in three ways. First, it is possible or even likely that microcrystalline goethite present in these sediments anomalously fit with ferrihydrite due to spectral similarities. Second, ferrihydrite, which often has measured surface areas in excess of $200 \text{ m}^2 \text{ g}^{-1}$,²⁶ may form aggregates or have blocked surfaces that make them less reactive. Indeed, the mole ratio of As effectively scavenged by fresh ferrihydrite is about 1:50,³¹ consistent with a surface area of about $\sim 60 \text{ m}^2 \text{ g}^{-1}$, and more aged forms could be less reactive. Third, the XAS data were collected in the middle of the dry season (January) during groundwater discharge. Fresh precipitates formed in such environments could indeed contain ferrihydrite that could be converted to more stable forms, catalyzed by their partial reduction²⁸ during the following rainy season. Other sediments collected in this area have also exhibited widely different sorption capacities, possibly due to different Fe mineral composition.⁴ The As sorption capacity of 13–23 mg kg^{-1} established by the batch sorption experiment using suboxic brown sediment collected in Nov 2007 is considerably lower than that observed for suboxic sediment collected in Jan 2003¹⁸ or Jan 2006 (Figure 1), which had P-extractable sediment As concentrations as high as 100–1000 mg kg^{-1} ; this difference is also attributed to the seasonal transformation of Fe minerals. The extent of such seasonal transformations needs to be assessed in the future.

Desorption of Arsenic and Iron Mineralogy. Desorption experiments performed in the absence of lactate (types I, II, and III) released measurable As into solution within 2 days, its concentration reaching $604 \pm 57 \mu\text{g L}^{-1}$, primarily as As(III) (Figure 3, Table S7). This As release was decoupled from a rise of Fe concentration, with 1237, <10, and $18 \mu\text{g L}^{-1}$ for types I, II, and III, respectively. Over the next 28 days, dissolved As concentrations decreased by 15–31% and it was oxidized to nearly 100% As(V) (Table S7). The oxidation of As is also observed in the sediment, with P-extractable As(III) decreasing from 57 mg kg^{-1} to finally 15, 32, and 22 mg kg^{-1} for types I, II, and III, respectively, while the total pool of adsorbed As remained constant, with the nearly unchanged total P-extractable (80–90 mg/kg) and HCl leachable As ($\sim 70 \text{ mg/kg}$) concentrations (Figure 3B, Table S7). Although we are unsure of the responsible oxidants, limited introduction of




Figure 3. Fe and As in aqueous phase (A) and solid phase (B) during desorption experiment using RS19-4 sediment core collected in Jan 2006. Type I: nanopure water + kanamycin (50 mg/L); Type II: artificial groundwater + kanamycin (50 mg/L); Type III: artificial groundwater; Type IV: artificial groundwater + lactate (1 mM). In the panel showing HCl-leachable Fe (B right), solid lines with large symbols represent Fe(III) concentration, and dotted lines with small symbols are Fe(III)/Fe ratio. Surface complexation modeling of As desorption in type III without lactate and type IV with lactate are indicated by black dashed line and red dashed line, respectively.

oxygen ($\sim 0.5 \text{ mg L}^{-1}$ dissolved O_2) at the beginning of the experiment, or another oxidant such as Mn(IV) oxides in the sediments, are possibilities.¹⁴ The HCl leachable Mn concentration is $\sim 100 \text{ mg kg}^{-1}$, but the mineralogy of Mn phase has not been confirmed in these sediments (Table S7). The overall oxidation state of sediment Fe was constant as evidenced by a constant Fe(III)/Fe ratio throughout the experiment (0.5–0.55), while the HCl leachable Fe(III) decreased from its initial concentration of 3500 to $\sim 2700 \text{ mg kg}^{-1}$ over the course of the experiment (Figure 3B). This could take place when some of the Fe(III) oxyhydroxides are transformed to more crystalline Fe(III) phases (e.g., goethite, hematite), resulting in the decreased leachability by HCl with little change in overall oxidation state.

Sediments incubated with 1 mM lactate (type IV) released similar quantities of As into solution ($\sim 630 \mu\text{g L}^{-1}$, Figure 3A) on the second day and also without Fe ($< 10 \mu\text{g L}^{-1}$), but As release was much more appreciable between the 5th and the 30th day, at which point the dissolved As reached $12\,081 \mu\text{g L}^{-1}$. This high As concentration is equivalent to mobilization of 64 mg kg^{-1} sediment As, or 70% of the initial P-extractable As. Concurrent decreases in P-extractable and HCl-leachable As concentrations were also observed, as was the nearly complete conversion of dissolved As to As(III) (Figure 3B, Table S7). The release of dissolved Fe resembled that of As except for the 17th day. In the first 5 days, Fe release was low and comparable to type I, with $482 \mu\text{g L}^{-1}$ dissolved Fe, equivalent to 2.5 mg kg^{-1} sediment Fe. It then increased to 3 mg L^{-1} , and finally to 8 mg L^{-1} , equivalent to 40 mg kg^{-1} sediment Fe. The HCl-leachable Fe(III) decreased from 3500 to $\sim 2450 \text{ mg kg}^{-1}$ and sediment Fe(III)/Fe ratio displayed a measurable and consistent decrease from 0.54 to 0.44 between the 10th and the 30th day (Figure 3). The change in extractable Fe(III)/Fe in the solid phase indicates that a significant portion of the Fe(III) was reduced (and not just changes in leachability by HCl), and that much of the Fe(II) produced was conserved in the solid phase. The reduction of Fe and As in both solution

and sediment, as well as subsequent desorption of As, appear to have been caused by microbial activity³² stimulated with addition of labile organic carbon to sediment preserved at $4 \text{ }^\circ\text{C}$ for 11 months.¹⁵ This is different from the naturally occurring reduction of As during sorption experiment without addition of labile organic carbon perhaps because the sorption experiment was conducted using fresh sediment containing active microbial community immediately after sample collection.

Decrease in surface site density due to microbial transformation of Fe minerals stimulated by lactate was likely a major cause for extensive As desorption. Although no XAS data were available to ascertain the nature of iron mineralogy change during the desorption experiment, SCM modeling based on surface sites derived from HCl-leachable Fe(III) concentration simulated the evolution of solute (Figure 3A) and solid As (Figure 3B) reasonably well, although dissolved As is overestimated (Figure 3A). The overestimation may be due to surface site density of natural Fe(III) oxyhydroxides in sediment being higher than that of typical synthetic ferrihydrite (0.2 mol surface sites per mol Fe), or some As coprecipitated with Fe in sediment could not be mobilized during the experiment. The sediment Fe(III)/Fe decrease with lactate over 1 month was interpreted as a redox transformation of Fe minerals from an initially ferrihydrite-like Fe(III) mineral with 0.20 mol surface sites per mol Fe(III) to finally magnetite-like Fe(II, III) minerals with 0.01 mol surface sites per mol Fe(III) (same as the surface site concentration from sorption experiment with RS39 sediments collected in Nov 2007), while the surface site concentrations for day 5, 10, and 17 were adjusted to 0.20, 0.15, and 0.04 mol per mol Fe(III), respectively, for the best fit. For types without lactate, 0.20 mol surface sites per mol Fe(III) were used throughout. It is worth noting that color change of Bangladesh Holocene aquifer sediment from brown to gray corresponds to a threshold Fe(III)/Fe ratio of ~ 0.6 .³³ Therefore, the SCM modeling results suggest that ferrihydrite-like Fe(III) mineral or poorly crystalline goethite with higher surface site density is initially

the dominant Fe mineral in RS19 sediment collected in Jan 2006 and is supported by XAS data (Table 1), whereas microcrystalline goethite with lower surface site density is possibly dominant in RS39 sediment collected in Nov 2007. The results also suggest that Meghna Riverbank sediment As will remain reasonably immobile in the absence of labile organic carbon.

Redox-Dependent As Partitioning in Iron-Dominant System. Arsenic partitioning to aquifer sands appears to vary systematically with sediment redox state. The K_d values for As(III) at equilibrium with $100 \mu\text{g L}^{-1}$ As(III) observed in batch sorption experiments using sands from the redox transition zones of Meghna Riverbank in this study compare favorably with those determined within the Waquoit Bay aquifer,⁹ as well as Bangladesh Pleistocene orange sands.^{14,15} Arsenic adsorption is considerably greater in orange sands than gray sands. Interestingly, adsorption (as indicated by the effective K_d) is not clearly proportional to sediment Fe(III) concentration (Figure S4) probably due to mineralogical differences such as crystallinity and morphology, which affect the leachability of Fe minerals by HCl. Instead, the K_d and sediment Fe(III)/Fe ratio can be fitted with an exponential function ($R^2 = 0.97$, $n = 7$; Figure 4) excluding a black reducing




Figure 4. Correlation between HCl-leachable Fe(III)/Fe ratio and As partitioning coefficient (K_d) for a solute As of $100 \mu\text{g L}^{-1}$ at circumneutral pH. Data are from the Meghna riverbank (this study) and the Waquoit Bay coastal aquifer,⁹ and Pleistocene aquifer sands from Bangladesh.^{14,15} Langmuir isotherm fit was used to obtain K_d , except for Stollenwerk et al.,¹⁴ in which K_d was estimated based on SCM fit. The exponential fit excludes the K_d for a black Waquoit Bay sediment with porewater molar Fe/S of ~ 0.005 ($\text{Fe/S} < 1$, a sulfur-rich system). In Bangladesh aquifer, the Fe(III)/Fe ratio and K_d range from 0.37 to 0.90 and from 1 to 26 L kg^{-1} , respectively.

sediment from Waquoit Bay with Fe(III)/Fe of 0.26.⁹ The higher than expected K_d value of the black sediment from Waquoit Bay is attributed to As-sulfide precipitation¹¹ during sorption experiment because of high sulfur and low iron concentration in porewater (Fe/S molar ratio < 1).^{9,11,34} This relationship can be used to estimate K_d for aquifer sands where sulfur content is low (e.g., $\text{Fe/S} > 1$), sediment Fe(III)/Fe ratio is between 0.3–0.9, and As mobility is governed by interaction with Fe minerals. The nearly constant K_d values of 2–6 reported for reducing GBMD aquifer sediments with Fe(III)/Fe < 0.5 ^{8,12,13} is thus not surprising.

Despite a number of limitations such as unavailable Fe mineralogy data for desorption experiment, spatially and seasonally heterogeneous sediment composition, and lack of differentiation between adsorption and coprecipitation, the results from this coupled field, experimental, and modeling study underscore the importance of characterizing the mineralogy, redox state, and surface site density of sediments of aquifers that are affected by seasonal and spatial redox transition. Recharge and discharge zones of the floodplain aquifer of South Asia, where seasonal and spatial redox transition is considerable due to groundwater–surface water interaction,^{18,20,22} are likely most susceptible to redox-dependent As partitioning through precipitation/dissolution and transformation of Fe minerals causing the change of the surface sorption site density. The sediment Fe(III)/Fe ratio is a useful parameter to assess the mobility and transport of groundwater As in iron-rich and sulfur-poor aquifers with circumneutral pH, but will not apply in the iron-poor environment where sulfides dominate the mobility of As.^{11,34}

■ ASSOCIATED CONTENT

📄 Supporting Information

Detailed sampling and analysis methods, chemical composition of water and sediment, database for surface complexation modeling, batch sorption and desorption data, sampling location map. This information is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Phone: 718-997-3300; fax: 718-997-3299; e-mail: yan.zheng@qc.cuny.edu, yzheng@ldeo.columbia.edu.

Present Address

§Pacific Northwest National Laboratory, Richland, WA 99354, USA.

■ ACKNOWLEDGMENTS

Funding was provided by NIEHS SBPR 2 P42 ES10349 to Y.Z. H.B.J. received a University Fellowship and Mina Rees Dissertation Fellowship from the Graduate Center, CUNY. We are grateful to Mohammad W. Rahman, Mohammad M. Rahman, and Mohammad Rezaul Huq for assistance in the field.

■ REFERENCES

- (1) Raven, K. P.; Jain, A.; Loeppert, R. H. Arsenite and arsenate adsorption on ferrihydrite: Kinetics, equilibrium, and adsorption envelopes. *Environ. Sci. Technol.* **1998**, *32*, 344–349.
- (2) Goldberg, S. Competitive adsorption of arsenate and arsenite on oxides and clay minerals. *Soil Sci. Soc. Am. J.* **2002**, *66*, 413–421.
- (3) Manning, B. A.; Fendorf, S. E.; Bostick, B.; Suarez, D. L. Arsenic(III) oxidation and arsenic(V) adsorption reactions on synthetic birnessite. *Environ. Sci. Technol.* **2002**, *36*, 976–981.
- (4) Dixit, S.; Hering, J. G. Comparison of arsenic(V) and arsenic(III) sorption onto iron oxide minerals: Implications for arsenic mobility. *Environ. Sci. Technol.* **2003**, *37*, 4182–4189.
- (5) Manning, B. A.; Goldberg, S. Arsenic(III) and arsenic(V) adsorption on three California soils. *Soil Sci.* **1997**, *162*, 886–895.
- (6) Smith, E.; Naidu, R.; Alston, A. M. Chemistry of arsenic in soils: I. Sorption of arsenate and arsenite by four Australian soils. *J. Environ. Qual.* **1999**, *28*, 1719–1726.
- (7) Zhang, H.; Selim, H. M. Kinetics of arsenate adsorption-desorption in soils. *Environ. Sci. Technol.* **2005**, *39*, 6101–6108.

- (8) BGS and DPHE. *Arsenic Contamination of Groundwater in Bangladesh, Final Report*; British Geological Survey Report WC/00/19; Kinniburgh, D. G., Smedley, P. L., Eds.; British Geological Survey: Keyworth, 2001.
- (9) Jung, H. B.; Charette, M. A.; Zheng, Y. Field, Laboratory, and Modeling Study of Reactive Transport of Groundwater Arsenic in a Coastal Aquifer. *Environ. Sci. Technol.* **2009**, *43*, 5333–5338.
- (10) Ford, R. G.; Wilkin, R. T.; Hernandez, G. Arsenic cycling within the water column of a small lake receiving contaminated ground-water discharge. *Chem. Geol.* **2006**, *228*, 137–155.
- (11) O'Day, P. A.; Vlassopoulos, D.; Root, R. A.; Rivera, N. A. The influence of sulfur and iron on dissolved arsenic concentrations in the shallow subsurface under changing redox conditions. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, *101*, 13703–13708.
- (12) Harvey, C. F.; Swartz, C. H.; Badruzzaman, A. B. M.; Keon-Blute, N.; Yu, W.; Ali, M. A.; Jay, J.; Beckie, R.; Niedan, V.; Brabander, D.; Oates, P. M.; Ashfaq, K. N.; Islam, S.; Hemond, H. F.; Ahmed, M. F. Arsenic mobility and groundwater extraction in Bangladesh. *Science* **2002**, *298*, 1602–1606.
- (13) van Geen, A.; Zheng, Y.; Goodbred, S.; Horneman, A.; Aziz, Z.; Cheng, Z.; Stute, M.; Mailloux, B.; Weinman, B.; Hoque, M. A.; Seddique, A. A.; Hossain, M. S.; Chowdhury, S. H.; Ahmed, K. M. Flushing history as a hydrogeological control on the regional distribution of arsenic in shallow groundwater of the Bengal Basin. *Environ. Sci. Technol.* **2008**, *42*, 2283–2288.
- (14) Stollenwerk, K. G.; Breit, G. N.; Welch, A. H.; Yount, J. C.; Whitney, J. W.; Foster, A. L.; Uddin, M. N.; Majumder, R. K.; Ahmed, N. Arsenic attenuation by oxidized aquifer sediments in Bangladesh. *Sci. Total Environ.* **2007**, *379*, 133–150.
- (15) Dhar, R. K.; Zheng, Y.; Saltikov, C. W.; Radloff, K. A.; Mailloux, B. J.; Ahmed, K. M.; van Geen, A. Microbes Enhance Mobility of Arsenic in Pleistocene Aquifer Sand from Bangladesh. *Environ. Sci. Technol.* **2011**, *45*, 2648–2654.
- (16) Radloff, K. A.; Zheng, Y.; Michael, H. A.; Stute, M.; Bostick, B. C.; Mihajlov, I.; Bounds, M.; Huq, M. R.; Choudhury, I.; Rahman, M. W.; Schlosser, P.; Ahmed, K. M.; van Geen, A. Arsenic migration to deep groundwater in Bangladesh influenced by adsorption and water demand. *Nat. Geosci.* **2011**, *4*, 793–798.
- (17) Neumann, R. B.; Ashfaq, K. N.; Badruzzaman, A. B. M.; Ali, M. A.; Shoemaker, J. K.; Harvey, C. F. Anthropogenic influences on groundwater arsenic concentrations in Bangladesh. *Nat. Geosci.* **2010**, *3*, 46–52.
- (18) Datta, S.; Mailloux, B.; Jung, H. B.; Hoque, M. A.; Stute, M.; Ahmed, K. M.; Zheng, Y. Redox trapping of arsenic during groundwater discharge in sediments from the Meghna riverbank in Bangladesh. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 16930–16935.
- (19) Fendorf, S.; Michael, H. A.; van Geen, A. Spatial and Temporal Variations of Groundwater Arsenic in South and Southeast Asia. *Science* **2010**, *328*, 1123–1127.
- (20) Polizzotto, M. L.; Harvey, C. F.; Sutton, S. R.; Fendorf, S. Processes conducive to the release and transport of arsenic into aquifers of Bangladesh. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 18819–18823.
- (21) Postma, D.; Larsen, F.; Minh Hue, N. T.; Duc, M. T.; Viet, P. H.; Nhan, P. Q.; Jessen, S. Arsenic in groundwater of the Red River floodplain, Vietnam: Controlling geochemical processes and reactive transport modeling. *Geochim. Cosmochim. Acta* **2007**, *71*, 5054–5071.
- (22) Polizzotto, M. L.; Kocar, B. D.; Benner, S. G.; Sampson, M.; Fendorf, S. Near-surface wetland sediments as a source of arsenic release to ground water in Asia. *Nature* **2008**, *454*, 505–508.
- (23) O'Day, P. A.; Rivera, N.; Root, R.; Carroll, S. A. X-ray absorption spectroscopic study of Fe reference compounds for the analysis of natural sediments. *Am. Mineral.* **2004**, *89*, 572–585.
- (24) Dzombak, D. A.; Morel, F. M. M. *Surface Complexation Modeling: Hydrous Ferric Oxide*; Wiley-Interscience: New York, 1990.
- (25) Mathur, S. S.; Dzombak, D. A. Surface complexation modeling: goethite. In *Surface Complexation Modeling*; Lutzenkirchen, J., Ed.; Elsevier: Amsterdam, 2006.
- (26) Appelo, C. A. J.; Postma, D. *Geochemistry, Groundwater and Pollution*; A.A. Balkema Publishers: Leiden Netherlands, 2005.
- (27) Zachara, J. M.; Kukkadapu, R. K.; Fredrickson, J. K.; Gorby, Y. A.; Smith, S. C. Biomineralization of poorly crystalline Fe(III) oxides by dissimilatory metal reducing bacteria (DMRB). *Geomicrobiol. J.* **2002**, *19*, 179–207.
- (28) Hansel, C. M.; Benner, S. G.; Neiss, J.; Dohnalkova, A.; Kukkadapu, R. K.; Fendorf, S. Secondary mineralization pathways induced by dissimilatory iron reduction of ferrihydrite under advective flow. *Geochim. Cosmochim. Acta* **2003**, *67*, 2977–2992.
- (29) Cornell, R. M.; Schwertmann, U. *The Fe Oxides: Structure, Properties, Reactions, Occurrences, and Uses*; VCH Weinheim: Federal Republic of Germany, 1996.
- (30) Sverjensky, D. A.; Fukushi, K. A predictive model (ETLM) for As(III) adsorption and surface speciation on oxides consistent with spectroscopic data. *Geochim. Cosmochim. Acta* **2006**, *70*, 3778–3802.
- (31) Berg, M.; Luzi, S.; Trang, P. T. K.; Viet, P. H.; Giger, W.; Stuben, D. Arsenic Removal from Groundwater by Household Sand Filters: Comparative Field study, Model Calculations, and Health Benefits. *Environ. Sci. Technol.* **2006**, *40*, 5567–5573.
- (32) Islam, F. S.; Gault, A. G.; Boothman, C.; Polya, D. A.; Charnock, J. M.; Chatterjee, D.; Lloyd, J. R. Role of metal-reducing bacteria in arsenic release from Bengal delta sediments. *Nature* **2004**, *430*, 68–71.
- (33) Horneman, A.; van Geen, A.; Kent, D. V.; Mathe, P. E.; Zheng, Y.; Dhar, R. K.; O'Connell, S. O.; Hoque, M. A.; Aziz, A.; Shamsuddhau, M.; Seddique, A. A.; Ahmed, K. M. Decoupling of As and Fe release to Bangladesh groundwater under reducing conditions. Part I: Evidence from sediment profiles. *Geochim. Cosmochim. Acta* **2004**, *68*, 3459–3473.
- (34) Keimowitz, A. R.; Zheng, Y.; Chillrud, S. N.; Mailloux, B.; Jung, H. B.; Stute, M.; Simpson, H. J. Arsenic Redistribution between Sediments and Water near a Highly Contaminated Source. *Environ. Sci. Technol.* **2005**, *39*, 8606–8613.