

Posttraumatic Stress Disorder and HIV Risk Among Poor, Inner-City Women Receiving Care in an Emergency Department

Nabila El-Bassel, DSW, Louisa Gilbert, PhD, Danielle Vinocur, PhD, Mingway Chang, PhD, and Elwin Wu, PhD

The relationship between posttraumatic stress disorder (PTSD) and HIV risk behaviors remains relatively underresearched. However, several studies have shown that PTSD is associated with sexual HIV risk behaviors and HIV seropositive status.^{1–3} Emergency departments have been identified as the first and primary source of medical treatment of many women infected with or at high risk for HIV^{4–6} and for those with high rates of interpersonal violence and trauma, including those suffering from PTSD.^{7–11}

Hutton et al. found that, after adjustment for potentially confounding factors, a PTSD diagnosis was associated with engaging in anal intercourse and exchanging sex for money or drugs in a sample of 177 female inmates.¹² High rates of PTSD have also been found among HIV-positive women,^{3,13,14} many of whom have experienced repeated traumas associated with PTSD, such as childhood sexual abuse and intimate partner violence (IPV).^{3,13,14} In a study of HIV-positive women, 35% of those with a trauma history met the criteria for PTSD,¹⁵ a rate far exceeding both the lifetime PTSD rate (10.4%) among women in the general population¹⁶ and the PTSD rate (4.6%) in a nationally representative sample of female crime victims.¹⁷

The relationship between PTSD and HIV risk behaviors has been found to vary according to the presence of different PTSD symptoms (avoidance, hyperarousal, and re-experiencing trauma). In their study of 64 HIV-positive women and men, Gore-Felton and Koopman found that moderate to severe re-experiencing symptoms were associated with multiple sexual partners and unprotected sex during the preceding 3 months.¹⁸ Individuals with avoidant symptoms were less likely to have engaged in unprotected sex, possibly as a result of deficits in establishing and maintaining intimate partnerships.¹⁸ The presence of

Objectives. We examined the associations between posttraumatic stress disorder (PTSD) and HIV risk behaviors among a random sample of 241 low-income women receiving care in an urban emergency department.

Methods. We recruited participants from the emergency department waiting room during randomly selected 6-hour blocks of time. Multivariate analyses and propensity score weighting were used to examine the associations between PTSD and HIV risk after adjustment for potentially confounding sociodemographic variables, substance use, childhood sexual abuse, and intimate partner violence.

Results. A large majority of the sample self-identified as Latina (49%) or African American (44%). Almost one third (29%) of the participants met PTSD criteria. Women who exhibited symptoms in 1 or more PTSD symptom clusters were more likely than women who did not to report having had sex with multiple sexual partners, having had sex with a risky partner, and having experienced partner violence related to condom use in the preceding 6 months.

Conclusions. The high rate of PTSD found in this sample and the significant associations between PTSD symptom clusters and partner-related risk behaviors highlight the need to take PTSD into account when designing HIV prevention interventions for low-income, urban women. (*Am J Public Health.* 2011;101:120–127. doi:10.2105/AJPH.2009.181842)

hyperarousal symptoms may trigger individuals to seek sexual stimulation and engage in riskier sex, and they may experience difficulty in problem solving and negotiating safe sex.¹⁹

The research just described highlights mechanisms of how different PTSD symptom clusters may increase the likelihood of engaging in HIV risks. However, it should also be acknowledged that the relationship may be bidirectional: a traumatic experience (e.g., forced unprotected sex) associated with a risk of HIV may lead to PTSD.

Furthermore, research suggests that the relationship between PTSD and HIV risk may be mediated by several factors, including childhood sexual abuse, IPV, and substance abuse. Those who have experienced childhood sexual abuse are at increased risk of developing PTSD,^{20–22} engaging in subsequent sexual HIV risk behaviors, and HIV transmission.^{3,23} Similarly, IPV has been found to increase the risk of both developing PTSD and engaging in a range

of HIV risk behaviors, including unprotected sex,^{24–38} sexual practices leading to a high risk of sexually transmitted infections,^{6,32,39–42} sex with multiple partners,^{31,32,43} trading of sex for money or drugs,^{40,44} sex with risky partners,^{38,45} and sex with HIV-positive partners.³⁸ Finally, substantial research indicates that drug and alcohol dependencies are associated with both PTSD^{46,47} and engaging in a range of HIV risk behaviors.^{48–51}

We examined the relationship between PTSD (and the symptom clusters of avoidance, re-experiencing trauma, and hyperarousal) and sexual HIV risk behaviors in a random sample of 241 women attending an emergency department in a low-income neighborhood of the Bronx, New York. We hypothesized that women who met the criteria for PTSD and the symptom clusters of hyperarousal, re-experiencing trauma, or avoidance would be more likely than women who did not meet these criteria to engage in sexual HIV risk behaviors

after adjustment and matching for potentially confounding factors such as sociodemographic characteristics, childhood sexual abuse, substance abuse, and IPV.

METHODS

We collected data for this study in a Bronx, New York, hospital emergency department that serves a catchment area of 1.1 million residents. Participants were recruited during randomly selected 6-hour blocks of time from August 2001 through April 2003. In 2002, 18 045 unduplicated female patients aged 18 years or older visited the emergency department (unpublished data from the emergency department's 2002 records). A total of 215 blocks of time were selected as follows: 9 AM to 3 PM, 7 blocks (3%); 3 PM to 9 PM, 118 blocks (55%); 9 PM to 3 AM, 77 blocks (36%); and 3 AM to 9 AM, 13 blocks (6%).

Among the selected blocks, 29% occurred on a weekend. The probability of a specific block being selected was adjusted to match the proportion of patients seen in the emergency department (according to emergency department census data from the preceding year) based on the day of the week and time of day. Of the 6422 female patients admitted to the emergency department during the selected time blocks, 1251 were approached and 452 refused to participate. A sample of 799 women participated in the screening interview, of whom 396 met the study's eligibility criteria. Of these eligible women, 241 (61%) agreed to participate and completed the study questionnaire a week after being screened.

Recruitment procedures were based on success with similar studies in other emergency department settings.^{52–54} Female research assistants, fluent in Spanish and English, attempted to approach every woman who was admitted to the emergency department during the designated time blocks prior to or immediately after medical care. Interested potential participants provided informed consent before completing a face-to-face screening interview in a private room.

To be eligible, patients had to be older than 18 years, female, and admitted to the emergency department during one of the time blocks selected for study inclusion. In addition, they were required to have had a sexual

relationship (at least 3 months in duration) during the preceding year with someone they described as their boyfriend, ex-boyfriend, spouse, ex-spouse, the father of their children, or a regular nonpaying sexual partner, and they had to have had vaginal sex in the preceding 6 months with that partner. Women who were admitted for psychiatric emergencies and women who demonstrated severe cognitive or psychological impairments were excluded because of their inability to provide informed consent.

In the case of moderate or severe triage patients who required hospitalization, screening interviews were conducted within 2 weeks of their hospital admission. Eligible women were asked to participate in the study interview within 10 days of being screened.

Measurement

HIV risk behaviors. We used 2 continuous measures to capture the number of unprotected acts of vaginal or anal sex and the percentage of condom-protected sexual encounters across all partners (i.e., intimate, casual, paying) in the preceding 30 days. Also, we used several dichotomous measures to assess HIV risks in the preceding 6 months, including having had unprotected anal sex, having had multiple sexual partners, having had sex with a risky partner (i.e., a partner with HIV, an injection drug user, or a partner who had sex with multiple partners), having used alcohol or drugs immediately before or during sex, having experienced violence related to condom use (i.e., being forced to have sex without a condom), and having exchanged sex for money or drugs. To facilitate more accurate reporting of sexual behaviors, we used the timeline follow-back method, in which calendar techniques are used to enhance recall.

Posttraumatic stress disorder. We used the PTSD Checklist–Civilian (PCL-C), a 17-item self-report scale, to assess PTSD. Respondents were asked to use a 5-point scale to describe the extent to which they experienced each symptom during the time frame.

The PCL-C mirrors the criteria of the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders*,⁵⁵ which include the following: experiencing or witnessing a stressful life event involving a threat of serious injury or death or a threat to the physical integrity of oneself

or others, a diagnosis of reexperiencing symptoms (indicated by 1 or more of the 5 items of the PCL-C reexperiencing subscale), a diagnosis of avoidance symptoms (indicated by 3 or more of the 7 items of the PCL-C avoidance subscale), a diagnosis of arousal symptoms (indicated by 2 or more of the 5 items of the PCL-C arousal subscale), and symptoms of disturbance. With the exception of disturbance, each symptom was required not to have been present before the trauma; symptoms of disturbance had to have persisted for more than 1 month. The PCL-C has an alpha reliability of 0.97 and has been cross validated with related measures.⁵⁶

Covariates. Our data analyses included covariates that have been found to be associated with PTSD and HIV risk outcomes. In addition to collecting data on sociodemographic characteristics such as age, race/ethnicity, education level, homelessness, employment status in the preceding 6 months, and receipt of public assistance within 12 months of the screening interview, we assessed whether participants had a history of childhood sexual abuse, physical or injurious IPV, sexual IPV, illicit drug use, and heavy episode drinking.

We measured childhood sexual abuse using the Childhood Sexual Abuse Interview.^{57,58} This instrument evaluates history of sexual experiences during childhood and includes 2 subscales assessing touching: exposure (6 items) and penetration (3 items). Participants were classified as having been sexually abused if they reported that someone who was a relative or 5 or more years older than they were had subjected them to either of these forms of abuse before the age of 16 years or if the sexual experience was unwanted or coerced.

We assessed history of IPV using the minor and severe sexual, physical, and injurious subscales of the Revised Conflict Tactics Scales.⁵⁹ Sexual IPV scale items focus on coercive acts intended to engage a partner in unwanted sexual activity. Physical IPV scale items focus on physical assaults perpetrated against the partner, and injurious IPV scale items focus on the consequences of violence. The internal consistencies of the instrument's subscales are high, ranging between 0.79 and 0.95.⁵⁹

Drug use was assessed according to whether the participant reported having used crack, cocaine, heroin (not injected), or marijuana in the preceding 6 months. Participants who

reported having had 4 or more drinks on the same occasion in the preceding 6 months were categorized as having engaged in heavy episode drinking.

Data Analysis

We used multiple imputation via the multivariate imputation by chained equations module in Stata (StataCorp LP, College Station, TX) to reduce the potential bias resulting from missing data for 19 women who refused to answer PTSD questions.^{60–63} Multiple imputation procedures generated 5 imputed data sets to replace each missing value and account for the uncertainty about the missing values.

We used negative binomial regression to examine the associations between PTSD variables and occurrences of unprotected vaginal sex, multiple ordinary least square regression to assess the associations between PTSD variables and proportion of protected vaginal sex, and multiple logistic regression to examine the associations between PTSD and dichotomous HIV riskdependent variables. The adjusted covariates were age, ethnicity, homelessness, employment, years of schooling, heavy episode drinking, drug use, childhood sexual abuse, and history of physical or injurious IPV.

All of the multivariate linear models were estimated with propensity score weights, a well-established method of estimating treatment effects (i.e., the presence of PTSD in this study) in observational studies to reduce selection bias. This approach minimizes the likelihood that pretreatment covariates will confound the treatment effect by selecting groups that are similar on average with respect to all covariate values.^{64–68} The propensity score is the probability that a respondent is assigned to the pretreatment group (e.g., women who exhibited PTSD symptoms) rather than to the comparison group (e.g., women without PTSD symptoms). The propensity score is a 1-dimensional variable that represents the selected pretreatment covariates. Among respondents with a given propensity score, the distribution of these covariates is the same on average in the treatment and comparison groups.

Propensity score models can reduce selection bias in observational studies, especially when the variable of interest was not a key factor in sample selection.^{64–68} In our analyses, the PSMATCH2 (Stata shareware, StataCorp LP,

College Station, TX) module⁶⁹ was used in calculating propensity scores. Also, balance diagnostics were conducted for all covariates to verify that the treatment and comparison groups had similar distributions for the set of covariates.⁷⁰ We used propensity score weighting to reweight treatment and comparison observations so that they would be representative of the population of interest.

RESULTS

Figure 1 shows that 30% of the sample met PTSD criteria. Two thirds of the participants met the criteria for reexperiencing trauma, 40% met the criteria for avoidance symptoms, and 48% met the criteria for hyperarousal symptoms.

Table 1 presents data on sociodemographic characteristics, substance abuse history, and history of violence according to PTSD diagnosis. Participants ranged in age from 18 to 61 years, with a mean of 33 years (SD=10, median=32). The majority of the participants identified themselves as Latina or African American. About half had a high school diploma and had been employed in the preceding 6 months. Fewer than 20% had been homeless in the preceding 6 months. The only significant association found between PTSD criteria and sociodemographic characteristics


was that women who met all of the PTSD criteria were less likely to have been employed in the preceding 6 months than were women who did not meet any of the criteria.

One quarter of the women reported heavy episode drinking, and 43% had used drugs in the preceding 6 months. Women who met PTSD criteria were significantly more likely to report heavy episode drinking and drug use than were their counterparts who did not meet these criteria.

Two fifths of the women reported experiencing childhood sexual abuse. More than two thirds had experienced sexual, physical, or injurious IPV. Women who met PTSD criteria were more likely than those who did not to report experiencing childhood sexual abuse and sexual, physical, or injurious IPV.

HIV Risk Behaviors

On average, participants had used condoms 33% of the time during vaginal sex in the preceding 30 days, and only 25% of the women reported always using condoms during vaginal sex (Table 2). Only 10% reported unprotected anal sex in the preceding 30 days. More than half of the women indicated that they had not used condoms in the preceding 30 days. On average, the participants reported 10 acts of unprotected vaginal sex in the past 30 days. More than two fifths (43%) reported



Note. The sample size was n=241.

FIGURE 1—Posttraumatic stress disorder (PTSD) and symptom clusters.

TABLE 1—Sociodemographic Characteristics, Substance Abuse History, and History of Violence: Low-Income Women Seeking Emergency Care in the Bronx, NY, 2001–2003

	Total (n = 241)	Participants Not Meeting PTSD Criteria (n = 169)	Participants Meeting PTSD Criteria (n = 72)
Sociodemographic characteristics			
Age, y, mean (SD)	33 (10)	33 (10)	33 (10)
Race/ethnicity, no. (%)			
Latina	119 (49)	81 (48)	38 (53)
African American	105 (44)	75 (44)	30 (42)
Other	17 (7)	13 (8)	4 (6)
High school diploma, no. (%)	127 (53)	93 (55)	34 (47)
Employed in past 6 mo, no. (%)	111 (46)	86 (51)	25* (35)
Homeless in past 6 mo, no. (%)	38 (16)	23 (14)	15 (21)
Substance abuse in past 6 mo, no. (%)			
Heavy episode drinking	57 (24)	30 (18)	27** (38)
Illicit drug use	104 (43)	61 (36)	43** (60)
History of violence, no. (%)			
Childhood sexual abuse (before age 16 y)	99 (41)	50 (30)	49** (68)
Lifetime sexual IPV	167 (69)	107 (63)	60** (83)
Lifetime physical or injurious IPV	165 (68)	103 (61)	62** (86)

Note. IPV = intimate partner violence; PTSD = posttraumatic stress disorder.
*P < .05; **P < .01.

that either they or their partners were under the influence of drugs or alcohol during sexual encounters in the preceding 6 months. Almost one third (29%) of the women reported

having had sex with a risky partner. Twenty-two percent reported IPV related to condom use, and 18% reported more than 1 sexual partner in the preceding 6 months. Eight

women reported sex trading in the past 6 months.

Relative to women who did not meet PTSD criteria, those who did were more likely to report multiple sexual partners, having a risky partner, experiencing violence related to condom use, and trading sex in the preceding 6 months. Eight participants (3%) reported being HIV positive.

Multivariate Analyses

Table 3 shows the estimates of the effects of PTSD and the 3 symptom clusters on different HIV risk behaviors after adjustment for the study covariates (sociodemographic characteristics, substance abuse, childhood sexual abuse, and physical or injurious IPV). We estimated each multivariate model with propensity score weights.

The findings from the multivariate analyses showed that those who met PTSD criteria were more likely than those who did not to report having a risky partner (adjusted odds ratio [OR]=2.3) in the preceding 6 months. In comparison with their counterparts, women who met criteria for reexperiencing symptoms were more likely to report having multiple sexual partners (adjusted OR=5.9), having a risky partner (adjusted OR=4.7), and experiencing violence related to condom use (adjusted OR=3.4) in the preceding 6 months.

TABLE 2—Descriptive Statistics for HIV Risk Behaviors and Posttraumatic Stress Disorder (PTSD): Low-Income Women Seeking Emergency Care in the Bronx, NY, 2001–2003

	Total (n = 241)	Participants Not Meeting PTSD Criteria (n = 169)	Participants Meeting PTSD Criteria (n = 72)
No. of unprotected vaginal sex acts with all partners in past 30 d, mean (SD)	10 (15)	10 (15)	10 (16)
Percentage of protected vaginal sex acts with all partners in past 30 d, mean (SD)	33 (43)	33 (43)	35 (44)
Condom use during vaginal sex in past 30 d, no. (%)			
Never	138 (57)	99 (59)	39 (54)
Sometimes	43 (18)	29 (17)	14 (20)
Always	60 (25)	41 (24)	19 (26)
Unprotected anal sex in past 30 d, no. (%)	24 (10)	14 (8)	10 (14)
Multiple sex partners in past 6 mo, no. (%)	43 (18)	21 (12)	22** (31)
Risky partners in past 6 mo, no. (%)	70 (29)	40 (24)	30** (42)
Sex under the influence of drugs or alcohol in past 6 mo, no. (%)	103 (43)	66 (39)	37 (51)
Violence related to condom use in past 6 mo, no. (%)	52 (22)	25 (15)	27** (38)
Sex trading in past 6 mo, no. (%)	8 (3)	2 (1)	6** (8)

**P < .01.

TABLE 3—Associations Between Posttraumatic Stress Disorder (PTSD) and HIV Risk Behaviors: Low-Income Women Seeking Emergency Care in the Bronx, NY, 2001–2003

	No. of Unprotected Vaginal Sex Acts With All Partners ^a		Proportion of Protected Vaginal Sex Acts With All Partners ^a		Unprotected Anal Sex ^a		Multiple Sex Partners ^b		Risky Partners ^b		Violence Related to Condom Use ^b	
	Unadjusted IRR (95% CI)	Adjusted IRR (95% CI)	Unadjusted OLS Estimate (95% CI)	Adjusted OLS Estimate (95% CI)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
Model 1: PTSD	1.0 (0.9, 1.2)	1.0 (0.8, 1.3)	0.03 (-0.1, 0.2)	0.001 (-0.2, 0.2)	1.8 (0.8, 4.3)	1.0 (0.4, 2.7)	3.1** (1.6, 6.2)	2.1 (0.8, 5.5)	2.4** (1.3, 4.4)	2.3* (1.1, 4.8)	3.7** (1.9, 7.0)	2.3 (0.98, 5.2)
Model 2: reexperiencing	1.0 (0.8, 1.1)	0.9 (0.8, 1.1)	0.03 (-0.1, 0.2)	0.1 (-0.1, 0.2)	1.6 (0.6, 4.1)	1.0 (0.3, 3.2)	6.1** (2.1, 17.8)	5.9* (1.4, 24.1)	3.8** (1.8, 8.2)	4.7** (1.9, 12.0)	2.5* (1.1, 5.4)	3.4** (1.4, 8.6)
Model 3: avoidance	1.0 (0.8, 1.2)	1.0 (0.8, 1.1)	0.04 (-0.1, 0.2)	0.01 (-0.1, 0.2)	1.3 (0.6, 3.1)	0.6 (0.2, 1.7)	4.0** (2.0, 8.2)	4.3** (1.5, 12.7)	3.6** (2.0, 6.5)	3.7** (1.7, 7.8)	6.7** (3.3, 13.6)	7.3** (3.1, 17.2)
Model 4: arousal	0.9 (0.8, 1.1)	0.9 (0.8, 1.1)	0.04 (-0.1, 0.2)	0.003 (-0.1, 0.1)	1.9 (0.8, 4.6)	0.8 (0.3, 2.2)	5.4** (2.5, 11.9)	4.4** (1.6, 12.4)	2.5** (1.4, 4.7)	2.3* (1.1, 4.8)	3.1** (1.5, 6.3)	3.6** (1.5, 8.6)

Note. CI = confidence interval; IRR = incidence rate ratio; OLS = ordinary least square; OR = odds ratio. Sample size was n = 241. Values were derived from unadjusted and adjusted models with propensity score weighting. The adjusted covariates were age, ethnicity (Black, Latino, other), homelessness, employment, years of schooling, heavy episode drinking, illicit drug use, childhood sexual abuse, and physical or injurious intimate partner violence. The same covariates were used for propensity score weighting.

^aPast 30 days.
^bPast 6 months.
 *P < .05; **P < .01.

Women with avoidance symptoms were more likely to report having multiple sexual partners (adjusted OR=4.3), having a risky partner (adjusted OR=3.7), and experiencing violence related to condom use (adjusted OR=7.3) than women without these symptoms. Similarly, women with hyperarousal symptoms were more likely than their counterparts to report having multiple sexual partners (adjusted OR=4.4), having a risky partner (adjusted OR=2.3), and experiencing violence related to condom use (adjusted OR=3.6).

PTSD was not associated with the number or proportion of protected vaginal sex acts, having unprotected anal sex, or having sex under the influence of drugs or alcohol. Only 8 women reported sex trading in the preceding 6 months, so this variable was not included in the multivariate analyses.

DISCUSSION

We found remarkably high rates of PTSD in this representative sample of predominantly poor, minority women visiting an urban emergency department; almost one third met full PTSD criteria. We also found low rates of consistent condom use and high rates of HIV risk behaviors.

The multivariate analyses suggest a consistent pattern of associations between PTSD variables and having sex with risky partners. Meeting full PTSD criteria was found to be associated with an increased risk of having sex with a risky partner and experiencing IPV related to condom use. Similarly, the PTSD symptom clusters of hyperarousal, reexperiencing, and avoidance were independently associated with these partner risk factors as well as with participants' reports of having had sex with multiple partners, which is consistent with previous study findings.^{11,2,18} No significant associations were found between PTSD symptoms and condom use variables, contrary to some previous research.¹⁸

Specific mechanisms that may explain the relationships between PTSD and certain HIV risk behaviors remain to be researched. However, several factors may contribute to the increased likelihood of experiencing partner-related risks associated with different PTSD symptom clusters. Symptoms associated with reexperiencing trauma (flashbacks, detachment

from one's own body, emotional unresponsiveness) may result in difficulty in negotiating sexual encounters. As a result, women may acquiesce to a risky partner and have unsafe sex.

Women meeting criteria for the avoidance cluster are more likely to report numbing symptoms, emotional detachment, and feelings of disassociation, all of which may prevent them from establishing and maintaining monogamous intimate relationships and make them prone to engaging in concurrent or short-term serial relationships. They may also be more likely to seek out riskier sexual encounters to counter numbing symptoms. Another consequence of avoidance may be the loss of one's ability to recognize or appropriately respond to sexual risk situations.

Women who meet the criteria for hyperarousal are subject to physiological and emotional dysregulation and may seek sexual stimulation. This can lead to impulsive sexual behaviors and multiple partners. Furthermore, in sexual encounters that trigger additional physical stimulation and chronic arousal, women are likely to have difficulty accessing areas of the brain responsible for the higher order thinking needed to practice safe sex.¹⁹ An alternative explanation for the association between PTSD variables and HIV risk behaviors is that engaging in sex with multiple partners or risky partners increases the likelihood that women will be exposed to trauma that results in PTSD.

Limitations

This study involved several limitations. First, the exclusion of women who were admitted for psychiatric emergencies may have led to underestimations of the rate of PTSD, childhood sexual abuse, and IPV, which have been found to be associated with a range of mental health problems.⁷¹⁻⁷³ Second, the study's cross-sectional design and failure to control for all potentially confounding variables limits our ability to draw conclusions about causal relationships between PTSD and HIV risks. Third, the eligibility inclusion criterion requiring women to have had a minimum 3-month sexual relationship with their partner excluded sexually active women who had sex with only casual or paying partners. High rates of PTSD may be more likely among these women because, as a result of

their riskier encounters, they may experience more sex-related trauma than women in steady relationships.

Fourth, our data were collected between 2001 and 2003, which may limit the present-day generalizability of the findings. However, to our knowledge, this is the first study that has examined the relationship between PTSD and HIV risks in a random sample of poor, urban women seeking emergency department care, and thus it has the potential to address gaps in the research on this issue. Finally, although the use of a random sample that included time blocks encompassing all days of the week and times of day was a strength of the study, the 61% participation rate may limit the generalizability of our findings.

Implications and Conclusions

Despite its limitations, this study has important implications for future research and for HIV prevention interventions with poor, urban women. Future longitudinal research that examines the relationship between PTSD symptomatology and HIV risk behaviors over time is needed to help ascertain the temporal causality between these co-occurring problems.

Further research investigating the specific mechanisms linking different PTSD symptoms and a broad range of risk behaviors would also advance our understanding and help elucidate targeted prevention strategies that are likely to be effective with women who experience PTSD and engage in risky behaviors. Such strategies may synergistically address co-occurring PTSD and HIV risks by raising awareness of the relationship between PTSD symptoms and sexual risk behaviors, assessing patients for PTSD and referring them to clinical treatment services, and developing techniques to avoid PTSD-related triggers for unsafe sex.

Emergency departments remain a primary source of medical care for poor, urban women at risk for HIV as well as for women experiencing PTSD. Low-income women experiencing PTSD may not seek screening or treatment for HIV and sexually transmitted infections elsewhere because of high costs and inadequate access to services. Failure to detect PTSD among emergency department patients may increase the likelihood of repeated emergency department visits, as PTSD has been found to be associated with a broad range of physical

and mental health problems in addition to HIV transmission risks. ■

About the Authors

Nabila El-Bassel, Louisa Gilbert, Mingway Chang, and Elwin Wu are with the Social Intervention Group, Columbia University, New York, NY. Danielle Vinocur is with the Women's Health Project Treatment and Research Center, St. Luke's-Roosevelt Hospital Center, New York.

Correspondence should be sent to Nabila El-Bassel, DSW, Columbia University School of Social Work, 1255 Amsterdam Ave, New York, NY 10027 (e-mail: ne5@columbia.edu). Reprints can be ordered at <http://www.ajph.org> by clicking the "Reprints/Eprints" link.

This article was accepted March 26, 2010.

Contributors

N. El-Bassel wrote the first draft of the article. N. El-Bassel and L. Gilbert conceptualized the article, generated the analysis approach, and revised the final version of the article. D. Vinocur, M. Chang, and E. Wu reviewed and provided feedback on the article. M. Chang analyzed the data.

Acknowledgments

This study was funded by grant 5R01MH05829 from the National Institute of Mental Health to Nabila El-Bassel.

Human Participant Protection

This study was approved by Columbia University's institutional review board. Participants provided written informed consent.

References

1. Stiffman AR, Dore P, Earls F, Cunningham R. The influence of mental health problems on AIDS-related risk behaviors in young adults. *J Nerv Ment Dis*. 1992;180(5):314-320.
2. Wyatt GE, Myers HF, Ashing-Giwa KT, Durvasula RS. Dating patterns: sociocultural factors affecting sexual risk taking in African American men and women—results from two empirical studies. In: Staples R, ed. *The Black Family: Essays and Studies*. Belmont, CA: Wadsworth; 1998:45-57.
3. Wyatt GE, Myers HF, Williams JK, et al. Does a history of trauma contribute to HIV risk for women of color? *Am J Public Health*. 2002;92(4):660-665.
4. Haukoos JS, Hopkins E, Byyny RL. Patient acceptance of rapid HIV testing practices in an urban emergency department: assessment of the 2006 CDC recommendations for HIV screening in health care settings. *Ann Emerg Med*. 2008;51(3):303-309.
5. Rothman RE, Lyons MS, Haukoos JS. Uncovering HIV infection in the emergency department: a broader perspective. *Acad Emerg Med*. 2007;14(7):653-657.
6. El-Bassel N, Gilbert L, Krishnan S, et al. Partner violence and sexual HIV-risk behaviors among women in an inner-city emergency department. *Violence Vict*. 1998;13(4):377-393.
7. Houry D, Kaslow NJ, Kembell R, et al. Does screening in the emergency department hurt or help victims of intimate partner violence? *Ann Emerg Med*. 2008;51(4):433-442.

8. Hurley KF, Brown-Maher T, Campbell SG, Wallace T, Venugopal R, Baggs D. Emergency department patients' opinions of screening for intimate partner violence among women. *Emerg Med J*. 2005;22(2):97–98.
9. MacMillan HL, Wathen CN, Jamieson E, et al. Approaches to screening for intimate partner violence in health care settings: a randomized trial. *JAMA*. 2006; 296(5):530–536.
10. Wathen CN, Jamieson E, Wilson M, Daly M, Worster A, MacMillan HL. Risk indicators to identify intimate partner violence in the emergency department. *Open Med*. 2007;1(2):e113–e122.
11. Lipsky S, Field CA, Caetano R, Larkin GL. Post-traumatic stress disorder symptomatology and comorbid depressive symptoms among abused women referred from emergency department care. *Violence Vict*. 2005; 20(6):645–660.
12. Hutton HE, Treisman GJ, Hunt WR, et al. HIV risk behaviors and their relationship to posttraumatic stress disorder among women prisoners. *Psychiatr Serv*. 2001; 52(4):508–513.
13. Kalichman SC, Sikkema KJ, DiFonzo K, Luke W, Austin J. Emotional adjustment in survivors of sexual assault living with HIV/AIDS. *J Trauma Stress*. 2002; 15(4):289–296.
14. Simoni JM, Ng MT. Trauma, coping, and depression among women with HIV/AIDS in New York City. *AIDS Care*. 2000;12(5):567–580.
15. Kimerling R, Calhoun KS, Forehand R, Armistead L. Traumatic stress in HIV-infected women. *AIDS Educ Prev*. 1999;11(4):321–331.
16. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson C. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1995;52(12):1048–1060.
17. Resnick H, Kilpatrick D, Dansky B, Saunders B, Best C. Prevalence of civilian trauma and post-traumatic stress disorder in a representative national sample of women. *J Consult Clin Psychol*. 1993;61(6):984–991.
18. Gore-Felton C, Koopman C. Traumatic experiences: harbinger of risk behavior among HIV-positive adults. *J Trauma Dissociation*. 2002;3(4):121–135.
19. Hien DA, Nunes E, Levin F, Fraser D. PTSD and short-term outcome in early methadone treatment. *J Subst Abuse Treat*. 2000;19(1):31–37.
20. Classen C, Nevo R, Koopman C, et al. Recent stressful life events, sexual revictimization, and their relationship with traumatic stress symptoms among women sexually abused in childhood. *J Interpers Violence*. 2002;17(12):1274–1290.
21. Twaite JA, Rodriguez-Srednicki O. Childhood sexual and physical abuse and adult vulnerability to PTSD: the mediating effects of attachment and dissociation. *J Child Sex Abuse*. 2004;13(1):17–38.
22. Widom CS. Posttraumatic stress disorder in abused and neglected children grown up. *Am J Psychiatry*. 1999; 156(8):1223–1229.
23. Greenberg JB. Childhood sexual abuse and sexually transmitted diseases in adults: a review of and implications for STD/HIV programmes. *Int J STD AIDS*. 2001; 12(12):777–783.
24. Wingood GM, DiClemente RJ. The effects of an abusive primary partner on the condom use and sexual negotiation practices of African-American women. *Am J Public Health*. 1997;87(6):1016–1018.
25. Amaro H. Love, sex and power: considering women's realities in HIV prevention. *Am Psychol*. 1995; 50(6):437–447.
26. Fernandez MI. Latinas and AIDS: challenges to HIV prevention efforts. In: O'Leary A, Jemmott LS, eds. *AIDS Prevention and Mental Health*. New York, NY: Plenum Press; 1995:159–174.
27. Cunningham RM, Stiffman AR, Dore P, Earls F. The association of physical and sexual abuse with HIV risk behaviors in adolescence and young adulthood: implications for public health. *Child Abuse Negl*. 1994;18(3): 233–245.
28. Wyatt GE. Child sexual abuse and its effects on sexual functioning. *Annu Rev Sex Res*. 1991;2:249–266.
29. Amaro H, Fried LE, Cabral H, Zuckerman B. Violence during pregnancy and substance abuse. *Am J Public Health*. 1990;80(5):575–579.
30. Worth D. Sexual decision-making and AIDS: why condom promotion among vulnerable women is likely to fail. *Stud Fam Plann*. 1989;20(6):297–307.
31. Gilbert L, El-Bassel N, Rajah V, et al. The converging epidemics of mood-altering-drug use, HIV, HCV, and partner violence: a conundrum for methadone maintenance treatment. *Mt Sinai J Med*. 2000;67(5–6):452–464.
32. Wu E, El-Bassel N, Witte SS, Gilbert L, Chang M. Intimate partner violence and HIV risk among urban minority women in primary health care settings. *AIDS Behav*. 2003;7(3):291–301.
33. Tucker JS, Wenzel SL, Elliott MN, Marshall GN, Williamson S. Interpersonal violence, substance use, HIV-related behavior and cognitions: a prospective study of impoverished women in Los Angeles County. *AIDS Behav*. 2004;8(4):463–474.
34. Hamburger ME, Moore J, Koenig LJ, et al. Persistence of inconsistent condom use: relation to abuse history and HIV serostatus. *AIDS Behav*. 2004;8(3): 333–344.
35. Bassuk EL, Buckner JC, Weinreb LF, et al. Homelessness in female-headed families: childhood and adult risk and protective factors. *Am J Public Health*. 1997; 87(2):241–248.
36. Gielen AC, McDonnell KA, O'Campo PJ. Intimate partner violence, HIV status, and sexual risk reduction. *AIDS Behav*. 2002;6(2):107–116.
37. Saul J, Moore J, Murphy ST, Miller LC. Relationship violence and women's reactions to male- and female-controlled HIV prevention methods. *AIDS Behav*. 2004; 8(2):207–214.
38. El-Bassel N, Gilbert L, Wu E, Go H, Hill J. HIV and intimate partner violence among women on methadone. *Soc Sci Med*. 2005;61(1):171–183.
39. Gilbert L, El-Bassel N, Schilling RF, Catan V, Wada T. Partner abuse and sexual risk behavior among women on methadone. *Violence Vict*. 1998;13(4):1–17.
40. El-Bassel N, Gilbert L, Schilling RF, Wada T. Drug abuse and partner violence among women in methadone treatment. *J Fam Violence*. 2000;15(3):209–225.
41. Rodriguez MA, Szkupinski Quiroga S, Bauer HM. Breaking the silence: battered women's perspective on medical care. *Arch Fam Med*. 1996;5(3):153–158.
42. Hogben M, Gange SJ, Watts DH, Robison E, Young M, Richardson J. The effect of sexual and physical violence on risky sexual behavior and STDs among a cohort of HIV seropositive women. *AIDS Behav*. 2001; 5(4):353–361.
43. Rich JD, Dickinson BP, Macalino G, et al. Prevalence and incidence of HIV among incarcerated and re-incarcerated women in Rhode Island. *J Acquir Immune Defic Syndr*. 1999;22(2):161–166.
44. Beadnell B, Baker SA, Morrison DM, Knox K. HIV/STD risk factors for women with violent male partners. *Sex Roles*. 2000;42(7–8):661–690.
45. Gilbert L, El-Bassel N, Schilling R, Wada T, Bennet B. Partner violence and sexual HIV risk behaviors among women in methadone treatment. *AIDS Behav*. 2000; 4(3):261–269.
46. Dansky BS, Brady KT, Saladin ME. Untreated symptoms of PTSD among cocaine-dependent individuals: changes over time. *J Subst Abuse Treat*. 1998;15(6): 499–504.
47. Najavits L, Weiss R, Shaw S. The link between substance abuse and posttraumatic stress disorder in women: a research review. *Am J Addict*. 1997;6(4):273–283.
48. Adimora AA, Schoenbach VJ, Martinson FEA, et al. Heterosexually transmitted HIV infection among African Americans in North Carolina. *J Acquir Immune Defic Syndr*. 2006;41(5):616–623.
49. Miller M, Liao Y, Gomez AM, Gaydos CA, D'Mellow D. Factors associated with the prevalence and incidence of *Trichomonas vaginalis* infection among African American women in New York City. *J Infect Dis*. 2008; 197(4):503–509.
50. Plitt SS, Garfein RS, Gaydos CA, Strathdee SA, Sherman SG, Taha TE. Prevalence and correlates of *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, *Trichomonas vaginalis* infections, and bacterial vaginosis among a cohort of young injecting drug users in Baltimore, Maryland. *Sex Transm Dis*. 2005;32(7): 446–453.
51. Edlin BR, Irwin KL, Faruque S, et al. Intersecting epidemics: crack cocaine use and HIV infection in inner-city young adults. *N Engl J Med*. 1994;331(21):1422–1427.
52. Abbott J, Johnson R, Koziol-McLain J, Lowenstein SR. Domestic violence against women: incidence and prevalence in an emergency department population. *JAMA*. 1995;273(22):1763–1767.
53. Roberts GL, O'Toole BI, Lawrence JM, Raphael B. Domestic violence victims in a hospital emergency department. *Med J Aust*. 1993;159(5):307–310.
54. Roberts GL, O'Toole BI, Raphael B, Lawrence JM, Ashby R. Prevalence study of domestic violence victims in an emergency department. *Ann Emerg Med*. 1996; 27(6):741–753.
55. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*. Washington, DC: American Psychiatric Association; 1994.
56. Weathers FW, Blake DD, Krinsky DE, Haddad W, Huska JA, Keane TM. The Clinician-Administered PTSD Scale: reliability and construct validity. Paper presented at: annual meeting of Association for Advancement of Behavior Therapy, November 1992, Boston, MA.
57. Finkelhor D. *Sexually Victimized Children*. New York, NY: Free Press; 1979.
58. Sgroi SM. *Handbook of Clinical Intervention in Child Sexual Abuse*. Lexington, MA: DC Heath & Co; 1982.

59. Straus MA, Hamby SL, Boney-McCoy S, Sugarman DB. The revised Conflict Tactics Scales (CTS2): development and preliminary psychometric data. *J Fam Issues*. 1996;17(3):283–316.
60. Rubin DB. *Multiple Imputation for Nonresponse in Surveys*. New York, NY: John Wiley & Sons Inc; 1987.
61. Schafer JL. *Analysis of Incomplete Multivariate Data*. London, England: Chapman & Hall; 1997.
62. Royston P. Multiple imputation of missing values. *Stata J*. 2004;4(3):227–241.
63. van Buuren S, Boshuizen HC, Knook DL. Multiple imputation of missing blood pressure covariates in survival analysis. *Stat Med*. 1999;18(6):681–694.
64. Connors A, Speroff T, Dawson N, et al. The effectiveness of right heart catheterization in the initial care of critically ill patients. *JAMA*. 1996;276(11):889–897.
65. Dehejia R, Wahba S. Causal effects in non-experimental studies: re-evaluating the evaluation of training programs. *J Am Stat Assoc*. 1999;94(448):1053–1062.
66. Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika*. 1983;70(1):41–55.
67. Rosenbaum PR, Rubin DB. Constructing a control group using multivariate matched sampling methods that incorporate the propensity score. *Am Stat*. 1985;39(1):33–38.
68. Stenestrånd U, Wallentin L. Early statin treatment following acute myocardial infarction and one-year survival. *JAMA*. 2001;285(4):430–436.
69. Leuven E, Sianesi B. *PSMATCH2: Stata Module to Perform Full Mahalanobis and Propensity Score Matching, Common Support Graphing, and Covariate Imbalance Testing*. Boston, MA: Boston College Dept of Economics; 2003.
70. Berger LM, Hill J, Waldfogel J. Maternity leave, early maternal employment and child health and development in the US. *Econ J*. 2005;115(501):F29–F47.
71. Schiff M, El-Bassel N, Engstrom M, Gilbert L. Psychological distress and intimate physical and sexual abuse among women in methadone maintenance treatment programs. *Soc Serv Rev*. 2002;76(2):302–320.
72. Wathen CN, MacMillan HL. Interventions for violence against women: scientific review. *JAMA*. 2003;289(5):589–600.
73. Dansky BS, Byrne CA, Brady KT. Intimate violence and post-traumatic stress disorder among individuals with cocaine dependence. *Am J Drug Alcohol Abuse*. 1999;25(2):257–268.