

Brief Reports

Harmaline-Induced Tremor in Mice: Videotape Documentation and Open Questions About the Model

Melody M. Cheng¹, Guomei Tang¹ & Sheng-Han Kuo^{1*}

Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, New York, United States of America

Abstract

Background: Harmaline-induced tremor in rodents has been extensively used as an animal model for essential tremor (ET). However, there is no visual documentation in the published literature.

Methods: We injected mice subcutaneously with either 20 mg/kg of harmaline hydrochloride or saline and then videotaped the responses.

Results: Action and postural tremor in the mouse began 5 minutes after subcutaneous harmaline injection and peaked at approximately 30 minutes. The tremor involved the head, trunk, tail, and four limbs and lasted for approximately 2 hours. The forelimb tremor was postural or action tremor, similar to that observed in ET.

Discussion: This video segment provides the first visual documentation of the phenomenology of harmaline-induced tremor in a mouse. We also raise several unanswered questions regarding the use of harmaline-induced tremor to model ET.

Keywords: Essential tremor, harmaline, mouse, cerebellum, inferior olive

Citation: Cheng MM, Tang GM, Kuo SH. Harmaline-induced tremor in mice: videotape documentation and open questions about the model. Tremor Other Hyperkinet Mov 2013; 3: <http://tremorjournal.org/article/view/205>

*To whom correspondence should be addressed. E-mail: sk3295@columbia.edu

Editor: Elan D. Louis, Columbia University, United States of America

Received: October 8, 2013 **Accepted:** October 25, 2013 **Published:** December 9, 2013

Copyright: © 2013 Cheng et al. This is an open-access article distributed under the terms of the Creative Commons Attribution–Noncommercial–No Derivatives License, which permits the user to copy, distribute, and transmit the work provided that the original author(s) and source are credited; that no commercial use is made of the work; and that the work is not altered or transformed.

Funding: Dr. Kuo has received funding from an NINDS grant (K08 NS08738, principal investigator), the Louis V. Gerstner Jr. Scholar Award, the Parkinson's Disease Foundation, an American Academy of Neurology Research Fellowship, and the American Parkinson's Disease Association.

Financial disclosure: None.

Conflict of interest: The authors report no conflict of interest.

Introduction

Essential tremor (ET) is the most common type of tremor, characterized by action and postural tremor in the upper extremities and/or neck and voice tremor.¹ The underlying mechanisms of ET have been extensively investigated. Genetic factors are important in ET, as it is common in kindreds.² However, the concordance in monozygotic twins is only about 60–63%,³ indicating that environmental factors also play a role in ET.⁴

β -carboline alkaloids (BCAs), including norharman, harmine, harmaline, and ibogaine, are well known to induce action and postural tremor in mice, rats, rabbits, cats, and monkeys.^{5–9} Exposure to exogenous BCAs seems to be associated with ET, as ET patients have higher levels of harmine in the blood and brain as compared to healthy controls.^{10–15} Harmaline can induce tremor by the mechanism of synchronized firing in the inferior olivary nucleus (IO), leading to rhythmic activity within the olivocerebellar system, which has been hypothesized by some to be the

anatomical substrate of ET.¹⁶ Harmaline-induced tremor in animals is predominantly postural and action at the frequency of 8–14 Hz, similar to tremor in ET patients.^{17,18} Furthermore, treatments for ET, such as β -blockers and primidone, can dampen harmaline-induced tremor.¹⁸ These findings suggest that harmaline-induced rodent tremor models might be useful to investigate the mechanisms of ET.

Despite extensive research on harmaline-induced tremor in rodents, visual documentation of this tremor in published literature is lacking. The goal of the current study was to 1) provide video of harmaline-induced tremor in a mouse, 2) describe in detail the phenomenology of harmaline-induced tremor, 3) raise several unanswered questions regarding the use of harmaline-induced tremor to model ET.

Methods

We subcutaneously injected C57/BL6 mice with either harmaline 20 mg/kg or saline and observed tremor at rest, posture, and action.



Video 1. Harmaline-induced Tremor. Subcutaneous administration of 20 mg/kg harmaline induced whole-body tremor in a mouse, whereas the saline-injected control mouse had no tremor. The harmaline-treated mouse exhibited action tremor in the body and tail during locomotion, as well as forelimb postural and action tremor. The harmaline-treated mouse also had intermittent rest tremor. The control mouse did not have any action tremor or forelimb tremors.

Results

Five minutes after subcutaneous harmaline administration, we observed tremors at approximately 10–16 Hz, involving the head, trunk, and tail (Video). The tremor became more apparent during locomotion and occurred only intermittently at rest. The tremor lasted for approximately 2 hours. A head tremor was noted on elevating the head and was mainly seen as a vertical motion. The tail tremor was also mostly vertical. The forelimb tremor was seen when the animal reared and raised its forepaw. The hindlimbs also had tremor during locomotion. Control mice did not have any apparent tremors (Video). In addition, harmaline also caused slow locomotion, but these symptoms disappeared when the tremor resolved.

Discussion

We present the first visual documentation of harmaline-induced tremor in mice. We observed postural and action tremors, as previously reported.^{8,17} In addition, we observed prominent forelimb postural tremors in harmaline-treated mice, which have not been described in detail.

Acute harmaline-induced tremors in rodents are a very useful animal model for ET research as they recapitulate disease phenotypes, and this report further supports this notion. Acute harmaline or ibogaine exposure in rats can also induce cerebellar Purkinje cell (PC) loss,^{8,20} which is a major pathological feature of ET pathology.^{21,22}

Harmaline and other BCAs can cause synchronization of rhythms in the IO and climbing fibers, which subsequently leads to PC dysfunction. ET, along with action tremors due to other causes, such as post-traumatic tremor, tremor of hyperthyroidism, and valproate-induced tremor, may share common anatomical tremor circuits with

harmaline-induced tremor, and thus have similar pharmacological responses to treatment.¹⁹ We observed mild unsteadiness in a harmaline-treated mouse during the period of tremor. However, whether harmaline can cause ataxic gait in mice requires further investigation. In addition, harmaline-treated mice also exhibited slow movements and locomotion, but these resolved with the resolution of tremor and were likely a secondary response to the tremor rather than bradykinesia per se. Thus, it is more likely that slowness is an aversive response to action and postural tremor¹⁷ rather than a direct effect of harmaline on the basal ganglia.

There are several unanswered questions with regard to the use of harmaline-induced tremor to model ET. First, several postmortem studies found structural changes in the cerebellum, such as PC axonal torpedoes²² and other associated axonal pathology,²³ basket cell axonal process changes,^{24,25} and heterotopic PCs.²⁶ Acute effects of harmaline are unlikely to recapitulate these chronic pathological features of ET, although this remains to be determined. Second, the chronic effects of harmaline in terms of tremor characteristics and duration are poorly understood. Chronic administration of harmaline or other BCAs might be more useful in studying ET, especially for screening treatments that could prevent or reverse the pathological features of ET. The chronic effects of harmaline in mice have not been extensively investigated, and establishing a chronic harmaline tremor mouse model might be an important future research direction.

References

1. Louis ED. Clinical practice. Essential tremor. *N Engl J Med* 2001;345:887–891, doi: <http://dx.doi.org/10.1056/NEJMcp010928>.
2. Deng H, Le W, Jankovic J. Genetics of essential tremor. *Brain* 2007;130:1456–1464, doi: <http://dx.doi.org/10.1093/brain/awm018>.
3. Tanner CM, Goldman SM, Lyons KE, et al. Essential tremor in twins: an assessment of genetic vs environmental determinants of etiology. *Neurology* 2001;57:1389–1391, doi: <http://dx.doi.org/10.1212/WNL.57.8.1389>.
4. Louis ED. Environmental epidemiology of essential tremor. *Neuroepidemiology* 2008;31:139–149, doi: <http://dx.doi.org/10.1159/000151523>.
5. Du W, Aloyo VJ, Harvey JA. Harmaline competitively inhibits [3H]MK-801 binding to the NMDA receptor in rabbit brain. *Brain Res* 1997;770:26–29, doi: [http://dx.doi.org/10.1016/S0006-8993\(97\)00606-9](http://dx.doi.org/10.1016/S0006-8993(97)00606-9).
6. De Montigny C, Lamarre Y. Effects produced by local applications of harmaline in the inferior olive. *Can J Physiol Pharmacol* 1975;53:845–849, doi: <http://dx.doi.org/10.1139/y75-116>.
7. Bergström M, Westerberg G, Kihlberg T, Långström B. Synthesis of some 11C-labelled MAO-A inhibitors and their in vivo uptake kinetics in rhesus monkey brain. *Nucl Med Biol* 1997;24:381–388.
8. Miwa H, Kubo T, Suzuki A, Kihira T, Kondo T. A species-specific difference in the effects of harmaline on the rodent olivocerebellar system. *Brain Res* 2006;1068:94–101, doi: <http://dx.doi.org/10.1016/j.brainres.2005.11.036>.
9. Zetler G, Singbartl G, Schlosser L. Cerebral pharmacokinetics of tremor-producing harmala and iboga alkaloids. *Pharmacology* 1972;7:237–248, doi: <http://dx.doi.org/10.1159/000136294>.
10. Louis ED, Benito-León J, Moreno-García S, et al. Blood harmaline (1-methyl-9H-pyrido[3,4-b]indole) concentration in essential tremor cases in

- Spain. *Neurotoxicology* 2013;34:264–268, doi: <http://dx.doi.org/10.1016/j.neuro.2012.09.004>.
11. Louis ED, Zheng W, Applegate L, Shi L, Factor-Litvak P. Blood harmaline concentrations and dietary protein consumption in essential tremor. *Neurology* 2005;65:391–396, doi: <http://dx.doi.org/10.1212/01.wnl.0000172352.88359.2d>.
 12. Louis ED, Jiang W, Pellegrino KM, et al. Elevated blood harmaline (1-methyl-9H-pyrido[3,4-b]indole) concentrations in essential tremor. *Neurotoxicology* 2008;29:294–300, doi: <http://dx.doi.org/10.1016/j.neuro.2007.12.001>.
 13. Louis ED, Factor-Litvak P, Liu X, et al. Elevated brain harmaline (1-methyl-9H-pyrido[3,4-b]indole) in essential tremor cases vs. controls. *Neurotoxicology* 2013;38:131–135, doi: <http://dx.doi.org/10.1016/j.neuro.2013.07.002>.
 14. Louis ED, Zheng W, Jurewicz EC, et al. Elevation of blood beta-carboline alkaloids in essential tremor. *Neurology* 2002;59:1940–1944, doi: <http://dx.doi.org/10.1212/01.WNL.0000038385.60538.19>.
 15. Louis ED, Rios E, Pellegrino KM, et al. Higher blood harmaline (1-methyl-9H-pyrido[3,4-b]indole) concentrations correlate with lower olfactory scores in essential tremor. *Neurotoxicology* 2008;29:460–465, doi: <http://dx.doi.org/10.1016/j.neuro.2008.02.013>.
 16. Llinás R, Yarom Y. Oscillatory properties of guinea-pig inferior olivary neurones and their pharmacological modulation: an in vitro study. *J. Physiol* 1986;376:163–182.
 17. Miwa H. Rodent models of tremor. *Cerebellum* 2007;6:66–72, doi: <http://dx.doi.org/10.1080/14734220601016080>.
 18. Martin FC, Thu Le A, Handforth A. Harmaline-induced tremor as a potential preclinical screening method for essential tremor medications. *Mov Disord* 2005;20:298–305, doi: <http://dx.doi.org/10.1002/mds.20331>.
 19. Elble RJ. Tremor disorders. *Curr Opin Neurol* 2013;26:413–419, doi: <http://dx.doi.org/10.1097/WCO.0b013e3283632f46>.
 20. O’Hearn E, Molliver ME. The olivocerebellar projection mediates ibogaine-induced degeneration of Purkinje cells: a model of indirect, trans-synaptic excitotoxicity. *J Neurosci* 1997;17:8828–8841.
 21. Axelrad JE, Louis ED, Honig LS, et al. Reduced Purkinje cell number in essential tremor: a postmortem study. *Arch Neurol* 2008;65:101–107, doi: <http://dx.doi.org/10.1001/archneurol.2007.8>.
 22. Louis ED, Faust PL, Vonsattel JP, et al. Neuropathological changes in essential tremor: 33 cases compared with 21 controls. *Brain* 2007;130:3297–3307, doi: <http://dx.doi.org/10.1093/brain/awm266>.
 23. Babij R, Lee M, Cortes E, et al. Purkinje cell axonal anatomy: quantifying morphometric changes in essential tremor versus control brains. *Brain* 2013;136:3051–3061, doi: <http://dx.doi.org/10.1093/brain/awt238>.
 24. Erickson-Davis CR, Faust PL, Vonsattel JP, Gupta S, Honig LS, Louis ED. “Hairy baskets” associated with degenerative Purkinje cell changes in essential tremor. *J Neuropathol Exp Neurol* 2010;69:262–271, doi: <http://dx.doi.org/10.1097/NEN.0b013e3181d1ad04>.
 25. Kuo SH, Tang G, Louis ED, et al. Lingo-1 expression is increased in essential tremor cerebellum and is present in the basket cell pinceau. *Acta Neuropathol* 2013;125:879–889, doi: <http://dx.doi.org/10.1007/s00401-013-1108-7>.
 26. Kuo SH, Erickson-Davis C, Gillman A, Faust PL, Vonsattel JP, Louis ED. Increased number of heterotopic Purkinje cells in essential tremor. *J Neurol Neurosurg Psychiatry* 2011;82:1038–1040, doi: <http://dx.doi.org/10.1136/jnnp.2010.213330>.