

SYNTHESIS

Open Access

# Climate and health in Africa

Madeleine C Thomson<sup>1,2\*</sup>, Simon Mason<sup>1</sup>, Barbara Platzer<sup>3</sup>, Abere Mihretie<sup>4</sup>, Judy Omumbo<sup>5</sup>, Gilma Mantilla<sup>1,6</sup>, Pietro Ceccato<sup>1</sup>, Michel Jancloes<sup>7</sup> and Stephen Connor<sup>8</sup>

## Abstract

This paper describes the work of the International Research Institute for Climate and Society (IRI) and its partners towards the development of climate services for the health sector in Africa; integrating research, operational applications and capacity building alongside policy development and advocacy. It follows the evolution of IRI's health work from an initial focus on the use of seasonal climate forecasts to a wider agenda serving climate and environmental information needs to a broad range of health-related users. Recognizing that climate information must be relevant to the priority policy and programming needs of national and international health stakeholders, this review highlights an approach that has centered not only on the assessment and creation of evidence, but also on knowledge transfer through engagement with decision-makers. Current opportunities and priorities identified for the routine use of climate and environmental information in health in Africa include: i) understanding mechanisms by which climate impacts on transmission and occurrence of disease; ii) mapping populations at risk both in space and by season; iii) developing early warning systems; iv) understanding the contributions of climate to trends in disease incidence v) improving the evaluation of the impacts of climate-sensitive interventions.

While traditional metrics (e.g. peer review publications) have been important in establishing evidence for policy, the IRI's role as a knowledge broker (in research and professional capacity building, facilitation of communities of practice, and engagement in policy dialogue at local and global scale) has been critical to delivery of its mission.

**Keywords:** Climate; Health; Africa; Climate services; Risk Management; Interdisciplinary; Adaptation; Early Warning System

## Introduction

Although Sub-Saharan Africa is rapidly urbanizing, the continent remains substantially reliant on rain-fed agriculture and surface water resources (McCartney et al. 2013). Despite significant economic growth in several countries, there is still a high burden of climate-sensitive diseases, food insecurity and hydro-meteorological disasters (WHO-AFRO 2011). Health and well-being therefore remain highly climate dependent.

Health has been an important focus of IRI activities through much of the institute's history. In 1999 IRI supported its first workshop on "Climate Prediction and Diseases/Health", which took place in Bamako, Mali in

March/April 1999 (Tourre 1999). The objectives of the Bamako workshop were:

- To expose professionals of the climate and health sectors in Africa to state-of-the-art climate and health science.
- To explore the linkages between physical and biological mechanisms of disease transmission and epidemics relative to climate variability;
- To foster the application of regional climate prediction to preparedness and decision-making processes in the health sector;
- To provide cross-fertilization, exchange of ideas, expertise and analytical tools between climate and health researchers from other countries and continents.

\* Correspondence: mthomson@iri.columbia.edu

<sup>1</sup>International Research Institute for Climate and Society, PO Box 1000, Palisades, New York 10964, USA

<sup>2</sup>Mailman School of Public Health, Columbia University, New York, USA

Full list of author information is available at the end of the article

In this paper, we outline key developments and lessons learned in the health programme at IRI since the Bamako workshop.

### Development of a climate and health strategy

The initial emphasis at the IRI on *early warning systems* was a logical beginning for its health work as this emphasis provided a natural opportunity to introduce IRI's core strength in seasonal climate forecasts (Mason *et al.* 1999) into vector borne disease control planning. The approach taken was to drive biological models with global climate data (Hopp and Foley 2001) and then to compare the results with disease incidence (Hopp and Foley 2003). While this work provided an important starting point for the development of forecasting models, the models did not incorporate much of what is known about disease transmission dynamics (Morin *et al.* 2013) and were not focused on the operational needs of the health sector.

In order to better respond to potential user needs, the strategy adopted for the IRI's health programme in Africa since 2002 has been to align activities with the political, technical and economic priorities of the international health community. The overarching goal established was "to increase the well-being of people in Africa by reducing vulnerability to adverse climate impacts". The purpose was "to create the knowledge and capacity, and thus the opportunity for health organizations and their partners to predict, prevent and manage adverse climate-influenced health outcomes." To realise this knowledge and capacity, efforts were focused on diseases of major public health significance that impact the poor across the differing disease ecological strata. The choice of efforts was based on their potential to impact health policy rather than scientific curiosity. A set of *verifiable indicators* was created that included i) establishment of a coalition of partners who are able to identify and commission operational research activities ii) the provision of innovative mechanisms for health organizations to routinely access and use new knowledge to anticipate, prevent and manage climate-related adverse health outcomes; and iii) the creation of cadres of researchers focused on health – climate issues.

The following were identified as *key intermediary indicators* i) evidence on the role of climate in driving the multiple spatial and temporal determinants of health outcomes, which may include: infectious disease transmission, food security, livelihoods, poverty, disasters, socio-economic development; ii) the creation of tailored climate/environmental products that can be used both for research and operational activities, including near real-time monitoring products and downscaled climate forecasts; iii) research indicating the economic value of the use of tailored climate/environmental information in national, regional health decision-making in Africa.

An important part of the strategy to engage the global health community was the creation, in 2004, of the IRI-Pan American Health Organization-World Health Organization Collaborating Center (US 306) for "Early Warning Systems for Malaria and other Climate Sensitive Diseases." Since its inception, the collaborating center has provided a formal link with this key UN health agency. It maintains active engagement with the WHO/PAHO's Regional Offices in Africa and the Americas as well as with headquarters in Geneva, where the sponsoring department is the Global Malaria Programme (GMP).

This new strategy resulted in an explicit focus on the achievement of the Millennium Development Goals (MDGs) (UN 2013). As the MDGs became the focus of international development, scientists at IRI were concerned that climate variability and change might pose a risk to their achievement. As a result malaria was prioritized at IRI because control of this climate sensitive disease was seen as essential to the achievement of the MDGs (Gallup and Sachs 2001). The report "Sustainable Development in Africa – Is the Climate Right?" (IRI 2005) was designed to alert high-level policy-makers, involved in the Group of Eight (G8) summit at Gleneagles in 2005, to the challenge posed by climate variability and change. The report's specific objective was to link two expected outcomes of the G8 agenda: to increase overseas development assistance in support of MDG achievements in Africa, and to strengthen the continent's climate observing systems (G8 2005).

Following the IRI 2005 report, the UK Department for International Development (DFID) commissioned the Institute to undertake a "Gap Analysis for the Implementation of a Global Climate Observing System (GCOS) program in Africa" (IRI 2006). The objective was to better understand how improvements in climate data and services could have a positive impact on development outcomes – particularly on the MDGs. The "Gap Analysis" identified four major areas requiring investment in order to overcome barriers to the effective use of climate information in national decision-making for development. These were i) policy; ii) practice; iii) services; and iv) data. In short, to use climate information effectively policies need to be flexible in order to respond to changes in climatic risk, practitioners need to be able to use policy relevant climate information when and where cost-effective, the providers of climate information need to provide reliable, relevant, accurate and timely information, and the underlying observational data need to be quality-controlled, analysed and readily accessible (Connor *et al.* 2006).

The "Gap Analysis" was presented at a major stakeholder meeting in Addis Ababa in 2006 that launched Climate for Development in Africa (ClimDev-Africa); a multilateral initiative overseen by a Secretariat comprised of the African Union, the United Nations Economic

Commission for Africa (UNECA) and the African Development Bank. Following the Addis Ababa meeting, the IRI was invited to write its first *Climate and Society Report* to showcase examples of the effective use of climate information in development decision-making in Africa, including health (Hellmuth et al. 2007). One of the case studies focused on the development of Malaria Early Warning Systems (MEWS) in Botswana. Because much of the initiative originated from the regional health partners, this case study demonstrated the value of collaborations between climate and health sectors in the development of the MEWS.

A prerequisite to the effective use of new (in this case climate) information in health decision-making is the role of evidence (Kula et al. 2013). Evidence needs to be gathered in relation to two distinct issues. The first is the need to establish the degree to which climate is a significant driver of specific health-related outcomes (Kelly-Hope and Thomson 2008), and the second is the need to establish the utility (including cost-effectiveness) of using climate information in improving such outcomes. Demonstrating cost-effectiveness is easiest to do when a climate-informed intervention is compared with no such intervention (Worrall et al. 2008); it is much more challenging to compare the impact of a climate-informed intervention with a non-climate related intervention.

In a multi-disciplinary environment, agreeing on what constitutes evidence is not always simple since each community may approach the creation and acceptance of evidence with diverse criteria and perspectives. Getting evidence into policy and practice in the health sector is a widely identified challenge, both in the developed world (Nutley and Davies 2000; Lomas 2007) and globally (WHO 2004). Our aim has been to create policy-relevant evidence and transfer results to operational decision-making in both communities. We have used a number of approaches for knowledge transfer or 'brokering' (Ward et al. 2009) including traditional approaches for the development and diffusion of knowledge (publications, reports, presentations, conferences), fostering links between the producers and users of information (communities of practice, coalitions of partners) and targeted research and professional capacity building activities (multi-disciplinary training).

As indicated above, establishing and strengthening the evidence base for the role of climate as a significant driver of health outcomes has been a central part of IRI health work. For example, understanding the basic mechanism whereby climate affects infectious disease transmission is important for an improved scientific understanding of the disease dynamics and better targeting of interventions. In the context of malaria, the mechanisms (i.e. influences of rainfall and temperature on vector dynamics, pathogen development and human/vector interactions) are relatively well understood. Malaria is a climate-sensitive, vector-

borne disease that is transmitted in Africa by mosquitoes belonging to the *Anopheles gambiae s.l.* species complex, which breeds primarily in sun lit, rain-fed puddles and other still, clear water sources (Thomson et al. 1996). Laboratory studies confirm the importance of temperature on the development rates of both the malaria parasite and its mosquito vector, but the relationship is likely more complicated than initially thought (Paaijmans et al. 2012). Mechanisms underpinning many other climate-sensitive diseases or health outcomes are not necessarily well understood. Climate-driven statistical and process-based models are increasingly being used to refine our knowledge of these relationships and predict the effects of climate variability and climate change on disease incidence. However, results are not always consistent and may be of limited operational use (Morin et al. 2013) if different components of the system are not properly elaborated.

Early on in its history, the Roll Back Malaria Partnership recognized the potential to use climate information in Malaria Early Warning Systems (MEWS) to help predict and prevent malaria epidemics (WHO 2001). The MEWS approach was designed to include elements of vulnerability assessment, seasonal climate forecasting, weather/environmental monitoring and improved case surveillance within an integrated framework that could inform malaria control planning, enable preparedness, early detection and timely response. The use of seasonal climate forecasting was explored in depth through a collaboration between IRI and the European Union funded DEMETER (Development of a European Multi-model Ensemble Forecast System for Seasonal to Inter-annual Climate Prediction) project (Palmer et al. 2004). Forecasts were developed using a multi-model system permitting forecasts to be presented to decision-makers in such a way as to indicate the level of uncertainty in the predictions (Thomson et al. 2000). Probabilistic forecasting is intended to encourage the use of forecast information in regions, seasons and years when predictability is most likely to be high (e.g. the East African short rains during El Niño-Southern Oscillation events) and ignored when predictability is insufficient for improved outcomes, e.g. during the long rains in Kenya during March-May (Thomson et al. 2003).

Initial success of the MEWS approach was demonstrated at the regional level in the Southern Africa Development Community (SADC) (DaSilva et al. 2004). The approach built on lessons learned from the food security community (Buchanan-Smith and Davies 1995; Kuhn et al. 2005) and focused on data, methodologies and tools (Grover-Kopec et al. 2005; Grover-Kopec et al. 2006) that could be readily transferred with an emphasis on political engagement with key stakeholders (DaSilva et al. 2004). Regional support for the development of MEWS came from the WHO- Southern Africa Malaria Control

(SAMC) program supported by the RBM Inter-Country Team and SADC Drought Monitoring Center (DMC) based in Harare, Zimbabwe. Botswana was selected as an initial focus because of its long history of reliable, laboratory-confirmed epidemiological data. IRI worked with the Ministry of Health to determine the relationship between observed and forecasted rainfall and malaria incidence (Thomson *et al.* 2005; Thomson *et al.* 2006; Mason and Stephenson 2008). The strength of the observed relationship encouraged neighboring SADC countries to develop their own MEWS.

Between 2004 and 2007, the principal mechanism established for awareness-raising and information sharing between the climate and malaria control communities in Southern Africa was the seasonal Malaria Outlook Forum (MALOF), modeled on the traditional "Climate Outlook Forum" (Ogallo *et al.* 2008). The main purpose of the MALOF meeting was to integrate climate information into the routine planning cycle of national malaria control programmes. The meetings were timed for November, just prior to the onset of the annual rainy season when seasonal climate forecasts were of sufficient skill to enable proactive decisions to be made (DaSilva *et al.* 2004). The MALOF garnered great interest in its first two years of operation. The activity was considered a practical example of adaptation to climate change and the approach was seen to be applicable to other climate sensitive sectors (WMO 2007).

The impetus in this regional activity declined, however, once donor funding ended. Without the resources to enable inter-sectorial regional organization and implementation efforts, the health sector could no longer adequately support the initiative and the climate community was unable to pick up the cost. Currently, while some national level activities have continued in the SADC member states and the malaria epidemics of the past have not re-emerged at the scale seen historically, a lack of sustained, long-term investment in this regional climate service initiative has led to gaps in delivery and focus.

Despite this demise in the meetings, the MALOF has inspired continued support to malaria control from the regional meteorological agencies. In addition, there is anecdotal evidence from newspaper reports as well as ministerial, donor and non-governmental organization statements suggesting that epidemic early warning activities continue. Moving forward, early warning systems continue to be seen as a component of national malaria elimination strategies (Simon *et al.* 2013).

While exploring how sustainable early warning systems might be established, IRI has continued to conduct research on the various sources of environmental and climate data and products that might be incorporated into malaria risk assessments. These include rainfall, temperature, vegetation and extents of water bodies

derived from remotely-sensed images. New products provided by the National Oceanic and Atmospheric Administration (NOAA) and the National Aeronautics and Space Administration (NASA) have been routinely tested against field observations and new, refined products developed. For example, estimations of air temperature derived from Moderate-Resolution Imaging Spectro-radiometer (MODIS) Land Surface Temperature products (Vancutsem *et al.* 2010; Ceccato *et al.* 2010) were developed and provided to the user community to assess risks of malaria transmission in the highland regions of Africa. Maps of water bodies and vegetation using the MODIS sensor at 250 metre spatial resolution were developed and integrated into vector-borne disease risk models (Baeza *et al.* 2013; Ceccato *et al.* 2006). Additionally, a Vectorial Capacity Model (VCAP) has been designed to include the influence of rainfall and temperature variables on the potential for malaria transmission. The expanded model was tested in Eritrea and Madagascar using remote sensing data to check the viability of the approach. The analysis of climate driven VCAP and malaria incidence data in these countries showed that the model correctly tracked the risk of malaria in regions where rainfall and or temperature were the limiting factors (Ceccato *et al.* 2012).

In 2003, Eritrea was a primary recipient of World Bank and USAID funding for malaria control (Simon *et al.* 2011) and the IRI was invited to provide technical support to the Ministry of Health through the creation of a Malaria Early Warning System (MEWS) in this highly epidemic prone country. Analysis of climate and malaria data in Eritrea identified a potential limitation of the national malaria program's impact evaluation process. A more than 60% reduction in malaria incidence observed between 1998 and 2003 was widely lauded by the World Bank and others as a result of the scale up of control efforts such as indoor residual spraying and the use of insecticide impregnated bed nets (Barat 2006). However, during the intervention period, Eritrea experienced much drier climatic conditions compared with the baseline year of 1999. Without accounting for the impact of drought on the observed decline in malaria, the Ministry of Health risked overstating the effectiveness of their control program and thereby potentially underestimating the need for future investments. A careful analysis which accounted for rainfall variability in the assessment of the impact of anti-malarial interventions in Eritrea concluded that drought contributed to approximately 40% of the total decline in cases during the period of observation (Graves *et al.* 2008). This analysis alerted the IRI to the value of using climate information in malaria control impact assessments (Dinku *et al.* 2014b; Thomson *et al.* 2012).

Epidemics of malaria remain a major public health issue in East Africa where both desert-fringe and highland-fringe epidemics have occurred following unusual rainfall

and temperature conditions (Worrall *et al.* 2004). The highland areas are of particular concern due to their economic importance and dense populations. Significant epidemics in many highland regions in the 1990s prompted calls for improved epidemic early warning (Connor *et al.* 1999). Some considered these epidemics as evidence of climate change in the region (Patz *et al.* 2003). However, this perspective was strongly contested, largely on the erroneous basis that epidemics had emerged in the absence of a warming climate (Reiter *et al.* 2004).

The IRI's capacity to work with experts from both climate and sectorial communities has proven particularly valuable in helping to bridge the disciplinary divide at the country level. For example, in Kenya, the scientists worked closely with the Kenyan Meteorological Department to shed new light on the decade-long dispute on whether or not the highlands in Western Kenya have experienced a significant warming since the 1970's (Shanks *et al.* 2002; Hay *et al.* 2002; Pascual *et al.* 2006). A study by Omumbo and colleagues in 2011 using 30 years of quality-controlled daily meteorological data from a disputed highland study site (Kericho, Kenya) concluded definitively that significant warming has occurred and provided evidence that what was happening in the local climate in Kericho was intimately associated with what was happening at the regional level (Omumbo *et al.* 2011a). Year to year anomalies in minimum temperatures, for example, were found to be strongly correlated with equatorial land and sea surface temperatures.

Given that the climate-malaria linkages in the Kenyan highlands have been studied extensively by several research teams and that the results of different studies have been disseminated widely and cited in key reviews (Reiter 2008) and policy documents (IPCC 2007), it is disconcerting to note that it took over a decade to obtain a quality assured historical temperature record that could be used to identify these trends. This study and other observed challenges, with regard to availability and access of climate observations, spurred efforts at the IRI to overcome the policy and technical barriers to climate data that exist at the national level (Dinku *et al.* 2014a).

### **Getting evidence into policy and practice**

Early on in its development the IRI health programme identified the need for "coalitions of partners" and "communities of practice" in furthering its multi-disciplinary, research-to-practice mission. Building communities involving both researchers and policy makers and practitioners is challenging given their differing priorities (Lomas 1997). However, such communities can help researchers identify questions of direct relevance to operational needs while giving the opportunity for decision-makers to better understand the strengths and limitations of science (Sutherland *et al.* 2013). One such "coalition of partners" is the

Meningitis Environment Risk Information Technologies (MERIT) initiative, a multi-sectorial research-to-practice consortium designed to support current and future vaccination strategies for the prevention and control of meningococcal meningitis in the Sahel.

The history of MERIT, its membership and its processes are described in detail elsewhere (Thomson *et al.* 2013). The research supported by MERIT has targeted identified priorities for meningococcal meningitis control and the impact of serogroup A vaccination across the African meningitis belt (Greenwood 2013). Given the make-up of the consortium membership, the most significant research outputs have been largely focused on climate/environmental drivers of meningitis incidence (Yaka *et al.* 2008; Abdussalam *et al.* 2014; Dukic *et al.* 2012; Perez *et al.* 2014). In addition considerable advances have been made in understanding and monitoring of the climate of the Sahelian dry-season (Pérez *et al.* 2011). Although MERIT has advanced a research for policy and practice agenda, traditional differences between research and health services worlds remain and act as a constraint on effective knowledge transfer. Differences in the incentives for researchers (grants, publications) and decision-makers (actionable information) as well as timelines for expected delivery (years versus months) pose significant hurdles to the engagement of both communities (Lomas 2007), as does the time taken to engage in meaningful discourse (Ward *et al.* 2009).

In 2008, a multi-agency and cross-disciplinary community of practice "Climate and Health Working Group" (CHWG) was formed in Ethiopia with support from a Google.org funded project "Building Capacity to Produce and Use Climate and Environmental Information for Improving Health in East Africa". The project was designed to facilitate the articulation of climate information needs for improved control of climate-sensitive diseases (Ghebreyesus *et al.* 2008). The CHWG was composed of approximately 12 agency representatives and chaired by the Ministry of Health with the National Meteorological Agency (NMA) serving as co-chair. The remaining membership was drawn from agencies that broadly supported public health development in Ethiopia including: the Ethiopian offices of UN and other international organizations, the Ethiopian Health and Nutrition Research Institute, Addis Ababa University, the Association of Schools of Public Health and the Health-Development and Anti-Malaria Association (HDAMA; previously called the Anti-Malaria Association), a local community based-organization which also acted as the CHWG secretariat.

In the control and prevention of malaria, Ethiopia provides a striking example given its large population (80 million – the second largest in Africa only after Nigeria) and the fact that 75% of its inhabitants live in areas of unstable malaria transmission where year to year and

longer term changes in climate can have a major impact. It therefore provided an ideal environment to explore the use of climate information in malaria control decision-making.

In its first three years of operation, the Ethiopian CHWG, with the support of HDAMA, organized and supported cross-disciplinary training and research activities across these agencies at national and sub-national levels. These activities in turn led to elaboration of information needs and the development of operational information products designed to serve control initiatives for a number of climate-sensitive health issues (Teka 2009; Dinku 2010). This demonstration of cross-institutional facilitation from Ethiopia stimulated interest elsewhere and CHWGs (with a similar composition to Ethiopia's) were established in Madagascar with the support of the World Meteorological Organization (WMO), in Kenya with the support of WHO and in West Africa with the support of the African Center of Meteorological Applications for Development (ACMAD) and AEMET (the National Meteorological Agency of Spain). Based on these initial results, WMO has adopted CHWGs as a key strategy for the development of its engagement with the health community in Africa (Guillemot 2014).

A key outcome of the Ethiopian capacity building work has been the prioritization of improved historical and real-time climate data and information over climate forecasts. The observational data provide the basis for improved historical, current and future analyses, and since the opportunity to create useful forecast information is severely constrained where data are lacking the prioritization of the observational data is effectively unavoidable. The creation of "Enhanced National Climate Services" (ENACTS) products from quality controlled national observational data blended with appropriate satellite data (Dinku *et al.* 2011) has not only transformed the way the National Meteorological Agencies can serve the development community, but has focused attention on the need for robust historical information, with national coverage, to form the basis of climate-sensitive decision-making (Thomson *et al.* 2011). In response to demand from the national meteorological agencies and with support from WMO and regional partners, ENACTS have been developed at the regional level in West Africa and at the national level in Madagascar and Tanzania with other countries now seeking similar support. These new "gold standard" nationally owned climate resources have enabled, for the first time, detailed analyses of the impact of climate variability on malaria impact assessment by National Malaria Control Programmes (Dinku *et al.* 2014b).

Multi-disciplinary trainings (involving both the health and climate research and operational communities) are an important part of the capacity building portfolio of IRI and have been a central pillar of the CHWGs. Short training courses held in Ethiopia, Kenya and Madagascar

have been facilitated by alumni from the Summer Institute "Climate Information for Public Health" held in New York (Mantilla *et al.* 2014). In Madagascar, two training courses on the use of climate information for public health were conducted in October 2009 and March 2010. These courses were implemented and funded by the Public Weather Services (PWS) program of WMO as part of its new "Learning Through Doing" (LTD) approach to the delivery of climate services. The LTD effort implemented in Madagascar offered new opportunities for multi-disciplinary research and collaboration between climate experts and public health professionals (Cibrelus *et al.* 2010). The content of these courses was based on the IRI Climate Information for Public Health curriculum (Cibrelus and Mantilla 2010) and the courses were delivered in the local language with materials tailored to local needs, using national disease surveillance data and examples relevant to the areas of work of the participants.

As a boundary institute (both a WHO Collaborating Center and a source of expertise for many WMO programs, projects and processes), the IRI has been present and engaged in many key inter-institutional efforts including, for example, the World Climate Conference III held in Geneva in 2009 to launch the GFCS process (Guillemot 2014; Connor *et al.* 2010). To facilitate inter-sectorial dialogue and processes, a US 501c3 Non-Governmental Organization, the Health and Climate Foundation (HCF 2007) was created. IRI has partnered with HCF to strengthen its engagement in the policy arena.

In April 2011, the IRI and the Health and Climate Foundation, in partnership with key regional and international organizations initiated and facilitated the "Climate and Health in Africa – 10 Years On" multi-stakeholder workshop in Addis Ababa (Omumbo *et al.* 2011b). The workshop was designed to reflect on achievements in climate and health research, policy and practice in Africa since the Bamako meeting of 1999 and to chart a way forward for greater impact. Steering member organizations included WMO, UNECA's African Climate Policy Center (ACPC), the United Nations Development Program (UNDP), WHO-AFRO, the Met Office, Exeter University and HDAMA. With over 100 participants, many from National Meteorological Agencies or Ministries of Health, the workshop participants developed a series of key recommendations in the areas of policy, practice, climate services and data, research and education (Table 1) building on lessons learned in the "Gap Analysis". The recommendations testify to the emergence of a mature, experience-based, discourse between diverse stakeholders.

Engagement of WHO-AFRO in the workshop highlighted the importance of linking this effort to those emerging across the continent following the Libreville Declaration

(WHO-AFRO 2008). In response to the increasing need of Ministries of Health to adapt to climate change, WHO-AFRO has recently focused its efforts on the development of an informal international consortium for public health early warning and response, under the umbrella of ClimHealth-Africa (WHO-AFRO 2011). The IRI is currently supporting this effort.

While the African health community has been focused on the development of an enabling policy environment for the better management of climate-related health risks, progress towards the creation of a Global Framework of Climate Services (GFCS) under the leadership of WMO has been underway (Hewitt *et al.* 2012). The GFCS further reiterated the view that climate services must fully engage the user communities and be based on accurate, relevant and reliable observational data obtained in an effective, efficient, credible and transparent manner.

### Lessons learned

Through the evolution of the IRI's strategy on climate and health over the last fifteen years, the health programme at IRI has learned many key lessons some of which are described below.

#### With regard to evidence creation

Attempts to promote the application of seasonal climate forecasts within the health sector are unlikely to succeed unless the forecasts are packaged as only one component within a broader climate service system. This system should include historical and monitoring climate and environmental products as well as forecast information, which extends across timescales (from weeks to decades). This breadth of effort enables engagement in a much wider set of climate-sensitive development challenges (Hellmuth *et al.* 2007; Hellmuth *et al.* 2011; Hellmuth *et al.* 2009) and provides the health community with a wide-ranging new set of products and services. With this expanded range of information products we can improve our understanding of disease drivers; map populations at risk; investigate the seasonality and timing of interventions; monitor and forecast year-to-year variability and trends (including epidemics); and assess the contribution of climate as a confounder in an impact assessment of interventions.

To influence policy and practice, evidence needs to be built up in a cumulative and strategic manner so that new knowledge can be accepted by specific user communities and incorporated in their planning process. Developing sufficient evidence might require using a range of methods, repeating the analysis at multiple sites to ensure robust, locally relevant, results and ensuring that the information is presented in the 'language' of the specific community. The varying spatial and temporal uncertainty of forecast information presents a challenge

to decision-makers – as policies must often be consistent across administrative regions and seasons. However, by integrating seasonal climate forecasts into an early warning system that incorporates a range of information sources (such as a historical probability of climate suitability for epidemics, recent environmental observations, early case detection *etc.*), decision-makers may then access actionable information even when seasonal climate forecast signals are weak (Ceccato *et al.* 2007; Thomson 2013).

Seasonal climate forecast verification has been an important area of research at the IRI (Mason 2012). Ensuring that practitioners are aware of the limitations as well as the opportunities of climate forecasts is critical to their effective use (Mason and Chidzambwa 2008). Recent work includes establishing a verification framework for decadal prediction (Goddard *et al.* 2013). Such a framework is essential to establishing the potential skill of temperature and rainfall model outputs from Coupled Model Inter-comparison Project Phase 5 (CMIP5).

Once again evidence is required to establish the strength of the relationship between forecasts and outcomes as well as the value of the information to decision-making. Even if long term assessments were shown to have predictive skill, it is not clear how such information could be used in health decision-making. This is because most policy timelines for infectious disease prevention and control are relatively short when compared to climate change timelines of decades to centuries. Planning cycles in the Ministry of Health usually follow annual and political cycles, *i.e.* normally 1–4 years. Longer time-frames associated with programme development (e.g. roll-out of the malaria elimination strategy or new vaccines for meningococcal meningitis) may be on the order of 5–20 years. While climate scientists seek to find predictability in this time-frame, very little is known about the potential for decadal forecasts to have meaningful skill (Cane 2010).

Given these decision-making constraints, current opportunities and priorities identified by IRI work for the routine use of climate and environmental information in health in Africa are focused on: i) understanding mechanisms by which climate impacts on transmission and occurrence of disease; ii) mapping populations at risk both in space and by season; iii) developing early warning systems; iv) understanding the contributions of climate to trends in disease incidence; and v) improving the evaluation of the impacts of climate-sensitive interventions.

#### With regard to knowledge transfer

Knowledge transfer is complex and requires an understanding of the often misaligned agendas of research and decision-making communities (Lomas 2007; Tsui *et al.* 2006). Meaningful exchange can only be built on a

**Table 1 Consensus recommendations on policy from workshop held on “Climate and Health in Africa: 10 Years On” held in Addis Ababa, Ethiopia in April 2011**

Theme 1 Policy	Theme 2 Practice	Theme 3 Services and Data	Theme 4 Research and Education
<p><i>Support effective implementation of the Joint Statement on Climate Change and Health in Africa adopted by African Ministers of Health and Environment in Luanda, 2010, as an overarching platform for addressing climate and health issues to:</i></p>	<p>-Integrate climate health risk management into cross-sector planning and practice for adaptation to climate variability and change by developing climate services and products that address disease prevention at end-user level.</p>	<p>-Develop tailored services in partnerships with weather/ climate and health organisations. These should recognise that health forecasts, which are different from weather forecasts, should be well designed and understood by all. They should act as early warnings to users of differing types, that assist in the prediction of future health outcomes.</p>	<p>-Understand the relationships between climate and climate-sensitive diseases and health issues under different environmental conditions through interdisciplinary, multi-sectoral and multi-centre research.</p>
<p>-Bridge the gap between policies and practices through legislation and guidelines, appropriate planning, including relevant vulnerability assessments, programmatic support and multi-sectoral and participatory processes that are gender sensitive.</p>	<p>-Create a human resource center/ virtual hub where expertise is shared in order to develop the capacity of African health and climate communities, institutions, practitioners and negotiators to understand/integrate climate change challenges into policy, socio-economics, planning and programming by identifying institutions and organisations in Africa that can deliver training courses and conduct research on “Climate, Health and Prevention”.</p>	<p>-Improve existing data, for example through: the digitisation of historical health and climatic data; the increased use of metadata analyses and validation tools; the inclusion of aggregated health data at appropriate spatial and temporal scales; and the enhanced awareness of, and use of, observational and processed data, appropriate satellite, and climate model data sources.</p>	<p>-Ensure that climate change mitigation and adaptation strategies are informed by multi-disciplinary research.</p>
<p>-Support countries to establish integrated health surveillance and climate observation and processing systems.</p>	<p>-Strengthen community-based organisations by liaising, in a gender-sensitive fashion, with their leaders to develop locally owned sustainable strategies for adaptation to climate change and/or variability in their communities taking account of local knowledge rooted in social history and disseminated by appropriate channels, including the mass media.</p>	<p>-Access and use data in a systematic manner in order to identify vulnerable groups and areas. This needs to involve: employing data strategically within and across sectors; considering trend and seasonality issues; using data to evaluate the success of interventions; and, importantly, understanding how communities cope.</p>	<p>-Develop capacity within Africa for the generation, interpretation and use of climate, health and other interdisciplinary data enabling informed, evidence-based decision making.</p>
<p>-Strengthen health systems using climate information tailored to decision needs at all relevant levels and time scales.</p>	<p>Define the different levels and needs (including learning outcomes) of health practitioners and stakeholders across different geographic scales, specifically researchers and teachers, graduate and undergraduate students, practitioners in the public health system, community opinion leaders, traditional healers, impacted communities and other special interest groups and develop appropriate curricula for adaptation to climate change and/or variability in the health sector.</p>	<p>-Incorporating other data into these health forecast services, for example population, rural vs. urban residence, migration, nutritional status, environmental and poverty data.</p>	<p>-Standardise and quality control data collection and storage, ensuring data are available on relevant temporal and spatial scales.</p>
<p>-Make evidence-based, sound climate-informed decisions to implement a set of preventive actions to reduce population vulnerability and lessen the additional burden imposed by climate-sensitive diseases and health issues according to their respective epidemiological circumstances.</p>	<p>-Promote a gender-sensitive approach to interventions on climate and health in cross-sectoral disaster risk reduction and preventive health strategies.</p>	<p>-Collaboration +: new, multi-disciplinary initiatives that involve communities beyond health and climate/ weather; build upon existing initiatives and progress; aim to meet emerging challenges; and communicate with end-users in appropriate ways.</p>	<p>-Enhance knowledge transfer and communication of information across disciplines and communities through existing networks, encouraging the introduction of climate and health into the curriculum at all levels of education.</p>



**Table 1 Consensus recommendations on policy from workshop held on “Climate and Health in Africa: 10 Years On” held in Addis Ababa, Ethiopia in April 2011 (Continued)**

Anticipate, prepare for and respond to the health consequences of extreme weather events, particularly by strengthening the functioning of health systems and other relevant sectors.	-Commitment at all levels that brings climate and health communities together, clarifies responsibilities, builds capacity in the climate and health sectors to achieve these services, facilitates joint initiatives and ensures resources such as data are shared in a suitable way.	-Strengthen existing partnerships and collaborations while developing new groups and building links across disciplines.
Multilateral partners to consider the significant co-benefits of environment integrity, population health and consequent economic development that can result from mitigation and adaptation policies in the climate and health sectors and to support African countries in gaining access to resources under the various climate-related funds.		

confident and trusting relationship. Approaches to increasing uptake in policy and practice must happen at a number of levels. First of all, policies must be flexible enough to respond to new information. If they are not, then opportunities to change policies should be explored. Researchers must recognize that scientific evidence is only one piece of information that is used in the pragmatic and political process of decision-making. Even when policies favour the use of new information, practitioners may not adopt it in their decision-making processes without additional support. This might come in the form of formal or informal (including web-based) training, access to decision-relevant peer-review publications (e.g. open access) and technical reports and peer-to-peer communication.

The metrics developed and described earlier in this document have acted as a guide to programme development rather than an explicit evaluation tool. Nonetheless, significant advances in the creation of evidence for policy, “support to communities of practice” and “coalitions of partners” at national and international levels have demonstrated the value of this approach.

The challenge of designing and implementing rigorous impact evaluation for in-country use of climate information remains. The effort needed to undertake significant impact evaluations, which could connect climate information to development outcomes, is not only challenging methodologically but requires resources beyond those available through conventional research funding. In addition, basic capacities in the delivery of climate information in Africa at the national level are only now emerging, making it difficult to undertake proper assessments of the value of the information in a ‘real world’ context.

### Opportunities going forward

IRI has long proposed the incorporation of climate information into routine epidemiological surveillance systems (Jima *et al.* 2012) for climate-sensitive diseases (Thomson and Mantilla 2011). To achieve this integration requires new and innovative mechanisms for strengthening observations, data management and sharing, development of relevant climate services, inter-sectorial collaboration, training and capacity building, all within an enabling policy environment. The premise is that improved management of health risks associated with climate variability (such as the heat early warning systems recently established in Europe and North America) increases adaptive capacity of the public health sector to longer-term climate change.

In response to this need, the IRI has invested considerable effort in training and research capacity development in the area of climate and health through its Climate Information for Public Health curriculum and associated initiatives (Mantilla *et al.* 2014). Going forward, a broader health agenda has to be identified, which connects climate and environmental information to the needs of communities dealing with infectious disease, public health outcomes of hydro-meteorological disasters and nutrition. Central to this strategy is a focus on availability, access and use of historical, current and forecast data at temporal and spatial scales appropriate for decision-making (Dinku *et al.* 2011). In addition there is an urgent need to develop research and operational capacity to use the information in decision-making.

The “Climate and Health in Africa - 10 years On” workshop held in Addis Ababa in April 2011 demonstrated the dramatic change in perception of the value of climate information in health decision-making in Africa

since the Bamako workshop of 1999. This change is fostered by new resources for climate change adaptation, policy developments at the global and regional scale, donor and government requirements for more targeted health interventions and the need for better assessments of the impact of climate sensitive interventions. The specific role of IRI's contribution to mainstreaming climate information into health policy and practice in Africa cannot be fully assessed in the short term. However, we believe that the current and former staff and alumni of the IRI have played a significant role in advancing the concept and implementation of a 'climate-smart' health sector in Africa.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

All authors contributed to this review. MT was responsible for lead authorship and assumed final word on all writing. SC provided substantial contributions to conception and overview, with BP providing critical early stage revisions and content. JO, SM and GM are listed as co-authors due to their indispensable effort in reflecting on and capturing climate and health activities at IRI and launched with partners in Africa and their diligent editing advice. PC provided critical feedback on the remote sensing contributions of IRI's work to framing the health approach and new product development driven by user needs and MJ contributed to the discussions on MERIT and the broader policy implications of IRI work while AM initiated the development of Climate and Health Working Groups in Africa and reported on their development. All authors read and approved the final manuscript.

#### Acknowledgements

We would like to thank Google.org, the Health and Climate Foundation, WMO, WHO, USAID, NASA and NOAA for financial support to the projects identified in the paper, along with countless regional partners for their dedicated engagement on collaborations in this area. We also acknowledge the scientific and technical support of many IRI staff who have contributed to the health work over the past decade.

Responsible editor: Vivek Singh

#### Author details

<sup>1</sup>International Research Institute for Climate and Society, PO Box 1000, Palisades, New York 10964, USA. <sup>2</sup>Mailman School of Public Health, Columbia University, New York, USA. <sup>3</sup>Columbia Global Center-Africa, PO Box 51412-00100, Nairobi, Kenya. <sup>4</sup>Health, Development and Anti-Malaria Association, P. O Box 27279, Addis Ababa, Ethiopia. <sup>5</sup>Kenya Medical Research Institute-Wellcome Trust Research Programme, PO Box 43640-00100, Nairobi, Kenya. <sup>6</sup>Centro de Estudios e Investigacion Fundacion Santa Fe de Bogota, Calle 123 #7B-90, Bogota, Colombia. <sup>7</sup>Health and Climate Foundation, M25 K Street NW, Suite 350, Washington, DC 20005, USA. <sup>8</sup>School of Environmental Sciences, University of Liverpool, Liverpool L69 3BX, UK.

Received: 1 October 2013 Accepted: 22 April 2014

Published: 17 June 2014

#### References

- Abdussalam AF, Monaghan AJ, Dukić VM, Hayden MH, Hopson TM, Leckebusch GC, Thomes JE (2014) Climate Influences on Meningitis Incidence in Northwest Nigeria. *Weather Climate Soc* 6:62-76. doi.org/10.1175/WCAS-D-13-00004.1
- Anon (2013) Clim-Health Africa. *Bull World Meteorol Org* 62(Special Issue):35-40
- Baeza A, Bouma MJ, Dhiman R, Baskerville E, Ceccato P, Yadav RS, Pascual M (2013) Long lasting transition towards sustainable elimination of desert malaria under irrigation development in India. *Proc Natl Acad Sci* 110:15157-15162. doi:10.1073/pnas.1305728110
- Barat L (2006) Four malaria success stories: how malaria burden was successfully reduced in Brazil, Eritrea, India and Vietnam. *Am J Trop Med Hyg* 74(1):12-16
- Buchanan-Smith M, Davies S (1995) *Famine Early Warning and Response: The Missing Link*. Intermediate Technology Publications, London, p 228pp
- Cane M (2010) Climate science: decadal predictions in demand. *Nat Geosci* 3:231-232
- Ceccato P, Bell MA, Blumenthal B, Connor SJ, Dinku T, Grover-Kopec EK, Ropelewski CF, Thomson MC (2006) "Use of Remote Sensing for Monitoring Climate Variability for Integrated Early Warning Systems: Applications for Human Diseases and desert Locust Management". IGARSS Denver, July 31 2006-Aug. 4 2006. IEEE International Conference on Geoscience and Remote Sensing Symposium, Los Alamitos, California. doi.org/10.1109/IGARSS.2006.74
- Ceccato P, Ghebremeskel T, Jaiteh M, Graves PM, Levy M, Ghebreselassie S, Ogbamariam A, Barnston AG, Bell M, del Corral J, Connor SJ, Fesseha I, Brantly EP, Thomson MC (2007) Malaria stratification, climate, and epidemic early warning in Eritrea. *Am J Trop Med Hyg* 77(6):61-68. <Go to ISI>://000252212600013
- Ceccato P, Vancutsem C, Temimi M (2010) "Monitoring Air and Land Surface Temperatures from Remotely Sensed Data for Climate-Human Health Applications". International Geoscience and Remote Sensing Symposium (IGARSS) Honolulu, Hawaii, 25-30 July 2010. IEEE, pp 178-180. doi:10.1109/IGARSS.2010.5649810
- Ceccato P, Vancutsem C, Klaver R, Rowland J, Connor SJ (2012) A Vectorial Capacity Product to Monitor Changing Malaria Transmission Potential in Epidemic Regions of Africa. *J Trop Med* 2012(595948). doi: 10.1155/2012/595948
- Cibrelus L, Mantilla G (2010) *Climate Information for Public Health: A Curriculum for Best Practices - Putting Principles to Work*. IRI, Palisades, New York, p 137pp. <http://iri.columbia.edu/publications/id=1044>
- Cibrelus L, Raholijao N, Raelina Y, Rakoaivony MC, Tuseo L, Cousin R, Kootval H (2010) *Training Courses on Climate Information for Public Health in Madagascar, 2009 and 2010: Synthesis Report*. IRI, Palisades, New York, p 66pp. [http://iri.columbia.edu/docs/publications/TR10-12\\_MadagascarNov1.pdf](http://iri.columbia.edu/docs/publications/TR10-12_MadagascarNov1.pdf)
- Connor S, Thomson M, Molyneux D (1999) Forecasting and prevention of epidemic malaria: new perspectives on an old problem. *Parasitologia* 41:439-448
- Connor SJ, Ceccato P, Dinku T, Omumbo JA, Grover-Kopec EK, Thomson MC (2006) Using Climate Information for Improved Health in Africa: Relevance, Constraints, and Opportunities. *Geospat Health* 1(1):17-31
- Connor SJ, Omumbo J, DaSilva J, Green C, Mantilla G, Delacollette C, Hales S, Rogers D, Thomson MC (2010) *Health and Climate - Needs*. Proc Environ Sci 1:27-36. World Climate Conference - 3
- DaSilva J, Garanganga B, Teveredzi V, Marx S, Mason SJ, Connor SJ (2004) Improving Epidemic Malaria Planning, Preparedness and Response in Southern Africa. *Malar J* 3(1):37. <http://www.malariajournal.com/content/3/1/37>
- Dinku T (2010) The need for national centres for climate and development in Africa. *Climate Dev* 2:1-5
- Dinku T, Hilemariam K, Grimes D, Kidane A, Connor S (2011) Improving availability, access and use of climate information. *WMO Bull* 60(2). [http://www.wmo.int/pages/publications/bulletin\\_en/archive/60\\_2\\_en/60\\_2\\_Tufa\\_en.html](http://www.wmo.int/pages/publications/bulletin_en/archive/60_2_en/60_2_Tufa_en.html)
- Dinku T, Block P, Sharoff J, Hilemariam K, Osgood D, Del Corral J, Cousin R, Thomson MC (2014a) Bridging Critical Gaps in Climate Service and Applications in Africa. The International Research Institute for Climate & Society: Shaping the Landscape of Climate Services. *Earth Perspect* 1(1):15. <http://www.earth-perspectives.com/content/1/1/15>
- Dinku T, Kanemba A, Platzer B, Thomson MC (2014b) Leveraging the Climate for Improved Malaria Control in Tanzania. IEEE. <http://www.earthzine.org/2014/02/15/leveraging-the-climate-for-improved-malaria-control-in-tanzania/>
- Dukić V, Hayden M, Adams Forgor A, Hopson T, Akweongo P, Hodgson A, Wiedinmyer C, Yoksas T, Thomson M, Trzaska S, Pandya R (2012) The role of weather in meningitis outbreaks in Navrongo, Ghana: A Generalized Additive Modeling Approach. *J Agric Biol Environ Stat* 17(3):442-460. doi:10.1007/s13253-012-0095-9
- G8 (2005) *Gleneagles Plan of Action*. Climate Change, Clean Energy and Sustainable Development. Gleneagles, Scotland. <http://www.number10.gov.uk/Page7882>
- Gallup JL, Sachs JD (2001) The economic burden of malaria. *Am J Trop Med Hyg* 64(1,2):85-96
- Ghebreyesus TA, Tadese Z, Jima D, Bekele E, Mihretie A, Yihdego YY, Dinku T, Connor SJ, Rogers DP (2008) Public health and weather services-climate information for the health sector. *Bull World Meteorol Org* 57(4):256-261
- Goddard L, Kumar A, Solomon A, Smith D, Boer G, Gonzalez P, Khariin V, Meryfield W, Deser C, Mason SJ, Kirtman BP, Msadek R, Sutton R, Hawkins E,

- Fricker T, Hegerl G, Ferro CAT, Stephenson DB, Meehl GA, Stockdale T, Burgman R, Greene AM, Kushnir Y, Newman M, Carton J, Fukumori I, Delworth T (2013) A verification framework for interannual-to-decadal predictions experiments. *Climate Dynam* 40(1–2):245–272
- Graves PM, Osgood DE, Thomson MC, Sereke K, Araia A, Zerom M, Ceccato P, Bell M, del Corral J, Ghebreselassie S, Brantly EP, Ghebremeskel T (2008) Effectiveness of malaria control during changing climate conditions in Eritrea, 1998–2003. *Trop Med Int Health* 13(2):218–228. doi: 10.1111/j.1365-3156.2007.01993.x
- Greenwood B (2013) Priorities for research on meningococcal disease and the impact of serogroup A vaccination in the African meningitis belt. *Vaccine* 31(11):1453–1457. doi: 10.1016/j.vaccine.2012.12.035. Epub 2012 Dec 27
- Grover-Kopec E, Kawano M, Klaver RWW, Blumenthal B, Ceccato P, Connor SJ (2005) An online operational rainfall-monitoring resource for epidemic malaria early warning systems in Africa. *Malar J* 4(6). doi:10.1186/1475-2875-4-6
- Grover-Kopec E, Blumenthal B, Ceccato P, Dinku T, Omumbo J, Connor S (2006) Web-Based Climate Information Resources for Malaria Control in Africa. *Malar J* 5:38. doi: 10.1186/1475-2875-4-6
- Guillemot J (2014) Health Exemplar: Annex to the Implementation Plan for the Global Framework for Climate Services. Geneva. [http://webcache.googleusercontent.com/search?q=cache:XfnXMuFyoQJftp://ftp.wmo.int/Documents/gfcs/ImplementationPlan/Exemplars/GFCS-HEALTH-EXEMPLAR-14152\\_en\\_proof-read.doc+&cd=1&hl=en&ct=clnk&gl=us&client=safari](http://webcache.googleusercontent.com/search?q=cache:XfnXMuFyoQJftp://ftp.wmo.int/Documents/gfcs/ImplementationPlan/Exemplars/GFCS-HEALTH-EXEMPLAR-14152_en_proof-read.doc+&cd=1&hl=en&ct=clnk&gl=us&client=safari)
- Hay SI, Rogers DJ, Randolph SE, Stern DI, Cox J, Shanks GD, Snow RW (2002) Hot topic or hot air? Climate change and malaria resurgence in East African highlands. *Trends Parasitol* 18(12):530–534
- HCF (2007). "<http://hc-foundation.org>."
- Hellmuth ME, Moorhead A, Thomson MC, Williams J (ed) (2007) *Climate Risk Management in Africa: Learning from Practice*. Climate and Society Series. IRI, Palisades, New York
- Hellmuth ME, Osgood DE, Hess U, Moorhead A, Bhojwani H (ed) (2009) *Index Insurance and Climate Risk: Prospects for Development and Disaster Management*. Climate and Society Report/Climate and Society Report. IRI, Palisades, New York
- Hellmuth ME, Mason SJ, Vaughan C, van Aalst MK, Choularton R (2011) *A Better Climate for Disaster Risk Management*. Climate and Society Report Series. IRI, Palisades, New York
- Hewitt C, Mason S, Walland D (2012) *The Global Framework for Climate Services*. *Nat Clim Chang* 2:831–832
- Hopp MJ, Foley JA (2001) Global-scale relationships between climate and the dengue fever vector, *Aedes aegypti*. *Clim Change* 48(2–3):441–463
- Hopp MJ, Foley JA (2003) Worldwide fluctuations in dengue fever cases related to climate variability. *Climate Res* 25(1):85–94
- IPCC (2007) *IPCC Third Assessment Report 2007 (TAR) Climate Change: Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC Assessment Report. Cambridge University Press, Cambridge, UK
- IRI (2005) *Sustainable development in Africa: is the climate right?* IRI, Palisades, New York, p 40pp
- IRI (2006) *A Gap Analysis for the Implementation of the Global Climate Observing System Programme in Africa*. IRI, Palisades, New York, p 06–01. 39pp
- Jima D, Wondabeku M, Alemu A, Teferra A, Awel N, Deressa W, Adissie A, Tadesse Z, Gebre T, Mosher A, Richards F, Graves P (2012) Analysis of malaria surveillance data in Ethiopia: what can be learned from the integrated disease surveillance and response system? *Malar J* 5(17):330. doi: 10.1186/1475-2875-11-330
- Kelly-Hope LA, Thomson MC (2008) *Climate and Infectious Disease*. In: Thomson MC, Garcia-Herrera R, Beniston M (ed) *Seasonal Forecasts, Climatic Change, and Human Health*. Springer Science+Business Media, Dordrecht, pp 31–70. doi:10.1007/978-1-4020-6877-5
- Kuhn K, Campbell-Lendrum D, Haines A, Cox J (2005) *Using Climate to Predict Infectious Disease Epidemics*. WHO, Geneva, Switzerland, p 54pp
- Kula N, Haines A, Fryatt R (2013) *Reducing Vulnerability to Climate Change in Sub-Saharan Africa: The Need for Better Evidence*. *PLoS Med* 10(1). doi:10.1371/journal.pmed.1001374
- Lomas J (1997) *Improving research dissemination and uptake in the health sector: Beyond the sound of one hand clapping*. McMaster University Centre for Health Economics and Policy Analysis C97-1, Ontario, p 42pp. [http://www.cfhi-fcass.ca/migrated/pdf/mythbusters/handclapping\\_e.pdf](http://www.cfhi-fcass.ca/migrated/pdf/mythbusters/handclapping_e.pdf)
- Lomas J (2007) *The in-between world of knowledge brokering*. *Br Med J* 334(7585):129–132
- Mantilla G, Thomson C, Sharoff J, Barnston AG, Curtis A (2014) *Capacity building through the sharing of climate information with diverse user communities - The International Research Institute for Climate & Society: Shaping the Landscape of Climate Services*. *Earth Perspect* 1(1).
- Mason SJ (2009) *Recommended Procedures for the Verification of Operational Seasonal Climate Forecasts*. WMO, Geneva, Switzerland
- Mason SJ (2012) *Seasonal and longer-range forecasts*. In: Jolliffe IT, Stephenson DB (ed) *Forecast Verification: A Practitioner's Guide in Atmospheric Science*. Wiley, Chichester, pp 203–220
- Mason SJ, Chidzambwa S (2008) *Verification of RCOF Forecasts*. IRI, Palisades, New York, p 09–02. 23pp
- Mason SJ, Stephenson DB (2008) *How can we know whether the forecasts are any good?* In: Troccoli A, Harrison MSJ, et al. (ed) *Seasonal Climate Variability: Forecasting and Managing Risk*. Springer Academic Publishers, Dordrecht, pp 3–11
- Mason SJ, Goddard L, Graham NE, Yulaeva E, Sun L, Arkin PA (1999) *The IRI Seasonal Climate Prediction System and the 1997/1998 El Nino Event*. *Bull Am Meteorol Soc* 80(9):1853–1874
- McCartney M, Rebelo L-M, Xenarios S, Smakhtin V (2013) *Agricultural Water Storage in an Era of Climate Change: Assessing Need and Effectiveness in Africa*. IWMI, Colombo, Sri Lanka, p 26pp
- Morin CW, Comrie AC, Ernst K (2013) *Climate and Dengue Transmission: Evidence and Implications*. *Environ Health Perspect* 121:1264–1272. doi: 10.1289/ehp.1306556
- Nutley S, Davies HTO (2000) *Making a reality of evidence-based practice: Some lessons from the diffusion of innovations*. *Public Money Manag* 20(4):35–42
- Ogallo L, Bessemoulin P, Ceron J-P, Mason S, Connor SJ (2008) *Adapting to climate variability and change: the Climate Outlook Forum process*. *Bull World Meteorol Org* 57(2):93–102
- Omumbo J, Lyon B, Waweru SM, Connor S, Thomson MC (2011a) *Raised temperatures over the Kericho tea estates: revisiting the climate in the East African highlands malaria debate*. *Malar J* 10:12. doi: 10.1186/1475-2875-10-12
- Omumbo J, Platzer B, Girma A, Connor SJ (2011b) *Climate and Health in Africa: 10 Years On Workshop*. Addis Ababa, Ethiopia. IRI, Palisades, New York, p 102pp
- Paaijmans KP, Blanford S, Chan BHK, Thomas M (2012) *Warmer temperatures reduce the vectorial capacity of malaria mosquitoes*. *Biol Letters* 8(3):465–468. doi: 10.1098/rsbl.2011.1075
- Palmer T, Alessandri A, Andersen U, Cantelaube P, Davey M, Décléclue P, Dequé M, Díez E, Doblaz-Reyes F, Feddersen H, Graham R, Gualdi S, Guérémy J, Hagedorn R, Hoshen M, Keenlyside N, Latif M, Lazar A, Maisonnave E, Marletto V, Morse A, Orfila B, Rogel P, Terres J, Thomson M (2004) *Development of a European multi-model ensemble system for seasonal to inter-annual prediction*. *Bull Am Meteorol Soc* 85:853–872
- Pascual M, Ahumada JA, Chaves LF, Rodo X, Bouma M (2006) *Malaria resurgence in the East African Highlands: Temperature trends revisited*. *Proc Natl Acad Sci* 103:5829–5834
- Patz JA, Githeko AK, McCarty JP, Hussein S, Confalonieri U, de Wet N (2003) *Climate Change and Infectious Disease*. In: McMichael AJ, Campbell-Lendrum D, Corvalan C, et al. (ed) *Climate Change and Human Health: Risks and Responses*. WHO, Geneva, pp 103–127
- Pérez C, Haustein K, Janjic Z, Jorba O, Huneus N, Baldasano JM, Black T, Basart S, Nickovic S, Miller RL, Perlwitz JP, Schulz M, Thomson M (2011) *Atmospheric dust modeling from meso to global scales with the online MMB/BSC-Dust model — Part 1: Model description, annual simulations and evaluation*. *Atmos Chem Phys* 11:13001–13027. doi: 10.5194/acp-11-13001-2011
- Perez CP, Thomson MC, Staton M, Diggle P, Hopson T, Miller R, Hugonnet S (2014) *Meningitis and Climate: from Science to Practice*. The International Research Institute for Climate & Society: Shaping the Landscape of Climate Services. *Earth Perspect* 1(1):14
- Reiter P (2008) *Global warming and malaria: knowing the horse before hitching the cart*. *Malar J* 7((Suppl 1)(S3)). doi:10.1186/1475-2875-7-S1-S3
- Reiter P, Thomas CJ, Atkinson PM, Hay SI, Randolph SE, Rogers DJ, Shanks GD, Snow RW, Spielman A (2004) *Global warming and malaria: a call for accuracy*. *Lancet Infect Dis* 4(6):323–324
- Shanks GD, Hay SI, Stern DI, Biomndo K, Snow RW (2002) *Meteorologic influences on Plasmodium falciparum malaria in the highland tea estates of Kericho, Western Kenya*. *Emerg Infect Dis* 8(12):1404–1408
- Simon J, Yeboah-Antwi K, Schapira A, Cham MK, Barber-Madden R, Brooks MI (2011) *External evaluation of the President's Malaria Initiative final report*. T. G. H. T. A. Project, Washington, p 92pp
- Simon C, Moakofhi K, Mosweunyane T, Baba Jibril HB, Nkomo B, Motlaleng M, Ntebela DS, Chanda E, Haque U (2013) *Malaria control in Botswana, 2008–*

- 2012: the path towards elimination. *Malar J* 12:458. <http://www.malariajournal.com/content/12/1/458>
- Sutherland WJ, Spiegelhalter D, Burgman M (2013) Policy: Twenty tips for interpreting scientific claims. *Nature* 503:225–337. doi: 10.1038/503335a
- Teka H (2009) Health and Climate Experts Must Jointly Tackle Disease. *Sci Dev Net*. <http://www.scidev.net/en/opinions/health-and-climate-experts-must-jointly-tackle-disease.html>
- Thomson MC (2013) Chapter 2: Climate change and disaster risk management: challenge and opportunities. In: Palutikof J, Karoly D, Boulter S (ed) *Natural Disasters and Adaptation to Climate Change*. Cambridge University Press, Cambridge UK, pp 8–32
- Thomson MC, Mantilla G (2011) Integrating Climate Information into Surveillance Systems for Infectious Diseases: New Opportunities for Improved Public Health Outcomes in a Changing Climate". *Emerging Persistent Infectious Diseases: Focus on Surveillance*, Airlie Conference Center, Warrenton, Virginia, Oct. 17–20, 2010. ISGP, Washington, pp 47–54. <http://www.scienceforglobalpolicy.org/LinkClick.aspx?fileticket=ve893KJC650=&tabid=133>
- Thomson MC, Connor SJ, Milligan PJM, Flasse SP (1996) The ecology of malaria - As seen from Earth-observation satellites. *Ann Trop Med Parasitol* 90(3):243–264
- Thomson MC, Palmer T, Morse AP, Cresswell M, Connor SJ (2000) Forecasting disease risk with seasonal climate predictions. *Lancet* 355(9214):1559–1560
- Thomson MC, Indeje M, Connor SJ, Dilley M, Ward N (2003) Malaria early warning in Kenya and seasonal climate forecasts. *Lancet (Letter)* 362:580
- Thomson MC, Mason SJ, Phindela T, Connor SJ (2005) Use of rainfall and Sea Surface Temperature monitoring for Malaria Early Warning in Botswana. *Am J Trop Med Hyg* 73(1):214–221
- Thomson MC, Doblaz-Reyes FJ, Mason SJ, Hagedorn R, Connor SJ, Phindela T, Morse AP, Palmer TN (2006) Malaria early warnings based on seasonal climate forecasts from multi-model ensembles. *Nature* 439:576–579
- Thomson MC, Connor SJ, Zebiak SE, Jancoles M, Mihretie A (2011) Africa needs climate data to fight disease. *Nature* 471:440–442
- Thomson MC, Zdravetz F, Lyon B, Mantilla G, Willis D, Ceccato P, Dinku T (2012) President's Malaria Initiative-USAID Report: Development of Climate Analysis Section for the President's Malaria Initiative Impact Evaluation: Reports for Ethiopia and Tanzania. IRI, Palisades, New York, p 62pp
- Thomson MC, Firth E, Jancoles M, Mihretie A, Onoda M, Nickovic S, Broutin H, Sow S, Perea W, Bertherat E, Hugonnet S (2013) A climate and health partnership to inform the prevention and control of meningococcal meningitis in sub-Saharan Africa: The MERIT initiative. In: Hurrell J, Arsar G (ed) *Priorities in Climate Research, Analysis and Prediction*. Springer, Netherlands, pp 459–484
- Tourre YM (1999) *Climate Prediction and Diseases/Health in Africa*. IRI, Palisades, New York, p 73pp
- Tsui L, Chapman SA, Schnirer L, Stewart S (2006) "A Handbook on Knowledge Sharing: Strategies and Recommendations for Researchers, Policymakers, and Service Providers". Y. Community-University Partnership for the Study of Children, and Families, Edmonton, Alberta, p 35pp. [http://www.uws.edu.au/\\_data/assets/pdf\\_file/0018/405252/Knowledge\\_Sharing\\_Handbook.pdf](http://www.uws.edu.au/_data/assets/pdf_file/0018/405252/Knowledge_Sharing_Handbook.pdf)
- UN (2013) *The Millennium Development Goals Report*. UN, New York, p 59pp. <http://www.un.org/millenniumgoals/pdf/report-2013/mdg-report-2013-english.pdf>
- Vancutsem C, Ceccato P, Dinku T, Connor SJ (2010) Evaluation of MODIS Land surface temperature data to estimate air temperature in different ecosystems over Africa. *Remote Sens Environ* 114:449–465
- Ward V, House A, Hamer S (2009) Knowledge Brokering: The missing link in the evidence to action chain? *Evid Policy* 5(3):267–279
- WHO (2001) *Malaria Early Warning Systems, Concepts, Indicators, and Partners: A Framework for Field Research in Africa*. WHO. WHO/CDS/RBM/2001.32, Geneva, p 80pp. <http://www.who.int/iris/handle/10665/66848>
- WHO (2004) *World Health Organization World Report on Knowledge for Better Health: Strengthening Health Systems*. WHO, Geneva, p 145pp
- WHO-AFRO (2008) *The Libreville Declaration on Environment and Health in Africa*. WHO-AFRO, Brazzaville, p 12pp. [http://www.ehrn.co.za/download/libreville\\_declaration.pdf](http://www.ehrn.co.za/download/libreville_declaration.pdf)
- WMO (2007) *Climate information for adaptation and development needs*. W. M. Organization, Geneva, p 44
- Worrall E, Rietveld A, Delacollette C (2004) The burden of malaria epidemics and cost-effectiveness of interventions in epidemic situations in Africa. *Am J Trop Med Hyg* 71(Suppl 2):136–140

Worrall E, Connor SJ, Thomson MC (2008) Improving the cost-effectiveness of IRS with climate informed health surveillance systems. *Malar J* 7:263. doi:10.1186/1475-2875-7-263

Yaka P, Sultan B, Broutin H, Janicot S, Philippon S, Fourquet N (2008) Relationships between climate and year-to-year variability in meningitis outbreaks: A case study in Burkina Faso and Niger. *Int J Health Geogr* 7:34. doi: 3410.1186/1476-072x-7-34

doi:10.1186/2194-6434-1-17

**Cite this article as:** Thomson *et al.*: Climate and health in Africa. *Earth Perspectives* 2014 1:17.

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)