



An update to the Surface Ocean CO₂ Atlas (SOCAT version 2)

D. C. E. Bakker¹, B. Pfeil^{2,3}, K. Smith^{4,5}, S. Hankin⁴, A. Olsen^{2,3,6}, S. R. Alin⁴, C. Cosca⁴, S. Harasawa⁷, A. Kozyr⁸, Y. Nojiri⁷, K. M. O'Brien^{4,5}, U. Schuster^{9,*}, M. Telszewski¹⁰, B. Tilbrook^{11,12}, C. Wada⁷, J. Akl¹¹, L. Barbero¹³, N. R. Bates¹⁴, J. Boutin¹⁵, Y. Bozec^{16,17}, W.-J. Cai¹⁸, R. D. Castle¹⁹, F. P. Chavez²⁰, L. Chen^{21,22}, M. Chierici^{23,24}, K. Currie²⁵, H. J. W. de Baar²⁶, W. Evans^{4,27}, R. A. Feely⁴, A. Fransson²⁸, Z. Gao²¹, B. Hales²⁹, N. J. Hardman-Mountford³⁰, M. Hoppema³¹, W.-J. Huang¹⁸, C. W. Hunt³², B. Huss¹⁹, T. Ichikawa³³, T. Johannessen^{2,3,6}, E. M. Jones³¹, S. D. Jones³⁴, S. Jutterström³⁵, V. Kitidis³⁶, A. Körtzinger³⁷, P. Landschützer¹, S. K. Lauvset^{2,3}, N. Lefèvre^{38,39}, A. B. Manke⁴, J. T. Mathis⁴, L. Merlivat¹⁵, N. Metzl¹⁵, A. Murata⁴⁰, T. Newberger⁴¹, A. M. Omar^{6,3,2}, T. Ono³³, G.-H. Park⁴², K. Paterson¹¹, D. Pierrot¹³, A. F. Ríos⁴³, C. L. Sabine⁴, S. Saito⁴⁴, J. Salisbury³², V. V. S. S. Sarma⁴⁵, R. Schlitzer³¹, R. Sieger³¹, I. Skjelvan^{6,2,3}, T. Steinhoff³⁷, K. F. Sullivan¹³, H. Sun²¹, A. J. Sutton^{4,5}, T. Suzuki⁴⁶, C. Sweeney⁴¹, T. Takahashi⁴⁷, J. Tjiputra^{6,3}, N. Tsurushima⁴⁸, S. M. A. C. van Heuven⁴⁹, D. Vandemark³², P. Vlahos⁵⁰, D. W. R. Wallace⁵¹, R. Wanninkhof¹⁹, and A. J. Watson^{9,*}

¹Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich, UK

²Geophysical Institute, University of Bergen, Bergen, Norway

³Bjerknes Centre for Climate Research, Bergen, Norway

⁴Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, Seattle, Washington, USA

⁵Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, Washington, USA

⁶Uni Climate, Uni Research, Bergen, Norway

⁷National Institute for Environmental Studies, Tsukuba, Japan

⁸Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁹College of Life and Environmental Sciences, University of Exeter, Exeter, UK

¹⁰International Ocean Carbon Coordination Project, Institute of Oceanology of the Polish Academy of Sciences, Sopot, Poland

¹¹CSIRO Marine and Atmospheric Research, Hobart, Australia

¹²Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Australia

¹³Cooperative Institute for Marine and Atmospheric Studies, Rosenstiel School for Marine and Atmospheric Science, University of Miami, Miami, Florida, USA

¹⁴Bermuda Institute of Ocean Sciences, Ferry Reach, Bermuda

¹⁵Sorbonne Universités (UPMC, Univ Paris 06), CNRS, IRD, MNHN, LOCEAN Laboratory, Paris, France

¹⁶CNRS, UMR 7144, Equipe Chimie Marine, Station Biologique de Roscoff, Roscoff, France

¹⁷Sorbonne Universités, UPMC Université Paris 06, UMR7144, Adaptation et Diversité en Milieu Marin, SBR, Roscoff, France

¹⁸School of Marine Science and Policy, University of Delaware, Newark, Delaware, USA

¹⁹Atlantic Oceanographic and Meteorological Laboratory, National Oceanic and Atmospheric Administration, Miami, Florida, USA

²⁰Monterey Bay Aquarium Research Institute, Moss Landing, California, USA

²¹Key Laboratory of Global Change and Marine-Atmospheric Chemistry, Third Institute of Oceanography, State Oceanic Administration, Xiamen, P. R. China

²²Chinese Arctic and Antarctic Administration, Beijing, P. R. China

- ²³Institute of Marine Research, Tromsø, Norway
- ²⁴Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg, Sweden
- ²⁵National Institute of Water and Atmospheric Research, Dunedin, New Zealand
- ²⁶Royal Netherlands Institute for Sea Research, Texel, the Netherlands
- ²⁷Ocean Acidification Research Center, University of Alaska Fairbanks, Fairbanks, Alaska, USA
- ²⁸Norwegian Polar Institute, Fram Centre, Tromsø, Norway
- ²⁹College of Earth, Ocean and Atmospheric Sciences, Oregon State University, Corvallis, Oregon, USA
- ³⁰CSIRO Marine and Atmospheric Research, Floreat, WA, Australia
- ³¹Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany
- ³²Ocean Process Analysis Laboratory, University of New Hampshire, Durham, New Hampshire, USA
- ³³National Research Institute for Fisheries Science, Fisheries Research Agency, Yokohama, Japan
- ³⁴Tyndall Centre for Climate Change Research, University of East Anglia, Norwich Research Park, Norwich, UK
- ³⁵IVL Swedish Environmental Research Institute, Gothenburg, Sweden
- ³⁶Plymouth Marine Laboratory, Plymouth, UK
- ³⁷GEOMAR, Helmholtz Centre for Ocean Research, Kiel, Germany
- ³⁸IRD, Sorbonne Universités (UPMC, Univ Paris 06), CNRS, MNHN, LOCEAN Laboratory, Paris, France
- ³⁹LaboMar, Universidade Federal do Ceará, Fortaleza-Ceará, Brazil
- ⁴⁰Japan Agency for Marine-Earth Science and Technology, Yokosuda, Japan
- ⁴¹Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, Colorado, USA
- ⁴²East Sea Research Institute, Korea Institute of Ocean Science and Technology, Uljin, Korea
- ⁴³Instituto de Investigaciones Marinas de Vigo, Consejo Superior de Investigaciones Científicas, Vigo, Spain
- ⁴⁴Marine Division, Global Environment and Marine Department, Japan Meteorological Agency, Tokyo, Japan
- ⁴⁵National Institute of Oceanography, Regional Centre, Visakhapatnam, India
- ⁴⁶Marine Information Research Center, Japan Hydrographic Association, Tokyo, Japan
- ⁴⁷Lamont Doherty Earth Observatory, Columbia University, Palisades, New York, USA
- ⁴⁸National Institute of Advanced Industrial Science and Technology, Tsukuba, Japan
- ⁴⁹Centre for Isotope Research, University of Groningen, Groningen, the Netherlands
- ⁵⁰Department of Marine Sciences, University of Connecticut, Groton, Connecticut, USA
- ⁵¹Department of Oceanography, Dalhousie University, Halifax, Canada
- *formerly at: School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich, UK

Correspondence to: D. C. E. Bakker (d.bakker@uea.ac.uk)

Received: 25 June 2013 – Published in Earth Syst. Sci. Data Discuss.: 16 August 2013

Revised: 12 January 2014 – Accepted: 24 January 2014 – Published:

Abstract. The Surface Ocean CO₂ Atlas (SOCAT), an activity of the international marine carbon research community, provides access to synthesis and gridded *f*CO₂ (fugacity of carbon dioxide) products for the surface oceans. Version 2 of SOCAT is an update of the previous release (version 1) with more data (increased from 6.3 million to 10.1 million surface water *f*CO₂ values) and extended data coverage (from 1968–2007 to 1968–2011). The quality control criteria, while identical in both versions, have been applied more strictly in version 2 than in version 1. The SOCAT website (<http://www.socat.info/>) has links to quality control comments, metadata, individual data set files, and synthesis and gridded data products. Interactive online tools allow visitors to explore the richness of the data. Applications of SOCAT include process studies, quantification of the ocean carbon sink and its spatial, seasonal, year-to-year and longer-term variation, as well as initialisation or validation of ocean carbon models and coupled climate-carbon models.

Data coverage

Repository-References: Individual data set files and synthesis product: doi:10.1594/PANGAEA.811776

Gridded products:

doi:10.3334/CDIAC/OTG.SOCAT_V2_GRID

Available at: <http://www.socat.info/>

Coverage: 79° S to 90° N; 180° W to 180° E

Location Name: Global Oceans and Coastal Seas

Date/Time Start: 16 November 1968

Date/Time End: 26 December 2011

1 Introduction

Human activity is releasing large quantities of the greenhouse gas carbon dioxide (CO₂) into the atmosphere. As a result, the atmospheric CO₂ mole fraction has increased from 280 μmol mol⁻¹ in pre-industrial times (Jansen et al., 2007) to 397 μmol mol⁻¹ in April 2013 (Tans and Keeling, 2014).

The rapid, ongoing change in the atmospheric composition by greenhouse gas emissions has been predicted to increase global mean temperature by 1.5 °C to 5.0 °C by the end of the century (Peters et al., 2013). Such warming would be accompanied by sea level rise, increased storm frequency, melting of ice caps and sea ice, changes in precipitation patterns and ocean acidification (Solomon et al., 2007), to name only the most prominent examples. Already many changes in the Earth's climate are apparent, such as the decline in Arctic sea ice extent (Stroeve et al., 2007), and warming in Alaska, near the Antarctic Peninsula (Vaughan et al., 2003; Mulvaney et al., 2012) and of the upper ocean (Levitus et al., 2005).

The oceans absorb a substantial part of the CO₂ emissions by human activity, thereby mitigating climate change. From pre-industrial times to 1994 the oceans have taken up 118 ± 19 Pg C from the atmosphere (Sabine et al., 2004). This is equivalent to roughly 50 % of CO₂ emissions from fossil fuel burning and cement production or 30 % of the total anthropogenic emissions, if CO₂ emissions from land use change are included. Recent estimates indicate that the oceans are a contemporary sink for roughly 27 % of the annual CO₂ emissions by fossil fuel combustion, cement production and land use change (Le Quéré et al., 2013). Uncertainty in the land use change emissions leads to a large error estimate for the proportion of the anthropogenic emissions taken up by the oceans.

There is uncertainty on how much CO₂ the oceans will absorb in a warming climate of the future (e.g. Jones et al., 2013). Considerable year-to-year, decadal and longer-term variation of CO₂ uptake is apparent in the North Atlantic Ocean (Corbière et al., 2007; Schuster and Watson, 2007; Thomas et al., 2008; Schuster et al., 2009; McKinley et al., 2011), the North Sea (Thomas et al., 2007), the North Pacific Ocean (Takamura et al., 2010), the equatorial Pacific Ocean (Feely et al., 2002, 2006; Ishii et al., 2004, 2009; Park et al., 2006, 2012) and the Southern Ocean (Le Quéré et al., 2007; Metzl, 2009), with large differences between ocean regions (Le Quéré et al., 2010; Lenton et al., 2012).

Measurements of CO₂ in the surface oceans (generally expressed as the mole fraction of CO₂ ($x\text{CO}_2$), partial pressure ($p\text{CO}_2$), or fugacity ($f\text{CO}_2$)) enable estimation of CO₂ air–sea fluxes and their variability. The fugacity can be measured underway on the surface water supply of ships. This method is used on a variety of ships, including ships of opportunity on commercial routes. The number of CO₂ measurements has greatly increased over the past four decades (Fig. 1) (Sabine et al., 2010). Data collection started in the late 1960s and 1970s, increased in the 1980s and intensified from the 1990s onwards. Roughly four times more data have been collected during the 2000s than in the 1990s. The growth in data collection has partly resulted from large international research programmes, for example JGOFS (Joint Global Ocean Flux Study) and WOCE (World Ocean Circulation Experiment), and regional funding initiatives. The development of autonomous instrumentation for the contin-




Figure 1. (a) The number of surface water $f\text{CO}_2$ values per year in SOCAT versions 1 and 2 and (b) per region per year in version 2. The SOCAT operationally defined region names are the Coastal Seas, the Arctic Ocean, the North Pacific Ocean, the Tropical Pacific Ocean, the North Atlantic Ocean, the Tropical Atlantic Ocean, the Indian Ocean and the Southern Ocean (Fig. 5, Table 5). These data points originate from data sets with flags of A, B, C or D and have a WOCE flag of 2. The subsequent figures only show $f\text{CO}_2$ values with these characteristics.

uous measurement of surface water $f\text{CO}_2$ (e.g. Körtzinger et al., 1996; Cooper et al., 1998; Pierrot et al., 2009), the inter-comparison of such instrumentation at sea (Körtzinger et al., 1996, 2000) and its installation on ships of opportunity (e.g. Cooper et al., 1998; Lüger et al., 2004; Schuster and Watson, 2007; Watson et al., 2009; Takamura et al., 2010; Lefèvre et al., 2013) and on moorings and drifters (e.g. Hood et al., 1999; Emerson et al., 2011) have played an important role in the increase in data collection.

Quantification of global and regional, annual mean ocean CO₂ uptake requires observations of surface water $f\text{CO}_2$ with adequate spatial and temporal coverage (Sweeney et al., 2000; Lenton et al., 2006). Studies of year-to-year, decadal and longer-term trends in air–sea CO₂ uptake necessitate consistent, multi-decade data records of surface ocean $f\text{CO}_2$ (e.g. Schuster and Watson, 2007; Park et al., 2012). Statistical techniques and modelling have been developed to infer basin-wide distributions of surface water $f\text{CO}_2$ from limited observations, for example a diffusion–advection based interpolation scheme (Takahashi et al., 1997, 2009), (multiple) linear regression (e.g. Boutin et al., 1999; Sarma et al., 2006), neural network approaches (e.g. Lefèvre et al., 2005) and a diagnostic ocean mixed layer model (Rödenbeck et al., 2013).

Uniform procedures for the collection, reporting, processing and archiving of CO₂ data, as well as public release of data, are essential for creating global and regional, long-term, consistent surface ocean $f\text{CO}_2$ synthesis products. Takahashi




Figure 2. The global distribution of surface water $f\text{CO}_2$ values in SOCAT version 2: (a) for 1968 to 2011 and (b) for 2008 to 2011.

and co-workers have constructed an impressive series of surface ocean CO₂ climatologies, the most recent one for the climatological year 2000 (Takahashi et al., 2009), and now provide annual updates to their global surface ocean $p\text{CO}_2$ data set (Takahashi et al., 2013). The Surface Ocean CO₂ Atlas (SOCAT) (Bakker et al., 2012; Pfeil et al., 2013; Sabine et al., 2013) complements this work. The SOCAT and Takahashi data sets benefit from standardisation and intercomparison of measurement and reporting protocols, as well as discussions between data providers and quality controllers on reporting standards and data quality (Dickson et al., 2007; IOCCP, 2008; SOCAT, 2011; Wanninkhof et al., 2013a). Both data sets contribute towards more rapid availability of ocean carbon data for synthesis products and policy-related assessments.

SOCAT is an international activity of ocean carbon scientists. It aims to create, make publicly available and archive the following (IOCCP, 2007):

- A 2nd level quality-controlled, global surface ocean $f\text{CO}_2$ data set following internationally agreed-upon procedures and regional review;
- A gridded data product of mean monthly surface water $f\text{CO}_2$ on a 1° latitude by 1° longitude grid with minimal temporal or spatial interpolation using the 2nd level quality-controlled, global surface ocean $f\text{CO}_2$ data set.

The first SOCAT release was made public as versions 1.4 and 1.5, here jointly referred to as version 1, in September 2011 (Bakker et al., 2012). SOCAT version 1 contains 6.3 million surface $f\text{CO}_2$ data points from 1851 data sets in the global oceans and coastal seas between 1968 and 2007 (Fig. 1, Table 1) (Pfeil et al., 2013; Sabine et al., 2013). Version 2 is presented here.

2 SOCAT version 2

2.1 An update of version 1

Version 2 is an update of version 1 with 60 % more data and 4 years extra data coverage. SOCAT version 2 contains 10.1 million surface $f\text{CO}_2$ values from 2660 data sets for the global oceans and coastal seas between November 1968 and December 2011 (Figs. 1 and 2). Version 2 was made public on 4 June 2013 at the 9th International Carbon Dioxide Conference in Beijing, China (SOCAT, 2013b).

SOCAT data products provide surface water $f\text{CO}_2$ values at sea surface temperature ($f\text{CO}_{2\text{rec}}$, with “rec” indicating recommended $f\text{CO}_2$), which have been (re-)calculated from the original CO₂ values reported by the data provider, following a strict calculation protocol. Sea surface temperature refers to the temperature at the seawater intake, often at about 5 m depth on ships. The procedures for data retrieval, for data entry, for the (re-)calculation of surface water $f\text{CO}_2$, for quality control, and for the creation of data products in version 2 are analogous to those used in version 1 (Pfeil et al., 2013; Sabine et al., 2013) and are described in Sects. 2.2, 2.3 and 2.4. The sections also highlight where version 2 differs from version 1 (Table 1).

Version 2 has three data products (Tables 2 and 3):

1. Individual data set files of surface water $f\text{CO}_2$ in a uniform format which have been subject to 2nd level quality control;
2. A synthesis data set of surface water $f\text{CO}_2$ for the global oceans and coastal seas;
3. Global gridded products of surface water $f\text{CO}_2$ means.

These data products are much the same as those for version 1 (Sect. 2.4) (Pfeil et al., 2013; Sabine et al., 2013). The SOCAT website (<http://www.socat.info/>) provides access to the data products together with online visualisation tools, data documentation, quality control comments, meeting reports, publications and a list of contributors (Tables 4, 5 and 6).

2.2 Data assembly and (re-)calculation of $f\text{CO}_2$ in version 2

2.2.1 Data origin

New data sets were either submitted directly to SOCAT or were retrieved from public websites hosted by the

Table 1. Key differences between SOCAT versions 1 (released as versions 1.4 and 1.5) and 2. Further details are in the text.

	Version 1	Version 2
Description	Pfeil et al. (2013); Sabine et al. (2013).	This study.
Data coverage	1968 to 2007, 6.3 million surface water <i>f</i> CO ₂ values, 1853 data sets.	1968 to 2011, 10.1 million surface water <i>f</i> CO ₂ values, 2660 data sets.
Time stamp	The time stamp did not contain seconds. Multiple entries for the same time stamp were reported in individual data set files (version 1.4), but were averaged in the synthesis files (version 1.5).	The time stamp includes seconds for all new and updated data sets. Seconds were added to time stamps for version 1 data sets to avoid concurrent entries. Artificial times with tenths and hundreds of a second have been generated for a dozen historical data sets reported at midnight or with few decimals in the time stamp.
Version numbers	Version numbers 1.4 and 1.5 highlighted the different treatment of multiple entries for the same time stamp.	Version 2 only.
Expocode	Expocodes were not used for moored and drifting buoys.	Expocodes are used for moored and drifting buoys.
Arctic region	Arctic data were included under the North Atlantic, North Pacific and coastal regions.	An Arctic region has been defined as all open ocean and coastal waters north of 70° N for 100° W to 43° E and north of 66° N elsewhere.
Identification of outliers	No systematic search was carried out for outliers and unrealistic values.	A systematic search for outliers and unrealistic values has been carried out. In total 154 data sets have been suspended.
Suspension of data sets	Data sets part of version 1.	Suspension of 70 data sets included in version 1 upon identification of unrealistic values.
WOCE flags of 2 (good), 3 (questionable), 4 (bad)	Virtually all <i>f</i> CO ₂ values were reported with a WOCE flag of 2.	WOCE flags of 2, 3 and 4 have been assigned to <i>f</i> CO ₂ values. Flags of 3 and 4 given during version 1 quality control (0.2 % of data) have been reinstated. A total of 20 850 <i>f</i> CO ₂ values (0.2 %) has been given a flag of 3 or 4 in version 2.
Parameters in the individual and synthesis files	Atmospheric CO ₂ mole fractions were from GLOBALVIEW-CO ₂ (2008). The files downloadable from the Cruise Data Viewer contained more parameters than the synthesis files.	Atmospheric CO ₂ mole fractions are from GLOBALVIEW-CO ₂ (2012). New parameters are the data set quality control flags of A to D and distance to a major land mass (Table 3). The parameters in files downloadable via the Cruise Data Viewer as “All Variables” and “Current Variable” match those in the synthesis files at CDIAC.
Gridded Data Viewer	Available	The capabilities of the Gridded Data Viewer have been expanded.
Release notes	None	Release notes document problems in version 2 data sets and data products.

Carbon Dioxide Information Analysis Center (CDIAC), PANGAEA®, institutions and research projects. Version 2 has an additional 3.8 million surface *f*CO₂ values and 807 data sets relative to version 1, mostly from 2006 to 2011 (Fig. 1, Table 1). Figure 3 shows the number of *f*CO₂ values from the 30 ships, including 1 ship-based time series, with the most intense data collection effort. The data sets in version 2 originate from 107 different ships, plus 3 ship-based

time series, 13 mooring-based time series and 3 drifters (Table 7). This study will adopt the term “data set” rather than “cruise” for individual data sets to reflect the presence of mooring and drifter data (0.7 % of *f*CO₂ values in version 2). Tools and parameters available online will be referred to by their name, e.g. “Cruise Data Viewer” (Sect. 2.4.5) and “cruise-weighted means” (Sect. 2.4.6).

Table 2. Key characteristics of the three SOCAT data products for surface ocean $f\text{CO}_2$ values in version 2 (Sect. 2.4). The synthesis product is available as synthesis files and as subsets of the global synthesis data set. The table lists whether the data products include only $f\text{CO}_2$ data with a WOCE flag of 2 (good) or also with flags of 3 (questionable) and 4 (bad). Information on access to metadata and quality control comments is provided. All data products can be accessed via the SOCAT website (<http://www.socat.info>) and via the links in the table.

	Characteristics	WOCE flag	Metadata	QC entries	Access and format
Individual data set files	The files contain all original CO ₂ measurements and $f\text{CO}_2$ values with a flag of 2, 3 and 4 for data sets with flags of A, B, C or D. Metadata accompany the files.	2, 3, 4	Yes	No	Text files at Pangaia ¹
Synthesis data set	Synthesis files consist of data sets with flags of A, B, C and D and contain $f\text{CO}_2$ values with a flag of 2. The global synthesis data set is available as global ^{2,3} and regional files ² .	2 only	No	No	Zip text files ² and in Ocean Data View format ³
Subset of synthesis data set (i)	Subset of the synthesis data set containing $f\text{CO}_2$ values with a flag of 2. Selection of “Include SOCAT invalids” gives access to $f\text{CO}_2$ values with a flag of 2, 3 and 4.	2 as default; 2, 3, 4 upon request	No	No	Text files via Cruise Data Viewer ⁴
Subset of synthesis data set (ii)	Subset of the synthesis data set containing original CO ₂ measurements and $f\text{CO}_2$ values with a flag of 2, 3 and 4. Metadata and quality control entries are available.	2, 3, 4	Yes	Yes	Text files via Table of Cruises ⁴
Gridded files	Gridded means of $f\text{CO}_2$ values on a 1° × 1° grid with minimal interpolation. Means are per year, monthly per year, monthly per decade and per climatological month from 1970 to 2011. A monthly 0.25° × 0.25° data set is available for coastal regions.	2 only	No	No	NetCDF files ⁵ , in Ocean Data View format ³ , and via Gridded Data Viewer ⁶

¹ doi:10.1594/PANGAEA.811776, ² <http://cdiac.ornl.gov/ftp/oceans/SOCATv2/>,

³ http://odv.awi.de/en/data/ocean/socat_fco2_data, ⁴ http://ferret.pmel.noaa.gov/SOCAT2_Cruise_Viewer/,

⁵ http://cdiac.ornl.gov/ftp/oceans/SOCATv2/SOCATv2_Gridded_Dat/, doi:10.3334/CDIAC/OTG.SOCAT_V2_GRID,

⁶ http://ferret.pmel.noaa.gov/SOCAT_gridded_viewer/

As in version 1 (Sect. 3.1 in Pfeil et al., 2013), most surface water CO₂ values have been measured by equilibration of a headspace with seawater and subsequent analysis of the CO₂ content of the headspace. Historical measurements generally used gas chromatographic analysis, while more recent measurements are based on infrared detection. SOCAT versions 1 and 2 include a small number of historical, discrete surface water $f\text{CO}_2$ measurements. SOCAT products do not include $f\text{CO}_2$ calculated from other carbon parameters, such as pH, alkalinity or dissolved inorganic carbon. A small percentage of the $f\text{CO}_2$ values (0.2% in version 2) is from measurements by a spectrophotometric method using a pH-sensitive dye (Table 7) (e.g. Hood et al., 1999).

2.2.2 Data entry

The data were assembled in a uniform file format, as in version 1 (Sect. 3.2 in Pfeil et al., 2013). Key differences in data entry between versions 1 and 2 relate to the time stamp, version numbering and an exocode for moorings and drifters, as described in Sect. 2.2.3.

Primary quality control was carried out at this stage. Primary quality control included identification of basic problems in the data, for example unrealistic positions, times and orders of magnitude. Additional basic problems were identified during secondary quality control (Sect. 2.3).

2.2.3 Key differences with version 1 in data entry

Time stamp and version numbering: the time stamp for SOCAT version 1 products did not contain seconds (Table 1) (Pfeil et al., 2013). In some cases this resulted in multiple entries for a given time stamp. Such multiple entries were averaged in the synthesis files (version 1.5), but not in the individual data set files (version 1.4). Two version numbers (version 1.4 and 1.5) highlight the different treatment of multiple entries for the same time stamp in the version 1 data products (Table 1).

SOCAT version 2 products include seconds, as reported by the data contributor, in the time stamp for all new and updated data sets (Table 1). However, a time stamp including seconds is not available for version 1 data sets. For these

Table 3. Content of the individual data set files (IF) and the synthesis files in SOCAT version 2. The global synthesis product is available as zip text files at CDIAC (ZIP) and in Ocean Data View (ODV) format. Subsets of the global synthesis data set can be created via the Cruise Data Viewer for All Variables (CDV_AV), Current Variable (CDV_CV) and via the Table of Cruises (CDV_TC). The first column (“Notation”) lists column headers for the parameters in the files.

Notation	IF	ZIP, CDV_AV	CDV_CV	ODV	CDV_TC	Unit	Description
Exocode	–	✓	✓	✓	✓	–	12-character exocode
SOCAT_DOI	–	✓	✓	✓	✓	–	Digital object identifier for the individual data set and metadata
QC_ID	–	✓	²	–	✓	–	Data set quality control flag with 11 for A, 12 for B, 13 for C and 14 for D
Date/Time	✓	–	✓	–	–	–	yyyy-mm-dd/hh:mm:ss (ISO8859)
yr	–	✓	–	✓	✓	Year	Year (UTC) ¹
mon	–	✓	–	✓	✓	Month	Month (UTC) ¹
day	–	✓	–	✓	✓	Day	Day (UTC) ¹
hh	–	✓	–	✓	✓	Hour	Hour (UTC) ¹
mm	–	✓	–	✓	✓	Minute	Minute (UTC) ¹
ss	–	✓	–	✓	✓	Seconds	Seconds (may include decimals) ¹
Time	–	–	²	–	–	Hour	Hours since 1970
Day of Year	–	–	²	–	–	Day of Year	Day of Year (UTC) with 1 January 00:00 as 1.0.
Longitude	✓	✓	✓	✓	✓	° E	Longitude (0 to 360) ¹
Latitude	✓	✓	✓	✓	✓	° N, ° S	Latitude (–90 to 90) ¹
Sample_depth/Depth water	✓	✓	²	✓	✓	m	Water sampling depth ^{1,3}
Sal	✓	✓	²	✓	✓	–	Salinity on Practical Salinity Scale ¹
Temp/SST	✓	✓	²	✓	✓	°C	Sea surface temperature ¹
Tequ	✓	✓	²	✓	✓	°C	Equilibrator chamber temperature ¹
PPPP	✓	✓	²	✓	✓	hPa	Atmospheric pressure ¹
Pequ	✓	✓	²	✓	✓	hPa	Equilibrator chamber pressure ¹
WOA_SSS/Sal interp	✓	✓	²	✓	✓	–	Salinity from WOA (2005) ⁴
NCEP_SLP/PPPP interp	✓	✓	²	✓	✓	hPa	NCEP Atmospheric pressure ⁵
ETOPO2_depth/Bathy depth interp	✓	✓	²	✓	✓	m	ETOPO2 Bathymetry ⁶
Distance/d2l	✓	✓	²	–	✓	km	Distance to major land mass
GVCO2/xCO ₂ air_interp	✓	✓	²	✓	✓	μmol mol ^{–1}	Atmospheric xCO ₂ from GLOBALVIEW-CO ₂ (2012)
xCO ₂ water_equ_dry	✓	–	–	–	✓	μmol mol ^{–1}	xCO ₂ (water) at equilibrator temperature (dry air) ¹
fCO ₂ water_SST_wet	✓	–	–	–	✓	μatm	fCO ₂ (water) at sea surface temperature (air at 100 % humidity) ¹
pCO ₂ water_SST_wet	✓	–	–	–	✓	μatm	pCO ₂ (water) at sea surface temperature (air at 100 % humidity) ¹
xCO ₂ water_SST_dry	✓	–	–	–	✓	μmol mol ^{–1}	xCO ₂ (water) at sea surface temperature (dry air) ¹
fCO ₂ water_equ_wet	✓	–	–	–	✓	μatm	fCO ₂ (water) at equilibrator temperature (air at 100 % humidity) ¹
pCO ₂ water_equ_wet	✓	–	–	–	✓	μatm	pCO ₂ (water) at equilibrator temperature (air at 100 % humidity) ¹
fCO ₂ rec/fCO ₂ water_SST_wet	✓	✓	²	✓	✓	μatm	Recommended fCO ₂ calculated following the SOCAT protocol
fCO ₂ rec_src/Algorithm	✓	✓	²	✓	✓	–	Algorithm for calculating fCO ₂ rec (0: not generated; algorithm 1–14 in Table 8)
fCO ₂ rec_flag/Flag	✓	✓	²	–	✓	–	WOCE flag for fCO ₂ rec (2: good, 3: questionable, 4: bad) ⁷

¹ Data reported by the data originator.

² Available upon selection of parameter.

³ If the intake depth has not been reported by the data originator, an intake depth of 5 m has been assumed.

⁴ Sea surface salinity on the Practical Salinity Scale interpolated from the World Ocean Atlas (WOA) 2005 (Antonov et al., 2006), available at: http://www.nodc.noaa.gov/OC5/WOA05/pr_woa05.html (last access: 1 May 2013).

⁵ Atmospheric pressure interpolated from the NCEP/NCAR (National Centers for Environmental Prediction/National Center for Atmospheric Research) 40-Year Reanalysis Project on a 6-hourly, global, 2.5° latitude by 2.5° longitude grid (Kalnay et al., 1996), available at: <http://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.surface.html> (last access: 1 May 2013).

⁶ Bathymetry interpolated from ETOPO2 (2006) 2-minute Gridded Global Relief Data, available at: <http://www.ngdc.noaa.gov/mgg/global/etopo2.html> (last access: 1 May 2013).

⁷ Individual data set files contain all fCO₂rec data. Synthesis files at CDIAC and via ODV only contain fCO₂rec data with a WOCE flag of 2 (Table 2).

Table 4. Activities and key participants in SOCAT versions 2 and 3 to date. Regional group leads are in Table 5.

Activity	Key Participants
Global group for coordination	Bakker (chair), Hankin, Kozyr, Metzl, Olsen, Pfeil, Pierrot, Telszewski
Data retrieval, data entry, (re-)calculation of $f\text{CO}_2$	Pfeil, Olsen
Quality control	Alin, Bakker, Barbero, Castle, Cosca, Evans, Hales, Harasawa, Hoppema, Huang, Hunt, Huss, Park, Paterson, Pierrot, Schuster, Skjelvan, Steinhoff, Suzuki, Tilbrook, Van Heuven, Vlahos, Wada, Wanninkhof
Live Access Server	Hankin, O'Brien, Smith
Individual data set files, synthesis products and gridded products	Pfeil, Smith, Manke, Hankin
Ocean Data View	Schlitzer
Matlab files	Pierrot, Landschützer
SOCAT website	Pfeil
Data archiving and online access	Pfeil, Sieger, Kozyr, Smith, Manke, Hankin
Meetings	Alin, Bakker, Hales, Hankin, Nojiri, Telszewski
Alternative sensors (version 3)	Wanninkhof, Steinhoff, Bakker, Bates, Boutin, Olsen, Sutton
Automation (versions 3 to 4)	Hankin, S. Jones, Kozyr, O'Brien, Pfeil, Smith, Bakker, Olsen, Schweitzer

data sets, seconds were added artificially to the time stamp to avoid the problem of multiple entries. The next version of SOCAT will include seconds, as reported by the data contributor, for all data sets.

The CO₂ measurements for a dozen historical data sets are listed at midnight or their time stamp in fractional days contains insufficient decimals for retrieving minutes and seconds. Artificial seconds, in some cases including tenths or hundreds of a second, were generated for these valuable data, such that they can remain in SOCAT version 2. Every effort will be made to retrieve a more adequate time stamp for these data sets for future versions. Unlike version 1, which has version 1.4 and 1.5 data products, version 2 data products have a single version number (Table 1).

Expocode for moorings and drifters: SOCAT uses twelve character expocodes (Swift, 2008) as stable and unique data set identifiers. For example, 49P120101218 indicates a cruise on the Japanese (49) ship of opportunity Pyxis (P1) with the first day of the cruise on 18 December 2010. In contrast to version 1, expocodes have been assigned for moorings and drifters in version 2, by registering a “vessel code” at International Council for the Exploration of the Sea (ICES) in collaboration with the National Oceanographic Data Center (NODC) and the British Oceanographic Data Centre (BODC) (Table 1).

2.2.4 (Re-)calculation of recommended $f\text{CO}_2$

Surface water $f\text{CO}_2$ values at sea surface temperature (or intake temperature), also known as recommended $f\text{CO}_2$ ($f\text{CO}_2\text{rec}$), have been recalculated, analogous to version 1 (Sect. 3.3 in Pfeil et al., 2013). A single set of equations and a strict order of preference for the CO₂ input parameter has been used (Table 8) (Pfeil et al., 2013). Six different CO₂ parameters were reported in the original data files, notably $x\text{CO}_2$, $p\text{CO}_2$ and $f\text{CO}_2$, either at the equilibration temperature (Tequ) or at the sea surface temperature (SST).

The (re-)calculation procedure of $f\text{CO}_2$ has the following philosophy (Pfeil et al., 2013):

1. Whenever possible, (re-)calculate $f\text{CO}_2$;
2. The favourite starting point for the calculations is $x\text{CO}_2$, next $p\text{CO}_2$, followed by $f\text{CO}_2$;
3. Minimise the amount of external data required for the calculations.

Table 8 lists surface water CO₂ parameters and ancillary parameters used for calculation of recommended $f\text{CO}_2$ in version 2 in order of preference with algorithm 1 as the favourite (analogous to Table 4 in Pfeil et al., 2013). The algorithm is provided in the output files (Table 3). Equations recommended by Dickson et al. (2007) have been used for the conversion of the dry CO₂ mole fraction to $p\text{CO}_2$, for the calculation of the water vapour pressure and for the conversion of $p\text{CO}_2$ to $f\text{CO}_2$, similar to version 1 (Sect. 3.3 of Pfeil et al., 2013). As in version 1, the correction of Takahashi et al. (1993) has been applied for temperature change between the seawater intake and the site of equilibration:

$$f\text{CO}_2^{\text{SST}} = f\text{CO}_2^{\text{equT}} \exp(0.0423(\text{SST} - \text{Tequ})). \quad (1)$$

Climatological values of salinity and atmospheric pressure from reanalysis have been used in the calculation of recommended $f\text{CO}_2$ (Table 8), if the data contributor did not report in situ salinity and atmospheric pressure, following Pfeil et al. (2013).

Table 5. Regions and regional group leads in SOCAT version 2 (Fig. 5).

Region	Definition	Lead(s)
Coastal Seas	Less than 400 km from land; between 30° S and 70° N for 100° W to 43° E; between 30° S and 66° N elsewhere	Alin, Cai, Hales
Arctic Ocean	North of 70° N for 100° W to 43° E; north of 66° N elsewhere, including coastal waters	Mathis
North Atlantic	30° N to 70° N	Schuster
Tropical Atlantic	30° N to 30° S	Lefèvre
North Pacific	30° N to 66° N	Nojiri
Tropical Pacific	30° N to 30° S	Cosca
Indian Ocean	North of 30° S	Sarma
Southern Ocean	South of 30° S, including coastal waters	Tilbrook, Metzl

Table 6. Meetings for SOCAT versions 2 and 3 to date.

Timing	Meeting description	Location	Reference
09/2011	Public release of version 1. Session on future SOCAT.	UNESCO, Paris, France	(SOCAT, 2011)
05/2012	Automation planning meeting	NOAA-PMEL, Seattle, USA	(SOCAT, 2012a)
07/2012	SOCAT progress meeting	Epochal Centre, Tsukuba, Japan	(SOCAT, 2012b)
10/2012	Coastal and Arctic SOCAT quality control workshop	NOAA-PMEL, Seattle, USA	(IOCCP, 2012)
06/2013	SOCAT side event at the 9th International Carbon Dioxide Conference. Public release of version 2.	Beijing International Convention Center, Beijing, China	(SOCAT, 2013b)

2.3 Secondary quality control in version 2

2.3.1 Secondary quality control criteria

Criteria for 2nd level quality control have been defined in a series of workshops (IOCCP, 2008, 2009, 2010; Pfeil et al., 2013). Second level quality control consists of assigning a quality control flag to each data set and a WOCE flag to individual surface water $f\text{CO}_2$ values. The criteria for quality control are identical in versions 1 (Sect. 4.1 in Pfeil et al., 2013) and 2.

Only data sets with a quality control flag of A, B, C and D are included in SOCAT version 1 and 2 data products (Table 9) (Pfeil et al., 2013). The data set quality control flags (formerly known as “cruise flags”) in versions 1 and 2 have been developed for automated shipboard measurement of surface water $f\text{CO}_2$, mainly by infrared detection and frequent at sea standardisation using calibration gases with a range of CO₂ concentrations (IOCCP, 2008; Pfeil et al., 2013). Much weight is put on whether approved methods or standard operating procedures (SOP) (AOML, 2002; Dickson et al., 2007; Pierrot et al., 2009) were followed by making this a prerequisite for flags of A and B. Citing Pfeil et al. (2013):

“Seven SOP criteria need to be fulfilled for a cruise (or data set) flag of A or B in SOCAT:

1. The data are based on $x\text{CO}_2$ analysis, not $f\text{CO}_2$ calculated from other carbon parameters, such as pH, alkalinity or dissolved inorganic carbon;
2. Continuous CO₂ measurements have been made, not discrete CO₂ measurements;
3. The detection is based on an equilibrator system and is measured by infrared analysis or gas chromatography;
4. The calibration has included at least 2 non-zero gas standards, traceable to World Meteorological Organization (WMO) standards;
5. The equilibrator temperature has been measured to within 0.05 °C accuracy;
6. The intake seawater temperature has been measured to within 0.05 °C accuracy;
7. The equilibrator pressure has been measured to within 0.5 hPa accuracy.”

The $f\text{CO}_2$ values from data sets with flags of A and B are judged to have an accuracy of $\pm 2 \mu\text{atm}$ or better. Criterion 1




Figure 3. The number of surface water $f\text{CO}_2$ values obtained on the 30 ships, including 1 ship-based time series, hosting the most intense data collection effort in SOCAT version 2.

also needs to be met for flags of C and D, similar to version 1 (Sect. 4.1 in Pfeil et al., 2013). Complete metadata documentation is required for data set quality control flags of A, B and C. Comparison to other data is carried out, if possible. The overall quality of the data needs to be deemed acceptable for flags of A, B, C and D (Table 9) (Pfeil et al., 2013).

The Southern and Indian Ocean groups (Table 5) have applied three additional quality control criteria for the temperature change between the seawater intake and the equilibrator in versions 1 and 2 (IOCCP, 2010), citing Pfeil et al. (2013):

- “Warming should be less than 3 °C;
- Warming rate should be less than 1 °C h⁻¹, unless a rapid temperature front is apparent;
- Warming outliers should be less than 0.3 °C, compared to background data.”

In addition:

- Cooling between the seawater intake and the equilibrator is unlikely in high-latitude oceans for an indoor measurement system;
- Zero or constant temperature change may indicate absence of SST values.

The above five guidelines have been applied widely in version 2 for open ocean data away from sea ice, as part of a systematic search for unrealistic data and outliers (Sect. 2.3.3). Such a systematic search has not been carried out for version 1 (Table 1).

These quality control criteria (Table 9) have also been applied for quality control of surface water CO₂ measurements from moorings and drifters in versions 1 and 2 (Table 7).

Individual $f\text{CO}_2$ values are assigned WOCE flags: 2 (good), 3 (questionable) or 4 (bad) with 2 being the default




Figure 4. The number of quality controllers in SOCAT versions 1 and 2 based in Europe, the USA, Asia and Australia, respectively. The figure demonstrates the international character of the quality control effort in SOCAT.

setting (Sect. 4.1.3 in Pfeil et al., 2013). Outliers in parameters required for the timing, position and (re-)calculation of $f\text{CO}_2$ values are given flags of 3 and 4. Thus, flags of 3 and 4 might indicate an erroneous time or position stamp, an unrealistic seawater temperature, strong warming between the seawater intake and the equilibrator or a large pressure difference between the equilibrator and the atmosphere. Data sets with a large number of flags of 3 and 4 (> 50, as a guide line) are suspended, as was also the case for version 1 (Pfeil et al., 2013).

2.3.2 Secondary quality control in practice

Secondary quality control for version 2 has been carried out by 24 marine carbon scientists from eight countries (Fig. 4, Table 4). Quality control in SOCAT is carried out by groups organised according to region. These regions have been operationally defined and do not necessarily follow common oceanographic definitions. Regions for version 2 are the Coastal Seas, the North Atlantic, Tropical Atlantic, North Pacific, Tropical Pacific, Indian Ocean and Southern Oceans and a newly defined Arctic region (Fig. 5, Table 5). The Arctic region includes both coastal seas and the deep ocean. It encompasses all waters north of 70° N for 100° W to 43° E (Atlantic sector) and north of 66° N elsewhere (Table 1) (Sect. 2.3.3) (SOCAT, 2012b).

The regional group members assign data set quality control flags, WOCE flags and enter quality control comments. The Live Access Server (<http://ferret.pmel.noaa.gov/LAS>) is used for quality control, as in version 1 (Sect. 4 in Pfeil et al., 2013). Quality control comments are available via the Table of Cruises on the Cruise Data Viewer (Table 2) (Sect. 2.4.5).

All new and updated data sets are subject to quality control. Each data set is assigned a flag of A, B, C, D, S (Suspend) or X (Exclude) for each region it crosses. As a final step the quality controllers need to resolve any “conflicting”

Table 7. Drifters and time series in SOCAT version 2 with the location, year(s) of operation, platform type, CO₂ instrument type, algorithm used for (re-)calculation of *f*CO₂ (Table 8), number of data sets, number of *f*CO₂ values with a WOCE flag of 2, and reference.

Drifters and time series	Location	Year(s)	Platform type	CO ₂ instrument type	Algorithm	Number data sets	Number <i>f</i> CO ₂ values	Reference
CARIOCA	75.0° N 3.0° W	1996–1997	Drifter	Membrane spectrophotometer	6	1	2668	H1999
CARIOCA	0.4° S 7.8° W	1997	Drifter	Membrane spectrophotometer	6	1	1964	B2001
CARIOCA	40.1° S 15.8° E	2005	Drifter	Membrane spectrophotometer	6	1	1451	BM2009
PIRATA_10W_6S	6° S 10° W	2006–2007	Mooring	Membrane spectrophotometer	6	2	11 820	L2008
Papa_145W_50N	50.1° N 144.8° W	2007	Mooring	Equilibrator-IR	6	1	4987	J2010
JKEO_147E_38N	37.9° N 146.6° E	2007	Mooring	Equilibrator-IR	6	1	927	J2010
KEO_145E_32N	32.3° N 144.5° E	2007–2008	Mooring	Equilibrator-IR	6	2	4740	J2010
MOSEAN_158W_23N	22.8° N 158.1° W	2004–2007	Mooring	Equilibrator-IR	6	5	6034	J2010
WHOTS_158W_23N	22.7° N 158.1° W	2007	Mooring	Equilibrator-IR	6	1	4750	J2010
CRIMP1	21.4° N 157.8° W	2005	Mooring	Equilibrator-IR	6	1	1993	J2010
TAO_170W_0	0.0° S 170.0° W	2005	Mooring	Equilibrator-IR	6	1	2577	J2010
TAO_155W_0	0.0° W 155.0° W	2005	Mooring	Equilibrator-IR	6	1	2198	J2010
TAO_140W_0	0.0° N 139.8° W	2004–2007	Mooring	Equilibrator-IR	6	5	5253	J2010
TAO_125W_0	0.2° S 124.4° W	2004–2007	Mooring	Equilibrator-IR	6	4	3686	J2010
BTM_64W_32N	31.8° N 64.2° W	2005–2007	Mooring	Equilibrator-IR	6	3	5095	J2010
Stratus_85W_20S	19.7° S 85.6° W	2006	Mooring	Equilibrator-IR	6	1	9466	J2010
Station M	66.0° N 2.0° E	2006–2007	Ship	Equilibrator-IR	1	19	159 671	WT1993
Station P	50° N 145° W	1973–1976	Ship	Equilibrator-IR	6	12	4158	None
Western Channel Observatory	50.1° N 4.3° W	2007–2009	Ship	Equilibrator-IR	1	1	899	HM2008, K2012

References are: B2001 – Bakker et al. (2001); BM2009 – Boutin and Merlivat (2009); H1999 – Hood et al. (1999); HM2008 – Hardman-Mountford et al. (2008); J2010 – Johengen (2010); K2012 – Kitidis et al. (2012); L2008 – Lefèvre et al. (2008); WT1993 – Wanninkhof and Thoning (1993).




Figure 5. Quality control regions for SOCAT version 2 (Table 5). White shading corresponds to the coastal region. The regions have been defined for operational reasons and do not necessarily reflect common oceanographic definitions.

data set flags between regions and decide on the “agreed” flag for a data set. The data set quality control flag has been added as a parameter in the synthesis files (Tables 1 and 3) (Sects. 2.4.4 and 2.4.7).

Overall data quality and reporting of metadata has improved from version 1 to version 2, which we attribute to the SOCAT effort. In version 1, 41 % of the data sets were assigned a flag of A or B, 22 % obtained a flag of C and, 37 % received a flag of D. Version 2 has a larger proportion of data

sets with flags of A or B (48 %) and smaller proportions of data sets with a flag of C (18 %) and D (33 %).

2.3.3 Key differences with version 1 in secondary quality control

This section identifies key differences in secondary quality control between versions 1 and 2.

Creation of an Arctic regional designation: in version 1 Arctic data were part of the North Pacific and North Atlantic Oceans and the Coastal Region (Sect. 2.2 in Pfeil et al., 2013). Given the importance of Arctic research and the rapid increase in the quantity of Arctic *f*CO₂ values, an Arctic region has been defined for version 2 (Figs. 5 and 6; Table 1) (Sect. 2.3.2) (SOCAT, 2012b).

Identification of unrealistic values: in version 2 a systematic search has been carried out for unrealistic values or patterns in all data relevant for the timing, position or (re-)calculation of *f*CO₂ (Sect. 2.3.1). This activity considered all the data sets submitted to version 2, including data sets previously included in the version 1 data release. The search applied to the ship’s cruise track, position, time, atmospheric pressure, equilibrator pressure, salinity, sea surface temperature, equilibrator temperature, and temperature change between the seawater intake and the equilibrator. This helped locate problems with data entry, e.g. overlap between data sets, reversal of hours and minutes, reversal of SST and salinity, presence of undefined values (e.g. –999, –99, –9.999, –9.99, –9.9, –9, 0), identification of unrealistic values (e.g. an atmospheric pressure of 780 mbar) and problems with water flow (absolute temperature change between intake and equilibrator exceeding 3 °C). Depending on the nature of

Table 8. Surface water CO₂ parameters used for the calculation of recommended $f\text{CO}_2$ ($f\text{CO}_{2\text{rec}}$) at sea surface temperature in version 2 (after Table 4 in Pfeil et al., 2013). The parameters are listed in order of preference (with algorithm 1 as the favourite). The algorithm is provided in the output files (Table 3). In cases of incomplete data reporting, these ancillary parameters have been used for atmospheric pressure and salinity: NCEP (National Centers for Environmental Prediction) atmospheric pressure (Kalnay et al., 1996) and WOA (World Ocean Atlas) salinity (Antonov et al., 2006) (Sect. 2.2.4).

Algorithm	Reported CO ₂ parameter	Unit	Data Percentage (%)	Extra variable
1	$x\text{CO}_2\text{water_equi_dry}$	$\mu\text{mol mol}^{-1}$	66.7	–
2	$x\text{CO}_2\text{water_SST_dry}$	$\mu\text{mol mol}^{-1}$	4.5	–
3	$p\text{CO}_2\text{water_equi_wet}$	μatm	4.5	–
4	$p\text{CO}_2\text{water_SST_wet}$	μatm	2.6	–
5	$f\text{CO}_2\text{water_equi}$	μatm	0.2	–
6	$f\text{CO}_2\text{water_SST_wet}$	μatm	10.8	–
7	$p\text{CO}_2\text{water_equi_wet}^1$	μatm	0.3	NCEP Pressure
8	$p\text{CO}_2\text{water_SST_wet}^1$	μatm	8.3	NCEP Pressure
9	$x\text{CO}_2\text{water_equi_dry}^2$	$\mu\text{mol mol}^{-1}$	0.2	WOA Salinity
10	$x\text{CO}_2\text{water_SST_dry}^2$	$\mu\text{mol mol}^{-1}$	0.7	WOA Salinity
11	$x\text{CO}_2\text{water_equi_dry}^1$	$\mu\text{mol mol}^{-1}$	0.01	NCEP Pressure
12	$x\text{CO}_2\text{water_SST_dry}^1$	$\mu\text{mol mol}^{-1}$	1.0	NCEP Pressure
13	$x\text{CO}_2\text{water_equi_dry}^{1,2}$	$\mu\text{mol mol}^{-1}$	0.01	NCEP Pressure, WOA Salinity
14	$x\text{CO}_2\text{water_SST_dry}^{1,2}$	$\mu\text{mol mol}^{-1}$	0.1	NCEP Pressure, WOA Salinity

¹ Atmospheric pressure was not reported in the original data file.

² Salinity was not reported in the original data file.

the problem, this resulted in suspension of a data set (Table 10) or assignation of a WOCE flag of 3 (questionable) or 4 (bad) to individual $f\text{CO}_2$ values.

In total, 154 data sets have been suspended, of which 70 had previously been included in the version 1 release (Table 1). Table 10 lists grounds for suspension of data sets. These include absence of CO₂ values (14 %), a data entry problem (10 %), use of a constant atmospheric pressure or salinity in the calculation of $f\text{CO}_2$ (45 %), absence of SST (8 %), and concerns on the quality of $f\text{CO}_2$ (3 %), temperature (14 %), or atmospheric pressure (2 %). In case of a data entry problem, data sets will be re-entered into the SOCAT quality control system for version 3. In other cases, data sets may need revision before resubmission to SOCAT. Finally, six data sets (4 %) made by a spectrophotometric method were suspended, as the data set flags of A to D were not deemed appropriate by the quality controller. In response, a new data set flag of E has been defined for use in version 3 (Sect. 4.2) (Wanninkhof et al., 2013a).

Suspension of 70 data sets included in version 1: 70 data sets previously included in version 1 were suspended from version 2 upon identification of data quality concerns (Tables 1 and 10), as discussed above. Most of these (59) were suspended as a constant atmospheric pressure had been used in the calculation of $f\text{CO}_2$. These 59 data sets have since been revised and resubmitted to SOCAT for version 3. Six data sets were suspended for a data entry problem, while three data sets lacked surface water CO₂ values. Concerns on

the quality of a temperature or atmospheric pressure reading were grounds for suspension of a further two data sets.

WOCE flags of 3 and 4: WOCE flags of 2 (good), 3 (questionable) and 4 (bad) have been assigned to all $f\text{CO}_2$ values in version 2, including for data sets part of the version 1 release. During version 1 quality control, 0.2 % of the $f\text{CO}_2$ values had been assigned a flag of 3 or 4. However, these flags were accidentally reset to a flag of 2 prior to the version 1 release, such that most $f\text{CO}_2$ values in version 1 were reported with a flag of 2. The initial flags of 3 and 4 set during version 1 quality control have been reinstated in version 2. In version 2, a total of 20 850 $f\text{CO}_2$ values (0.2 %) has been given a flag of 3 or 4.

2.4 Version 2 data policy and data products

2.4.1 Data policy

Users of the SOCAT data products are requested to do the following (SOCAT, 2013a, b):

1. Recognise the contribution of SOCAT data contributors and quality controllers in the form of invitation to co-authorship or citation of relevant scientific articles by data contributors;
2. Cite all SOCAT data products by reference to publications documenting SOCAT;

Table 9. Criteria for assigning data set quality control flags based on the expected quality of the recommended $f\text{CO}_2$ values (per Table 6 in Pfeil et al., 2013). All criteria need to be met for assigning a data set flag. Only data sets with a quality control flag of A, B, C and D are included in version 1 and 2 data products. SOP is Standard Operating Procedures (Dickson et al., 2007); QC is quality control.

Data set flag (ID)	Criteria
A (11)	1. Followed approved methods or SOP criteria and 2. Metadata documentation complete and 3. Extended QC was deemed acceptable and 4. A comparison with other data was deemed acceptable.
B (12)	1. Followed approved methods or SOP criteria and 2. Metadata documentation complete and 3. Extended QC was deemed acceptable.
C (13)	1. Did not follow approved methods or SOP criteria but 2. Metadata documentation complete and 3. Extended QC was deemed acceptable (including comparison with other data if possible).
D (14)	1. Did or did not follow approved methods or SOP criteria and 2. Metadata documentation incomplete but 3. Extended QC was deemed acceptable (including comparison with other data if possible).
S (Suspend)	1. Did or did not follow methods or SOP criteria and 2. Metadata documentation complete or incomplete and 3. Extended QC revealed non-acceptable data but 4. Data are being updated.
X (15) (Exclude)	The data set duplicates another data set in SOCAT.
N (No flag)	No data set flag has yet been given to this data set.
U (Update)	The data set has been updated. No data set flag has yet been given to the revised data.

3. Send references of publications using SOCAT products to submit@socat.info.

2.4.2 SOCAT data products

The SOCAT data products provide access to recommended surface ocean $f\text{CO}_2$ values in a uniform format for the global oceans and coastal seas. Three different SOCAT data products are available: individual data set files, synthesis files and gridded files. User-defined subsets of the synthesis files are available via the Cruise Data Viewer. The Gridded Data Viewer facilitates querying of the gridded data products. All data products can be accessed via the SOCAT website (<http://www.socat.info/>) or via the web-links provided below. Table 2 identifies the key characteristics of the SOCAT data products, while Table 3 lists the contents of downloadable files. The version 2 data products resemble those for version 1 (Pfeil et al., 2013; Sabine et al., 2013), apart from further standardisation and extra parameters. The data products and tools are discussed below, followed by a description of key differences between version 1 and 2 (Table 1) (Sect. 2.4.7).

2.4.3 Individual data set files

Individual data set files provide surface water $f\text{CO}_2$, the parameters used to (re-)calculate $f\text{CO}_2$ and the original CO_2 parameter(s) reported by the data contributor for data sets with a flag of A, B, C or D (Table 2). The files include all surface water $f\text{CO}_2$ values with WOCE flags of 2, 3 and 4. Individual data set files are archived at PANGAEA® (doi:10.1594/PANGAEA.811776). Each data set has a digital object identifier (doi) (Table 3). Metadata provided by the data contributor accompany the data set files. As in version 1, the individual data set and synthesis files include the climatological values of salinity and atmospheric pressure from reanalysis. The files also contain values for the water depth, the distance to a major land mass and the atmospheric CO_2 mole fraction interpolated from GLOBALVIEW-CO2 (2012). Via PANGAEA®, version 2 is made available to the World Data System (WDS) of the International Council for Science (ICSU), to the Group of Earth Observations (GEO) Portal and to the Global Earth Observation System of Systems (GEOSS).

Table 10. Grounds for suspension of data sets from SOCAT version 2. A distinction is made between data sets previously included in version 1 and new data sets in version 2. Abbreviations are SST for sea surface temperature, Tequ for equilibrator temperature and dT for the difference between the equilibrator temperature and the sea surface temperature.

Ground for suspension	Number data sets version 1	Number data sets version 2	Percentage of total (%)
Overlap with other data set	0	1	1
Data entry problem (incomplete data set, time, position, SST, salinity)	6	9	10
Constant atmospheric pressure in calculation of $f\text{CO}_2\text{rec}$	59	5	42
Constant salinity (0 or -999) in calculation of $f\text{CO}_2\text{rec}$	0	5	3
No $x\text{CO}_2$, $p\text{CO}_2$ or $f\text{CO}_2$ reported	3	19	14
No SST reported	0	13	8
Concerns on quality of $f\text{CO}_2\text{rec}$	0	4	3
Concerns on quality of SST, Tequ or dT	1	20	14
Concerns on quality of atmospheric pressure	1	2	2
No appropriate sensor flag	0	6	4
Total	70	84	100

2.4.4 Global synthesis product

The global synthesis data set consists of individual data sets with flags of A, B, C and D and contains $f\text{CO}_2$ values with a WOCE flag of 2 (Table 2). The synthesis files do not contain the original CO₂ values (Table 3). Each line in the files lists the doi-number of the corresponding individual data set file at PANGAEA® (Sect. 2.4.3). The synthesis data set is available as global and regional files for the SOCAT regions (Fig. 5). The regional files only contain data from within that region, so that data from most cruises are split between several regional files. The global and regional files are publicly available as compressed zip text files via CDIAC (<http://cdiac.ornl.gov/ftp/oceans/SOCATv2/>). Matlab code is available for reading these synthesis files. The global synthesis product is also available in Ocean Data View format (http://odv.awi.de/en/data/ocean/socat_fCO2_data).

2.4.5 Subsetting the global synthesis product

The Cruise Data Viewer (http://ferret.pmel.noaa.gov/SOCAT2_Cruise_Viewer/), an interactive tool on the Live Access Server, enables searching and subsetting the global synthesis data set by year, month, day, region, parameter, expocode, cruise name, vessel, and data set quality control flag. One may define search limits, for example salinity below 32. The user can create property-property plots and download data. The default setting is access to $f\text{CO}_2$ values with a WOCE flag of 2 (Table 2). However, the user can include data with flags of 3 (questionable) and 4 (bad) by selecting “Include SOCAT invalids”. Figures 2 and 7 have been made with the Cruise Data Viewer.

The Table of Cruises, available via the pull-down menu “Tables” on the Cruise Data Viewer, enables the user to find metadata and read quality control comments for specific data sets (Table 2). Files downloadable via the Table of Cruises contain $f\text{CO}_2$ values with WOCE flags of 2, 3 and 4 and the original CO₂ data (Table 3).

2.4.6 Gridded products

Sabine et al. (2013) detail the gridding of the $f\text{CO}_2$ values on a 1° latitude by 1° longitude grid with a higher 0.25° latitude by 0.25° longitude resolution product for the coastal seas in version 1. The procedures for gridding the data are identical in versions 1 and 2.

Several gridded products of surface ocean $f\text{CO}_2$ means with minimal interpolation are available (doi:10.3334/CDIAC/OTG.SOCAT_V2_GRID). Surface water $f\text{CO}_2$ values with a flag of 2 have been put on a 1° latitude by 1° longitude grid in four ways: per year, monthly per year, monthly per decade, and per climatological month from 1970 to 2011 (Table 2). A higher resolution of 0.25° latitude by 0.25° longitude is available as monthly means per year for the coastal region (Fig. 5).

Gridded $f\text{CO}_2$ values are reported as unweighted means and as cruise-weighted (or data set-weighted) means (Sabine et al., 2013). In an unweighted mean all the recommended $f\text{CO}_2$ values in a grid cell have been given equal weight in calculating the mean. In a cruise-weighted mean, first averages of the recommended $f\text{CO}_2$ values per cruise (or data set) have been calculated within a grid cell, before averages of the cruise means have been determined. Grid cells without $f\text{CO}_2$ values are empty. No correction has been made for the expected long-term increase in surface water $f\text{CO}_2$,




Figure 6. The density of surface water $f\text{CO}_2$ values for each region in SOCAT versions 1 and 2. Regions are the Coastal Seas, the Arctic Ocean, the North Pacific Ocean, the Tropical Pacific Ocean, the North Atlantic Ocean, the Tropical Atlantic Ocean, the Indian Ocean and the Southern Ocean (Fig. 5, Table 5). In version 1, Arctic data were included in the North Pacific, North Atlantic and Coastal Regions.

something users of the monthly climatological and decadal gridded products should keep in mind. Furthermore, the gridded products may have a temporal bias in grid cells with uneven temporal data coverage. For example, an annual gridded product may have a strong seasonal bias, if only summertime $f\text{CO}_2$ values are available.

Gridded $f\text{CO}_2$ products can be accessed as NetCDF files from CDIAC (http://cdiac.ornl.gov/ftp/oceans/SOCATv2/SOCATv2_Gridded_Dat/), in Ocean Data View format (http://odv.awi.de/en/data/ocean/socat_fCO2_data) and via the Gridded Data Viewer (http://ferret.pmel.noaa.gov/SOCAT_gridded_viewer/) (Table 2). Matlab code is available for reading the NetCDF files.

The capabilities of the Gridded Data Viewer have been expanded in version 2. The number of different years is a new variable in the monthly climatological gridded data set. The Gridded Data Viewer now shows the 400 km continental margin mask at 1 min resolution used for defining the Coastal Region (Table 5) and the distance to the nearest major land mass from 0 to 1000 km at 20 min resolution. The Gridded Data Viewer has an option for animation of gridded products. The interface has a new comparison capability for up to four gridded data sets. This enables the user to visualise, for example, gridded data products in SOCAT versions 1 and 2 in a multiple-plot view.

2.4.7 Key differences with version 1 in the data products

This section identifies key differences between the data products for versions 1 and 2.

Parameters in the individual and synthesis data set files: the data set quality control flags of A to D have been added as numerical values 11 to 14 to the synthesis files in version 2 (Tables 1 and 3). The distance to a major land mass is a new parameter in the files. Atmospheric CO₂ mole fractions from the 2012 GLOBALVIEW-CO₂ are reported in version 2 output files; this represents an update from the 2008 GLOBALVIEW-CO₂ values which were reported for




Figure 7. Seasonal distribution of surface water $f\text{CO}_2$ values for 2000 to 2009 in SOCAT version 2 for (a) January to March and (b) July to September.

version 1. The number of parameters in the downloadable files available via the Cruise Data Viewer as “All Variables” and “Current Variable” has been strongly reduced to match those in the synthesis files at CDIAC (Tables 1 and 3).

Gridded Data Viewer: the number of different years has been added as a variable to the monthly climatological gridded data set (Sect. 2.4.6). Data sets for the 400 km continental margin mask and the distance to the nearest major land mass are now available. The visualisation tools of the Gridded Data Viewer have been expanded.

Release notes: release notes document issues identified with individual data sets or data products in version 2. The notes are available on the CDIAC (<http://cdiac.ornl.gov/ftp/oceans/SOCATv2/>) and SOCAT (<http://www.socat.info/access.html>) websites (Table 1).

3 Spatial and temporal data coverage

SOCAT version 2 includes surface ocean $f\text{CO}_2$ values collected between 1968 and 2011 for the global oceans and coastal seas (Figs. 1 and 2). Data availability has increased over time for most ocean regions (Fig. 1b). A notable exception is the Indian Ocean, for which data are available from the 1990s, but where few subsequent observations have been made. Marked increases in data collection are apparent in the Gulf of Mexico (not shown) and the Arctic Ocean (Fig. 1b).




Figure 8. The number of (a) months of the year and (b) total months with surface water $f\text{CO}_2$ values in each 1° latitude by 1° longitude grid cell from 1970 to 2011 in SOCAT version 2. Figure 8a updates a similar figure for version 1 in Sabine et al. (2013, Fig. 5).

For example, version 2 has a total of 40 data sets in the Arctic Ocean, of which 10 data sets were collected in 2011 alone. Data coverage remains sparse in large parts of the Southern Hemisphere oceans (Fig. 2).

On average 3.4 surface water $f\text{CO}_2$ values have been collected per 100 km^2 between 1968 and 2011 in the global oceans and coastal seas. Data density ranges widely from 0.8 $f\text{CO}_2$ values per 100 km^2 in the Indian Ocean to 6.7 values per 100 km^2 in the North Atlantic Ocean (Fig. 6). Data density in the Southern Ocean appears somewhat high with 2.6 values per 100 km^2 relative to the North Pacific, the Tropical Pacific and the Tropical Atlantic Oceans. However, the Southern Ocean includes coastal waters with higher than average data density (Fig. 5), while the other three open ocean regions do not. Five of the ten most “productive” ships in terms of data collection are active south of 30° S , notably the *Nathaniel B. Palmer*, the *Laurence M. Gould*, the *Southern Surveyor*, the *Polarstern* and the *Aurora Australis* (Fig. 3).

The seasonal distribution of surface water $f\text{CO}_2$ values is shown in Fig. 7 for the period 2000 to 2009. The maps demonstrate the near absence of wintertime data in the high-latitude regions. The Ross Sea (Southern Ocean) has about 20 months of observations spanning five months from austral spring to autumn (Fig. 8).

The installation of automated $f\text{CO}_2$ systems on ships of opportunity and Antarctic supply ships has greatly improved the data availability along shipping routes and including for coastal seas near major ports (Fig. 9). For example, between 2000 and 2009 more than 40 individual ship visits have been made to the 1° latitude by 1° longitude grid boxes in the




Figure 9. Number of data sets (colour bar on top of subplots) with surface water $f\text{CO}_2$ measurements per 1° latitude by 1° longitude grid cell for 2000 to 2009 for (a) the northwest Atlantic Ocean and the Caribbean Sea, (b) the northeast Atlantic Ocean and European shelf seas, (c) the northwest Pacific Ocean and (d) Drake Passage in the Southern Ocean. The presence of repeated $f\text{CO}_2$ observations made on ships of opportunity and research supply ships is clearly visible in coastal seas and the open ocean.

Florida Straits, the English Channel, off the coast of Japan and near the Antarctic Peninsula.

The numbers of unique months and total months with $f\text{CO}_2$ values per 1° latitude by 1° longitude grid cell shed light on data collection activities for 1970 to 2011 (Fig. 8). High data density along shipping routes highlights the repeated $f\text{CO}_2$ observations. For example, numerous grid boxes east of Japan have observations in all months of the year for about 50 months in total, reflecting an intense CO_2 observational effort over a large number of years. This ongoing data collection effort is critical for the quantification of the variability and trends in CO_2 air–sea exchange.

4 Future plans

4.1 Progress towards version 3

Surface water CO_2 values and accompanying metadata can be submitted to CDIAC in the IOCCP-recommended formats (<http://cdiac.ornl.gov/oceans/submit.html>) at all times. Ideally data are submitted as they become available. The SOCAT global group sets deadlines for consideration of data in specific SOCAT versions; for example, the deadline for submission to SOCAT version 3 was 28 February 2014. Version 3 quality control is scheduled to take place during the

summer and autumn of 2014 with the release of version 3 planned for mid-2015.

4.2 Quality control flags for alternative sensors on a range of platforms

The SOCAT data quality control flags have been primarily designed for shipboard, continuous surface water CO₂ measurements by gas chromatography or infrared detection (Pfeil et al., 2013). Since the definition of these flags, high-precision and stable cavity ring-down spectroscopy (CRDS) has become available for surface water CO₂ measurements (Friedrichs et al., 2010; Becker et al., 2012). The quality control criteria in SOCAT are deemed adequate for the measurements by CRDS. Measurements made by CRDS will be included in future SOCAT versions, provided calibrations have been carried out at least daily (SOCAT, 2012b).

The quality control criteria, as used in versions 1 and 2, need revision for *f*CO₂ values from sensors on surface moorings, surface drifters and self-propelled surface vehicles (SOCAT, 2012b). These measurements do not follow all the standard operating procedures and at-sea calibration of such *f*CO₂ measurements is often infrequent or non-existent. Also, the sensors tend to use fewer gas standards than on ships due to logistical and power constraints. A working group on alternative sensors (Table 4) has developed a vision on how to include such *f*CO₂ values, as measured for example by infrared analysis and spectrophotometry, in future SOCAT versions (Wanninkhof et al., 2013a). The working group has determined which quality control criteria should apply to surface water CO₂ data from these new sensors and platforms. The term “data set quality control flag” replaces “cruise quality control flag”. The accuracy of data with data set flags of C and D has been specified as 5 µatm. A new data set quality control flag of “E” with an accuracy better than 10 µatm has been defined. The platform and the CO₂ instrument type will be introduced as parameters. These quality control criteria and other recommendations of the working group will be adopted for SOCAT version 3.

4.3 Automation

The large effort for data entry and quality control is a major obstacle for regular and prompt SOCAT updates, especially with more data becoming available each year. The need for automating SOCAT was formally recognised in September 2011 (SOCAT, 2011) and an automation team was created (Table 4). The automation vision proposed by the team was accepted by regional and global group leads (SOCAT, 2012a, b). The automation system will allow the data provider to upload, review and submit data and metadata. It will calculate surface water *f*CO₂. The automation system will provide a single portal for data providers, data managers and quality controllers. Manual data entry by the SOCAT data managers will be strongly reduced. Regular, prompt releases of SOCAT

will be more straightforward once the automation system is fully operational. The automation system is expected to become the primary mode of data submission from version 4 onwards.

5 Scientific applications of SOCAT

Several scientific studies have used SOCAT data products. The global synthesis product is the most popular SOCAT product in scientific publications. Both files in zip text format (Lourantou and Metzl, 2011; Tjiputra et al., 2012; Nakaoka et al., 2013; Rödenbeck et al., 2013; Wanninkhof et al., 2013b) and the Ocean Data View collection (Chierici et al., 2012) are used for data access. Two studies utilise a global gridded product (Landschützer et al., 2013; Schuster et al., 2013).

Scientific applications of SOCAT include:

- Visualisation of surface ocean *f*CO₂ data coverage (Chierici et al., 2012) and data requirements (Wanninkhof et al., 2013b);
- Use of the SOCAT continental margin mask (Evans and Mathis, 2013);
- Process studies (Lourantou and Metzl, 2011);
- Creation and validation of surface water *f*CO₂ and CO₂ air–sea flux maps by a variety of techniques, including multiple linear regression (Schuster et al., 2013), neural network approaches (Landschützer et al., 2013; Nakaoka et al., 2013) and an ocean mixed layer model (Rödenbeck et al., 2013);
- Quantification of the annual mean ocean carbon sink (Schuster et al., 2013);
- Studies of variation in the ocean carbon sink on seasonal (Rödenbeck et al., 2013), year-to-year (Landschützer et al., 2013) and decadal timescales (Lourantou and Metzl, 2011);
- Initialisation and validation fields for ocean carbon cycle models (Tjiputra et al., 2012).

These applications highlight the importance of SOCAT for regional and global air–sea CO₂ flux assessments, process studies and ocean carbon modelling.

6 Conclusions

SOCAT version 2 represents a 44 yr record of surface water *f*CO₂ values from 1968 to 2011 for the global oceans and coastal seas (Figs. 1 and 2). Version 2 extends version 1 by four years, while also adding more *f*CO₂ values for the years 2006 and 2007. The data are in a uniform format and have been subject to documented quality control. The quality

of data and of data reporting has improved in version 2 relative to version 1. The temporal data distribution partly reflects activities in large international research programmes. Over time, data coverage in all ocean regions has increased, with the exception of the Indian Ocean. Data coverage has increased four-fold from the 1990s to the 2000s, thus providing much better seasonal and spatial coverage for large parts of the Northern Hemisphere oceans and coastal seas. Data coverage remains sparse in large parts of the Southern Hemisphere and the Indian Ocean.

The international importance of SOCAT is evident from recent scientific articles using SOCAT data products for quantification of the ocean carbon sink, process studies and ocean carbon modelling. Regular updates to SOCAT will extend the SOCAT data record and ensure that new data are promptly made available for flux assessments and modelling. Future plans include use of the revised quality control criteria for $f\text{CO}_2$ values from alternative sensors and platforms, as well as automation.

Acknowledgements. The International Ocean Carbon Coordination Project (IOCCP), the Surface Ocean Lower Atmosphere Study (SOLAS), and the Integrated Marine Biogeochemistry and Ecosystem Research program (IMBER) promote SOCAT. We thank both referees, anonymous referee 1 and D. Woolf, for their constructive and detailed comments, which have considerably improved the Cruise Data Viewer and this manuscript. Support for SOCAT has been received from the University of East Anglia (UK), the Bjerknes Centre for Climate Research, the Geophysical Institute at the University of Bergen (Norway), the Climate Observation Division of the Climate Program Office of the US National Oceanic and Atmospheric Administration, the University of Washington, Oak Ridge National Laboratory (USA), PANGAEA® Data Publisher for Earth and Environmental Data (Germany), the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research (Germany), the National Institute for Environmental Studies (Japan), the Korean Institute of Ocean Science and Technology, the European Union projects CarboChange (FP7 264879) and GEO-CARBON (FP7 283080), the UK Ocean Acidification Research Programme (NE/H017046/1; funded by the Natural Environment Research Council, the Department for Energy and Climate Change and the Department for Environment, Food and Rural Affairs), the Research Council of Norway (CARBON-HEAT), the Scientific Committee on Oceanic Research (SCOR, US, OCE-0938349), the US National Science Foundation (OCE-1068958), Swedish Research Council projects (2009–2994; 2008–6228; 2009–2994) and the Swedish National Space Board (RESCUE project; REmote Sensing ocean Carbon UptakE).

Edited by: F. Schmitt

References

- Antonov, J. I., Locarnini, R. A., Boyer, T. P., Mishonov, A. V., and Garcia, H. E.: World Ocean Atlas 2005, in: Volume 2: Salinity, edited by: Levitus, S., NOAA Atlas NESDIS 62, US Government Printing Office, Washington, DC, 182 pp., 2006.
- AOML: Underway $p\text{CO}_2$ systems workshop, Atlantic Oceanographic and Meteorological Laboratory (AOML) National Oceanic and Atmospheric Administration, Miami, 2–3 October 2002, <http://www.aoml.noaa.gov/ocd/gcc/uwpc02/workshops/> (last access: 13 March 2012), 2002.
- Bakker, D. C. E., Etcheto, J., Boutin, J., Dandonneau, Y., and Merlivat, L.: Variability of surface water $f\text{CO}_2$ during seasonal upwelling in the equatorial Atlantic Ocean as observed by a drifting buoy, *J. Geophys. Res.*, 106, 9241–9253, doi:10.1029/1999JC000275, 2001.
- Bakker, D. C. E., Pfeil, B., Olsen, A., Sabine, C., Metzl, N., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., and Telszewski, M.: Global data products help assess changes to the ocean carbon sink, *Eos, Transactions American Geophysical Union*, 93, 125–126, doi:10.1029/2012EO120001, 2012.
- Becker, M., Andersen, N., Fiedler, B., Fietzek, P., Körtzinger, A., Steinhoff, T., and Friedrichs, G.: Using cavity ringdown spectroscopy for continuous monitoring of $\delta^{13}\text{C}(\text{CO}_2)$ and $f\text{CO}_2$ in the surface ocean, *Limnol. Oceanogr.-Meth.*, 10, 752–766, doi:10.4319/lom.2012.10.752, 2012.
- Boutin, J., Etcheto, J., Dandonneau, Y., Bakker, D. C. E., Feely, R. A., Inoue, H. Y., Ishii, M., Ling, R. D., Nightingale, P. D., Metzl, N., and Wanninkhof, R.: Satellite sea surface temperature: a powerful tool for interpreting in situ $p\text{CO}_2$ measurements in the equatorial Pacific Ocean, *Tellus*, 51B, 490–508, 1999.
- Boutin, J. and Merlivat, L.: New in situ estimates of carbon biological production rates in the Southern ocean from CARIOCA drifter measurements, *Geophys. Res. Lett.*, 36, L13608, doi:10.1029/2009GL038307, 2009.
- Chierici, M., Signorini, S. R., Mattsdotter-Björk, M., Fransson, A., and Olsen, A.: Surface water $f\text{CO}_2$ algorithms for the high-latitude Pacific sector of the Southern Ocean, *Remote Sens. Environ.*, 119, 184–196, doi:10.1016/j.rse.2011.12.020, 2012.
- Cooper, D. J., Watson, A. J., and Ling, R. D.: Variations of $p\text{CO}_2$ along a North Atlantic shipping route (UK to the Caribbean): A year of automated observations, *Mar. Chem.*, 60, 147–164, 1998.
- Corbière, A., Metzl, N., Reverdin, G., Brunet, C., and Takahashi, T.: Interannual and decadal variability of the carbon dioxide sink in the North Atlantic subpolar gyre, *Tellus*, 59B, 168–178, doi:10.1111/j.1600-0889.2006.00232.x, 2007.
- Dickson, A. G., Sabine, C. L., and Christian, J. R. (Eds.): Guide to best practices for ocean CO₂ measurements, PICES Special Publication 3, IOCCP Report 8, 191 pp., 2007.
- Emerson, S., Sabine, C., Cronin, M. F., Feely, R., Cullison Gray, S. E., and DeGrandpré, M.: Quantifying the flux of CaCO₃ and organic carbon from the surface ocean using in situ measurements of O₂, N₂, $p\text{CO}_2$, and pH, *Global Biogeochem. Cy.*, 25, GB3008, doi:10.1029/2010GB003924, 2011.
- ETOPO2: 2-minute Gridded Global Relief Data (ETOPO2v2), National Geophysical Data Center, National Oceanic and Atmospheric Administration, US Dept. of Commerce, available at: <http://www.ngdc.noaa.gov/mgg/global/etopo2.html> (last access: 1 May 2013), 2006.
- Evans, W. and Mathis, J. T.: The Gulf of Alaska coastal ocean as an atmospheric CO₂ sink, *Cont. Shelf Res.*, 65, 52–63, doi:10.1016/j.csr.2013.06.013, 2013.
- Feely, R. A., Boutin, J., Cosca, C. E., Dandonneau, Y., Etcheto, J., Inoue, H. Y., Ishii, M., Le Quééré, C., Mackey, D. J., McPhaden, M., Metzl, N., Poisson, A., and Wanninkhof, R.: Seasonal and

- interannual variability of CO₂ in the equatorial Pacific, *Deep-Sea Res. Pt. II*, 49, 2442–2469, 2002.
- Feely, R. A., Takahashi, T., Wanninkhof, R., McPhaden, M. J., Cosca, C. E., and Sutherland, S. C.: Decadal variability of the air-sea CO₂ fluxes in the Equatorial Pacific Ocean, *J. Geophys. Res.*, 111, CO8S90, doi:10.1029/2005JC003129, 2006.
- Friedrichs, G., Bock, J., Temps, F., Fietzek, P., Körtzinger, A., and Wallace, D. W. R.: Toward continuous monitoring of seawater¹³CO₂/¹²CO₂ isotope ratio and *p*CO₂: Performance of cavity ringdown spectroscopy and gas matrix effects, *Limnol. Oceanogr.-Meth.*, 8, 539–551, doi:10.4319/lom.2010.8.539, 2010.
- GLOBALVIEW-CO2: Cooperative Atmospheric Data Integration Project – Carbon Dioxide, 2008 version, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, Colorado, USA, available at: <http://www.esrl.noaa.gov/gmd/ccgg/globalview/> (last access: 26 June 2013), 2008.
- GLOBALVIEW-CO2: Cooperative Atmospheric Data Integration Project – Carbon Dioxide, 2012 version, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, Colorado, USA, available at: <http://www.esrl.noaa.gov/gmd/ccgg/globalview/> (last access: 26 June 2013), 2012.
- Hardman-Mountford, N. J., Moore, G. F., Bakker, D. C. E., Watson, A. J., Schuster, U., Barciela, R., Hines, A., Moincoiffé, G., Brown, J., Dye, S., Blackford, J., Somerfield, P. J., Holt, J., Hydes, D. J., and Aiken, J.: An operational monitoring system to provide indicators of CO₂-related variables in the ocean, *ICES J. Mar. Sci.*, 65, 1498–1503, doi:10.1093/icesjms/fsn110, 2008.
- Hood, E. M., Merlivat, L., and Johannessen, T.: Variations of *f*CO₂ and air-sea flux of CO₂ in the Greenland Sea gyre using high-frequency time series data from CARIOCA drifting buoys, *J. Geophys. Res.* 104, 20571–20583, doi:10.1029/1999JC900130, 1999.
- IOCCP: Surface Ocean CO₂ Variability and Vulnerabilities Workshop, UNESCO, Paris, France, 11–14 April 2007, IOCCP Report 7, available at: <http://www.ioccp.org/> (last access: 1 May 2013), 2007.
- IOCCP: Surface Ocean CO₂ Atlas Project, UNESCO, Paris, France, IOCCP Report 9, available at: <http://www.ioccp.org/> (last access: 1 May 2013), 2008.
- IOCCP: SOCAT Atlantic and Southern Oceans Regional Meeting, University of East Anglia, Norwich, UK, 25–26 June 2009, IOCCP Report 13, available at: <http://www.ioccp.org/> (last access: 1 May 2013), 2009.
- IOCCP: Southern and Indian Ocean SOCAT Workshop, CSIRO Marine Laboratories, Hobart, Tasmania, Australia, 16–18 June 2010, IOC Workshop Report 234, IOCCP Report 21, available at: <http://www.ioccp.org/> (last access: 1 May 2013), 2010.
- IOCCP: Coastal and Arctic Surface Ocean CO₂ Atlas (SOCAT) Quality Control Workshop, NOAA-PMEL, Seattle, USA, 2–4 October 2012, IOCCP Report 4/2012, available at: <http://www.ioccp.org/> (last access: 1 May 2013), 2012.
- Ishii, M., Saito, S., Tokieda, T., Kawano, T., Matsumoto, K., and Inoue, H. Y.: Variability of surface layer CO₂ parameters in the Western and Central Equatorial Pacific, in: *Global environmental change in the ocean and on land*, edited by: Shiyomi, M., Kawahata, H., Koizumi, H., Tsuda, A., and Awaya, Y., TERRAPUB, Tokyo, Japan, 2004.
- Ishii, M., Inoue, H. Y., Midorikawa, T., Saito, S., Tokieda, T., Sasano, D., Nakadate, A., Nemoto, K., Metzl, N., Wong, C. S., and Feely, R. A.: Spatial variability and decadal trend of the oceanic CO₂ in the western equatorial Pacific warm/fresh water, *Deep-Sea Res. Pt. II*, 56, 591–606, 2009.
- Jansen, E., Overpeck, J., Briffa, K. R., Duplessy, J.-C., Joos, F., Masson-Delmotte, V., Olago, D., Otto-Bliesner, B., Peltier, W. R., Rahmstorf, S., Ramesh, R., Raynaud, D., Rind, D., Solomina, O., Villalba, R., and Zhang, D.: Palaeoclimate, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- Johengen, T.: Performance demonstration statement PMEL MAPCO2/Battelle Seaology *p*CO₂ monitoring system, Alliance for Coastal Technologies report, ACT DS10-02, available at: http://www.act-us.info/Download/Evaluations/pCO2/PMEL_MAPCO2_Battelle_Seaology/ (last access: 10 January 2014), 2010.
- Jones, C., Robertson, E., Arora, V., Friedlingstein, P., Shevliakova, E., Bopp, L., Brovkin, V., Hajima, T., Kato, E., Kawamiya, M., Liddicoat, S., Lindsay, K., Reick, C. H., Roelandt, C., Segsneider, J., and Tjiputra, J.: Twenty-first century compatible CO₂ emissions and airborne fraction simulated by CMIP5 Earth System models under 4 Representative Concentration Pathways, *J. Climate*, 26, 4398–4413, doi:10.1175/JCLI-D-12-00554.1, 2013.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–470, 1996.
- Kitidis, V., Hardman-Mountford, N. J., Litt, E., Brown, I., Cummings, D., Hartman, S., Hydes, D., Fishwick, J. R., Harris, C., Martinez-Vicente, V., Woodward, E. M. S., and Smyth, T. J.: Seasonal dynamics of the carbonate system in the Western English Channel, *Cont. Shelf Res.*, 42, 30–40, 2012.
- Körtzinger, A., Thomas, H., Schneider, B., Gronau, N., Mintrop, L., and Duinker, J. C.: At-sea intercomparison of two newly designed underway *p*CO₂ systems – Encouraging results, *Mar. Chem.*, 52, 133–145, 1996.
- Körtzinger, A., Mintrop, L., Wallace, D. W. R., Johnson, K. M., Neill, C., Tilbrook, B., Towler, P., Inoue, H., Ishii, M., Shaffer, G., Torres, R. F., Ohtaki, E., Yamashita, E., Poisson, A., Brunet, C., Schauer, B., Goyet, C., and Eiseid, G.: The international at-sea intercomparison of *f*CO₂ systems during the R/V *Meteor* cruise 36/1 in the North Atlantic Ocean, *Mar. Chem.*, 72, 171–192, 2000.
- Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U., Nakaoka, S., Payne, M. R., Sasse, T. P., and Zeng, J.: A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink, *Biogeosciences*, 10, 7793–7815, doi:10.5194/bg-10-7793-2013, 2013.
- Lefèvre, N., Watson, A. J., and Watson, A. R.: A comparison of multiple regression and neural network techniques for mapping in situ *p*CO₂ data, *Tellus*, 57B, 375–384, 2005.

- Lefèvre, N., Guillot, A., Beaumont, L., and Danguy, T.: Variability of $f\text{CO}_2$ in the Eastern Tropical Atlantic from a moored buoy, *J. Geophys. Res.*, 113, C01015, doi:10.1029/2007JC004146, 2008.
- Lefèvre, N., Caniaux, G., Janicot, S., and Gueye, A. K.: Increased CO₂ outgassing in February–May 2010 in the tropical Atlantic following the 2009 Pacific El Niño, *J. Geophys. Res.–Oceans*, 118, 1645–1657, doi:10.1002/jgrc.20107, 2013.
- Lenton, A., Matear, R. J., and Tilbrook, B.: Design of an observational strategy for quantifying the Southern Ocean uptake of CO₂, *Global Biogeochem. Cy.*, 20, GB4010, doi:10.1029/2005GB002620, 2006.
- Lenton, A., Metzl, N., Takahashi, T., Kuchinke, M., Matear, R. J., Roy, T., Sutherland, S. C., Sweeney, C., and Tilbrook, B.: The observed evolution of oceanic $p\text{CO}_2$ and its drivers over the last two decades, *Global Biogeochem. Cy.*, 26, GB2021, doi:10.1029/2011GB004095, 2012.
- Le Quéré, C., Rödenbeck, C., Buitenhuis, E. T., Conway, T. J., Lagenfelds, R., Gomez, A., Labuschagne, C., Ramonet, M., Nakazawa, T., Metzl, N., Gillett, N., and Heimann, M.: Saturation of the Southern Ocean CO₂ sink due to recent climate change, *Science*, 316, 1735–1738, 2007.
- Le Quéré, C., Takahashi, T., Buitenhuis, E., Rödenbeck, C., and Sutherland, S. C.: Impact of climate change and variability on the global oceanic sink of CO₂, *Global Biogeochem. Cy.*, 24, GB4007, doi:10.1029/2009GB003599, 2010.
- Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters, G. P., van der Werf, G. R., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C., Enright, C., Friedlingstein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein Goldewijk, K., Levis, S., Levy, P., Lomas, M., Poulter, B., Raupach, M. R., Schwinger, J., Sitch, S., Stocker, B. D., Viovy, N., Zaehle, S., and Zeng, N.: The global carbon budget 1959–2011, *Earth Syst. Sci. Data*, 5, 165–185, doi:10.5194/essd-5-165-2013, 2013.
- Levitus, S., Antonov, J., and Boyer, T.: Warming of the world ocean, 1955–2003, *Geophys. Res. Lett.*, 32, L02604, doi:10.1029/2004GL021592, 2005.
- Lourantou, A. and Metzl, N.: Decadal evolution of carbon sink within a strong bloom area in the subantarctic zone, *Geophys. Res. Lett.*, 38, L23608, doi:10.1029/2011GL049614, 2011.
- Lüger, H., Wallace, D. W. R., Körtzinger, A., and Nojiri, Y.: The $p\text{CO}_2$ variability in the midlatitude North Atlantic Ocean during a full annual cycle, *Global Biogeochem. Cy.*, 18, GB3023, doi:10.1029/2003GB002200, 2004.
- McKinley, G. A., Fay, A. R., Takahashi, T., and Metzl, N.: Convergence of atmospheric and North Atlantic carbon dioxide trends on multidecadal timescales, *Nat. Geosci.*, 4, 606–610, doi:10.1038/NGEO1193, 2011.
- Metzl, N.: Decadal increase of oceanic carbon dioxide in Southern Indian Ocean surface waters (1991–2007), *Deep-Sea Res. Pt. II*, 56, 607–619, 2009.
- Mulvaney, R., Abram, N. J., Hindmarsh, R. C. A., Arrowsmith, C., Fleet, L., Triest, J., Sime, L. C., Alemany, O., and Foord, S.: Recent Antarctic Peninsula warming relative to Holocene climate and ice-shelf history, *Nature*, 489, 141–145, doi:10.1038/nature11391, 2012.
- Nakaoka, S., Telszewski, M., Nojiri, Y., Yasunaka, S., Miyazaki, C., Mukai, H., and Usui, N.: Estimating temporal and spatial variation of ocean surface $p\text{CO}_2$ in the North Pacific using a self-organizing map neural network technique, *Biogeosciences*, 10, 6093–6106, doi:10.5194/bg-10-6093-2013, 2013.
- Park, G.-H., Lee, K., Wanninkhof, R., and Feely, R. A.: Empirical temperature-based estimates of variability in the oceanic uptake of CO₂ over the past 2 decades, *J. Geophys. Res.*, 111, C07S07, doi:10.1029/2005JC003090, 2006.
- Park, G.-H. and Wanninkhof, R.: A large increase of the CO₂ sink in the western tropical North Atlantic from 2002 to 2009, *J. Geophys. Res.*, 117, C08029, doi:10.1029/2011JC007803, 2012.
- Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré, C., Marland, G., Raupach, M. R., and Wilson, C.: The challenge to keep global warming below 2 °C, *Nature Clim. Change*, 3, 4–6, doi:10.1038/nclimate1783, 2013.
- Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N., Sabine, C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales, B., Hardman-Mountford, N., Heinze, C., Hood, M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Jones, S. D., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A uniform, quality controlled Surface Ocean CO₂ Atlas (SOCAT), *Earth Syst. Sci. Data*, 5, 125–143, doi:10.5194/essd-5-125-2013, 2013.
- Pierrot, D., Neill, C., Sullivan, K., Castle, R., Wanninkhof, R., Lüger, H., Johannessen, T., Olsen, A., Feely, R. A., and Cosca, C. E.: Recommendations for autonomous underway $p\text{CO}_2$ measuring systems and data reduction routines, *Deep-Sea Res. Pt. II*, 56, 512–522, doi:10.1016/j.dsr2.2008.12.005, 2009.
- Rödenbeck, C., Keeling, R. F., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global surface-ocean $p\text{CO}_2$ and sea-air CO₂ flux variability from an observation-driven ocean mixed-layer scheme, *Ocean Sci.*, 9, 193–216, doi:10.5194/os-9-193-2013, 2013.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T. H., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.
- Sabine, C. L., Ducklow, H., and Hood, M.: International carbon coordination. Roger Revelle's legacy in the Intergovernmental Oceanographic Commission, *Oceanography*, 23, 48–61, 2010.
- Sabine, C. L., Hankin, S., Koyuk, H., Bakker, D. C. E., Pfeil, B., Olsen, A., Metzl, N., Kozyr, A., Fassbender, A., Manke, A., Malczyk, J., Akl, J., Alin, S. R., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Feely, R. A., González-Dávila, M., Goyet, C., Hardman-Mountford, N., Heinze, C., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton,

- A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Pater-son, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Sal-isbury, J., Santana-Casiano, J. M., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Stein-hoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Vandemark, D., Veness, T., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: Surface Ocean CO₂ Atlas (SOCAT) gridded data products, *Earth Syst. Sci. Data*, 5, 145–153, doi:10.5194/essd-5-145-2013, 2013.
- Sarma, V. V. S. S., Saino, T., Sasaoka, K., Nojiri, Y., Ono, T., Ishii, M., Inoue, H. Y., and Matsumoto, K.: Basin-scale *p*CO₂ distribu-tion using satellite sea surface temperature, Chl *a*, and climato-logical salinity in the North Pacific in spring and summer, *Global Biogeochem. Cy.*, 20, GB3005, doi:10.1029/2005GB002594, 2006.
- Schuster, U. and Watson, A. J.: A variable and decreasing sink for atmospheric CO₂ in the North Atlantic, *J. Geophys. Res.*, 112, C11006, doi:10.1029/2006JC003941, 2007.
- Schuster, U., Watson, A. J., Bates, N. R., Corbière, A., González-Dávila, M., Metzl, N., Pierrot, D., and Santana-Casiano, M.: Trends in North Atlantic sea-surface *f*CO₂ from 1990 to 2006, *Deep-Sea Res. Pt. II*, 56, 620–629, 2009.
- Schuster, U., McKinley, G. A., Bates, N., Chevallier, F., Doney, S. C., Fay, A. R., González-Dávila, M., Gruber, N., Jones, S., Krijnen, J., Landschützer, P., Lefèvre, N., Manizza, M., Mathis, J., Metzl, N., Olsen, A., Rios, A. F., Rödenbeck, C., Santana-Casiano, J. M., Takahashi, T., Wanninkhof, R., and Watson, A. J.: An assessment of the Atlantic and Arctic sea-air CO₂ fluxes, 1990–2009, *Biogeosciences*, 10, 607–627, doi:10.5194/bg-10-607-2013, 2013.
- SOCAT: Future for the Surface Ocean CO₂ Atlas: Data quality, management and products, Data2Flux Workshop, Unesco, Paris, 12–13 September 2011, SOCAT Report, available at: <http://www.socat.info/meetings.html> (last access: 24 June 2013), 2011.
- SOCAT: Surface Ocean CO₂ Atlas (SOCAT) automation planning meeting, NOAA-PMEL, Seattle, USA, 10–11 May 2012, SO-CAT Report, available at: <http://www.socat.info/meetings.html> (last access: 1 May 2013), 2012a.
- SOCAT: Surface Ocean CO₂ Atlas (SOCAT) progress meeting, Epochal Tsukuba, Tsukuba, Japan, 3–5 July 2012, SOCAT Re- port, available at: <http://www.socat.info/meetings.html> (last ac- cess: 1 May 2013), 2012b.
- SOCAT: Data policy for the SOCAT public release, available at: http://www.socat.info/SOCAT_data_policy_public_release_v2.htm (last access: 1 May 2013), 2013a.
- SOCAT: Surface Ocean CO₂ Atlas (SOCAT) side event: Release of version 2 and science highlights, SOCAT side event during the 9th International Carbon Dioxide Conference, Beijing Inter- national Convention Center, Beijing, China, 4 June 2013, SO- CAT Report, available at: <http://www.socat.info/meetings.html> (last access: 29 November 2013), 2013b.
- Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L. (Eds.): *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- Stroeve, J., Holland, M. M., Meier, W., Scambos, T., and Serreze, M.: Arctic sea ice decline: Faster than forecast, *Geophys. Res. Lett.*, 34, L09501, doi:10.1029/2007GL029703, 2007.
- Sweeney, C., Takahashi, T., and Wanninkhof, R.: Spatial and tem- poral variability of surface water *p*CO₂ and sampling strategies, Report presented at the Advisory Meeting for the Sea-Air CO₂ Flux Program, National Oceanic and Atmospheric Administra- tion, Boulder, Colorado, USA, 2000.
- Swift, J.: A guide to submitting CTD/hydrographic/tracer data and associated documentation to the CLIVAR and Carbon Hydro- graphic Data Office, version of 22 April 2008, Scripps Institu- tion of Oceanography, University of California, San Diego, 37 pp., 2008.
- Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S. C.: Seasonal variation of CO₂ and nutrients in the high-latitude surface oceans: a comparative study, *Global Bio- geochem. Cy.*, 7, 843–878, 1993.
- Takahashi, T., Feely, R. A., Weiss, R. F., Wanninkhof, R. H., Chip- man, D. W., Sutherland, S. C., and Takahashi, T. T.: Global air- sea flux of CO₂: An estimate based on measurements of sea-air *p*CO₂ difference, *P. Natl. Acad. Sci. USA*, 94, 8292–8299, 1997.
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C. L., Watson, A. J., Bakker, D. C. E., Schuster, U., Metzl, N., Inoue, H. Y., Ishii, M., Midorikawa, T., No- jiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, J. M. J., Olaf- sson J., Arnarson, T. S., Tilbrook, B., Johannessen T., Olsen, A., Bellerby, R. G. J., Wong, C. S., Delille B., Bates, N. R., and De Baar, H. J. W.: Climatological mean and decadal change in surface ocean *p*CO₂, and net sea-air CO₂ flux over the global oceans, *Deep-Sea Res. Pt. II*, 56, 544–577, doi:10.1016/j.dsr2.2008.12.009, 2009.
- Takahashi, T., Sutherland, S. C., and Kozyr, A.: Global ocean surface water partial pressure of CO₂ database: Measurements performed during 1957–2012 (Version 2012), ORNL/CDIAC-160, NDP-088(V2012), Carbon Dioxide Infor- mation Analysis Center, Oak Ridge National Labora- tory, US Department of Energy, Oak Ridge, Tennessee, doi:10.3334/CDIAC/OTG.NDP088(V2012), available at: http://cdiac.ornl.gov/oceans/LDEO_Underway_Database/ (last access: 30 May 2013), 2013.
- Takamura, T. R., Inoue, H. Y., Midorikawa, T., Ishii, M., and Nojiri, Y.: Seasonal and inter-annual variations in *p*CO₂^{sea} and air-sea CO₂ fluxes in mid-latitudes of the western and eastern North Pacific during 1999–2006: Recent results utilizing vol- untary observation ships, *J. Meteorol. Soc. Jpn.*, 88, 883–898, doi:10.2151/jmsj.2010-602, 2010.
- Tans, P. and Keeling, R.: Trends in atmospheric carbon dioxide, available at: http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html#global_data (last access: 7 January 2014), 2014.
- Thomas, H., Prowe, A. E. F., Heuven, S. van, Bozec, Y., Baar, H. J. W. de, Schiettecatte, L. S., Suykens, K., Koné, M., Borges, A. V., Lima, I. D., and Doney, S. C.: Rapid decline of the CO₂ buffering capacity in the North Sea and implications for the North Atlantic Ocean, *Global Biogeochem. Cy.*, 21, GB4001, doi:10.1029/2006GB002825, 2007.
- Thomas, H., Prowe, A. E. F., Lima, I., Doney, S. C., Wanninkhof, R., Greatbatch, R. J., Schuster, U., and Corbière, A.: Changes in the North Atlantic Oscillation influence CO₂ uptake in the North

- Atlantic over the past 2 decades, *Global Biogeochem. Cy.*, 22, GB4027, doi:10.1029/2007GB003167, 2008.
- Tjiputra, J. F., Olsen, A., Assmann, K., Pfeil, B., and Heinze, C.: A model study of the seasonal and long-term North Atlantic surface $p\text{CO}_2$ variability, *Biogeosciences*, 9, 907–923, doi:10.5194/bg-9-907-2012, 2012.
- Vaughan, D. G., Marshall, G. J., Connolley, W. M., Parkinson, C., Mulvaney, R., Hodgson, D. A., King, J. C., Pudsey, C. J., and Turner, J.: Recent rapid regional climate warming on the Antarctic Peninsula, *Climatic Change*, 60, 243–274, Kluwer Academic Publishers, the Netherlands, 2003.
- Wanninkhof, R. H. and Thoning, K.: Measurement of fugacity of CO₂ in surface water using continuous and discrete sampling methods, *Mar. Chem.*, 44, 183–204, 1993.
- Wanninkhof, R., Bakker, D. C. E., Bates, N., Olsen, A., Steinhoff, T., and Sutton, A. J.: Incorporation of alternative sensors in the SOCAT database and adjustments to dataset quality control flags, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, doi:10.3334/CDIAC/OTG.SOCAT_ADQCF, 2013a.
- Wanninkhof, R., Park, G.-H., Takahashi, T., Sweeney, C., Feely, R., Nojiri, Y., Gruber, N., Doney, S. C., McKinley, G. A., Lenton, A., Le Quéré, C., Heinze, C., Schwinger, J., Graven, H., and Khatiwala, S.: Global ocean carbon uptake: magnitude, variability and trends, *Biogeosciences*, 10, 1983–2000, doi:10.5194/bg-10-1983-2013, 2013b.
- Watson, A. J., Schuster, U., Bakker, D. C. E., Bates, N. R., Corbière, A., González-Dávila, M., Friedrich, T., Heinze, C., Johannessen, T., Körtzinger, A., Metzl, N., Olafsson, J., Olsen, A., Oschlies, A., Padin, X. A., Pfeil, B., Santana-Casiano, J. M., Steinhoff, T., Telszewski, M., Ríos, A. F., Wallace, D. W. R., and Wanninkhof, R.: A network to accurately estimate the North Atlantic sink for CO₂, *Science*, 326, 1391–1393, 2009.