

CATACLYSMIC VARIABLES FROM THE SLOAN DIGITAL SKY SURVEY. I. THE FIRST RESULTS¹

PAULA SZKODY,² SCOTT F. ANDERSON,² MARCEL AGÜEROS,² RICARDO COVARRUBIAS,² MISTY BENTZ,² SUZANNE HAWLEY,²
 BRUCE MARGON,^{2,3} WOLFGANG VOGES,⁴ ARNE HENDEN,⁵ GILLIAN R. KNAPP,⁶ DANIEL E. VANDEN BERK,⁷ ARMIN REST,²
 GAJUS MIKNAITIS,² EUGENE MAGNIER,² J. BRINKMANN,⁸ I. CSABAI,⁹ M. HARVANEK,⁸ R. HINDSLEY,¹⁰ G. HENNESSY,¹⁰
 Z. IVEZIC,⁶ S. J. KLEINMAN,⁸ D. Q. LAMB,¹¹ D. LONG,⁸ P. R. NEWMAN,⁸ E. H. NEILSEN,⁷ R. C. NICHOL,¹²
 A. NITTA,⁸ D. P. SCHNEIDER,¹³ S. A. SNEDDEN,⁸ AND D. G. YORK¹¹

Received 2001 July 3; accepted 2001 October 1

ABSTRACT

The commissioning year of the Sloan Digital Sky Survey (SDSS) has demonstrated that many cataclysmic variables (CVs) have been missed in previous surveys with brighter limits. We report the identification of 22 CVs, of which 19 are new discoveries and three are known systems (SW UMa, BH Lyn, and OU Vir). A compendium of positions, colors, and characteristics of these systems obtained from the SDSS photometry and spectroscopy is presented, along with data obtained during follow-up studies with the Apache Point Observatory and Manastash Ridge Observatory telescopes. We have determined orbital periods for three of the new systems, two show dwarf nova outbursts, and the third is a likely magnetic system with eclipses of its region of line emission. Based on these results, we expect the completed survey to locate at least 400 new CVs. Most of these will be faint systems with low accretion rates that will provide new constraints on binary evolution models.

Key words: novae, cataclysmic variables — techniques: photometric — techniques: spectroscopic

On-line material: color figure

1. INTRODUCTION

Cataclysmic variables (CVs) comprise all the close binaries that contain a white dwarf accreting material transferred from a companion, usually a late main-sequence star (see review book by Warner 1995). CVs include novae, dwarf novae, and nova-likes, which are distinguished by their amplitudes and timescale for variability. A nova outburst due to a thermonuclear runaway event creates a rise in brightness of 10–20 mag, a dwarf nova undergoes 2–7 mag outbursts from a normal quiescent state on a quasi-periodic timescale ranging from weeks to years (usually attributed to a disk instability), and a nova-like exhibits random variability of several magnitudes likely related to changes in mass transfer causing states of low or high accretion. A large

range in orbital period (from just under 80 minutes to over 2 days, with most being under 2 hr) and in magnetic field strength of the white dwarf (from less than 1 to 240 MG) results in a wide variety of characteristic behavior and various observational selection effects. While most of the nearly 1000 known or suspected CVs (Downes et al. 2001, hereafter D01) are faint at quiescence and have late-type M dwarf secondaries, most of the early discoveries from past surveys were blue, bright objects with dominant accretion disks or outbursts that brought them to high luminosities, thus making them accessible to detection.

Evolutionary models for close binaries (Howell, Rappaport, & Politano 1997) predict CVs should reach a period minimum near 80 minutes in the lifetime of the Galaxy and have very cool, optically unobservable secondaries with very low mass transfer rates. Surveys that have the capability to reach faint magnitudes (greater than 16) should be able to find this population of faint, old systems. By determining the numbers and characteristics of these faint systems, we can constrain evolutionary theories and determine the true distribution of CV types in the Galaxy. We expect these results to be quite different from the bright-end surveys comprised of accretion disk-dominated systems, e.g., the Palomar-Green survey (Green et al. 1982).

The Sloan Digital Sky Survey (SDSS; York et al. 2000) is ideal for finding these faint blue and red objects. The photometry of 10^4 deg² of sky in five filters to greater than 20th magnitude allows a selection of all types of CV systems (blue ones dominated by disks or white dwarfs as well as those that are both blue and red if the accretion disk is negligible and the primary white dwarf and M dwarf secondary are the main sources of light). In addition to the photometry, SDSS spectroscopy allows unambiguous identification of a CV from the strong hydrogen Balmer and helium emission lines that typically signify ongoing accretion. The strength and width of these lines provide clues to the inclination, orbital period and mass accretion rate (Patterson 1984;

¹ Based on observations obtained with the Sloan Digital Sky Survey and with the Apache Point Observatory 3.5 m telescope, which are owned and operated by the Astrophysical Research Consortium.

² Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195.

³ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218.

⁴ Max-Planck-Institute für extraterrestrische Physik, Geissenbachstrasse 1, D-85741 Garching bei München, Germany.

⁵ Universities Space Research Association/US Naval Observatory, Flagstaff Station, P.O. Box 1149, Flagstaff, AZ 86002-1149.

⁶ Princeton University Observatory, Peyton Hall, Princeton, NJ 08544.

⁷ Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, IL 60510.

⁸ Apache Point Observatory, P.O. Box 59, Sunspot, NM 88349-0059.

⁹ Department of Physics and Astronomy, John Hopkins University, 3400 North Charles Street, Baltimore, MD 21218.

¹⁰ US Naval Observatory, 3450 Massachusetts Avenue, NW, Washington, DC 20392-5420.

¹¹ Department of Astronomy and Astrophysics and Enrico Fermi Institute, 5640 South Ellis Avenue, Chicago, IL 60637.

¹² Department of Physics, Carnegie Mellon University, 5000 Forbes Avenue, Pittsburgh, PA 15232.

¹³ Department of Astronomy and Astrophysics, 525 Davey Laboratory, Pennsylvania State University, University Park, PA 16802.

Warner 1995). We report here the results we have obtained for objects we could clearly identify as CVs during the SDSS commissioning year (data obtained prior to 2001 January 1) and the implications for the total number of CVs that will be found by the completion of the SDSS project.

2. FINDING CV CANDIDATES

The SDSS imaging data are obtained on a dedicated 2.5 m telescope at Apache Point Observatory with a mosaic CCD camera (Gunn et al. 1998), which observes in five bands: u , g , r , i , and z (Fukugita et al. 1996; see Lupton, Gunn, & Szalay 1999 for a discussion of the modified magnitude scale). Until the final SDSS photometric system is defined, the preliminary magnitudes are denoted by u^* , g^* , r^* , i^* , and z^* , with the errors in the calibration estimated to be about 5% at 20th magnitude. The data are reduced by a photometric pipeline (Lupton et al. 2001), which separates stars from galaxies, using their surface brightness distribution and computes several types of magnitude, including the point source magnitudes that we use in this paper. The astrometric accuracy is $0''.10$. Spectra of objects selected from the photometric data are observed for typically 45 minutes with two fiber-fed CCD spectrographs (on the 2.5 m telescope) covering the range of 3900–6200 Å (blue beam) and 5800–9200 Å (red beam) at a resolving power of ~ 1800 (York et al. 2000). The spectra are wavelength- and flux-calibrated, and atmospheric absorption bands are corrected using sdF stars. The resulting spectra are then classified as stars, galaxies, or quasars, and redshifts are determined. Of the 640 fibers used in each 3° field, 32 are used for sky measurements, ~ 10 for standards, and the rest are primarily for galaxies and quasars. Stellar projects typically target about 10 fibers per field, with CVs having approximately one of these fibers.

During the commissioning year, a number of photometric selection criteria were attempted in order to maximize the chance of finding new CVs. We first tried the range of colors found by Krisciunas, Margon, & Szkody (1998), who used filters close to those of SDSS to observe known CVs of various types. The CVs easily separate from main-sequence stars in $u^* - g^*$ but overlap with quasars (Fan 1999) and white dwarfs. In addition, the large color range possible for redder colors does not allow a unique discriminator. Thus, selecting by the blue colors resulted in spectra that turned out to be primarily quasars and single white dwarfs. When we tried simultaneous blue and red criteria ($u^* - g^* < 0.45$, $g^* - r^* < 0.7$, $r^* - i^* > 0.30$, and $i^* - z^* > 0.4$) to select for CVs without much disk contribution, we were successful in obtaining mostly noninteracting WD plus M binaries with a few CVs. These criteria remain for the project color selection; in addition to the CVs found, the WD plus M binaries are being used in follow-up studies to search for systems that are close to starting mass transfer (i.e., pre-CVs) and to compare magnetic activity levels for single M stars with those in close binaries (Raymond, Szkody, & Hawley 2002).

The best success at finding the blue CVs came from using the fact that QSOs are primary targets for spectroscopic fibers (hence more are observed than stars) and some quasars overlap the colors of CVs. Thus, we could search the spectra obtained of quasar candidates for those that turned out to be nonquasars, i.e., those having zero redshift and broad hydrogen Balmer and helium emission lines. Several

fibers for extreme blue objects obtained under a serendipity classification also yielded blue CVs.

Because the commissioning year involved changing color selections for all targets, this is not a uniform sample. Nor will the final sample of CVs be complete, since there are not sufficient fibers for stars to target all possible CVs. However, since the QSO, serendipity, and CV selection criteria will not undergo further large changes, the results from this past year cover a representative sample of what will be found and represent a lower limit to the actual number of CVs that can be located by combining the SDSS database with follow-up spectroscopic and variability studies. Our results reported here include spectra obtained from fields that cover an effective area of about 600 deg^2 . The spectroscopic identification of 22 CVs in this area means that we found about 0.04 deg^{-2} . Thus, we can expect about 400 new CVs in the entire survey from the spectra alone. Follow-up time-resolved spectra can be used to determine the nature of these objects, as well as to identify pre-CVs from the WD plus M binary sample. Wide-field, time-resolved photometry on selected fields will be necessary to identify a complete sample of CVs using the SDSS colors combined with variability information.

3. RESULTS

The 22 objects that we identify as CVs are listed in Table 1, along with the date of the SDSS spectrum and the g^* magnitude and colors (with no correction for reddening). The naming convention is SDSSp Jhhmmss.ss+ddmms.s, where the coordinate equinox is J2000.0 and the “p” refers to the preliminary astrometry. Throughout the rest of this paper, we will abbreviate the names to SDSS hhmm for convenience. Objects that have data publicly available through the SDSS early data release (Stoughton et al. 2002) are marked. Finding charts from the Palomar Digitized Sky Survey (DSS) will be available from the on-line CV catalog (D01). The last column of Table 1 gives brief comments, which are elaborated on in the sections below.

Figure 1 shows the colors of the 22 objects in the SDSS color-color diagrams along with the stellar locus. As noted above, the contributions of the accretion disk and white dwarf generally result in a separation of the CVs from the main sequence, especially in the bluest colors. In this sample, about 90% of the objects were found by QSO and serendipity selection criteria. Two CVs (SDSS 0155 and 0813) satisfied the selection criteria of blue and red colors given in § 2.

Figure 2 shows the SDSS spectra for all systems except for the previously known ones, BH Lyn and SW UMa, which have published spectra available (Hoard & Szkody 1997; Shafter, Szkody, & Thorstensen 1986), and SDSS 0155, which was saturated in the fixed exposure time for its field. Table 2 lists the equivalent widths and fluxes of the prominent hydrogen Balmer and helium lines. In this table, the object is identified both by its abbreviated SDSS name and the plate and fiber numbers, where the plate number refers to the catalog number of the SDSS plug plate used to observe each field. Of the 20 objects in Table 2, 13 (65%) have $H\beta$ equivalent widths greater than 20 Å, and 10 (50%) have values greater than 50 Å. According to the approximate relation between $H\beta$ EW and \dot{M} given in Patterson (1984), these numbers indicate that we are finding the low accretion rate systems (\dot{M} of 10^{-10} to $10^{-12} M_\odot \text{ yr}^{-1}$), i.e.,

TABLE 1
SUMMARY OF CVs WITH SDSS SPECTRA^a

SDSSp	Date ^b	g^*	$u^* - g^*$	$g^* - r^*$	$r^* - i^*$	$i^* - z^*$	Comments ^c
J015151.87 + 140047.2	2000 Nov 29	20.26	-0.29	0.24	0.42	0.46	High i
J015543.40 + 002807.2*	2000 Nov 23	15.39	0.67	0.05	-0.23	-0.17	H, L, He II, e
J023322.61 + 005059.5*	2000 Oct 3	19.94	-0.30	0.26	-0.25	0.01	
J072910.68 + 365838.3	2000 Nov 29	20.38	-0.07	0.34	0.26	0.05	He II
J074716.81 + 424849.0	2000 Dec 7	17.10	-0.02	-0.09	-0.09	-0.06	NL, He II
J081321.91 + 452809.4	2000 Nov 29	18.29	-0.05	0.62	0.46	0.22	Long P , K-M sec.
J081610.84 + 453010.2	2000 Nov 29	20.08	-0.41	0.54	0.47	0.50	M sec.
J082236.05 + 510524.6	2000 Dec 4	15.34	0.02	0.15	-0.02	-0.01	BH Lyn
J082409.73 + 493124.4	2000 Nov 21	19.28	-0.39	0.44	0.16	0.16	
J083642.80 + 532838.1	2000 Dec 5	16.87	0.03	-0.06	-0.07	0.14	SW UMa
J083845.23 + 491055.5	2000 Nov 25	19.59	-0.40	0.20	-0.02	0.30	
J125641.29 - 015852.0	2000 May 30	20.12	0.25	-0.06	-0.28	-0.15	
J143500.22 - 004606.2*	2000 Apr 3 ^d	18.59	-0.14	0.17	-0.02	0.10	OU Vir
J155531.99 - 001055.0	2000 May 28	19.36	-0.26	0.23	0.06	0.31	High i
J155644.24 - 000950.2	2000 May 29	18.05	0.24	0.12	0.24	0.44	
J161033.64 - 010223.3	2000 May 26	19.07	0.19	0.02	-0.13	0.13	
J161332.56 - 000331.0	2000 May 29	18.63	-0.01	0.66	0.52	0.31	
J163722.21 - 001957.1 ^e	2000 May 7	16.56	0.15	-0.05	-0.14	-0.08	H, L
J171247.71 + 604603.3* ^f	2000 May 31 ^g	19.95	0.92	0.91	0.59	0.45	
J172601.96 + 543230.7*	2000 Sep 26	20.52	-0.26	-0.04	-0.02	0.02	
J173008.38 + 624754.7*	2000 May 30	15.92	0.10	-0.25	-0.11	0.00	DN
J230351.64 + 010651.0	2000 Sep 5	19.05	-0.38	0.30	0.06	0.26	DN

^a Objects marked with asterisk are publicly available in the SDSS early data release.

^b UT date of spectrum.

^c “DN” is a dwarf nova, “e” is eclipsing, “NL” is a nova-like, “H, L” means shows high and low brightness states.

^d This object also had spectra obtained on 2000 May 26.

^e The correct object is the northern one of a close pair.

^f The correct object is the southern one of a close pair.

^g This object also had spectra obtained on 2000 August 24.

the objects are faint not because they are observed at a large distance, but because they are intrinsically faint due to low mass transfer rates.

For several systems, we conducted follow-up photometric observations using the University of Washington (UW) 0.76 m telescope in eastern Washington at Manastash Ridge Observatory (MRO) equipped with a 1024×1024 CCD, and spectroscopic observations with the UW share of the 3.5 m telescope in New Mexico at Apache Point Observatory (APO) equipped with a Double Imaging Spectrograph. The spectrograph was used in high-resolution mode (resolution about 3 Å) with a 1"5 slit covering the wavelength regions of 4200–5000 Å in the blue and 6300–7300 Å in the red. Sporadic spectra on different nights were obtained to look for changes in the spectral lines and intensities that would identify a dwarf nova outburst or the high/low state of a nova-like object. For the brightest systems, time-resolved spectra over several hours were used to determine the orbital period and to measure radial velocities. The orbital period can also be an indicator of accretion rate, since the lowest rates generally exist in disk systems at quiescence that have orbital periods under 2 hr (Patterson 1984; Warner 1995), although low accretion rates can also occur in systems with longer orbital period when mass transfer sporadically turns off. The dates and modes of follow-up observations are listed in Table 3.

3.1. Previously Known Systems

There were three previously known CVs for which SDSS spectra were obtained, since their colors matched the selection criteria of quasars or serendipity. Two are relatively

well-known systems with published photometry and spectroscopy (SW UMa and BH Lyn) listed in the D01 catalog, as well as the Ritter & Kolb (1998) catalog. The third object (OU Vir, originally listed as Vir4 in the D01 catalog) is a recently discovered eclipsing dwarf nova with an orbital period of 1.75 hr (Vanmunster, Velthuis, & McCormick 2000). Two SDSS spectra of this object at quiescence were obtained (April and May; Table 1) and three APO spectra were obtained in May (Table 3). Since there is only one published spectra of OU Vir (Berg et al. 1992), we show its spectrum (SDSS 1435) in Figure 1. All the spectra of OU Vir reveal deep central absorption in the Balmer emission lines, which increases up the Balmer series. This feature is typical of high-inclination, eclipsing systems (Vogt 1982) and is postulated to be caused by low-velocity, cool outer disk material in the line of sight.

3.2. High-Inclination Systems

Two other systems also reveal the deep central absorption that is likely an indication of high inclination (SDSS 1555 and 0151). Confirmation of high inclination from an eclipse awaits time-resolved photometry or spectroscopy with larger aperture telescopes or observations during an outburst when the system is brighter.

3.3. Dwarf Novae

Since the SDSS photometry provides a separate epoch of observation from the SDSS spectroscopy and the Palomar DSS provides an additional observation, there are several instances in which the brightness of the system can be veri-




FIG. 1.—SDSS color-color plots of the 22 objects in Table 1. The filled circles are the CVs, while the dots are stars defining the stellar locus.

fied to be different, signifying either outbursts (in the case of dwarf novae) or high versus low states for nova-like objects. To distinguish these cases requires spectra at the high and low brightness states.

In the case of SDSS 1637, the SDSS photometry reveals a bright source ($g^* = 16.6$), while the spectra are consistent with a 20th magnitude object. In addition, the SDSS photometry reveals two close (but separated) stars, the northern object being the bright one, while the DSS shows it fainter than its southern neighbor. This fact is consistent with the northern object being caught at an outburst during the SDSS photometry. Several random follow-up spectra at APO (Table 3) and other quick-look images during the course of 3 months all found the system at its quiescent state, so the outbursts are likely infrequent or of short duration. As the quiescent spectra show none of the high-excitation lines of a nova-like system, this object is very likely a dwarf nova.

In two instances (SDSS 1730 and 2303), the APO spectra revealed objects undergoing a dwarf nova outburst, during which the Balmer emission lines turned into absorption lines when the systems were several magnitudes brighter than quiescence. On 2000 October 2, we obtained a total of 20 time-resolved spectra of SDSS 1730 at outburst and two

spectra, three nights later, when the system was at quiescence. Figure 3 shows typical outburst spectra and the smoothed mean of the two quiescent spectra for the blue and red wavelength regions. At outburst, the overall flux increased by a factor of 20 in the blue and four in the red, with noticeable, broad absorption features surrounding the $H\beta$ and $H\gamma$ emission lines. Figure 4 shows a close-up of the $H\beta$ line throughout the orbit. While inspection of the line suggests the possible presence of a narrow component passing from blue (phase 0.48) to red (phase 0.57), careful analysis of the velocity of the peak component did not produce an unambiguous feature that could be associated with the orbital period.

The presence of the underlying absorption features at $H\beta$ and $H\gamma$ in SDSS 1730 made it difficult to obtain radial velocities for these two lines, and a reliable solution was obtained only for the $H\alpha$ data. We obtained values for the central peak of the $H\alpha$ line by using IRAF's¹⁴ fitting rou-

¹⁴ IRAF (Image Reduction and Analysis Facility) is distributed by the National Optical Astronomy Observatory, which is operated by Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.




FIG. 2.—SDSS spectra of the newly discovered CVs. The flux scale is in units of flux density $10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$.




FIG. 2.—Continued

tines in SPLIT. As the line shows a fair amount of structure at outburst, the centroid “e” method returned more consistent answers than did Gaussian fitting. An IDL routine, CURVEFIT, was used to fit the radial velocities with a sinusoidal function of the form $v = \gamma - K \sin [2\pi(t - t_0)/P]$, where γ (systemic velocity), K (semiamplitude), P (orbital period), and t_0 (time of conjunction) were left as free parameters (see Table 4 for the best-fit parameters). Figure 5 shows the best fit to the $H\alpha$ radial velocity curve. Four spectra showing extreme deviations were not included in the fitting and are indicated in Figure 5 as open circles. We studied these spectra for evidence of shifts in the lines and for other potential sources of error but were unable to detect any cause. Nevertheless, they provided velocities that were clearly inconsistent with the rest of our measurements. The velocities for these four spectra shown in Figure 5 are from measurements of the midpoint of the extremes of the $H\alpha$ line done by eye. We tested our fitting program while includ-

ing these spectra, and found that the best fit obtained for the orbital period (117 ± 5 minutes) was consistent with the fit excluding these points.

Finally, we used the period obtained from the spectra to look for evidence of variability in the photometry of this system obtained at MRO. We folded the differential magnitudes on each of the two longest nights on the orbital period but found no evidence for any significant orbital variability. An average of the two nights’ data produces an upper limit for any such variability of 0.04 mag. The absence of variability, along with the lack of line doubling, suggests that SDSS 1730 is a system with a fairly low inclination (Warner 1995).

For SDSS 2303, we obtained a total of nine time-resolved spectra at quiescence on 2000 October 2 and 11 spectra at outburst three nights later. Figure 6 shows typical outburst and quiescent spectra for both nights. At outburst, the overall flux increases by a factor of 120 in the blue and 60 in the

TABLE 2
SDSS SPECTRAL LINE FLUXES AND EQUIVALENT WIDTHS^a

SDSS	PLATE, FIBER	H γ		H β		H α		He λ 4471		He II λ 4686	
		Flux	EW	Flux	EW	Flux	EW	Flux	EW	Flux	EW
0151	430, 430	0.9	15	1.7	36	2.6	77
0155	403, 423 ^b	59.7	10	63.4	11	37.8	13	12.2	2	44.5	8
0233	407, 592	1.9	43	2.5	74	4.5	303	0.2	5
0729	431, 316	0.5	17	0.6	24	0.8	35	0.07	3	0.5	19
0747	434, 430	0.4	0.8	0.6	1.2	0.9	3.3	0.8	1.5
0813	439, 582	1.6	15	1.9	14	2.8	15
0816	439, 624	3.5	43	3.8	53	5.3	91	0.9	12
0824	443, 360	1.0	17	1.5	26	1.9	46
0838	445, 89	2.9	65	3.2	87	5.1	211	0.4	8
1256	338, 382	0.4	11	0.4	14	0.7	52
1435 ^c ...	306, 4	2.9	19	2.5	23	4.4	64
1435 ^d ...	306, 20	22.4	59	18.3	66	19.0	105	3.9	11
1555	343, 198	4.9	33	5.9	62	10.1	195	0.8	5.5
1556	344, 315	1.7	6	2.8	14	6.2	62
1610	345, 138	0.9	5	1.9	16	5.1	86
1613	346, 263	5.1	11	5.4	14	8.3	27
1637	348, 103	2.6	80	3.0	129	3.4	186	0.6	21
1712	351, 17	6.0	100	6.8	79	8.3	90	2.0	28	2.0	23
1726	357, 51	1.4	56	1.6	94	1.3	158	0.3	13	0.25	12
1730	352, 26	16.1	58	15.4	63	17.5	118	3.4	14	1.3	6
2303	380, 575	2.8	52	3.3	74	5.1	193	0.5	10

^a Fluxes are in units of 10^{-15} ergs cm^{-2} s^{-1} , equivalent widths are in units of \AA .

^b Measurements from APO spectra as SDSS spectrum saturated.

^c Measured from spectrum on 2000 April 3.

^d Measured from spectrum on 2000 May 26.




FIG. 3.—APO spectra showing the outburst and quiescence of SDSS 1730. The fluxes have units of 10^{-16} ergs cm^{-2} s^{-1} \AA^{-1} .

TABLE 3
 FOLLOW-UP DATA FROM MRO AND APO

UT Date	SDSS	Site	Interval (hr)	Data Obtained
2000 May 6	1435	APO	0.3	2 spectra at quiescence
2000 May 15...	1435	APO	0.17	1 spectrum at quiescence
2000 May 28...	1637	APO	0.17	1 spectrum at quiescence
2000 Jul 3	1637	APO	0.25	1 spectrum at quiescence
2000 Jun 5	1730	MRO	5	V photometry
2000 Jun 16	1730	MRO	1.75	V photometry
2000 Jul 16	1730	MRO	2.4	V photometry
2000 Aug 7	1730	MRO	4.8	V photometry
2000 Aug 8	1730	MRO	7.2	B photometry
2000 Aug 25 ...	1730	MRO	2	B photometry
2000 Oct 3	1730	APO	2.2	20 spectra at outburst
2000 Oct 6	1730	APO	0.25	2 spectra at quiescence
2000 Oct 3	2303	APO	1.5	9 spectra at quiescence
2000 Oct 6	2303	APO	1.9	10 spectra at outburst
2000 Dec 29....	0155	APO	3	17 spectra at high state
2001 Jan 15	0747	APO	0.17	1 spectrum at high state

red, with broad absorption features surrounding the $H\beta$ and $H\gamma$ emission lines.

We obtained the velocities for the $H\alpha$ and $H\beta$ emission lines by using the IRAF fitting routines, with the Gaussian fitting routine “k” providing the best fit to a velocity curve (lowest residuals). Once again, the CURVEFIT routine was used to fit the radial velocity data with a sinusoidal function. The lines of $H\alpha$ and $H\beta$ from the quiescent data provided the most consistent solutions with the lowest errors. These solutions are given in Table 4, and the fit to the velocity curve for $H\alpha$ is shown in Figure 7. The outburst data resulted in consistent period and velocity curves. Using the best-fit period of 100 minutes, the phases for $H\beta$ were computed with the time of the red-to-blue crossing of the emission lines for $H\alpha$ as phase zero. The absence of any eclipse during quiescence or outburst suggests that SDSS 2303 is a system with a fairly low inclination.

To obtain further information about the disk at quiescence, we computed a Doppler tomogram. Doppler tomography (Horne 1991) is a technique that allows imaging of the line-forming regions in a binary system by combining the velocity profile information obtained at all binary phases. To construct our tomogram, we used the Fourier-filtered back projection program provided by Keith Horne and modified for our computers and plotting needs by Donald Hoard. Since we have no fiducial phasing from an eclipse or lines from the secondary, we used the phasing from the red-to-blue crossing of the $H\alpha$ line solution (Table 4) at quiescence. This assumes that the emission lines origi-




FIG. 4.—APO spectra showing the variation of the $H\beta$ line profile of SDSS 1730 throughout an orbital period during outburst.

nate near the white dwarf and that zero phase is inferior conjunction of the secondary. Figure 8 shows the tomograms for $H\alpha$ and $H\beta$ with a mass ratio of $q = 0.25$ used to compute the Roche lobe and stream. While a phase resolution of 0.1 is not suitable to determine any detailed disk structure, it can provide information on the disk extent and the stream. The tomograms of SDSS 2303 are consistent with a low mass transfer rate system with minimal disk emission and no evidence for a luminous hot spot near the stream location. There is some evidence for emission regions near the locations of the secondary ($V_y = +150$, $V_x = 0$) and of the white dwarf ($V_y = -150$, $V_x = 0$), but better data are required to confirm this.

3.4. Nova-likes with Strong He II

The three systems SDSS 0155, 0729, and 0747 reveal strong emission in He II $\lambda 4686$. While SDSS 0729 is very faint and has very narrow lines, indicative of a low mass transfer rate (possibly a magnetic) system, SDSS 0747 shows the bright blue continuum and weak Balmer emission of an old nova with a high accretion rate.

The system SDSS 0155 caught our attention for further study since the SDSS photometry and spectra indicated a very bright object, but the Palomar DSS showed a much

 TABLE 4
 RADIAL VELOCITY SOLUTIONS

SDSS	Line	P (minutes)	γ	K (km s $^{-1}$)	ϕ	σ
1730	$H\alpha$	117 ± 5	23 ± 3	74 ± 6	0.0 ± 0.02	19
2303	$H\alpha$	100 ± 14	9.3 ± 0.1	53 ± 7	0.0 ± 0.02	14
2303	$H\beta$	100	55.4 ± 0.1	49 ± 5	0.11 ± 0.02	9
0155	$H\alpha$	86 ± 2	89 ± 8	359 ± 29	0.09 ± 0.01	64
0155	$H\beta$	87 ± 2	135 ± 5	406 ± 20	0.05 ± 0.01	45
0155	He II $\lambda 4686$	88 ± 2	153 ± 4	404 ± 18	0.00 ± 0.01	38




FIG. 5.—Best fit to the radial velocity curve of SDSS 1730 at outburst using the emission line of $H\alpha$. The open circles produced large deviations to a sine fit and were thus left out of the best solution, which is shown as the solid line.

fainter star. The presence of strong emission at the bright state, combined with the presence of $He\ II$, suggests either an AM Her system with a magnetic white dwarf or a high accretion rate SW Sex star (characteristics of these types of CVs are reviewed by Warner 1995).

Our follow-up spectroscopy at APO, 1 month after the SDSS spectrum, showed the system also in its high state. Time-resolved spectra throughout 3 hr showed asymmetries and narrow component structure in the Balmer emission lines. Most important, an eclipse of the Balmer lines was evi-

dent (Fig. 9). When this occurred, the Balmer and $He\ I$ lines went into absorption while $He\ II$ remained visible but weaker, and the continuum did not appear to be eclipsed. This is opposite to the behavior typically seen in an eclipsing SW Sex system (e.g., BH Lyn; Hoard & Szkody 1997), where the $He\ II$ line (usually originating near the white dwarf) is deeply eclipsed along with the continuum light, while the Balmer emission lines (from the outer disk area) remain largely visible through the eclipse. Unfortunately, rising humidity levels necessitated a dome closure, which caused a 70 minute gap in the data sequence. The data sets before and after the dome closure cover intervals of 100 and 60 minutes, respectively.

Despite the gap, the velocities show a large and repetitive variation, allowing the determination of the orbital period to be 87 minutes. This short period also argues against an interpretation as an SW Sex system since those objects usually have orbital periods between 3–4 hr. The best fit to the radial velocity curve for $He\ II$ is shown in Figure 10, and the solutions are given in Table 4 for that line, as well as $H\beta$ and $H\alpha$. The very large (400 km s^{-1}) semi-amplitude of the velocity curves of SDSS 0155 compared with the other disk systems in Table 4 is typical of AM Her systems (Warner 1995) and strengthens the case for a magnetic interpretation for SDSS 0155.

Using the convention of the red-to-blue crossing of the $He\ II$ line as phase zero, the eclipse of the lines occurs at phase 0.7, which is maximum redshift. The most reasonable explanation for the lack of continuum eclipse is that the eclipse is of a line-forming area in a magnetic CV system. Because the lines are affected through more than one 10




FIG. 6.—Same as Fig. 3, but for SDSS 2303




Fig. 7.—Same as Fig. 5, but for SDSS 2303 at quiescence

minute exposure, the area must be larger than just a spot on the white dwarf. The emission region may be eclipsed by the secondary or self-eclipsed by the white dwarf. To test the geometry further, we constructed tomograms for the He II and H β lines, trying a phasing based on a zero of the red-to-blue crossing of the He II line (if its origin were on the white dwarf) and, alternatively, considering zero phase to be the eclipse. Figure 11 shows the results for the phasing of red-blue crossing, which produced tomograms that are consistent with those of AM Her stars (Hoard 1999; Schwöpe et al. 1999) with the brightest emission zones associated with the magnetic settling region (lower left quadrant) and the ballistic stream as drawn for a mass ratio of $\frac{1}{3}$. Unfortunately, the usual prominent areas due to the white dwarf and the irradiated secondary are not apparent, and so the actual phasing is ambiguous. Data obtained during a low

state should allow the absolute phasing to be determined by using the velocities of spectral lines from both the secondary star and the white dwarf.

3.5. Systems Showing the Secondary Star

Inspection of the spectra of Figure 2 shows that there are two systems that have spectral signatures of the secondary. SDSS 0813 shows the spectrum of a K5–M0 star (Mg band near 5200 Å and TiO at 7100 Å), similar to the long-period system TT Crt (Szkody et al. 1992). Thus, SDSS 0813 is likely to also have a long orbital period. The other system, SDSS 0816, also shows evidence for the TiO band at 7100 Å. The appearance of these features indicates a relatively low contribution of the accretion disk and, hence, a low mass transfer rate system.

3.6. ROSAT, 2MASS, and LONEOS Correlations

The newly identified CVs were cross-checked against the X-ray *ROSAT* All Sky Survey (RASS; Voges et al. 1999, 2000), the infrared Two Micron All Sky Survey (2MASS), and the optical Lowell Observatory Near-Earth Object Survey (LONEOS; Rest et al. 2002).

Besides the previously known bright system that is a known X-ray source (SW UMA; Shafter et al. 1986), five of the new CVs are detected in the RASS bandpass of 0.1–2.4 keV. Table 5 lists the X-ray count rates of these five sources. The two objects identified as dwarf novae (SDSS 1730 and 2303) are both detected, as well as the system with strong He II that is likely to be magnetic (SDSS 0155). As about half the known dwarf novae are detected by the RASS (Verbunt et al. 1997, hereafter V97), it is not so surprising that SDSS 1730 and 2303 are detected. However, it is unusual that the RASS count rates are identical for these two systems, while their optical fluxes differ by 3 mag. Since V97 found that a higher accretion rate corresponds to a lower X-ray to optical flux ratio, this would argue that the fainter system SDSS 2303 has the lower accretion rate (consistent with its shorter orbital period) compared with SDSS 1730,




Fig. 8.—Doppler tomograms of the H α (left) and H β (right) lines of SDSS 2303 at quiescence. The oval marks the secondary, the open circle below the origin marks the white dwarf, and the dotted line marks the mass transfer stream for a mass ratio of 0.25.




FIG. 9.—APO spectra of the He II $\lambda 4686$ and $H\beta$ lines throughout the orbital period in the likely magnetic system SDSS 0155 during a high state.




FIG. 10.—Best fit to the He II radial velocity curve of SDSS 0155

and that the lower optical flux is not just due to a greater distance for SDSS 2303 than SDSS 1730. In addition, it is remarkable that the two dwarf novae are brighter than the likely magnetic system SDSS 0155, as V97 and others have found that magnetic systems stand out as the brightest soft X-ray sources among CVs. However, this can be explained if SDSS 0155 was in a low state of mass accretion at the time of the RASS observations. We searched the existing X-ray data on the objects in Table 5 for any evidence of time variability and found a result only for SDSS 0233. In this case, the long pointing revealed variability of at least a factor of 4 on a timescale of 500 s. However, the low statistics and sampling do not allow any conclusion about periodicity.

Table 6 lists the 2MASS detections among the 19 new CVs (the known system SW UMa is also detected). The bright dwarf nova SDSS 1730 is among the detections and shows IR colors typical of accretion disks (Szkody 1985). The system showing a K–M dwarf secondary in the SDSS spectrum (SDSS 0813) is also detected and has red colors consistent with a K star. SDSS 1712 appears to have red col-




FIG. 11.—Same as Fig. 8, but for He II $\lambda 4686$ (left) and $H\beta$ (right) for SDSS 0155. The secondary, white dwarf, and mass transfer stream are shown for a mass ratio of 0.33.

TABLE 5
ROSAT DETECTIONS

SDSS	ROSAT (counts s ⁻¹) ^a	Exposure (s)	Observation ^b
0155	0.04 ± 0.01	394	S
0233	0.0058 ± 0.0005	28220	P
0838	0.017 ± 0.003	4569	P + S
1730	0.060 ± 0.006	2644	S
2303	0.061 ± 0.015	348	S

^a For a 2 keV bremsstrahlung spectrum, 1 counts s⁻¹ corresponds to a 0.1–2.4 keV flux of about 7×10^{-12} ergs cm⁻² s⁻¹.

^b “S” signifies a survey observation, “P” a pointed observation.

ors (also evident in the optical; Table 1) but does not show any spectral features from a late secondary (Fig. 2). Thus, this faint object may be at a large distance and/or may have significant reddening.

While the primary targets of the LONEOS project are moving objects, the existing database of a large fraction of the sky imaged in multiple epochs and calibrated to USNO-A2.0 red magnitudes provides a good resource for time-variable objects such as CVs. We searched the LONEOS database for our new SDSS CVs and found observations for the objects listed in Table 7 (data on the known system BH Lyn also exist). This table summarizes the range of variability evident along with the number of measurements made and the number of nights on which observations were obtained. Two of the four objects show evidence of large variability (more than 1 mag). The dwarf nova SDSS 2303 was near quiescence on two of the six measurements and at brighter magnitudes on the other four observations. Considering that one of our two APO measurements was also at outburst, the outburst timescale for this dwarf nova is likely to be very short. The other object showing large variability is the eclipsing, likely magnetic, system SDSS 0155. This object shows 1 mag variations among observations obtained on a single night (likely due to the eclipse), as well as 3 mag variations in observations separated by months (likely due to high and low states of mass transfer). The LONEOS database can ultimately be matched to the SDSS photometric fields to help identify new CV candidates by their variability.

4. CONCLUSIONS

The commissioning year of SDSS operation has shown that a wide variety of interesting CVs may be easily identi-

TABLE 6
2MASS DETECTIONS

SDSS	<i>J</i>	<i>H</i>	<i>K</i>
0747	16.59 ± 0.13	16.31 ± 0.22	15.5 ^a
0813	15.99 ± 0.09	15.30 ± 0.09	15.19 ± 0.13
1712	16.80 ± 0.16	15.94 ± 0.19	15.5 ^a
1730	15.30 ± 0.05	15.20 ± 0.09	15.28 ± 0.19

^a Detection only.

TABLE 7
LONEOS OBSERVATIONS

SDSS	No. of Observations	Days	Mag. Range ^a	Error
0155	21	8	14.7–17.6	0.05
0747	6	4	17.04–17.10	0.15
0824	7	2	17.60–17.75	0.1
2303	14	6	17.3–18.6	0.1

^a Calibration from USNO-A2.0 red magnitudes.

fied. Included among these are a few bright systems missed in past surveys, as well as members of the fainter population that models predict. This year of partial operation has led to the discovery of 19 new CVs and the recovery of three known systems. While the objects fainter than 20th mag at short orbital periods will need their orbital characteristics determined by 8–10 m class telescopes, the 15th–19th magnitude ones can be followed-up with 4 m class or smaller apertures. Using sporadic and time-resolved data, we have been able to identify two dwarf novae and one interesting, high-excitation, eclipsing system that is likely to be a magnetic CV. The fact that the orbital periods of these three systems are all short (under 2 hr) and that more than half of all the CVs found have large H β equivalent widths argue that we are finding lower accretion rate systems compared with previous surveys. These preliminary results show that the SDSS will be highly successful at identifying new CVs of a variety of types, with at least 400 total expected from the completed survey. Most of these should be faint, low accretion rate systems, which will provide insight into the nature and numbers of the true CV population in our galaxy.

The Sloan Digital Sky Survey (SDSS) is a joint project of the University of Chicago, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Max-Planck-Institut für Astronomie (MPIA), the Max-Planck-Institut für Astrophysik (MPA), New Mexico State University, Princeton University, the US Naval Observatory, and the University of Washington. Apache Point Observatory, site of the SDSS telescopes, is operated by the Astrophysical Research Consortium (ARC). Funding for the project has been provided by the Alfred P. Sloan Foundation, the SDSS member institutions, the National Aeronautics and Space Administration, the National Science Foundation, the US Department of Energy, the Japanese Monbukagakusho, and the Max Planck Society. The SDSS Web site is <http://www.sdss.org/>.

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center, funded by the National Aeronautics and Space Administration and the National Science Foundation.

The authors are grateful to the Lowell Observatory Near-Earth Object Search for making their data available for the CVs on our list.

REFERENCES

- Berg, C., Wegner, G., Folte, C. B., Chaffee, F.H., & Hewett, P. C. 1992, *ApJS*, 78, 409
- Downes, R. A., Webbink, R. F., Shara, M. M., Ritter, H., Kolb, U., & Dürbeck, H. W. 2001, *PASP*, 113, 764 (D01)
- Fan, X. 1999, *AJ*, 117, 2528
- Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
- Green, R. F., Ferguson, D. H., Liebert, J., & Schmidt, M. 1982, *PASP*, 94, 560
- Gunn, J. E., et al. 1998, *AJ*, 116, 3040
- Hoard, D. W. 1999, in *ASP Conf. Ser. 157, Annapolis Workshop on Magnetic Cataclysmic Variables*, ed. C. Hellier & K. Mukai (San Francisco: ASP), 201
- Hoard, D. W., & Szkody, P. 1997, *ApJ*, 481, 433
- Horne, K. 1991, in *Fundamental Properties of Cataclysmic Variable Stars*, ed. A. Shafter (San Diego: San Diego State Univ.), 23
- Howell, S. B., Rappaport, S., & Politano, M. 1997, *MNRAS*, 287, 929
- Krisciunas, K., Margon, B., & Szkody, P. 1998, *PASP*, 110, 1342
- Lupton, R. H., Gunn, J. E., Ivezić, Z., Knapp, G. R., Kent, S. M., & Yasuda, N. 2001, *ASP Conf. Ser. 238, Astronomical Data Analysis Software and Systems*, ed. F. R. Harnden, Jr., F. A. Primiini, & H. E. Payne (San Francisco: ASP), 269
- Lupton, R. H., Gunn, J. E., & Szalay, A. 1999, *AJ*, 118, 1406
- Patterson, J. 1984, *ApJS*, 54, 443
- Raymond, S., Szkody, P., & Hawley, S. L. 2002, *Proc. 12th Cambridge Workshop Cool Stars, Stellar Systems, and the Sun*, in press
- Rest, A., Miknaitis, G., Miceli, A., & Stubbs, C. 2002, in preparation
- Ritter, H., & Kolb, U. 1998, *A&AS*, 129, 83
- Schwöpe, A. D., Schwarz, R., Staude, A., Heerlein, C., Horne, K., & Steeghs, D. 1999, *ASP Conf. Ser. 157, Annapolis Workshop on Magnetic Cataclysmic Variables*, ed. C. Hellier & K. Mukai (San Francisco: ASP), 71
- Shafter, A. W., Szkody, P., & Thorstensen, J. R. 1986, *ApJ*, 308, 765
- Stoughton, C., et al. 2002, *AJ*, submitted
- Szkody, P. 1985, in *Cataclysmic Variables and Low-Mass X-Ray Binaries*, ed. D. Q. Lamb & J. Patterson (Dordrecht: Reidel), 385
- Szkody, P., Williams, R. E., Margon, B., Howell, S. B., & Mateo, M. 1992, *ApJ*, 387, 357
- Vanmunster, T., Velthuis, F., & McCormick, J. 2000, *Inf. Bull. Variable Stars*, No. 4955
- Verbunt, F., Bunk, W. H., Ritter, H., & Pfeffermann, E. 1997, *A&A*, 327, 602 (V97)
- Voges, W., et al. 1999, *A&A*, 349, 389
- , 2000, *IAU Circ.*, 7432
- Vogt, N. 1982, *ApJ*, 252, 653
- Warner, B. 1995, in *Cataclysmic Variable Stars* (Cambridge: Cambridge Univ. Press)
- York, D. G., et al. 2000, *AJ*, 120, 1579