

RESEARCH ARTICLE

Multiple Paternity in a Reintroduced Population of the Orinoco Crocodile (*Crocodylus intermedius*) at the El Frío Biological Station, Venezuela

Natalia A. Rossi Lafferriere^{1,2}*, Rafael Antelo^{3,4,5}, Fernando Alda^{4,6}, Dick Mårtensson⁷, Frank Hailer^{8,9}, Santiago Castroviejo-Fisher¹⁰, José Ayarzagüena^{5†}, Joshua R. Ginsberg^{1,11}, Javier Castroviejo^{5,12}, Ignacio Doadrio⁴, Carles Vilá¹³, George Amato²

1 Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, New York, United States of America, **2** Sackler Institute of Comparative Genomics, American Museum of Natural History, New York, New York, United States of America, **3** Fundación Palmarito Casanare, Bogotá, Colombia, **4** Dpto. Biodiversidad y Biología Evolutiva, Museo Nacional de Ciencias Naturales, CSIC, Madrid, Spain, **5** Estación Biológica El Frío, Apure, Venezuela, **6** LSU Museum of Natural Science, Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana, United States of America, **7** Department of Evolutionary Biology, Uppsala University, Uppsala, Sweden, **8** School of Biosciences, Cardiff University, Cardiff, CF10 3AX, Wales, United Kingdom, **9** Center for Conservation and Evolutionary Genetics, Smithsonian Conservation Biology Institute, National Zoological Park, Washington, DC, United States of America, **10** Lab. de Sistemática de Vertebrados, Pontificia Universidade Católica do Rio Grande do Sul (PUCRS), Porto Alegre, Brasil, **11** Cary Institute of Ecosystem Studies, Millbrook, New York, United States of America, **12** Asociación Amigos de Doñana, Seville, Spain, **13** Conservation and Evolutionary Genetics Group, Estación Biológica de Doñana (EBD-CSIC), Seville, Spain

† Deceased.

* These authors contributed equally to this work.

* nar2118@columbia.edu



CrossMark
click for updates

 OPEN ACCESS

Citation: Rossi Lafferriere NA, Antelo R, Alda F, Mårtensson D, Hailer F, Castroviejo-Fisher S, et al. (2016) Multiple Paternity in a Reintroduced Population of the Orinoco Crocodile (*Crocodylus intermedius*) at the El Frío Biological Station, Venezuela. PLoS ONE 11(3): e0150245. doi:10.1371/journal.pone.0150245

Editor: Samuel Rezende Paiva, Embrapa, BRAZIL

Received: September 27, 2015

Accepted: February 11, 2016

Published: March 16, 2016

Copyright: © 2016 Rossi Lafferriere et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: We have made our data available at the Dryad Digital Repository under the DOI: <http://dx.doi.org/10.5061/dryad.n2114>.

Funding: Funders included the Department of Evolutionary biology, Uppsala University, Uppsala, Sweden (DM), to support fieldwork and laboratory work; Dpto. Biodiversidad y Biología Evolutiva, Museo Nacional de Ciencias Naturales, CSIC, Madrid, Spain (RA FA SCF ID), to support fieldwork and laboratory work; Sackler Institute of Comparative Genomics at the American Museum of Natural

Abstract

The success of a reintroduction program is determined by the ability of individuals to reproduce and thrive. Hence, an understanding of the mating system and breeding strategies of reintroduced species can be critical to the success, evaluation and effective management of reintroduction programs. As one of the most threatened crocodile species in the world, the Orinoco crocodile (*Crocodylus intermedius*) has been reduced to only a few wild populations in the Llanos of Venezuela and Colombia. One of these populations was founded by reintroduction at Caño Macanillal and La Ramera lagoon within the El Frío Biological Station, Venezuela. Twenty egg clutches of *C. intermedius* were collected at the El Frío Biological Station for incubation in the lab and release of juveniles after one year. Analyzing 17 polymorphic microsatellite loci from 335 hatchlings we found multiple paternity in *C. intermedius*, with half of the 20 clutches fathered by two or three males. Sixteen mothers and 14 fathers were inferred by reconstruction of multilocus parental genotypes. Our findings showed skewed paternal contributions to multiple-sired clutches in four of the clutches (40%), leading to an overall unequal contribution of offspring among fathers with six of the 14 inferred males fathering 90% of the total offspring, and three of those six males fathering more than 70% of the total offspring. Our results provide the first evidence of multiple

History (NARL), to support data analysis and preparation of manuscript.

Competing Interests: The authors have declared that no competing interests exist.

paternity occurring in the Orinoco crocodile and confirm the success of reintroduction efforts of this critically endangered species in the El Frío Biological Station, Venezuela.

Introduction

Successful reproduction is critical to the recovery of endangered species. Many factors may limit reproduction, but in a small, newly established population knowledge of a species' mating system can be of particular importance to understanding the effective population size and the factors that drive reproductive success more generally [1]. Mating systems and their potential influence in the effective population size are also of critical importance for the design of *ex situ* conservation strategies for the recovery of endangered species [2]. Reintroduction of captive populations has proven challenging with approximately one-third of reintroductions failing [3,4] due to one or a combination of factors including poor habitat quality [5], altered behaviors (e.g. nest site selection, courtship rituals), and depleted genetic diversity [6].

For many species it is difficult, if not impossible, to regularly observe matings. Over the last thirty years, genetic tools have provided an increasingly accurate, and subtle, means to assess mating systems and to examine the result of differential mating success among both males and females [7]. Genetic tools have revealed that, throughout the animal kingdom, females of many species pursue a variety of alternative reproductive strategies exercising a more active role in mate choice [8,9], often mating with more than one male [10,11]. The result of this discovery of widespread polyandry is a better understanding of the frequency of extra pair paternity [12], female cryptic choice [13], and multiple paternity of a clutch or litter of offspring [11].

Multiple paternity of a single clutch, whereby offspring in that clutch are fathered by different males, can be the result of multiple mating within a single breeding season [13] or fertilizations from sperm stored from matings in previous seasons [14]. The evolution of female multiple mating (polyandry), and multiple paternity, can be driven by direct and indirect benefits. Direct benefits include provisioning of resources to the mother and paternal care of offspring [15,16]. Indirect benefits that may accrue include: improved quality of mate resulting in transfer of "good genes" or offspring of higher fitness [17,18], maximization of genetic diversity among the offspring resulting in bet-hedging against variation in the environment in successive generations [19,20], and improved genetic compatibility between mating pairs [21,22]. Mate encounter rates within a reproductive season also have been shown to influence both the occurrence and frequency of multiple paternity [23]. By increasing the number of males that contribute offspring to successive generations, and by changing the frequency of such contributions multiple paternity has also been shown, both theoretically [24] and empirically [25], to increase the effective population size [24], potentially increasing the overall genetic diversity of a population [26].

As a group, crocodylians were long believed to be polygynous and monandrous, with dominant males establishing breeding territories, excluding other adult males, and mating with multiple females [27–29]. This belief was primarily based on observations of crocodylians in captivity, supported with a few, mostly anecdotal, accounts of wild animals [30]. Multi-locus, highly variable, codominant markers such as microsatellites are a powerful genetic tool for parentage analysis [31]. They can be used for reconstruction of full-sibling and half-sibling families and inference of parental genotypes using maximum likelihood frameworks [32]. The use of molecular markers has shown multiple paternity to be an ubiquitous phenomenon [33–39].

Using these methods and data, evidence of multiple paternity has been observed in many crocodylian species, including members of families Crocodylidae [34–37] and Alligatoridae

[38,39]. The frequency of multiple-sired nests ranges from 30% to 90% [34–37]. Across studies, multiply sired clutches were fathered by two to four males [34–37]. Paternal contributions to multiply sired clutches were found to be skewed in the American alligator (*Alligator mississippiensis*) [38] and spectacled caiman (*Caiman crocodilus*) [37], with some males significantly contributing more than others to the total reproductive output. Overall, across all species of Crocodylia, these studies reveal high variability both in the prevalence of multiple paternity, measured as the percentage of multiple-sired clutches, and in the relative paternal contribution of males. This variability suggests that the incidence of multiple paternity is likely influenced by a variety of environmental factors as well as species behavior and life history traits.

Captive breeding and reintroduction has been used worldwide as a tool to recover critically endangered species of crocodylians including the Chinese alligator (*Alligator sinensis*) [40], the Cuban crocodile (*Crocodylus rhombifer*) [41], the Philippine crocodile (*Crocodylus mindorensis*) [42], the Siamese crocodile (*Crocodylus siamensis*) [43,44], the gharial (*Gavialis gangeticus*) [45], and the Orinoco crocodile (*C. intermedius*) [46]. With the exception of *A. sinensis* [40], information about mating systems and occurrence of multiple paternity based on genetic data is lacking. Genetic investigations of reproductive success could improve our understanding of crocodylian reproductive ecology and greatly aid *ex-situ* and *in-situ* conservation strategies, in particular the demographic and population structure of crocodylian reintroduction efforts.

One of the most threatened crocodile species in the world, the Orinoco crocodile [46,47] once inhabited large areas of the Llanos of Venezuela and Colombia within the Orinoco basin [48,49]. Extensive hunting until the 1960s and persistent collection of eggs for local consumption decimated its populations [50,51], with only few wild populations remaining in both countries [52]. In Venezuela, these include principally natural populations at Capanaparo, Cojedes and Manapire river systems [52], and a reintroduced population at Caño Macanillal and La Ramera lagoon within the El Frío Biological Station (EFBS) (Fig 1), and the adjacent Caño Guaritico Wildlife Refuge [53]. The original population at EFBS was extirpated during the 20th century. From 1990–2006 efforts were made to re-establish this population by the reintroduction of more than 2000 crocodiles raised in four captive breeding centers in Venezuela [53]. Post reintroduction, several management interventions were made to improve the survival of the *C. intermedius* at EFBS: river sand was supplemented along the shores of rivers and lagoons to facilitate the excavation of nests by females, eggs were collected and artificially incubated to prevent losses from nest predation, with the primary nest predator being the Tegú lizard (*Tupinambis teguixin*), juveniles were retained in captivity in a "head-start" program and released as one year-old juveniles to avoid the highest mortality rates which occur during the first year of life [46,53,54].

The occurrence of male hierarchies in Orinoco crocodiles during the breeding season was observed in previous studies [55,56]. However, both the extent to which males are able to monopolize breeding, and the occurrence of multiple mating, were unknown. Here, we use 17 polymorphic microsatellite loci to determine the occurrence of multiple paternity in 20 clutches of *C. intermedius* collected in EFBS, Venezuela. Specifically, we investigated the incidence of multiple paternity over three breeding seasons, used a maximum likelihood approach for sibship and parental inference, and assessed the degree of reproductive skew among inferred fathers.

Materials and Methods

Sampling

Twenty clutches of *C. intermedius* were collected during three consecutive years (n = 1 in 2004; n = 7 in 2005; and n = 12 in 2006) along artificial nesting sites placed in the wetland shorelines



Fig 1. Location of the El Frío Biological Station (EFBS), and three additional localities where last remaining populations of *Crocodylus intermedius* are found in Venezuela.

doi:10.1371/journal.pone.0150245.g001

of Caño Macanillal (7° 50' 11.45"N, 68° 55' 43.33"W) and La Ramera lagoon (7° 50' 5.65"N, 68° 55' 38.14"W) in the EFBS, Venezuela (Fig 1). Clutches were transported to the EFBS and incubated in artificially constructed nests as part of a captive breeding and reintroduction program for this species. Upon hatching, skin tissue samples were taken from all hatched crocodiles (n = 335). Samples were removed from the dorsal section of the base of the tail and stored at 8°C in 95% ethanol, following sampling protocols described in [36].

The Oficina Nacional de Diversidad Biológica, Ministerio del Ambiente y de los Recursos Naturales de Venezuela, granted tissue sample collection permits (Permit number 5-0358) and endorsed all other research activities of this project. CITES export permit number 1395/VE9120190 allowed *C. intermedius* skin tissue samples to be transported to Uppsala University, Sweden, and Museo Nacional de Ciencias Naturales (CSIC) in Madrid, Spain, to conduct genetic laboratory work.

Genotyping

Genomic DNA was extracted from preserved scales using the DNeasy blood and tissue kit (Qiagen, Valencia, CA) following manufacturer protocols. A panel of 17 microsatellite markers previously developed for *Crocodylus*, including C391, Cj16, Cj18, Cj101, Cj122, Cj127, CUJ131, Cu5123 [57], Cj109 [58], CpDi13, CpP302, CpP305, CpP314, CpP801, CpP1409, CpP1610, CpP3216 [59], was used to genotype all samples. Microsatellites were amplified in 4 multiplex PCRs (Mix1: CpP302, CpP305, CpP314, CpP1409, CpP3216, CpP1610; Mix2: C391, Cj16, Cj122, CUJ-131; Mix3: Cj18, Cj109, Cu5-123; Mix4: Cj101, Cj127, Cp801, CpDi13) using the Qiagen Multiplex PCR kit (Qiagen, Valencia, CA) for 30 cycles and two different annealing temperatures (57°C for Mix1, Mix2 and Mix4, and 60°C for Mix3). Reactions were prepared in a final volume of 12.5 μ L including: 6.25 μ L of Qiagen 2X PCR Master Mix, 2.5 μ L of 10X primer mix (final concentration: 2 μ M each, except for Cj16, Cj127 and CpP801, which were used at 4 μ M), 2 μ L DNA (ca. 10 ng/ μ L) and 1.75 μ L of RNase-free H₂O. Fluorescently labeled PCR products were electrophoresed on an ABI 3730 XL DNA analyzer with GS500 (-250) ROX size standard (Applied Biosystems, Foster City, CA) and allele sizes were determined using GeneMapper 3.7 software (Applied Biosystems Foster City, CA). Genetic laboratory work was conducted at Uppsala University, Sweden, and Museo Nacional de Ciencias Naturales, Spain.

The number of alleles, observed and expected heterozygosities, polymorphic information content, exclusion probabilities, allele frequencies, and tests for presence of null alleles were calculated in CERVUS 3.0.7 [60] for a sub-sample of one randomly selected individual per nest ($n = 20$). CERVUS 3.0.7 [60] uses genetic data from codominant markers to estimate allele frequencies and calculate various summary statistics and exclusion probabilities for each locus, assuming autosomal markers, linkage equilibrium between genetic markers, and diploid species. Conformity to Hardy-Weinberg (HW) expectations for each locus and genotypic linkage disequilibrium (LD) between pairs of loci were tested in GENEPOP 4.3 [61]. Significance levels ($p = 0.05$) for departure from HW and LD were corrected for multiple comparisons with a sequential Bonferroni correction [62].

Contamination and primer-site mutations may result in the incorrect assignment of microsatellite genotypes (genotyping errors), potentially biasing the results of parentage analysis [63]. Additionally, DNA degradation, low DNA concentration, and primer-site mutations can potentially cause non-amplified alleles (null alleles), which have proven to introduce errors in parentage analysis leading to high frequencies of erroneous parentage exclusions [64]. Potential genotyping inconsistencies including the presence of null alleles, large allele dropout, scoring of stutter peaks and typographic errors, were assessed in Micro-checker [65] and specific allele dropout rates were estimated in Microdrop 1.10 [66].

Paternity analysis

Sibship and parental inference analyses were conducted using a maximum likelihood approach implemented in COLONY 2.0 [32]. COLONY assigns individuals into full-sib and half-sib families using multi-locus genetic marker data with or without the imputing of parental information [32], and allowing for the incorporation of various types of genotyping errors [67]. By clustering offspring into full- and half-sib families COLONY is able to infer parental genotypes, when unknown, and calculate paternity assignment likelihoods, assuming Mendelian segregation and no maximum limit on the numbers of contributing parents. Ten replicate runs of “long” length and “high” likelihood precision were carried out in COLONY using the same dataset assuming an error rate of 0.04 for allelic dropout and 0.05 for genotyping error (based on the highest error rate per locus obtained in Microdrop 1.10 [68] (S1 Table) and suggested genotyping error rates by [67]). Polygamy was assumed for males and females and a full

Table 1. Characterization of 17 microsatellite loci in *Crocodylus intermedius* at the El Frío Biological Station, Venezuela.

Locus	n	Allele size range (bp)	No. Alleles	H _o	H _E	PIC	PE	PrDM
CpP1409	20	231–238	4	0.464	0.371	0.310	0.073	
CpP1610	20	274–278	2	0.143	0.135	0.124	0.012	
CpP302	20	168–204	10	0.821	0.862	0.829	0.559	
CpP305	20	160–182	5	0.536	0.531	0.484	0.109	
CpP314	20	236–250	6	0.500	0.592	0.516	0.184	
CpP3216	20	121–125	2	0.464	0.503	0.372	0.120	
C391	20	164–196	11	0.815	0.807	0.762	0.362	
CUJ131	20	202–210	4	0.536	0.551	0.474	0.154	
Cj122	20	392–406	4	0.667	0.713	0.645	0.250	
Cj16	20	160–194	7	0.679	0.722	0.660	0.264	
Cj109	20	380–408	11	0.750	0.774	0.733	0.317	
Cj18	20	225–235	5	0.571	0.573	0.481	0.167	
Cu5123	20	200–218	4	0.286	0.249	0.215	0.030	
Cj101	20	378–388	4	0.464	0.551	0.442	0.119	
Cj127	20	351–357	3	0.179	0.166	0.149	0.024	
CpDi13	20	342–348	3	0.500	0.499	0.370	0.085	
CpP801	20	164–184	5	0.538	0.651	0.571	0.270	
Overall				0.524	0.544	0.479	0.976	0.999/0.876

H_o: observed heterozygosity; H_E: expected heterozygosity; PIC: polymorphic information content; PE: probability of excluding a potential parent when neither parent is known; PrDM: probability of detecting multiple paternity given 17 progeny sampled per clutch and two fathers contributing 50:50 / 90:10.

doi:10.1371/journal.pone.0150245.t001

likelihood method was implemented. Offspring genotypes and known maternal sibship for each clutch were inputted into COLONY.

We used GERUD 2.0 [69] as alternative analysis package for the detection of multiple paternity. GERUD uses an exhaustive algorithm to determine the minimum number of fathers that can explain a full or half-sib progeny array when one or neither parent is known [69]. Because GERUD can only incorporate up to 10 loci, we used the 10 loci with highest exclusion probabilities (Table 1).

Probabilities of detecting multiple paternity were estimated using the PrDM software [70] (<http://publish.uwo.ca/~bneff/software.html>) for scenarios of equal (50:50) or skewed (90:10) paternal contribution by two males, averaged across 10 replicate runs, and using an average of 17 hatchlings per clutch sampled in this study. The degree of reproductive skew (B) was assessed for clutches exhibiting multiple paternity by calculating the binomial skew index B [71–73] in SKEW CALCULATOR 2003 (<https://www.eeb.ucla.edu/Faculty/Nonacs/PI.html>). Significant levels of B were calculated by simulation with 100,000 permutations. Significant positive values of B indicate a skewed paternal contribution to a given clutch, significant negative values of B indicate an excessively equal paternal contribution, and non significant values do not show differences from a random paternal contribution [71]. A Pearson's product moment correlation test between number of alleles per locus and exclusion probabilities, clutch size and number of fathers, and clutch size and probability of detecting multiple paternity (PrDM) was conducted using the package stats in R version 3.1.2 [74].

Results

Characterization of microsatellite markers

We identified a total of 90 alleles averaging 5.3 alleles per locus among a subsample of 20 Orinoco crocodile individuals, taking a random individual from each clutch. Observed and

expected heterozygosities ranged from 0.143 and 0.135 for CpP1610, respectively, to 0.821 and 0.862 for CpP302, exhibiting overall values of 0.524 and 0.544, respectively. Our analysis did not detect significant ($p < 0.05$) deviations from Hardy-Weinberg expectations and linkage equilibrium at any locus (S1 Table), or evidence of null alleles.

The discriminating power of the 17 markers combined was high (Table 1), with a probability of excluding a potential parent when neither parent is known (PE) of 0.976 and probabilities of detecting multiple paternity (PrDM) given 17 offspring sampled per clutch and two fathers contributing 50:50 and 90:10 of 0.999 and 0.876, respectively. However, exclusion probabilities (PE) of individual markers were mostly low, ranging from 0.012 to 0.559, and were highly correlated with the number of alleles per locus ($r = 0.831$).

On a clutch-by-clutch basis, probabilities of detecting multiple paternity (PrDM) remained relatively high ($p > 0.830$) for all clutch sizes under the equal paternal contribution scenario, and ranged from 0.587 to 0.852 under a skewed paternal contribution (90:10) (Table 2).

Paternity analysis

We obtained complete multilocus genotypes for all of the 17 loci and for more than 95% of the total individuals (335) sampled within a total of 20 clutches. Genotype data can be accessed at the Dryad Digital Repository under the DOI: [10.5061/dryad.n2114](https://doi.org/10.5061/dryad.n2114). Multiple paternity and parental reconstruction results were consistent over 10 COLONY replicate runs, assuming error rates of 0.04 for allelic dropout and 0.05 for genotyping error, and using different random seed numbers for each run. Our analyses revealed evidence of multiple paternity, with 10 (50%) out of 20 clutches fathered by two or three males (Table 2). We found multiple paternity in two of the three breeding seasons analyzed (2005 and 2006), with four out of seven clutches (57%) in 2005, and six out of 12 clutches (50%) in 2006 being fathered by two or three males. The only nest from 2004 was single-sired. GERUD and COLONY inferred multiple fathers for the same 10 clutches. The number of fathers remained the same using both programs for all but one clutch (C6), for which GERUD determined a minimum of three fathers whereas COLONY determined a minimum of two fathers. Due to the different number of markers used (a maximum of 10 loci for GERUD and all 17 for COLONY), the estimate offered by COLONY is considered more reliable.

For 90% of clutches, configurations of full-sib families inferred by COLONY exhibited high probabilities of both including all full-sib individuals in a given paternal family (Prob Inc.), and excluding all non full-sibs from each paternal family (Prob Exc.). Clutches 5 and 20 showed the lowest probabilities for full-sib paternal family configurations. Potential causes of these low values may be associated with mislabeling of hatchlings or mutations at one or several of the microsatellite loci that might have generated mismatches among sib hatchling genotypes.

Of the 10 multiple-sired clutches, four (40%) were significantly skewed from equal paternal contributions (Table 2). Four additional clutches (C6, C7, C14, C20) failed to show significant skew between fathers, which may be explained by small clutch size ($n < 10$) hindering the degree of reproductive skew. We did not find significant correlations between clutch size and number of fathers ($t = -1.639$, $df = 21$, $p = 0.116$), and clutch size and probability of multiple paternity (PrDM) ($t = -1.328$, $df = 21$, $p = 0.199$).

COLONY reconstructed 16 maternal and 14 paternal multilocus genotypes (S2 Table). Four of the 16 inferred mothers laid eggs on consecutive nesting seasons (M1, M5, M6, and M7), with two of these (M1 and M6) returning to the same geo-referenced nesting site (data not shown). Full siblings were observed in different nests when females mated with the same male in successive years. Some of these females laying eggs over consecutive seasons mated exclusively with the same male (M5), but the majority mated with more than one male in either one (M1, M7) or two (M6) consecutive breeding seasons.

Table 2. Paternity analyses for 20 clutches of *Crocodylus intermedius* from the El Frio Biological Station, Venezuela.

Clutch ID	Year	N	PrDM	GERUD		COLONY							p
				Number of inferred fathers	Inferred mother ID	Number of inferred fathers	Paternal sibship	Inferred father ID	N full siblings	Prob (Inc.)	Prob (Exc.)	B value*	
C1	2004	3	0.747/ 0.546	1	M1	1	1A	F1	3	1.000	1.000		
C2	2005	16	0.952/ 0.788	2	M2	2	2A	F2	15	1.000	1.000	0.3516	0.0005
							2B	F3	1	1.000	1.000		
C3	2005	32	0.925/ 0.744	1	M3	1	3A	F4	32	1.000	1.000		
C4	2005	19	0.983/ 0.801	2	M4	2	4A	F5	13	1.000	1.000	0.0416	0.1679
							4B	F1	6	0.986	0.986		
C5	2005	19	0.972/ 0.776	1	M5	1	5A	F2	19	0.604	0.604		
C6	2005	8	0.944/ 0.734	3	M1	2	6A	F1	7	1.000	1.000	0.2188	0.0684
							6B	F6	1	1.000	1.000		
C7	2005	8	0.971/ 0.768	2	M6	2	7A	F2	7	0.999	0.999	0.2188	0.0708
							7B	F3	1	1.000	1.000		
C8	2005	42	0.996/ 0.852	1	M7	1	8A	F2	42	1.000	1.000		
C9	2006	14	0.972/ 0.971	3	M8	3	9A	F7	9	0.949	0.904		
							9B	F8	4	0.811	0.797		
							9C	F2	1	1.000	1.000		
C10	2006	24	0.957/ 0.743	1	M9	1	10A	F2	24	1.000	1.000		
C11	2006	8	0.980/ 0.798	1	M5	1	11A	F2	8	1.000	1.000		
C12	2006	22	0.931/ 0.706	2	M10	2	12A	F9	21	1.000	1.000	0.3905	0.0000
							12B	F10	1	1.000	1.000		
C13	2006	23	0.940/ 0.718	1	M11	1	13A	F7	23	1.000	1.000		
C14	2006	3	0.929/ 0.746	2	M6	2	14A	F2	2	0.999	0.999	-0.1111	1.0000
							14B	F6	1	1.000	1.000		
C15	2006	13	0.983/ 0.807	3	M7	3	15A	F2	11	1.000	1.000	0.3432	0.0007
							15B	F3	1	1.000	1.000		
							15C	F11	1	1.000	1.000		
C16	2006	19	0.897/ 0.653	1	M12	1	16A	F12	19	0.998	0.998		
C17	2006	16	0.954/ 0.745	1	M13	1	17A	F7	16	1.000	1.000		
C18	2006	23	0.902/ 0.677	3	M14	3	18A	F13	21	1.000	1.000	0.4751	0.0000
							18B	F7	1	1.000	1.000		
							18C	F14	1	1.000	1.000		

(Continued)

Table 2. (Continued)

Clutch ID	Year	N	PrDM	GERUD		COLONY							p
				Number of inferred fathers	Inferred mother ID	Number of inferred fathers	Paternal sibship	Inferred father ID	N full siblings	Prob (Inc.)	Prob (Exc.)	B value*	
C19	2006	16	0.950/ 0.738	1	M15	1	19A	F7	16	1.000	1.000		
C20	2006	7	0.830/ 0.587	2	M16	2	20A	F10	4	0.706	0.706	-0.0612	1.0000
							20B	F11	3	0.549	0.549		

N: number of offspring used in paternity analysis; PrDM: probability of detecting multiple paternity given two fathers contributing 50:50/90:10, averaged over 10 replicate simulations; Prob (Inc.): probability that all individuals of a given full-sib paternal family are full-sibs; Prob (Exc.): probability that no other individuals are full-sibs within this paternal family; B value: binomial skew index; p: significance level of B value.

*Significant values of B are represented in bold.

doi:10.1371/journal.pone.0150245.t002

Shared paternity across clutches was inferred by COLONY, with highly skewed contributions among identified males (Fig 2). Six of the 14 inferred males (F2, F4, F7, F9, F12 and F13) fathered 90% of the total offspring, and three of those six males (F2, F4 and F7) fathered more than 70% of the total offspring.

Discussion

Our results report for the first time a polygamous mating system for *C. intermedius*, with females mating with more than one male in 10 out of 20 clutches studied. *Crocodylus*




Fig 2. Paternal contributions from 14 fathers inferred by COLONY for 335 hatchlings of *Crocodylus intermedius*.

doi:10.1371/journal.pone.0150245.g002

intermedius at the EFBS exhibited both single-fathered and multiple-fathered clutches, indicating that females can employ different reproductive strategies. Results of this study accord with other studies of multiple paternity in crocodylians including 30% multiple-sired clutches in *A. sinensis* [39], 32% in *A. mississippiensis* [38], 50% in *Crocodylus moreletii* [33], 50% in *Caiman latirostris* [34], 69% in *Crocodylus porosus* [36], 90% in *Melanosuchus niger* [35], and 95% in *C. crocodilus* [37]. In these studies, two to four males fathered multiply sired clutches.

Observed differences in the incidence of multiple paternity within and among crocodylian species could be influenced by one or a combination of factors, including: density of breeding individuals, mate encounter rates, sex ratio of sexually mature individuals, differences in mating behavior and reproductive strategies (e.g. ability of males to monopolize matings, male territory defense during the breeding season, male harassment, female choice, timing of copulation), and habitat type and spatial configuration of the environment (e.g. mangrove swamps, internal rivers and lagoons, small streams, ponds). Relatively small sample sizes, and a dearth of comparative data, make it difficult to identify which of these factors come in to play between and among these studies.

Multiple paternity in *C. intermedius* could have resulted from multiple mating within a single breeding season and/or fertilizations from sperm stored in females' reproductive tract within one or more breeding seasons. Sperm storage has proven an effective strategy to ensure fertilization and lengthen the breeding season in taxa where rates of encounter between the sexes are low, multiple clutches are laid within a single reproductive season, and/or where high asynchrony in gonadal cycles between males and females occurs [38,75,76]. Although sperm storage has not been reported in *C. intermedius*, sperm was found in the oviducts of female *A. mississippiensis* [77] and possibly occurred in a captive female of *Paleosuchus palpebrosus* which laid fertile eggs a long time after being separated from a male [78]. The absence of sperm from the oviduct of female American alligators during non-reproductive periods, however, suggested that alligator sperm may be retained within a reproductive season but is unlikely to be retained from one year to the next [77]. Female crocodylians generally congregate in breeding areas with groups of males during the reproductive season [79–81], lay one clutch per reproductive season on an annual or bi-annual basis [30], and some species showed a degree of synchrony in gonadal cycles between males and females (*A. mississippiensis* [82] and *C. niloticus* [83]). Given low population numbers, female sperm storage in *C. intermedius* (if it does occur) would most likely be associated with low frequency of mate encounter and low mate availability. Female cryptic choice (female selection for a particular male's sperm [84]) and the assurance of paternal diversity within clutches [85] could also potentially result in sperm storage in this species.

Our results, and observations from previous studies documenting male hierarchies in Orinoco crocodiles during the breeding season [55,56], suggest unequal distribution of matings. In this study shared paternity across clutches revealed highly skewed contributions among identified males. Similarly, paternal contributions of multiply sired clutches were skewed in *A. mississippiensis* [86] and *C. crocodilus* [37]. Skewed paternal contributions to clutches may arise by one or a combination of different mechanisms, such as: female preference for a "dominant" male, higher success for copulations occurring closest to female ovulation, sperm storage and female cryptic choice, and competition among male's sperm in female's reproductive tract (sperm competition [87]).

Four of the 16 identified mothers returned to lay eggs in two consecutive years, with two of them laying eggs in the same nesting site on the second year. In this recently reintroduced population, because of a lack of appropriate habitat, nesting sites have been artificially supplemented by providing river sand along the shores of rivers and lagoons to facilitate the excavation of nests by females. Despite the artificial construction of nest sites, two females

showed site fidelity, a phenomenon not uncommon to crocodylians [44]. Of the four returning females, two exhibited a switch from a single-sired clutch in one breeding season to a multiple-sired clutch in the subsequent breeding season, whereas the other two maintained either single-sired or multiple-sired clutches in both breeding seasons (Table 2). Hence, in this small sample, mating strategy, nesting site, and site fidelity, do not appear correlated, although further research is required to parse out the reasons for site fidelity.

A recent study of *C. acutus* [88] reporting nests with full siblings in successive mating seasons suggests the occurrence of mate fidelity in this species. In this study, we found that all four females that laid clutches on two consecutive seasons mated with at least one of the same males. Of these, only one mated exclusively with a single male, whereas the other three had multiple mates in either one or two breeding seasons. These findings could be a consequence of female choice and/or male dominance within this population.

Inference of parental genotypes from offspring multilocus marker data and derived interpretations needs to be carefully assessed. Maximum likelihood approaches for sibship and parental inference implemented by COLONY 2.0 [32] have been shown to overestimate parent numbers in both empirical [89] and simulated marker datasets [90] with low to intermediate levels of polymorphism. Nonetheless, parentage studies and family reconstructions of simulated datasets identified number and diversity of loci as main factors determining accuracy of results, with COLONY outperforming most available methods of parentage analysis [91]. In this study, the resolving power of the combined marker set was high. In addition, convergence of multiple runs in COLONY and high values of full-sib family exclusion and inclusion probabilities for the majority of the clutches suggested that our marker set could be sufficient for accurate parental reconstruction. The alternative method used to assess multiple paternity (GERUD) accorded with COLONY and confirmed the robustness of our analysis.

Information on the mating system and breeding strategies of reintroduced populations is critical to their success, evaluation and effective management [1]. The occurrence of multiple paternity may accelerate the recovery and resilience of reintroduced populations of *C. intermedius* by increasing genetic diversity among siblings, potentially increasing the effective population size [24], and overall genetic diversity of the population [26]. However, the dominance of certain males could potentially have the opposite effect by reducing the number of contributing males and increasing the variance in their contribution to the gene pool of next generations. In this study, we report heterozygosity values similar to those of other crocodylian populations in the wild [92–94]. The latter suggests that the genetic makeup and/or rapid population growth after individuals were reintroduced may have quickly compensated any potential founder effect, despite biases in the contribution of the different males.

Before the first reintroductions of *C. intermedius*, the species had been depleted from the El Frío Biological Station and surrounding areas [53] that fall within the historical range of the species [48,49]. After first reintroduction efforts took place in 1990, a combination of management interventions resulted in a successful population recovery with increasing nesting numbers adding up to at least 93 nests between 1996 and 2007 [53]. Nesting at the EFBS lays within nesting numbers previously reported at other sites: in the Capanaparo river system there were 11 nests in 1991 [81], 14 nests in 2001, 20 nests in 2002 [47], and 25 nests in 2013 [95]; during the 2009 nesting season 13 nests were laid in the Cojedes [52] and four nests in the Manapire river systems [47]. In the context of the few remaining natural populations of *C. intermedius* in Venezuela, the EFBS reproductive output makes it a very important population for the recovery of the species.

Successful reproduction is a necessary first step in the recovery of the Orinoco crocodile. However, other conservation actions will be critical to the continued and sustained growth of this and other populations. Conservation actions include mitigating the threat of poaching and

habitat protection [50]. Continued involvement and support from the governmental nature conservation authorities is essential for the long-term sustainability of these initiatives. Political changes in Venezuela led to a reduction in conservation and research activities at the El Frío Biological Station in 2009. Since then, the fate of the reintroduced Orinoco crocodile population, one of the last populations of *C. intermedius*, is little known. The limited information published in the popular press in recent years suggests that this population, and the species more generally, continues to be in need of urgent conservation actions [96].

Future research should aim to better elucidate the factors that influence multiple paternity in crocodilian species, looking at environmental (e.g. latitude, temperature, habitat type and configuration), demographic and species-specific traits. Research on specific mechanisms involved in producing multiple-fathered clutches, potential long-term adaptive advantages of multiple paternity for crocodilians, and impacts of multiple paternity on the overall genetic diversity of populations will further our knowledge on crocodilian mating systems and can aid conservation strategies for endangered crocodilian species.

Supporting Information

S1 Table. Additional information on 17 microsatellite loci used in *Crocodylus intermedius* paternity analysis.

(DOC)

S2 Table. Parental genotypes reconstructed for 20 *Crocodylus intermedius* clutches across 17 polymorphic microsatellite loci.

(DOC)

Acknowledgments

We are especially grateful to the staff of the El Frío Biological Station, for their support in the field and dedication, and the family Maldonado for their conservation efforts at Hato El Frío. We thank Dr. Josefa Señaris for her invaluable support in project development and processing of permits; staff of Uppsala University and Museo Nacional de Ciencias Naturales, Madrid, for their support in laboratory work; the Laboratorio de SIG y Teledetección (LAST) of Estación Biológica de Doñana, in Sevilla, Spain, for their GIS support; Angelica Menchaca Rodriguez, Nicole Mihnovets, Georgina Espinosa López, Dustin Rubenstein, Kent Vliet, and Martin Mendez for critical review of earlier versions of the manuscript.

Author Contributions

Conceived and designed the experiments: NARL RA FA DM FH SCF JA CV JC ID. Performed the experiments: NARL RA FA DM FH SCF JA CV JC ID. Analyzed the data: NARL RA. Contributed reagents/materials/analysis tools: NARL RA FA DM FH SCF CV JA JG JC ID GA. Wrote the paper: NARL RA FA FH SCF CV JG GA.

References

1. Schmidt F, Franke F, Shirley M, Vliet K, Villanova V (2015) The importance of genetic research in zoo breeding programmes for threatened species: the African dwarf crocodiles (genus *Osteolaemus*) as a case study. *International Zoo Yearbook* 49: 125–136.
2. Willoughby JR, Fernandez NB, Lamb MC, Ivy JA, Lacy RC, DeWoody J (2015) The impacts of inbreeding, drift and selection on genetic diversity in captive breeding populations. *Molecular Ecology* 24: 98–110. doi: [10.1111/mec.13020](https://doi.org/10.1111/mec.13020) PMID: [25443807](https://pubmed.ncbi.nlm.nih.gov/25443807/)
3. Fischer J, Lindenmayer DB (2000) An assessment of the published results of animal relocations. *Biological Conservation* 96: 1–11.

4. Pérez I, Anadón JD, Díaz M, Nicola GG, Tella JL, Giménez A (2012) What is wrong with current translocations? A review and a decision-making proposal. *Frontiers in Ecology and the Environment* 10: 494–501.
5. Moorhouse T, Gelling M, Macdonald D (2009) Effects of habitat quality upon reintroduction success in water voles: evidence from a replicated experiment. *Biological Conservation* 142: 53–60.
6. Jamieson IG, Lacy RC (2012) Managing genetic issues in reintroduction biology. *Reintroduction Biology: Integrating Science and Management* 12: 441.
7. DeWoody J, Avise J (2001) Genetic perspectives on the natural history of fish mating systems. *Journal of Heredity* 92: 167–172. PMID: [11396575](#)
8. Clutton-Brock T, McAuliffe K (2009) Female mate choice in mammals. *The Quarterly Review of Biology* 84: 3–27. PMID: [19326786](#)
9. Jennions MD, Petrie M (1997) Variation in mate choice and mating preferences: a review of causes and consequences. *Biological Reviews* 72: 283–327. PMID: [9155244](#)
10. Reynolds JD (1996) Animal breeding systems. *Trends in Ecology and Evolution* 11: 68–72. PMID: [21237764](#)
11. Taylor ML, Price TA, Wedell N (2014) Polyandry in nature: a global analysis. *Trends in Ecology and Evolution* 29: 376–383. doi: [10.1016/j.tree.2014.04.005](#) PMID: [24831458](#)
12. Adams EM, Jones AG, Arnold SJ (2005) Multiple paternity in a natural population of a salamander with long-term sperm storage. *Molecular Ecology* 14: 1803–1810. PMID: [15836651](#)
13. Canal D, Jovani R, Potti J (2012) Multiple mating opportunities boost protandry in a pied flycatcher population. *Behavioral Ecology and Sociobiology* 66: 67–76.
14. Brante A, Fernández M, Viard F (2011) Microsatellite evidence for sperm storage and multiple paternity in the marine gastropod *Crepidula coquimbensis*. *Journal of Experimental Marine Biology and Ecology* 396: 83–88.
15. Richardson DS, Burke T, Komdeur J (2002) Direct benefits and the evolution of female-biased cooperative breeding in Seychelles warblers. *Evolution* 56: 2313–2321. PMID: [12487360](#)
16. Moller AP (2000) Male parental care, female reproductive success, and extrapair paternity. *Behavioral Ecology* 11: 161–168.
17. Jennions MD, Petrie M (2000) Why do females mate multiply? A review of the genetic benefits. *Biological Reviews* 75: 21–64. PMID: [10740892](#)
18. Yasui Y (1998) The 'genetic benefits' of female multiple mating reconsidered. *Trends in Ecology and Evolution* 13: 246–250. PMID: [21238286](#)
19. Reed DH, Frankham R (2003) Correlation between fitness and genetic diversity. *Conservation Biology* 17: 230–237.
20. Bach L, Dahllöf I (2012) Local contamination in relation to population genetic diversity and resilience of an arctic marine amphipod. *Aquatic Toxicology* 114: 58–66. doi: [10.1016/j.aquatox.2012.02.003](#) PMID: [22421731](#)
21. Petrie M, Doums C, Moller AP (1998) The degree of extra-pair paternity increases with genetic variability. *Proceedings of the National Academy of Sciences of the United States of America* 95: 9390–9395. PMID: [9689090](#)
22. Rubenstein DR (2007) Female extrapair mate choice in a cooperative breeder: trading sex for help and increasing offspring heterozygosity. *Proceedings of the Royal Society B: Biological Sciences* 274: 1895–1903. PMID: [17526455](#)
23. Boomer JJ, Harcourt RG, Francis MP, Walker TI, Braccini JM, Stow AJ (2013) Frequency of multiple paternity in gummy shark, *Mustelus antarcticus*, and rig, *Mustelus lenticulatus*, and the implications of mate encounter rate, postcopulatory influences, and reproductive mode. *Journal of Heredity* 104: 371–379. doi: [10.1093/jhered/est010](#) PMID: [23505312](#)
24. Sugg DW, Chesser RK (1994) Effective population sizes with multiple paternity. *Genetics* 137: 1147–1155. PMID: [7982568](#)
25. Martinez J, Moran P, Perez J, De Gaudemar B, Beall E, Garcia-Vazquez E (2001) Multiple paternity increases effective size of southern Atlantic salmon populations. *Molecular Ecology* 9: 293–298.
26. Rafajlović M, Eriksson A, Rimark A, Hintz-Saltin S, Charrier G, Panova M, et al. (2013) The effect of multiple paternity on genetic diversity of small populations during and after colonisation. *PloS One* 8: e75587. doi: [10.1371/journal.pone.0075587](#) PMID: [24204577](#)
27. Thorbjarnarson J (1989) Ecology of the American Crocodile, *Crocodylus acutus*. In *Crocodiles, their Ecology, Management and Conservation IUCN Publications New Series Gland, Switzerland*: 228–258.

28. Alonso-Tabet M (2009) Comportamiento del cocodrilo americano (*Crocodylus acutus*) en el Refugio de Fauna Monte Cabaniguan, Cuba. Republica de Cuba: Universidad de Alicante and Universidad de Pinar del Rio Hermanos Saiz Montes de Oca.
29. Grigg G, Kirshner D (2015) Biology and Evolution of Crocodylians: CSIRO PUBLISHING.
30. Thorbjarnarson JB (1996) Reproductive characteristics of the order Crocodylia. *Herpetologica*: 8–24.
31. Jones AG, Ardren WR (2003) Methods of parentage analysis in natural populations. *Molecular Ecology* 12: 2511–2523. PMID: [12969458](#)
32. Jones OR, Wang J (2010) COLONY: a program for parentage and sibship inference from multilocus genotype data. *Molecular Ecology Resources* 10: 551–555. doi: [10.1111/j.1755-0998.2009.02787.x](#) PMID: [21565056](#)
33. McVay JD, Rodriguez D, Rainwater TR, Dever JA, Platt SG, McMurry ST, et al. (2008) Evidence of multiple paternity in Morelet's crocodile (*Crocodylus moreletii*) in Belize, CA, inferred from microsatellite markers. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* 309A: 643–648.
34. Amavet P, Rosso E, Markariani R, Ignacio Pina C (2008) Microsatellite DNA Markers Applied to Detection of Multiple Paternity in *Caiman latirostris* in Santa Fe, Argentina. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* 309A: 637–642.
35. Muniz FL, Da Silveira R, Campos Z, Magnusson WE, Hrbek T, Farias IP (2011) Multiple paternity in the Black Caiman (*Melanosuchus niger*) population in the Anavilhanas National Park, Brazilian Amazonia. *Amphibia-Reptilia* 32: 428–434.
36. Lewis JL, FitzSimmons NN, Jamerlan ML, Buchan JC, Grigg GC (2013) Mating systems and multiple paternity in the estuarine crocodile (*Crocodylus porosus*). *Journal of Herpetology* 47: 24–33.
37. Oliveira DP, Marioni B, Farias IP, Hrbek T (2014) Genetic evidence for polygamy as a mating strategy in *Caiman crocodylus*. *Journal of Heredity* 105: 485–492.
38. Davis LM, Glenn TC, Eelsey RM, Dessauer HC, Sawyer RH (2001) Multiple paternity and mating patterns in the American alligator, *Alligator mississippiensis*. *Molecular Ecology* 10: 1011–1024. PMID: [11348507](#)
39. Hu Y, Wu X-B (2010) Multiple paternity in Chinese alligator (*Alligator sinensis*) clutches during a reproductive season at Xuanzhou Nature Reserve. *Amphibia-Reptilia* 31: 419–424.
40. Wang Z, Yao H, Ding Y, Thorbjarnarson J, Wang X (2011) Testing reintroduction as a conservation strategy for the critically endangered Chinese alligator: Movements and home range of released captive individuals. *Chinese Science Bulletin* 56: 2586–2593.
41. Rodríguez-Soberón R, Ramos R, McMahan W, Ross J (1996) Reintroduction of the Cuban crocodile on the Isle of Pines. *Crocodile Specialist Group Newsletter* 15: 10–11.
42. Van Weerd M, Guerrero J, Balbas M, Telan S, Ven W, Rodriguez D, et al. (2010) Reintroduction of captive-bred Philippine Crocodiles. *Oryx* 44: 13.
43. Daltry J, Starr A (2010) Development of a re-introduction and re-enforcement program for Siamese crocodiles in Cambodia. In: *Global Re-introduction Perspectives: Additional Case Studies from Around the Globe* (ed Soorae P), pp 118–123 Re-introduction Specialist Group and Environment Agency, Abu Dhabi, UAE: 118.
44. Eelsey RM, Trosclair PL III, Glenn TC (2008) Nest-site fidelity in American alligators in a Louisiana coastal marsh. *Southeastern Naturalist* 7: 737–743.
45. Ballouard JM, Priol P, Oison J, Ciliberti A, Cadi A (2010) Does reintroduction stabilize the population of the critically endangered gharial (*Gavialis gangeticus*, Gavialidae) in Chitwan National Park, Nepal? *Aquatic Conservation: Marine and Freshwater Ecosystems* 20: 756–761.
46. Antelo R, Ayarzagüena J, Castroviejo J (2010) Reproductive ecology of Orinoco crocodiles (*Crocodylus intermedius*) in a newly established population at El Frío Biological Station, Venezuela. *The Herpetological Journal* 20: 51–58.
47. Barros T, Jiménez-Oraá M, Heredia HJ, Seijas AE (2010) Artificial incubation of wild-collected eggs of American and Orinoco crocodiles (*Crocodylus acutus* and *C. intermedius*), Guárico and Zulia, Venezuela. *Conservation Evidence* 7: 111–115.
48. Gumilla J (1745) *El Orinoco Ilustrado y Defendido*, Historia Natural, Civil, y Geographica de este Gran Rio: Manuel Fernandez, editor.
49. Thorbjarnarson JB (1992) Recent investigations of the status and distribution of the Orinoco crocodile (*Crocodylus intermedius*) in Venezuela. *Biological Conservation* 62: 179–188.
50. Casal AC, Fornelino MM, Restrepo MFG, Torres MAC, Velasco FG (2013) Uso histórico y actual del caimán llanero (*Crocodylus intermedius*) en la Orinoquia (Colombia-Venezuela). *Biota Colombiana* 14: 65–82.

51. Medem F (1981) Los Crocodylia de Sur América, Vol. I. Los Crocodylia de Colombia Editorial Carrera 7a Ltda, Bogotá.
52. Espinosa-Blanco AS, Seijas AE, Hernández O (2013) Egg collection as a conservation tool of Orinoco crocodile (*Crocodylus intermedius*) in the Cojedes river system, Venezuela. *Interciencia* 38: 358–363.
53. Antelo R (2008) Biología del Cocodrilo o Caimán del Orinoco (*Crocodylus intermedius*) en la Estación Biológica El Frío, Estado Apure (Venezuela): Tesis Doctoral. Universidad Autónoma de Madrid, Spain.
54. Ramo C (1992) Breeding and rearing of the Orinoco crocodile (*Crocodylus intermedius*) in Venezuela. *Biological Conservation* 60: 101–108.
55. Seijas A, Chávez C (2002) Reproductive status and nesting ecology of the Orinoco crocodile (*Crocodylus intermedius*) in the Cojedes River System, Venezuela. *Vida Silvestre Neotropical* 11: 23–32.
56. Thorbjarnarson JB, Hernández G (1993) Reproductive ecology of the Orinoco Crocodile (*Crocodylus intermedius*) in Venezuela. I. Nesting ecology and egg and clutch relationships. *Journal of Herpetology*: 363–370.
57. FitzSimmons N, Tanksley S, Forstner M, Louis E, Daglish R, Gratten J, et al. Microsatellite markers for *Crocodylus*: new genetic tools for population genetics, mating system studies and forensics; 2001. Surrey Beatty & Sons. pp. 51–57.
58. Dever JA, Densmore LD (2001) Microsatellites in Morelet's crocodile (*Crocodylus moreletii*) and their utility in addressing crocodylian population genetics questions. *Journal of Herpetology* 35: 541–544.
59. Miles L, Isberg S, Moran C, Hagen C, Glenn T (2009) 253 Novel polymorphic microsatellites for the salt-water crocodile (*Crocodylus porosus*). *Conservation Genetics* 10: 963–980.
60. Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* 16: 1099–1106. PMID: [17305863](#)
61. Rousset F (2008) Genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106. doi: [10.1111/j.1471-8286.2007.01931.x](#) PMID: [21585727](#)
62. Rice WR (1989) Analyzing tables of statistical tests. *Evolution*: 223–225.
63. Hoffman J, Amos W (2005) Microsatellite genotyping errors: detection approaches, common sources and consequences for paternal exclusion. *Molecular Ecology* 14: 599–612. PMID: [15660949](#)
64. Dakin E, Avise J (2004) Microsatellite null alleles in parentage analysis. *Heredity* 93: 504–509. PMID: [15292911](#)
65. Van Oosterhout C, Hutchinson WF, Wills DP, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535–538.
66. Wang C, Schroeder KB, Rosenberg NA (2012) A maximum-likelihood method to correct for allelic dropout in microsatellite data with no replicate genotypes. *Genetics* 192: 651–669. doi: [10.1534/genetics.112.139519](#) PMID: [22851645](#)
67. Wang J (2004) Sibship reconstruction from genetic data with typing errors. *Genetics* 166: 1963–1979. PMID: [15126412](#)
68. Wang C, Rosenberg NA (2012) MicroDrop: a program for estimating and correcting for allelic dropout in nonreplicated microsatellite genotypes version 1.01.
69. Jones AG (2005) GERUD 2.0: a computer program for the reconstruction of parental genotypes from half-sib progeny arrays with known or unknown parents. *Molecular Ecology Notes* 5: 708–711.
70. Neff B, Pitcher T (2002) Assessing the statistical power of genetic analyses to detect multiple mating in fishes. *Journal of Fish Biology* 61: 739–750.
71. Nonacs P (2000) Measuring and using skew in the study of social behavior and evolution. *The American Naturalist* 156: 577–589.
72. Nonacs P (2003) Measuring the reliability of skew indices: is there one best index? *Animal Behaviour* 65: 615–627.
73. Liu J-X, Avise JC (2011) High degree of multiple paternity in the viviparous Shiner Perch, *Cymatogaster aggregata*, a fish with long-term female sperm storage. *Marine Biology* 158: 893–901. PMID: [24391263](#)
74. R Development Core Team (2010). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing
75. Pearse D, Avise J (2001) Turtle mating systems: behavior, sperm storage, and genetic paternity. *Journal of Heredity* 92: 206–211. PMID: [11396580](#)
76. Uller T, Olsson M (2008) Multiple paternity in reptiles: patterns and processes. *Molecular Ecology* 17: 2566–2580. doi: [10.1111/j.1365-294X.2008.03772.x](#) PMID: [18452517](#)

77. Gist DH, Bagwill A, Lance V, Sever DM, Eisey RM (2008) Sperm storage in the oviduct of the American alligator. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* 309: 581–587.
78. Davenport M (1995) Evidence of possible sperm storage in the caiman, *Paleosuchus palpebrosus*. *Herpetological Review* 26: 14–14.
79. Garrick LD, Lang JW (1977) Social signals and behaviors of adult alligators and crocodiles. *American Zoologist* 17: 225–239.
80. Vliet K (2001) Courtship behaviour of American alligators *Alligator mississippiensis*. *Crocodylian Biology and Evolution*: 383–408.
81. Thorbjarnarson JB, Hernández G (1993) Reproductive ecology of the Orinoco crocodile (*Crocodylus intermedius*) in Venezuela. II. Reproductive and social behavior. *Journal of Herpetology*: 371–379.
82. Lance VA (1989) Reproductive cycle of the American alligator. *American Zoologist* 29: 999–1018.
83. Kofron CP (1990) The reproductive cycle of the Nile crocodile (*Crocodylus niloticus*). *Journal of Zoology* 221: 477–488.
84. Birkhead T (2000) Defining and demonstrating postcopulatory female choice—again. *Evolution* 54: 1057–1060. PMID: [10937281](#)
85. Calsbeek R, Bonneaud C, Prabhu S, Manoukis N, Smith TB (2007) Multiple paternity and sperm storage lead to increased genetic diversity in *Anolis* lizards. *Evolutionary Ecology Research* 9: 495.
86. Davis LM (2001) Multiple paternity and mating patterns in the American alligator, *Alligator mississippiensis*. *Molecular Ecology* 10: 1011–1024. PMID: [11348507](#)
87. Andersson M, Iwasa Y (1996) Sexual selection. *Trends in Ecology & Evolution* 11: 53–58.
88. Budd KM, Spotila JR, Mauger LA (2015) Preliminary Mating Analysis of American Crocodiles, *Crocodylus acutus*, in Las Baulas, Santa Rosa, and Palo Verde National Parks, Guanacaste, Costa Rica. *South American Journal of Herpetology* 10: 4–9.
89. Jones B, Grossman GD, Walsh DC, Porter BA, Avise JC, Fiumera AC (2007) Estimating differential reproductive success from nests of related individuals, with application to a study of the mottled sculpin, *Cottus bairdi*. *Genetics* 176: 2427–2439. PMID: [17565960](#)
90. Sefc KM, Koblmüller S (2009) Assessing parent numbers from offspring genotypes: the importance of marker polymorphism. *Journal of Heredity* 100: 197–205. doi: [10.1093/jhered/esn095](#) PMID: [18984859](#)
91. Harrison HB, Saenz-Agudelo P, Planes S, Jones GP, Berumen ML (2013) Relative accuracy of three common methods of parentage analysis in natural populations. *Molecular Ecology* 22: 1158–1170. doi: [10.1111/mec.12138](#) PMID: [23278953](#)
92. Dever JA, Strauss RE, Rainwater TR, McMurry ST, Densmore LD III, Wood R (2002) Genetic diversity, population subdivision, and gene flow in Morelet's crocodile (*Crocodylus moreletii*) from Belize, Central America. *Copeia* 2002: 1078–1091.
93. Farias IP, Silveira RD, de Thoisy B, Monjeló LA, Thorbjarnarson J, Hrbek T (2004) Genetic diversity and population structure of Amazonian crocodylians. *Animal Conservation* 7: 265–272.
94. Milián-García Y, Venegas-Anaya M, Frias-Soler R, Crawford AJ, Ramos-Targarona R, Rodríguez-Soberón R, et al. (2011) Evolutionary history of Cuban crocodiles *Crocodylus rhombifer* and *Crocodylus acutus* inferred from multilocus markers. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* 315: 358–375.
95. Hernández O, Boede E, Amauci J (2014) Evaluation of the Orinoco crocodile nesting (*Crocodylus intermedius*) in Capanaparo River, Santos Luzardo National Park, Apure State, Venezuela *Boletín de la Academia de las Ciencias Físicas, Matemáticas y Naturales* 2: 53–59.
96. Neuman W, Ramon P (December 26 2013) Venezuela's Fitful Effort to Save a Scaly Predator. *The New York Times*. pp. A4.