



Nemoz, C., Ropars, V., Frit, P., Gontier, A., Drevet, P., Yu, J., ... Charbonnier, J. B. (2018). XLF and APLF bind Ku80 at two remote sites to ensure DNA repair by non-homologous end joining. *Nature Structural and Molecular Biology*, 25(10), 971-980. <https://doi.org/10.1038/s41594-018-0133-6>

Peer reviewed version

Link to published version (if available):  
[10.1038/s41594-018-0133-6](https://doi.org/10.1038/s41594-018-0133-6)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Springer Nature at <https://www.nature.com/articles/s41594-018-0133-6#Abs1>. Please refer to any applicable terms of use of the publisher.

## **University of Bristol - Explore Bristol Research**

### **General rights**

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/pure/about/ebr-terms>

1 **XLF and APLF bind to Ku80 on two remote sites to ensure DNA repair by**  
2 **non-homologous end-joining**

3  
4 **AUTHORS**

5 Clement Nemoz<sup>1\*</sup>, Virginie Ropars<sup>1\*</sup>, Philippe Frit<sup>2,3\*</sup>, Amandine Gontier<sup>1</sup>, Pascal Drevet<sup>1</sup>,  
6 Jinchao Yu<sup>1</sup>, Raphaël Guerois<sup>1</sup>, Aurelien Pitois<sup>1</sup>, Audrey Comte<sup>1</sup>, Christine Delteil<sup>2,3</sup>,  
7 Nadia Barboule<sup>2,3</sup>, Pierre Legrand<sup>4</sup>, Sonia Baconnais<sup>5</sup>, Yandong Yin<sup>6</sup>, Satish Tadi<sup>7</sup>,  
8 Emeline Barbet-Massin<sup>8</sup>, Imre Berger<sup>9</sup>, Eric Le Cam<sup>5</sup>, Mauro Modesti<sup>7</sup>, Eli Rothenberg<sup>6</sup>,  
9 Patrick Calsou<sup>2,3,§</sup>, Jean-Baptiste Charbonnier<sup>1,§</sup>

10  
11 **INSTITUTIONS**

12 <sup>1</sup> Institute for Integrative Biology of the Cell (I2BC), Institute Joliot, CEA, CNRS, Univ.  
13 Paris-Sud, Université Paris-Saclay, 91198, Gif-sur-Yvette cedex, France

14 <sup>2</sup> Institut de Pharmacologie et Biologie Structurale, IPBS, Université de Toulouse, CNRS,  
15 UPS, Toulouse, France

16 <sup>3</sup> Equipe Labellisée Ligue Contre le Cancer 2013 and 2018

17 <sup>4</sup> Synchrotron Soleil, L'Orme des Merisiers, Saint-Aubin - BP 48, 91192 Gif-sur-Yvette  
18 Cedex, France.

19 <sup>5</sup> Signalisations, Noyaux et Innovations en Cancérologie, UMR 8126, CNRS, Université  
20 Paris-Sud, Gustave Roussy, Université Paris-Saclay, 94805 Villejuif, France.

21 <sup>6</sup> New York University School of Medicine, Perlmutter Cancer Center, USA

22 <sup>7</sup> Cancer Research Center of Marseille, CNRS UMR7258, Inserm U1068, Institut Paoli-  
23 Calmettes, Aix-Marseille Université UM105, Marseille, France.

24 <sup>8</sup> Dynamic Biosensors GmbH, Martinsried, Germany

25 <sup>9</sup> BrisSynBio Centre, The School of Biochemistry, Faculty of Biomedical Sciences,  
26 University of Bristol, Bristol, United Kingdom.

27  
28 \* These authors contributed equally

29 § Corresponding authors : [jb.charbonnier@cea.fr](mailto:jb.charbonnier@cea.fr) and [calsou@ipbs.fr](mailto:calsou@ipbs.fr)

33 **ABSTRACT**

34 The Ku70-Ku80 (Ku) heterodimer binds rapidly and tightly to ends of DNA double-strand  
35 breaks and recruits several factors of the Non-Homologous End Joining (NHEJ) pathway  
36 through molecular mechanisms that remain unclear. Here, we describe the crystal structures  
37 of the Ku-binding motifs (KBM) of the NHEJ proteins APLF (A-KBM) and XLF (X-KBM)  
38 bound to a Ku-DNA complex. The two KBMs motifs bind on remote sites of Ku80  $\alpha/\beta$   
39 domain. The X-KBM occupies an internal pocket formed after an unprecedented large  
40 outward rotation of the Ku80  $\alpha/\beta$  domain. We reveal independent recruitment at laser-  
41 irradiated sites of the APLF-interacting protein XRCC4 and of XLF through the respective  
42 binding of A- and X-KBMs to Ku80. Finally, we show that mutations on the X-KBM and A-  
43 KBM binding sites in Ku80 compromises efficiency and accuracy of end-joining and cellular  
44 radiosensitivity. A- and X-KBMs may represent two initial anchorage points necessary to  
45 build the NHEJ intricate interactions network.

46

47 Keywords : DSB repair, NHEJ, X-ray crystallography, induced fit, laser micro-irradiation,  
48 super resolution microscopy, switchSENSE, microcalorimetry

49

50

51 **INTRODUCTION**

52

53 In mammals the majority of DNA double-strand breaks (DSBs) is repaired by the non-  
54 homologous end-joining (NHEJ) pathway <sup>1,2</sup>. The Ku70-Ku80 heterodimer (Ku) rapidly and  
55 tightly interacts as a preformed ring with DSBs extremities in a non-sequence specific manner  
56 <sup>3</sup>. Ku serves as a hub for the recruitment of several NHEJ factors <sup>4,5</sup>. Among them, Ku recruits  
57 the XRCC4-LIG4-XLF ligation complex through interaction with XRCC4-LIG4 <sup>6,7</sup> and with  
58 XLF, the latter relying on a Ku-binding motif (KBM) localized at the XLF extreme C-  
59 terminus (thereafter named X-KBM) <sup>8,9</sup> (Figure 1a and Supplementary Figure 1). The ligation  
60 complex organizes into filaments both *in vitro* and in cells <sup>10-14</sup>.

61

62 Interestingly, Ku interacts also with a number of accessory NHEJ factors. The APTX and  
63 PNKP-like factor (APLF) binds poly(ADP)-ribosylated proteins near DSBs sites <sup>15,16</sup>, and has  
64 been reported to have nuclease activity <sup>16,17</sup>. APLF tightly interacts with Ku through a KBM  
65 (thereafter named A-KBM) that is located in its central region <sup>9,18</sup> (Figure 1a and  
66 Supplementary Figure 1). This interaction has been mapped to the periphery of the Ku80 von  
67 Willebrand A domain (vWA) <sup>5</sup>. Ku-APLF interaction was shown to facilitate recruitment of  
68 the APLF-partner XRCC4 at damaged sites <sup>9</sup> and was proposed to stabilize the assembly of  
69 NHEJ factors around the DSB <sup>19</sup>. Notably, an A-KBM-like domain is present at the N-  
70 terminus of a recently identified inhibitor of the NHEJ pathway, CYREN(MRI), that also  
71 interacts with Ku80 <sup>20</sup> (Figure 1a). Ku also associates with the Werner syndrome protein  
72 (WRN) that is involved in many aspects of DNA metabolism including NHEJ <sup>21</sup>. Two motifs  
73 in the C-terminus of WRN cooperate for interaction with Ku, one being A-KBM like, and the  
74 other resembling the X-KBM present on XLF (Figure 1a). In addition, we and others showed  
75 recently that PAXX (Paralog of XRCC4 and XLF) interacts with the Ku70 subunit through a  
76 third type motif that is located in its C-terminus <sup>22,23</sup> (Figure 1a). Despite identification of  
77 KBMs in several NHEJ factors, their respective contribution to the efficiency of DSB repair is  
78 not fully understood. For example, the puzzling observations that KBM deletion in XLF or  
79 APLF depletion in human cells lead to null or intermediate repair defect deserve further  
80 investigations <sup>9,24-27</sup>.

81

82 The interactome of Ku thus defines a large ensemble of motifs and proteins that could  
83 potentially compete or act synergistically. However, despite important structural and  
84 biophysical studies on NHEJ complexes <sup>28</sup>, the absence of high resolution structures of Ku-

85 KBMs complexes limits our understanding of the roles and specificity of the different  
86 molecular interactions in the recruitment of NHEJ factors to DSBs. Mapping KBM-binding  
87 sites in structures is also needed to clarify potential competition of all the Ku interacting  
88 factors on limited positions (Supplementary figure 1).

89

90 Here, we establish the structural and functional basis of Ku unique modes of interaction with  
91 two factors within the NHEJ repair pathway. We describe the first crystal structures of Ku70-  
92 Ku80-DNA complex in interaction with the A-KBM (KBM of APLF) and with the X-KBM  
93 (KBM of XLF), revealing that the two KBMs occupy remote interaction sites on the Ku80  
94 vWA domain. The functional context of our structural data was determined using cell based  
95 assays to visualize the recruitment of wild-type and mutant X-KBM motifs or XLF proteins as  
96 well as of the APLF-partner XRCC4 to DSBs sites induced by micro-irradiation in wild-type  
97 or mutant Ku backgrounds. Our data provide new mechanistic insights on the function of XLF  
98 and APLF in the NHEJ process.

99

100

## 101 RESULTS

102

### 103 The KBM of APLF tightly interacts with a highly conserved site of Ku80 vWA domain.

104

105 The APLF factor contains a conserved Ku binding motif (A-KBM, aa 179-192) (Figure 1a).  
106 We co-crystallized a 18-mer A-KBM peptide (aa 174-191) with a Ku form deleted for the C-  
107 terminal regions (Ku<sub>cc</sub>), and an hairpin DNA (hDNA)<sup>3</sup> (Supplementary Figure 2a). In the  
108 crystal structure at 3.0 Å resolution (Table 1), the A-KBM peptide is well defined and is  
109 positioned at the periphery of the vWA domain of Ku80 (aa T61<sup>80</sup>-C156<sup>80</sup>) (thereafter,  
110 superscript<sup>80</sup> stands for Ku80) (Figure 1b, and Supplementary Figure 2b). It is located at  
111 more than 50 Å from the DNA binding site of Ku80. The A-KBM adopts an extended  
112 conformation in a pocket delineated by the helices  $\alpha 4$  and  $\alpha 5$  and the loop located between the  
113  $\beta$ -strands B and C of Ku80 (Figure 1c). The hydrophobic part of the A-KBM, located in the  
114 C-terminal part of the motif, is composed by the amino acids I<sub>185</sub>LPTWML<sub>191</sub> and is buried in  
115 a hydrophobic pocket formed by the Ku80 residues L68<sup>80</sup>, I112<sup>80</sup>, M115<sup>80</sup>, I149<sup>80</sup> and I150<sup>80</sup>  
116 (Figure 1d). The N-terminal part of the A-KBM contained a patch of three consecutive basic  
117 residues and an acid residue (E<sub>181</sub>RKR<sub>184</sub> in human sequence) (Figure 1e). It forms salt

118 bridges and charged hydrogen bonds with respectively the side chains of D106<sup>80</sup>, D109<sup>80</sup>,  
119 Q73<sup>80</sup> and S145<sup>80</sup>, and the main chain of K144<sup>80</sup> and S143<sup>80</sup>.

120

121 Isothermal Titration Calorimetry (ITC) showed that the interaction between A-KBM and Ku  
122 had a Kd of  $33 \pm 10$  nM (Table 2, and Supplementary Figure 2e). We measured a nanomolar  
123 Kd for the interaction between Ku and a 18bp DNA as already reported<sup>29</sup> and found a similar  
124 Kd for the interaction between the A-KBM and Ku alone or Ku bound to a 18bp DNA (Table  
125 2). The interaction of the A-KBM with Ku<sub>cc</sub> showed Kd and enthalpy values similar to full-  
126 length Ku (KuFL). Thus, the core heterodimeric region of Ku is likely sufficient for the  
127 interaction with the A-KBM. Notably, these affinities are stronger than the ones previously  
128 reported by fluorescence polarization with a labelled A-KBM peptide (Kd of 580 nM)<sup>9</sup>.

129

130 Mapping the conservation rate of the residues at the surface of Ku80 shows that this pocket is  
131 the main conserved pocket together with the DNA binding pocket (Figure 1f). The residues  
132 L68<sup>80</sup>, Y74<sup>80</sup> and I112<sup>80</sup> make tight interactions with the hydrophobic part of the A-KBM  
133 motif. Mutations at these positions were reported to greatly reduce or disrupt the interactions  
134 with APLF in yeast two-hybrid experiments or in EMSA<sup>9</sup>. We produced the Ku I112R  
135 mutant that, as expected had no residual interaction with the A-KBM motif by ITC (Table 2).

136

137

### 138 **X-KBM creates an outward rotation of the vWA domain and a large groove in Ku80.**

139

140 We then determined the crystal structures of the Ku70-Ku80-hDNA complex bound with  
141 peptides derived from the XLF X-KBM ((L281<sup>X</sup> to S299<sup>X</sup>) and (S287<sup>X</sup>-S299<sup>X</sup>) peptides)  
142 (Supplementary Figure 2c). The crystal structures at 2.8 and 2.9 Å resolution (Table 1) show  
143 an unprecedented large outward rotation of the Ku80 vWA (Figure 2a, b). This movement  
144 forms a large groove between the Ku80 vWA and the rest of the Ku heterodimer. We  
145 therefore termed this conformation the open state of Ku, in contrast to the closed state  
146 observed in the three other crystal structures reported (Ku alone (1JEQ), Ku-hDNA (1JEY)  
147 and Ku-hDNA-A-KBM (this study)). The conformational change of Ku80 vWA does not  
148 affect Ku interaction with the duplex DNA (Supplementary Figure 2d)

149

150 The X-KBM is located on the Ku80 vWA face of the newly created groove in a pocket  
151 delineated by 4 strands ( $\beta_A$ ,  $\beta_D$ ,  $\beta_E$  and  $\beta_{E'}$ ) and 3 helices ( $\alpha_2$ ,  $\alpha_7$ ,  $\alpha_7'$ ). The motif is positioned

152 closer to the DNA (at 12Å) than the A-KBM, though not in direct contact. We can model  
153 eight residues of the X-KBM (<sub>292</sub>KKPRGLFS<sub>299</sub>) in the crystal obtained with the 19mer or  
154 14mer X-KBM. These residues are the last eight residues of the XLF sequence. The residues  
155 <sub>296</sub>GLFS<sub>299</sub> of the X-KBM occupy a hydrophobic pocket delineated by Ku80 residues L12<sup>80</sup>,  
156 V37<sup>80</sup>, F41<sup>80</sup>, F135<sup>80</sup>, F164<sup>80</sup>, Y225<sup>80</sup>, and L234<sup>80</sup> (Figure 2c). In the closed state, these Ku  
157 residues are buried and mediate intramolecular contacts (Figure 2d). The X-KBM may thus  
158 stabilize a transient open conformation of Ku80 in equilibrium with the closed state under  
159 basal conditions. In addition, we observed that the Ku80 vWA opening comes along with  
160 important secondary structure changes in the linker region (R232<sup>80</sup>-E241<sup>80</sup>) that separates the  
161 vWA and the rest of the Ku heterodimer (Figure 2c, d).

162

163 ITC measurements showed a moderate affinity of X-KBM for Ku, with a K<sub>d</sub> of 4.4 ± 0.2 μM  
164 (about 200-fold weaker than the affinity of the A-KBM) (Table 2, and Supplementary Figure  
165 2f). We observed similar affinities and thermodynamic parameters for the X-KBM with a Ku-  
166 DNA complex or with Ku<sub>cc</sub> (Table 2). Notably, this micromolar interaction was not detected  
167 in a previous study using fluorescence polarization with labelled peptides<sup>9</sup>. We also measured  
168 similar affinities for the interactions between XLF homodimer and Ku alone or Ku bound to a  
169 18bp DNA (Table 2). These data show that in absence of DNA or with a short DNA protected  
170 by Ku ring, XLF and its X-KBM interact similarly with Ku.

171

172 EMSA analyses confirmed an interaction between XLF and Ku complexed with a 50bp DNA  
173 in the μM range (Supplementary Figure 3a-b). Competitions experiments showed that the X-  
174 KBM peptide (pXLF) competes in the μM range with the XLF protein and that the A-KBM  
175 peptide (pAPLF) does not displace XLF, supporting remote sites of interactions (Figure 2e  
176 and Supplementary Figure 3c, d). Also, the C-terminus of PAXX (pPAXX) does not compete  
177 with XLF binding, in agreement with previous studies that report an interaction between  
178 PAXX C-terminus and Ku70 subunit<sup>22,23</sup> (Supplementary Figure 3e).

179

180

181

182 **The outward rotation of the vWA domain of Ku80 is mediated by Glu133.**

183

184 To evaluate if the outward rotation of Ku80 observed in the crystal structure with the X-KBM  
185 peptide was present in solution, we performed SAXS analyses. The SAXS data with the A-

186 KBM peptide were comparable with those of the Ku-hDNA sample without any peptide  
187 (respective Rg of 53.1Å and 53.7) (Figure 2f). We measured an increase of the Rg to 59.0Å in  
188 presence of the X-KBM, corresponding to the opening of Ku80 observed in the crystal  
189 (Figure 2f). Comparison of the Dmax values provided further corroboration for the opening of  
190 the Ku molecule with an increase of 30 Å.

191

192 The acid residue E133<sup>80</sup> buried in Ku80 is well positioned to act as a spring facilitating the  
193 Ku80 opening (Figure 2c, d). The glutamate E133<sup>80</sup> is buried in the closed conformation of  
194 the Ku80 vWA and its pKa value is estimated by the PDB2PQR-2.0 server<sup>30</sup> at a value of 9.1,  
195 far from the normal pKa of 4.5 for a glutamate in solvent. The outward rotation of Ku80 vWA  
196 should be energetically facilitated by the solvation of this Glu following the displacement of  
197 Ku80 residues V236<sup>80</sup>, F237<sup>80</sup> and I240<sup>80</sup> away from E133<sup>80</sup> carboxylate function (Figure 2d).  
198 Multiple sequence alignments show that E133<sup>80</sup> position and the residues surrounding are  
199 well conserved in mammalian and saurian and that Ku70 has no equivalent buried acidic  
200 residue at this position (Supplementary Figure 4a-b).

201

## 202 **Molecular bases of the specificities of A-KBM and X-BKM motifs binding to Ku80.**

203

204 The A-KBM and X-KBM present sequence similarities with a basic patch in their N-terminus  
205 followed by a hydrophobic patch<sup>9</sup> (Supplementary Figure 1). Comparison of the crystal  
206 structures of Ku bound to these two motifs suggests that the high affinity of the A-KBM relies  
207 on the tryptophan W189<sup>A</sup> in place of the Leu297<sup>X</sup> in X-KBM. Thus, we used ITC to measure  
208 the interaction of the X-KBM motifs with the mutation L297W (LW) or with a non-  
209 conservative L297E mutation (LE). The (LW) peptide has a Kd of 0.12 ± 0.03 μM, an  
210 interaction 40-fold tighter than wild-type X-KBM (Table 2 and Supplementary figure 2g).  
211 The LE mutant presents no detectable interaction with Ku (Table 2). Competition experiments  
212 with Ku saturated with the A-KBM showed that the X-KBM (L297W) no longer interacts  
213 with Ku, suggesting that the sole L297W mutation is able to redirect the X-KBM towards the  
214 A-KBM binding site on Ku80 (Table 2).

215

216 Then live cell imaging was used to monitor the recruitment of CFP-fused A-KBM and X-  
217 KBM fragments to DSBs sites induced with laser micro-irradiation. Under conditions of  
218 similar damage yield (Supplementary Figure 5e), the A-KBM motif transfected in U2OS cells  
219 was strongly recruited but the W189G mutation impaired both its nuclear localization and



220 recruitment to laser sites (Figure 3a-b), as reported<sup>9,18</sup>. X-KBM live recruitment was impaired  
221 by L297E mutation but not L297W mutation (Figure 3c), corroborating ITC data. We then  
222 used U2OS cells expressing an inducible shRNA against Ku80<sup>31</sup> (Supplementary Figure 5a)  
223 that were complemented with wild-type or I112R mutant Ku80 (Supplementary Figure 5c-d).  
224 The I112R mutation impaired A-KBM recruitment, as expected, but not that of X-KBM  
225 (Figure 3d, e). Notably, I112R Ku80 mutant specifically lowered the recruitment of L297W  
226 X-KBM (compare Figures 3c and 3f). Conversely, APLF knock-down boosted the  
227 recruitment of the LW mutant peptide above that of wild-type X-KBM (Figure 3g and  
228 Supplementary Figure 5b for control of shAPLF efficiency). Together, these data in cells  
229 support that the LW mutation redirects the X-KBM fragment to the APLF-binding site in  
230 Ku80 and point out the W189 residue as a key determinant for APLF specific interaction with  
231 Ku80.

232

233

#### 234 **X-KBM mutations impair XLF recruitment and XRCC4-XLF filament stability.**

235

236 We then investigated the properties of the interaction between full-length XLF protein and  
237 Ku. We first used the SwitchSENSE approach<sup>32</sup> in which oligonucleotide nanolevers labelled  
238 with a fluorescence probe are bound to a gold surface (Supplementary Figure 3f). Ku bound  
239 onto 48bp DNA nanolevers with a Kd in the nM range as already reported<sup>29</sup> and a long  
240 dissociation time (Supplementary Figure 3g). Wild-type XLF onto the Ku-DNA complex  
241 showed a rapid  $k_{on}$  ( $4.7 \pm 1.7 \cdot 10^5 \text{ M}^{-1}\text{s}^{-1}$ ) followed by a rapid dissociation ( $k_{off} = 0,09 \pm 0,004$   
242  $\text{s}^{-1}$ ) and a corresponding Kd of  $0.19 \pm 0.07 \mu\text{M}$  (Figure 4a). This affinity is about 10 fold  
243 stronger than the one measured by ITC with a smaller DNA and may reflect additional  
244 interactions of XLF with DNA emerging from Ku ring as observed with PAXX<sup>22</sup>. LW and  
245 LE mutants showed a 2.3- and 5.1-fold weaker affinity than WT protein, respectively (Kd of  
246  $0.45 \pm 0.26 \mu\text{M}$  for LW and Kd of  $0.98 \pm 0.15 \mu\text{M}$  for LE) (Supplementary Figure 3h). As  
247 compared with ITC and recruitment data with X-KBM LW mutant peptide, this suggests that  
248 the LW mutation cannot redirect the XLF protein to the APLF binding site in Ku80.

249

250 We then measured the recruitment of the CFP-tagged full-length XLF protein (CFP-XLF)  
251 expressed in human XLF-defective BuS cells (SV40T-transformed, telomerase immortalized  
252 fibroblasts derived from the XLF-deficient P2 patient - homozygous C622T nonsense mutant  
253 (R178X))<sup>33</sup> (Figure 4b). Wild-type XLF showed a rapid mobilization to irradiated nuclear

254 sites, as reported <sup>34</sup>. Both LE and LW mutations impaired to various extents XLF protein  
255 recruitment ([Figure 4b](#)), again indicating that LW mutation cannot redirect the XLF protein to  
256 the APLF binding site in Ku80. In addition, we observed an important reduction of the  
257 recruitment of G296W, S299E or F298G XLF mutants in the extreme C-terminus  
258 ([Supplementary Figure 5f-h](#)). We also questioned the contribution of APLF-XRCC4 complex  
259 to XLF recruitment by expressing L115D XLF mutant unable to interact with XRCC4 <sup>35,36</sup>.  
260 We measured an efficient recruitment of L115D XLF to laser-induced DSBs that was  
261 insensitive to APLF knock-down but impaired with the XLF L115D/L297E XLF double  
262 mutant ([Supplementary Figure 5i](#)). Together, these data support a major role for Ku80  
263 interaction with L297 and extreme C-terminal residues for XLF recruitment at DSBs in cells.

264

265 Multi-color super-resolution localization microscopy (STORM) allows characterizing  
266 formation of XRCC4-XLF filaments close to Ku foci and DSBs ends <sup>14</sup>. On DSBs induction  
267 with the radiomimetic drug neocarzinostatin, extended XLF filaments close to Ku80 foci were  
268 observed in XLF complemented BuS cells, whereas cells harboring (LE) and (LW) mutants  
269 showed slightly smaller and more punctuated XLF structures ([Figure 4c](#) and [Figure 4d-e for](#)  
270 [quantification](#)). In contrast to these data, evaluation of the effect of mutations in the X-KBM  
271 on cell radiosensitivity showed that (LW) or (LE) XLF mutants were associated with  
272 respectively no or minor radiosensitization, compared to high radiosensitivity of BuS cells  
273 and full restoration of radioresistance on expression of wild-type XLF ([Figure 4f](#)).

274

275

#### 276 **APLF and XLF binding to Ku80 promote DSB repair and cell survival to IR.**

277

278 Finally, we questioned the discrepancy between subnormal cell survival and defective XLF  
279 recruitment and filaments formation associated with mutations in X-KBM. The outward  
280 rotation in Ku80 upon X-KBM binding more likely relies on E133<sup>80</sup>, the equivalent position  
281 of which in Ku70 is a methionine (M167) ([Supplementary Figure 4b](#)). Therefore, we designed  
282 E133M mutant in Ku80 and Q162E mutation that may alter the charge environment of E133  
283 ([Supplementary Figure 5c](#)). E133M or Q162E mutation in Ku80 negatively impacted X-KBM  
284 recruitment but not that of control A-KBM ([Supplementary Figures 6a and 6b](#)). This result  
285 further supports the independent binding sites on Ku80 of the A-KBM and X-KBM motifs  
286 and validates E133 and Q162 positions in Ku80 as key residues for X-KBM binding.

287

288 Then, mutations in the Ku80 binding sites for X- and A-KBM were combined in the  
289 I112R/E133M Ku80 double mutant (Supplementary Figure 5d). This combination of  
290 mutations clearly impaired the recruitment of both CFP-(X- and A-KBM) peptides (Figures  
291 5a and 5b) while the recruitment of the mutant CFP-Ku fusions was preserved  
292 (Supplementary Figure 6c-d). We also measured the recruitment of the APLF partner XRCC4  
293 and of XLF, expressed as CFP-tagged full-length proteins (Figure 5c-d). Strikingly, only the  
294 double E133M/I112R mutation strongly but not completely impaired the recruitment of both  
295 proteins (Figure 5c-d). This result supports the independent binding of APLF-XRCC4 and  
296 XLF proteins on Ku80.

297

298 Ku80 mutants expressed in U2OS cells had no detectable impact on XLF filaments  
299 (Supplementary Figure 6e-f) but lowered end-joining efficiency on a linear plasmid  
300 transfected in U2OS cells (Figure 5e). To assess repair accuracy, we used an assay in which  
301 GFP expression from a cassette integrated in cells relied on the loss of a DNA fragment  
302 between two cut sites (modified from <sup>37</sup>) (Figure 5f), reported to be favoured upon NHEJ  
303 inhibition <sup>38,39</sup>. Strikingly, we observed an increase in fragment loss with the three Ku80  
304 mutants (Figure 5f), indicating that loose APLF-XRCC4 or XLF interactions with Ku80  
305 promote genomic instability. Finally, the combination of E113M and I112R Ku80 mutations  
306 had a strong impact on cell radiosensitivity with an additive effect of both mutations (Figure  
307 5g). These results indicate that APLF-XRCC4 and XLF interactions with Ku80 cooperate to  
308 promote cell survival to IR.

309

310

## 311 **DISCUSSION**

312

313 Here, we show that each of the A- and X-KBM binds Ku independently of other APLF or  
314 XLF protein domains. This is in contrast with the absence of binding reported for the isolated  
315 X-KBM in recent studies with fluorescent polarization <sup>9,40</sup> that may be due to lower  
316 sensitivity, steric hindrance by the fluorescence probe or differences in the Ku constructs  
317 used. Therefore, the present results rule out the conclusion that Ku-XLF interaction  
318 necessarily needs XLF binding to DNA (even though DNA could stabilise the interaction, as  
319 deduced from our switchSENSE data) or to DNA-PKcs <sup>40</sup>.

320

321 Despite the A-KBM and X-KBM motifs have important sequence similarities ([Supplementary](#)  
322 [figure 1](#)), they target different regions of Ku80 with different modes of actions, a more rigid  
323 one for the A-KBM and an induced fit one for the X-KBM. In contrast to the L297W X-KBM  
324 peptide, the (LW) full-length XLF mutant protein cannot be redirected to the APLF-binding  
325 site on Ku80. This may be explained by steric constraints either intrinsic to the protein  
326 structure or due to interactions with other partners in the NHEJ complex. It was reported that  
327 a X-KBM in WRN protein functions cooperatively with an A-KBM located upstream (1403-  
328 1412) in binding Ku complexes <sup>40</sup> ([Figure 1a](#)). Our structural data allow proposing the first  
329 model of Ku interacting with the tandem sequence of the WRN A-KBM and X-KBM,  
330 including the central linker ([Supplementary Figure 4c](#)). This model now awaits validation by  
331 the crystal structure of the tandem motif of WRN on Ku.

332

333 XRCC4 and XLF organize into filaments both *in vitro* and in cells <sup>10-14</sup> and can also assemble  
334 as sliding sleeves-like structures on broken DNA *in vitro* even without Ku <sup>41</sup>. In this study,  
335 cells containing single or double mutations on the A-KBM and/or X-KBM binding sites of  
336 Ku80 show intact filaments while mutations on the X-KBM of XLF induce a slight reduction  
337 in the filaments size. These data suggest that interactions additional to Ku80-XLF may  
338 participate in filament formation, like the XLF C-terminal DNA-binding domain that spans  
339 the X-KBM <sup>42</sup>.

340

341 How is NHEJ ligation complex assembled at broken DNA? APLF FHA domain interacts with  
342 XRCC4 <sup>17,26</sup> and XLF establishes head to head contact with XRCC4 <sup>10-13</sup>. Here, we show that  
343 the two remote APLF- and XLF-KBM binding sites in Ku80 promote the independent  
344 recruitment of XRCC4 and XLF at broken DNA and that disruption of XRCC4-XLF  
345 interaction (through L115D mutation) does not compromise XLF recruitment. Destabilisation  
346 of APLF or XLF interactions with Ku80 impairs repair efficiency and cell survival to DSBs  
347 and also favours genome instability associated with distal end-joining. These effects are most  
348 likely explained by loose assembly of the NHEJ apparatus at break ends. Therefore, we  
349 propose a model in which APLF and XLF KBMs represent two initial anchorage points for  
350 the rapid and independent recruitment of APLF-XRCC4 and XLF on Ku ([Figure 5h](#)). After  
351 initial recruitment, interactions additional to Ku80-XLF contacts may stabilize XLF at DSB  
352 sites ([Figure 5h](#)): our recruitment data show a stable interaction of XLF at DSB sites in cells  
353 that differs from the rapid dissociation from Ku-DNA observed *in vitro* by switchSENSE;  
354 also, although XRCC4 is dispensable for XLF initial recruitment, it has been shown to

355 stabilize XLF at damaged sites <sup>34</sup>. Thus, additional contacts stabilizing XLF may include  
356 XLF-DNA <sup>34,42,43</sup> and/or XLF-XRCC4 interactions, including filaments formation <sup>10-14</sup> and/or  
357 interaction with the LIG4 BRCT1 domain <sup>44</sup>. In addition, after XRCC4 recruitment through  
358 APLF binding to Ku, XRCC4 may also stabilize in turn APLF at damaged sites since FHA  
359 mutants of APLF that do not interact with XRCC4 show a reduced retention after laser micro-  
360 irradiation <sup>18</sup>. Moreover, links with DNA-PK may further properly stabilize and/or position  
361 the NHEJ ligation complex at DSBs (Figure 5h): Ku directly interacts with the XRCC4-LIG4  
362 complex <sup>7</sup> through either XRCC4 <sup>45</sup> or LIG4 <sup>6</sup> and DNA-PKcs also directly contacts XRCC4  
363 <sup>46-48</sup>. Intimate links between the DNA-PK and ligation complexes are illustrated by the  
364 requirement of an intact XLF-XRCC4-LIG4 complex to ensure optimal DNA ends synapsis  
365 <sup>49,50</sup>.

366

367 The intricate network linking end-recognition and ligation NHEJ complexes may allow  
368 compensation of partially defective individual components. Indeed, we found that separate or  
369 even combined Ku80 mutations do not completely abolish XRCC4 and XLF recruitment and  
370 do not radiosensitize cells as much as XLF complete defect that abolishes all XLF functions  
371 in NHEJ <sup>51</sup>. In that view, only a mild, if any, repair defect in human cells has been associated  
372 with X-KBM deletion <sup>27</sup> or APLF depletion <sup>9,24-26</sup>: in case of individual absence of XLF C-  
373 terminal tail or of APLF, the other intact partner would still bind to Ku80 and be able,  
374 although with a slower kinetics, to recruit the other components to achieve ligation.

375

376 Finally, our present study adds a new aspect to the DNA-PK-ligation complex interaction  
377 network that is the swing of the Ku80 vWA domain upon XLF binding (Figure 5h). From our  
378 structural and mutagenesis approaches, we propose that the outward swing of the vWA  
379 domain of Ku80 is mainly dependent on the conserved acidic residue E133 <sup>80</sup>. Although this  
380 swing does not impact the A-KBM binding site nor the DNA binding domain, it exposes a  
381 large groove between the vWA and the ring domain of Ku80 that may non-exclusively  
382 reinforce XRCC4-LIG4 and/or XLF interaction with Ku or attract yet unknown components.  
383 Interestingly, XLF deficiency impacts on the activity of the Ku partner terminal  
384 deoxynucleotidyl transferase during V(D)J recombination <sup>52</sup>. Mutagenesis studies on the  
385 conserved positions that delineated the unmasked surface of this groove will help to define  
386 precisely the role of this swing in the NHEJ reaction.

387

388 In conclusion, the present data further substantiate the emerging model that several NHEJ  
389 factors bearing a limited repertoire of KBMs recognize a limited number of KBM-binding  
390 sites on Ku<sup>16</sup>. The complex regulation of Ku sites occupancy by NHEJ factors during the  
391 repair process deserves further investigations.

392

393 **ACCESSION CODES.** Crystal structures are deposited at the pdb with the following codes  
394 Ku-DNA-pAPLF (6ERF), Ku-DNA-pXLF (6ERH) and Ku-DNA-pXLFshort (6ERG)

395

### 396 **ACKNOWLEDGMENTS**

397

398 J-B.C is supported by ARC program (SLS220120605310), ANR (ANR-12-SVSE8-012),  
399 INCA DomRep (PLBIO 2012-280), CEFIPRA grant 5203C and by the French Infrastructure  
400 for Integrated Structural Biology (FRISBI) ANR-10-INBS-05. Work in E.R's laboratory is  
401 supported by National Institutes of Health Grants CA187612, GM108119, and the American  
402 Cancer Society RSG DMC-16-241-01-DMC. P.C's team is supported by the Ligue Nationale  
403 Contre Le Cancer (Equipe labellisée 2013 and 2018) and Electricité de France (EDF, Conseil  
404 de Radioprotection). J-B.C, M.M and P.C are supported by ANR (CE12 2017 NHEJLIG4  
405 grant). We thank K.W. Caldecott (University of Sussex, Brighton, UK) for the gift of anti-  
406 APLF antibody and J.M. Stark (City of Hope, Duarte, USA) for the gift of U2OS cells  
407 engineered with the integrated distal-end-joining reporter. We would like to thank the  
408 Imaging Core Facility TRI-IPBS, in particular S. Mazerès and R. Poincloux for maintenance  
409 of the live-cell microscopy equipment and E. Näser for the maintenance of the flow-  
410 cytometers. Flow-cytometry equipment was with the financial support from ITMO Cancer  
411 Aviesan (Alliance Nationale Pour les Sciences de la Vie et de la Santé, National Alliance for  
412 Life Science and Health) within the framework of Cancer Plan. We thank J-P. de Villartay  
413 and F. Theillet for careful reading of the manuscript.

414

### 415 **AUTHORS CONTRIBUTIONS**

416 P.C. and J-B.C conceived this study. C.N, V.R, A.G, A.P, A.C, S.B expressed and purified  
417 recombinant proteins with help from P.D, E.L.C and I.B. C.N and V.R produced crystals and  
418 collected crystallographic data with help from P.L. C.N, V.R, P.L and J-B.C carried out the  
419 crystallographic analysis and interpreted the results. C.N, A.G, S.B, E.L.C and J-B.C  
420 designed, performed, and analyzed microcalorimetry and biophysical experiments. E.B-M,  
421 A.C and J-B.C designed, performed, and analyzed switchSENSE experiments. S.T and M.M

422 designed, performed, and analyzed electromobility shift assays. P.F and P.C designed and  
423 constructed vectors and cell lines for life cell imaging and radiosensitivity. P.F, C.D, N.B and  
424 P.C designed, performed and analyzed western blot analyses of Ku variants, life cell imaging,  
425 DNA repair and radiosensitivity experiments. Y.Y and E.R designed, performed and analyzed  
426 the super resolution microscopy. J.Y and R.G performed bioinformatic analyses. All the  
427 authors discussed the data. C.N, P.C and J-B.C wrote the manuscript with input from V.R,  
428 P.F, P.D, R.G, M.M, E.B-M, Y.Y and E.R.

429

### 430 **COMPETING FINANCIAL INTERESTS**

431 The authors declare no competing financial interests.

432

433

### 434 **REFERENCES**

- 435 1. Lieber, M.R. The Mechanism of Double-Strand DNA Break Repair by the  
436 Nonhomologous DNA End-Joining Pathway. *Annu Rev Biochem* **79**, 181-211 (2010).
- 437 2. Chang, H.H.Y., Pannunzio, N.R., Adachi, N. & Lieber, M.R. Non-homologous DNA  
438 end joining and alternative pathways to double-strand break repair. *Nat Rev Mol Cell*  
439 *Biol* **18**, 495-506 (2017).
- 440 3. Walker, J.R., Corpina, R.A. & Goldberg, J. Structure of the Ku heterodimer bound to  
441 DNA and its implications for double-strand break repair. *Nature* **412**, 607-14 (2001).
- 442 4. Grundy, G.J., Moulding, H.A., Caldecott, K.W. & Rulten, S.L. One ring to bring them  
443 all-The role of Ku in mammalian non-homologous end joining. *DNA Repair (Amst)*  
444 **17**, 30-8 (2014).
- 445 5. Rulten, S.L. & Grundy, G.J. Non-homologous end joining: Common interaction sites  
446 and exchange of multiple factors in the DNA repair process. *Bioessays* **39**(2017).
- 447 6. Costantini, S., Woodbine, L., Andreoli, L., Jeggo, P.A. & Vindigni, A. Interaction of  
448 the Ku heterodimer with the DNA ligase IV/Xrcc4 complex and its regulation by  
449 DNA-PK. *DNA Repair (Amst)* **6**, 712-22 (2007).
- 450 7. Nick McElhinny, S.A., Snowden, C.M., McCarville, J. & Ramsden, D.A. Ku recruits  
451 the XRCC4-ligase IV complex to DNA ends. *Mol Cell Biol* **20**, 2996-3003 (2000).
- 452 8. Yano, K.I., Morotomi-Yano, K., Lee, K.J. & Chen, D.J. Functional significance of the  
453 interaction with Ku in DNA double-strand break recognition of XLF. *FEBS Lett* **585**,  
454 841-6 (2011).
- 455 9. Grundy, G.J. et al. APLF promotes the assembly and activity of non-homologous end  
456 joining protein complexes. *Embo J* **32**, 112-25 (2013).
- 457 10. Ropars, V. et al. Structural characterization of filaments formed by human Xrcc4-  
458 Cernunnos/XLF complex involved in nonhomologous DNA end-joining. *Proc Natl*  
459 *Acad Sci U S A* **108**, 12663-8 (2011).
- 460 11. Hammel, M. et al. XRCC4 interactions with XRCC4-like factor (XLF) create an  
461 extended grooved scaffold for DNA ligation and double-strand break repair. *J Biol*  
462 *Chem* **286**, 32638-50 (2011).
- 463 12. Andres, S.N. et al. A human XRCC4-XLF complex bridges DNA. *Nucleic Acids Res*  
464 **40**, 1868-78 (2012).

- 465 13. Wu, Q. et al. Non-homologous end-joining partners in a helical dance: structural  
466 studies of XLF-XRCC4 interactions. *Biochem Soc Trans* **39**, 1387-92 (2011).
- 467 14. Reid, D.A. et al. Organization and dynamics of the nonhomologous end-joining  
468 machinery during DNA double-strand break repair. *Proc Natl Acad Sci U S A* **112**,  
469 E2575-84 (2015).
- 470 15. Ahel, I. et al. Poly(ADP-ribose)-binding zinc finger motifs in DNA repair/checkpoint  
471 proteins. *Nature* **451**, 81-5 (2008).
- 472 16. Rulten, S.L., Cortes-Ledesma, F., Guo, L., Iles, N.J. & Caldecott, K.W. APLF  
473 (C2orf13) is a novel component of poly(ADP-ribose) signaling in mammalian cells.  
474 *Mol Cell Biol* **28**, 4620-8 (2008).
- 475 17. Kanno, S. et al. A novel human AP endonuclease with conserved zinc-finger-like  
476 motifs involved in DNA strand break responses. *Embo J* **26**, 2094-103 (2007).
- 477 18. Shirodkar, P., Fenton, A.L., Meng, L. & Koch, C.A. Identification and functional  
478 characterization of a Ku-binding motif in Aprataxin Polynucleotide  
479 kinase/phosphatase-Like Factor (APLF). *J Biol Chem* **288**, 19604-13 (2013).
- 480 19. Hammel, M. et al. An Intrinsically Disordered APLF Links Ku, DNA-PKcs, and  
481 XRCC4-DNA Ligase IV in an Extended Flexible Non-homologous End Joining  
482 Complex. *J Biol Chem* **291**, 26987-27006 (2016).
- 483 20. Arnoult, N. et al. Regulation of DNA repair pathway choice in S and G2 phases by the  
484 NHEJ inhibitor CYREN. *Nature* **549**, 548-552 (2017).
- 485 21. Rossi, M.L., Ghosh, A.K. & Bohr, V.A. Roles of Werner syndrome protein in  
486 protection of genome integrity. *DNA Repair (Amst)* **9**, 331-44 (2010).
- 487 22. Tadi, S.K. et al. PAXX Is an Accessory c-NHEJ Factor that Associates with Ku70 and  
488 Has Overlapping Functions with XLF. *Cell Rep* **17**, 541-555 (2016).
- 489 23. Ochi, T. et al. DNA repair. PAXX, a paralog of XRCC4 and XLF, interacts with Ku to  
490 promote DNA double-strand break repair. *Science* **347**, 185-8 (2015).
- 491 24. Bekker-Jensen, S. et al. Human Xip1 (C2ORF13) is a novel regulator of cellular  
492 responses to DNA strand breaks. *J Biol Chem* **282**, 19638-43 (2007).
- 493 25. Iles, N., Rulten, S., El-Khamisy, S.F. & Caldecott, K.W. APLF (C2orf13) is a novel  
494 human protein involved in the cellular response to chromosomal DNA strand breaks.  
495 *Mol Cell Biol* **27**, 3793-803 (2007).
- 496 26. Macrae, C.J., McCulloch, R.D., Ylanko, J., Durocher, D. & Koch, C.A. APLF  
497 (C2orf13) facilitates nonhomologous end-joining and undergoes ATM-dependent  
498 hyperphosphorylation following ionizing radiation. *DNA Repair (Amst)* **7**, 292-302  
499 (2008).
- 500 27. Malivert, L. et al. The C-terminal domain of Cernunnos/XLF is dispensable for DNA  
501 repair in vivo. *Mol Cell Biol* **29**, 1116-22 (2009).
- 502 28. Brosey, C.A., Ahmed, Z., Lees-Miller, S.P. & Tainer, J.A. What Combined  
503 Measurements From Structures and Imaging Tell Us About DNA Damage Responses.  
504 *Methods Enzymol* **592**, 417-455 (2017).
- 505 29. Blier, P.R., Griffith, A.J., Craft, J. & Hardin, J.A. Binding of Ku protein to DNA.  
506 Measurement of affinity for ends and demonstration of binding to nicks. *J Biol Chem*  
507 **268**, 7594-601 (1993).
- 508 30. Dolinsky, T.J. et al. PDB2PQR: expanding and upgrading automated preparation of  
509 biomolecular structures for molecular simulations. *Nucleic Acids Res* **35**, W522-5  
510 (2007).
- 511 31. Cheng, Q. et al. Ku counteracts mobilization of PARP1 and MRN in chromatin  
512 damaged with DNA double-strand breaks. *Nucleic Acids Res* **39**, 9605-9619 (2011).
- 513 32. Langer, A. et al. Protein analysis by time-resolved measurements with an electro-  
514 switchable DNA chip. *Nat Commun* **4**, 2099 (2013).



- 515 33. Buck, D. et al. Cernunnos, a novel nonhomologous end-joining factor, is mutated in  
516 human immunodeficiency with microcephaly. *Cell* **124**, 287-99 (2006).
- 517 34. Yano, K.I. & Chen, D.J. Live cell imaging of XLF and XRCC4 reveals a novel view  
518 of protein assembly in the non-homologous end-joining pathway. *Cell Cycle* **7**, 1321-5  
519 (2008).
- 520 35. Hammel, M., Yu, Y., Fang, S., Lees-Miller, S.P. & Tainer, J.A. XLF Regulates  
521 Filament Architecture of the XRCC4.Ligase IV Complex. *Structure* **18**, 1431-42  
522 (2010).
- 523 36. Malivert, L. et al. Delineation of the XRCC4 interacting region in the globular head  
524 domain of cernunnos/XLF. *J Biol Chem* **285**, 26475-83 (2010).
- 525 37. Bennardo, N., Cheng, A., Huang, N. & Stark, J.M. Alternative-NHEJ Is a  
526 Mechanistically Distinct Pathway of Mammalian Chromosome Break Repair. *PLoS*  
527 *Genet* **4**, e1000110 (2008).
- 528 38. Bennardo, N., Gunn, A., Cheng, A., Hasty, P. & Stark, J.M. Limiting the persistence  
529 of a chromosome break diminishes its mutagenic potential. *PLoS Genet* **5**, e1000683  
530 (2009).
- 531 39. Gunn, A., Bennardo, N., Cheng, A. & Stark, J.M. Correct end use during end joining  
532 of multiple chromosomal double-strand breaks is influenced by repair protein RAD50,  
533 DNA-dependent protein kinase DNA-PKcs, and transcription context. *J Biol Chem*  
534 **286**, 42470-82 (2011).
- 535 40. Grundy, G.J. et al. The Ku-binding motif is a conserved module for recruitment and  
536 stimulation of non-homologous end-joining proteins. *Nat Commun* **7**, 11242 (2016).
- 537 41. Brouwer, I. et al. Sliding sleeves of XRCC4-XLF bridge DNA and connect fragments  
538 of broken DNA. *Nature* **535**, 566-9 (2016).
- 539 42. Andres, S.N., Modesti, M., Tsai, C.J., Chu, G. & Junop, M.S. Crystal Structure of  
540 Human XLF: A Twist in Nonhomologous DNA End-Joining. *Mol Cell* **28**, 1093-101  
541 (2007).
- 542 43. Lu, H., Pannicke, U., Schwarz, K. & Lieber, M.R. Length-dependent binding of  
543 human XLF to DNA and stimulation of XRCC4: DNA ligase IV activity. *J Biol Chem*  
544 **282**, 11155-62 (2007).
- 545 44. Wu, P.Y. et al. Interplay between cernunnos-XLF and NHEJ proteins at DNA ends in  
546 the cell. *J Biol Chem* **282**, 31937-43 (2007).
- 547 45. Mari, P.O. et al. Dynamic assembly of end-joining complexes requires interaction  
548 between Ku70/80 and XRCC4. *Proc Natl Acad Sci U S A* **103**, 18597-602 (2006).
- 549 46. Hsu, H.L., Yannone, S.M. & Chen, D.J. Defining interactions between DNA-PK and  
550 ligase IV/XRCC4. *DNA Repair (Amst)* **1**, 225-35 (2002).
- 551 47. Leber, R., Wise, T.W., Mizuta, R. & Meek, K. The XRCC4 gene product is a target  
552 for and interacts with the DNA-dependent protein kinase. *J Biol Chem* **273**, 1794-801  
553 (1998).
- 554 48. Wang, Y.G., Nnakwe, C., Lane, W.S., Modesti, M. & Frank, K.M. Phosphorylation  
555 and regulation of DNA ligase IV stability by DNA-dependent protein kinase. *J Biol*  
556 *Chem* **279**, 37282-90 (2004).
- 557 49. Cottarel, J. et al. A noncatalytic function of the ligation complex during  
558 nonhomologous end joining. *J Cell Biol* **200**, 173-86 (2013).
- 559 50. Graham, T.G., Walter, J.C. & Loparo, J.J. Two-Stage Synapsis of DNA Ends during  
560 Non-homologous End Joining. *Mol Cell* **61**, 850-8 (2016).
- 561 51. Menon, V. & Povirk, L.F. XLF/Cernunnos: An important but puzzling participant in  
562 the nonhomologous end joining DNA repair pathway. *DNA Repair (Amst)* **58**, 29-37  
563 (2017).

564 52. Jspeert, H. et al. XLF deficiency results in reduced N-nucleotide addition during  
565 V(D)J recombination. *Blood* **128**, 650-9 (2016).  
566  
567

568

## 569 **FIGURE LEGENDS**

570

### 571 **Figure 1. Crystal structure of the APLF KBM (A-KBM) bound to the Ku80 vWA** 572 **domain.**

573 **(a)** Positions of the A-KBM (magenta) and X-KBM (blue) motifs in APLF, XLF, WRN and  
574 CYREN. The C-terminal domain of PAXX contains a P-KBM that interacts with Ku70  
575 subunit. NTD: N-terminal domain. **(b)** Overall view of the quaternary complex  
576 Ku70/Ku80/hDNA/(APLF peptide). The A-KBM (magenta) binds at the periphery of the  
577 Ku80 (light green) vWA domain. The Ku70 subunit and hDNA are represented respectively  
578 in orange and red. The hairpin part of the DNA has been removed for clarity. **(c)** The N-  
579 terminal part of the A-KBM motif has an extended conformation whereas the C-terminal  
580 residues form a turn. **(d-e)** Zoom of the interactions made by **(d)** the hydrophobic patch and  
581 **(e)** the basic patch of the A-KBM. **(f)** The A-KBM binding site is delineated by conserved  
582 residues of Ku80 vWA domain. The binding site is represented in surface mode with amino  
583 acids colored according to their conservation rate: red (highly conserved) to white (not  
584 conserved)). The conservation rate was measured using sequences of metazoan Ku80. The  
585 orientation is the same as in **(c)**.

586

### 587 **Figure 2. Crystal structure of the XPLF KBM (X-KBM) bound to the Ku80.**

588 **(a)** Crystal structure of the quaternary complex Ku70-Ku80-DNA-(X-KBM peptide). The X-  
589 KBM (blue) binds in an internal site of the Ku80 subunit created upon an outward rotation of  
590 the vWA domain. The Ku80 vWA opening creates a large groove between the Ku80 vWA  
591 and the rest of the heterodimer. **(b)** The crystal structure of Ku70/Ku80/DNA in presence of  
592 the A-KBM is shown with the same orientation. **(c-d)** Comparison of the X-KBM binding site  
593 in presence of X-KBM **(c)** or A-KBM **(d)** peptides. The X-KBM interacts with Ku80 residues  
594 involved in Ku intramolecular contacts in the closed state of Ku observed with the A-KBM or  
595 with no peptide. The last GLFS residues of the X-KBM interact with the bottom of the groove  
596 formed in the open state. The glutamic acid presents an atypical hydrophobic environment  
597 and could be at the origin of the vWA instability. The X-KBM residues occupy the position of

598 the helix 236-241 of Ku80 in the closed conformation and some X-KBM side chains (R295<sup>X</sup>,  
599 L297<sup>X</sup> and F298<sup>X</sup>) mimic the intramolecular interactions made by Ku80 residues with the  
600 vWA domain. **(e)** Gel shift assay with XLF and Ku in presence of a 50bp DNA with a FAM  
601 in 5' and competition with pXLF containing the X-KBM motif. The arrow indicates the XLF-  
602 Ku-DNA complex. Uncropped gel image is shown in Supplementary Data Set 1. **(f)** The pair  
603 distributions  $P(r)$  obtained in solution by SAXS analysis indicates an opening of the  
604 Ku70/Ku80/DNA complex with higher  $D_{max}$  and  $R_g$  in presence of the X-KBM (blue line)  
605 compared to the Ku/DNA complex without peptide (grey line) and to the A-KBM complex  
606 (magenta line). Values deduced from SAXS analysis are reported beside the curves.

607  
608

609 **Figure 3. Life cell imaging of A-KBM and X-KBM recruitment after nuclear micro-**  
610 **irradiation.**

611 **(a)** Wild-type (WT) and mutant CFP-(A-KBM) behaviour at 0 s and 50 s after laser nuclear  
612 micro-irradiation. The white rectangle and arrows mark irradiated areas. Magnification: X40.  
613 **(b)** Dynamics of wild-type and mutant CFP-(A-KBM) at laser-induced damage sites in U2OS  
614 cells. Mean values of relative fluorescence with s.e.m. were calculated from data obtained in  
615 several individual cells:  $n=23$  and  $19$  cells for WT and mutant A-KBM, respectively).  $p$   
616 values at last time point were calculated using unpaired two-tailed  $t$ -test: WT vs W189G  
617  $p<0.0001$ . **(c)** Dynamics of wild-type and mutant CFP-(X-KBM) at laser-damaged sites as in  
618 b).  $n=27$ ,  $21$ , and  $24$  cells for WT, L297E and L297W X-KBM, respectively.  $p$  values at last  
619 time point : WT vs L297W  $p=0.8574$ ; WT vs L297E  $p=0.0021$ . **(d-e)** Dynamics of CFP-(A-  
620 KBM) (d) and (X-KBM) (e) at laser damaged sites in cells expressing wild-type or I122R  
621 mutant Ku80 as in b).  $n=20$ ,  $13$  cells for A-KBM in WT or I122R Ku80, and  $n=48$  and  $39$   
622 cells for X-KBM in WT or I122R Ku80, respectively.  $p$  values at last time point : (d) WT vs  
623 I112R  $p=0.0002$ ; (e) WT vs I112R  $p=0.5692$  **(f-g)** Dynamics of wild-type and mutant CFP-  
624 (X-KBM) at laser-damaged sites in cells expressing I112R mutant Ku80 (f) or treated with a  
625 shAPLF (g) as in b).  $n=26$ ,  $28$ , and  $21$  cells for WT, L297E and L297W X-KBM in (f), and  
626  $n=15$  cells for each of WT, L297E and L297W X-KBM in (g).  $p$  values at last time point : (f)  
627 WT vs L297W  $p=0.023$ ; WT vs L297E  $p=<0.0001$ ; (g) WT vs L297W  $p=0.0144$ ; WT vs  
628 L297E  $p=0.2654$ .

629  
630

631 **Figure 4. Biophysical and cellular analyses of XLF mutants in X-KBM.**

632 **(a)** SwitchSENSE kinetic analysis of the WT XLF interaction with Ku-DNA complexes.  
633 Solid grey lines represent raw data (from 1 to 8  $\mu\text{M}$ ; light grey to dark grey; averages of  
634 triplicates). Global fitting was performed, following a single-exponential function (solid  
635 orange lines) yielding kinetic rate constants;  $k_{\text{ON}}=4.7 \pm 1.7 \cdot 10^5 \text{ M}^{-1}\text{s}^{-1}$  and  $k_{\text{OFF}} = 9.1 \pm 0.4 \cdot 10^{-2}$   
636  $\text{s}^{-1}$  for XLF(wt). **(b)** Dynamics of wild-type and mutant CFP-XLF at laser-damaged sites in  
637 BuS cells as in Figure 3b.  $n=20$  cells for WT, L297E and L297W XLF.  $p$  values at last time  
638 point : WT vs L297W  $p=0.0093$ ; WT vs L297E  $p<0.0001$ . **(c)** Representative super-resolution  
639 images of WT, L297E mutant, and L297W mutant BuS nucleus, with XLF and Ku displayed  
640 in green and magenta, respectively (scale = 2500 nm). Right: zoomed-in areas (scale = 250  
641 nm). **(d)** Representative pair correlation function calculated from the  $8 \times 8 \mu\text{m}^2$  center square of  
642 one XLF nucleus image of WT (green), L297E (red), and L297W (blue) mutants. WT XLF  
643 shows bigger correlation radius (arrow). **(e)** Statistics of XLF foci size. Each plot represents  
644 the average XLF foci size (indicated as radius translated from the correlation radius) in one  
645 nucleus. Box's height displays the s.d. with the mean value labelled in the middle.  $n=116, 95,$   
646  $104$  nuclei for WT, L297E, and L297W. The two-sample unpaired  $t$ -test between WT and  
647 L297E is  $p=10^{-13}$  while that between WT and L297W is  $p=0.03$ . **(f)** Cell survival of BuS cells  
648 complemented with vector (EV) or WT or mutated XLF.  $y$  axis is log scale. Error bars  
649 represent s.d.,  $n=5$  to 6 independent experiments.  $p$  values were calculated using unpaired  
650 two-tailed  $t$ -test: WT vs EV  $p=1.788\text{e-}06$  ; WT vs LW  $p=0.068$  ; WT vs LE  $p=0.021$ .  
651 Significant  $p$ -values are indicated as follows:  $*p<0.05$ ,  $**p<0.01$ ,  $***p<0.001$ .

652

653 **Figure 5. Effects of Ku80 mutations in APLF and XLF binding sites.**

654 **(a-b)** Dynamics of CFP-(A-KBM) (a) and (X-KBM) (b) at laser damaged sites as in Figure  
655 3b, in U2OS cells expressing wild-type or I112R/E133M mutant Ku80.  $n=20$  and 9 cells for  
656 WT and mutant Ku80 in (a) and  $n=48$  and 11 cells for WT and mutant Ku80 in (b).  $p$  values  
657 at last time point: (a) WT vs I112R/E133M  $p=0.001$ ; (b) WT vs I112R/E133M  $p=0.0111$ . **(c-**  
658 **d)** Dynamics of CFP-XRCC4 (c) and XLF (d) at laser-damaged sites in cells expressing wild-  
659 type, I112R, E133M or I112R/E133M mutant Ku80.  $n=38, 27, 28,$  and 24 cells for WT,  
660 E133M, I112R and I112R/E133M Ku80 conditions in (c) and  $n=24, 26, 20$  and 23 cells for  
661 I112R, WT, I112R/E133M and E133M Ku80 conditions in (d).  $p$  values at last time point: (c)  
662 WT vs E133M  $p=0.532$ ; WT vs I112R  $p=0.0133$ ; WT vs I112R/E133M  $p=0.0048$ ; (d) WT vs  
663 I112R  $p=0.246$ ; WT vs E133M  $p=0.0048$ ; WT vs I112R/E133M  $p=0.0248$ . **(e)** End-joining  
664 activity in U2OS cells expressing mutated or WT Ku80. Error bars represent s.d.,  $n=4$

665 independent experiments.  $p$  values were calculated using unpaired two-tailed  $t$ -test: WT vs  
666 E133M  $p=0.0004$ ; WT vs I112R  $p=0.0052$ ; WT vs I112R/E133M  $p=0.0002$ . **(f)** Distal end-  
667 joining in U2OS cells containing mutated or WT Ku80. Error bars represent s.d.,  $n=7$   
668 independent experiments.  $p$  values were calculated using unpaired two-tailed  $t$ -test: WT vs  
669 E133M  $p=7.49 \text{ e-}05$ ; WT vs I112R  $p=2.21 \text{ e-}06$ ; WT vs I112R/E133M  $p=4.05 \text{ e-}06$ . **(g)**  
670 Survival of U2OS cells expressing WT or mutated Ku80. y axis is log scale. Error bars  
671 represent s.d.,  $n=7$  to 10 independent experiments.  $p$  values were calculated using unpaired  
672 two-tailed  $t$ -test: WT vs I112R  $p=1.47 \text{ e-}06$ ; WT vs E133M  $p=6.32 \text{ e-}05$ ; WT vs  
673 I112R/E133M  $p=2.52 \text{ e-}13$ ; I112R vs E133M  $p=0.011$ . Significant  $p$ -values are indicated as  
674 follows:  $*p < 0.05$ ,  $**p < 0.01$ ,  $***p < 0.001$ . **(h)** Model for APLF and XLF KBMs function  
675 during NHEJ.

676 **Table 1 Data collection and refinement statistics**

	Ku-pAPLF (PDB 6ERF)	Ku-pXLF (PDB 6ERH)	Ku-pXLFs (PDB 6ERG)
<b>Data collection</b>			
Space group	P <sub>1</sub>	P2 <sub>1</sub>	P2 <sub>1</sub>
Cell dimensions :			
<i>a</i> , <i>b</i> , <i>c</i> (Å)	98.9, 140.8, 150.3	111.8, 118.9, 128.2	111.7, 114.3, 127.2
α, β, γ (°)	68.6, 80.8, 81.2	90.0, 93.1, 90.0	90.0, 93.1, 90.0
Resolution (Å)	49.5-3.0 (3.28-3.0)	49.01-2.8 (3.16-2.8)	50-2.9 (3.09-2.9)
Anisotropy resolution limits (Å) <sup>§</sup>	2.9, 3.9, 3.3	2.7, 4.4, 3.4	2.8, 3.8, 3.0
Resolution limit overall (Å) <sup>§</sup>	3.15	3.11	3.0
<i>R</i> <sub>merge</sub>	0.056 (0.79)	0.164 (1.35)	0.171 (2.48)
<i>R</i> <sub>meas</sub>	0.072 (0.93)	0.182 (1.47)	0.182 (2.56)
<i>R</i> <sub>pim</sub>	0.051 (0.66)	0.094 (0.76)	0.068 (0.94)
<i>I</i> /σ ( <i>I</i> )	11.8 (1.3)	9.1 (1.5)	9.7 (1.0)
<i>CC</i> <sub>1/2</sub>	0.99 (0.60)	0.997 (0.60)	0.996 (0.51)
Completeness (spherical, %) <sup>§</sup>	66.5 (14.3)	51.3 (8.5)	72.9 (21.0)
Completeness (ellipsoidal, %) <sup>§</sup>	92.1 (70.1)	92.4 (69.6)	94.6 (68.5)
Redundancy	3.4 (3.5)	7.1 (6.9)	14.0 (14.2)
<b>Refinement</b>			
Resolution (Å)	49.47-3.01	49.38-2.8	48.76-2.9
No. reflections	90993	41644	51644
<i>R</i> <sub>work</sub> / <i>R</i> <sub>free</sub>	0.209/0.227	0.225/0.252	0.218/0.244
No. Atoms	35410	18810	18989
Protein	33225	16649	16821
DNA	2178	2118	2133
SO <sub>4</sub> <sup>2-</sup>	0	10	15
Water	7	33	20
B factors (Å <sup>2</sup> )			
Protein	112	76	94
DNA	201	109	126
R.m.s. deviations			
Bond lengths (Å)	0.008	0.007	0.008
Bond angles (°)	0.96	0.95	0.95

677 \*Values in parentheses are for highest-resolution shell. § Values from STARANISO, Global Phasing Ltd.

678

679 **Table 2 : Interactions measured by microcalorimetry between Ku70/Ku80, A-KBM, X-KBM and XLF**

680

	<b>Protein in the measurement cell</b>	<b>Ligand in the syringe</b>	<b>K<sub>d</sub> (μM)</b>	<b>ΔH (kcal.M<sup>-1</sup>)</b>	<b>Remarks</b>
1	Ku <sub>FL</sub>	pAPLF	0.033 ± 0.01	-13.3 ± 0.1	A-KBM (APLF 174-191)
2	Ku <sub>FL</sub>	DNA 18bp	0.0041 ± 0.0007	+5.3 ± 0.2	DNA <sub>18bp</sub> versus Ku
3	Ku <sub>FL</sub> +DNA-18bp	pAPLF	0.023 ± 0.002	-16.0 ± 0.4	in presence of DNA <sub>18bp</sub>
4	Ku <sub>CC</sub>	pAPLF	0.020 ± 0.002	-18.4 ± 0.7	Ku without Cter domains
5	Ku <sub>FL</sub> I112R	pAPLF	NI <sup>(a)</sup>	NI	Ku80 mutant on APLF site
6	Ku <sub>FL</sub>	pXLF	4.4 ± 0.2	-2.8 ± 0.2	X-KBM (XLF 281-299)
7	Ku <sub>FL</sub> +DNA-18bp	pXLF	2.4 ± 0.1	-8.1 ± 0.6	in presence of DNA <sub>18bp</sub>
8	Ku <sub>CC</sub>	pXLF	2.2 ± 0.9	-3.1 ± 1.2	Ku without Cter domains
9	Ku <sub>FL</sub>	XLF	1.0 ± 0.1	-9.0 <sup>(b)</sup>	XLF versus Ku
10	Ku <sub>FL</sub> +DNA-18bp	XLF	2.35 ± 0.1	-8.1 ± 0.6	XLF versus Ku/DNA <sub>18bp</sub>
11	Ku <sub>FL</sub>	pXLF(LW)	0.12 ± 0.03	-12.1 ± 2.0	X-KBM (L297W)
12	Ku <sub>FL</sub>	pXLF(LE)	NI	NI	X-KBM (L297E)
13	Ku <sub>FL</sub> +pAPLF	pXLF(LW)	NI	NI	L297W in presence of A-KBM

681

682 (a) NI means no interaction; (b) The enthalpy value was deduced from the first injection point in absence of lower plateau;

683 The thermograms and isotherms of titration of the experiments corresponding to lines 1, 6 and 11 are presented in Supplementary Figure 2e-g.

684 Interactions in lines 1, 4, 6, 8, 9, 11 were measured in triplicate, and the mean value with standard deviation is reported. Interactions in lines 2, 3,

685 4, 6, 10 were measured in duplicate, and the mean value with variation between min and max values is reported.

686

## 687 **ONLINE METHODS**

688

### 689 **DNA preparation**

690 The 500 bp linear dsDNA molecules were amplified from the 3516-4016 region of pBR322  
691 plasmid with biotinylated primers (5'-bGGATCTCAACAGCGGTAA-3' and 5'  
692 bCTTTATCCGCCTCCATCC-3'). DNA fragments were purified on a MiniQ anion exchange  
693 column with a chromatography SMART system (GE Healthcare), ethanol precipitated and  
694 resuspended in a 10 mM Tris-HCl, pH 7.5, 1 mM EDTA buffer.

695

### 696 **Oligonucleotides for ITC and EMSA**

697 CN1 34bp : CGCGCCAGCTTTCCAGCTAATAAACTAAAAAC

698 CN2 21bp : GTTTTTAGTTTATTGGGCGCG

699 CN3 18bp up : GTTATCCGAGCGTGAGAC

700 CN4 18bp down : GTCTCACGCTCGGATAAC

701 NLB48 : TAG TCG TAA GCT GAT ATG GCT GAT TAG TCG GAA GCA TCG AAC

702 GCT GAT

703 MM 50bp up : FAM-TAAATGCCAATGCTGCTGATACGTA CTCTCGGACTGATTCGGA ACTGTAACG

704 MM 50bp do : CGTTACAGTTCCGAATCAGTCCGAGTACGTATCAGCAGCATTGGCATTTA

705

### 706 **Peptides and Proteins:**

707 The synthetic peptides containing the KBM and XLM motifs were purchased from Genecust  
708 at 95% purity, and the concentrations of the stock peptide solutions were determined by  
709 amino acid composition. The oligonucleotides used for ITC and crystallization were  
710 synthesized by Sigma-Aldrich and Eurogentec.

711 The full length Ku70(1-609)/Ku80(1-732) heterodimer and a truncated version of the  
712 heterodimer deleted Ku70(1-544)/Ku80(1-551) were cloned in the Multibac vectors with a  
713 10-His tag and a TEV site on the Ku80 N-terminus<sup>53</sup>. Each plasmid was integrated in a  
714 Yellow Green Protein (YFP) containing bacmid by transformation in EMBACY *E.coli* stain  
715 (kind gift from Imre Berger, Bristol University). The resulting recombinant bacmids were  
716 used to transfect Sf21 insect cells giving the V0 virus generation. After amplification, stocks  
717 of viruses were titrated by the dilution limit method using YFP as marker for infected cells  
718 and Mac Grady table. Production was initiated in Sf21 cells culture by infection with  
719 baculovirus at MOI of  $5 \times 10^{-3}$ . Insect cells were collected 5-6 days after the infection (3-4 days



720 after the proliferation arrest). Cells were sonicated and the supernatant was incubated with  
721 Benzonase (300 units for 30 min at 4°C). The Ku heterodimer was purified on a NiNTA-  
722 Agarose affinity column (Protino, Macherey Nagel) with a 1M NaCl wash step to remove  
723 DNA excess. The eluted Ku was then bound onto an anion exchange column (Resource Q,  
724 GE Healthcare) equilibrate with buffer Q (20 mM Tris pH 8.0, 50 mM NaCl, 50 mM KCl, 10  
725 mM  $\beta$ -mercaptoethanol). Final yield of the Ku heterodimer was typically 35 mg of purified  
726 heterodimer by liter of culture. The full length 10-His tagged XLF protein was produced in  
727 insect cells with a similar protocols with a yield of 50mg of purified protein by L of culture. A  
728 truncated version of XLF(1-224) was also produced in *E. coli* .

729

### 730 **Crystallization of the Ku70/Ku80/hDNA complexes with A-KBM and X-KBM peptides**

731 The DNA used in this study is the hairpin DNA previously used by Walker et al <sup>3</sup>. It was  
732 obtained using HPLC-purified oligonucleotide of 34 and 21 nucleotides (see oligonucleotides  
733 above). The oligonucleotides were annealed and added in 1.1-fold molar excess to Ku  
734 heterodimer. The peptides containing the A-KBM motif (18mer, 174-192), the long X-KBM  
735 motif (19mer, 281-299) or the short X-KBM motif (13mer 287-299) were added respectively  
736 with a 1.1, 2 and 2 fold excess. Crystallization screenings on the Ku<sub>cc</sub>-hDNA-peptides were  
737 performed on the HTX platform (EMBL, Grenoble) with an automatic visualization at 4°C.  
738 The crystals of Ku<sub>cc</sub>-hDNA-pAPLF were reproduced and optimized in the laboratory at 20 °C  
739 using the sitting drop method by mixing 1.5 $\mu$ L of the 20 mg/mL Ku-DNA-peptides  
740 complexes with 1.5  $\mu$ L of the solution containing 13% polyethylene glycol (PEG) 3350, 150  
741 mM NaNO<sub>3</sub>, and 100 mM Bis-Tris-Propan (pH 6.5). The crystals (100x150x1000 $\mu$ m) grew in  
742 5-6 hours and were frozen in a solution of the mother liquor with 20% glycerol. The Ku<sub>cc</sub>-  
743 hDNA-pXLF and Ku<sub>cc</sub>-DNA-pXLFs complexes were crystallized at 20 °C by mixing 1.5 $\mu$ L  
744 of the 7 mg/mL complex solution with 1.5  $\mu$ L of a solution containing 18% polyethylene  
745 glycol (PEG) 3350, 150 mM Na<sub>2</sub>SO<sub>4</sub>, and 100 mM Bis-Tris-Propane (pH 8,5). The crystals  
746 (100x200x50 $\mu$ m) grew in 5-6 days and were frozen with 20% glycerol.

747

### 748 **Determination of the crystal structures**

749 Diffraction data were collected at the Proxima 1 and Proxima 2 beamlines at the *synchrotron*  
750 SOLEIL. The datasets were indexed and integrated using the XDS package <sup>54</sup>, the XDSME  
751 package (XDS Made Easier, <https://github.com/legrandp/xdsme>) and the CCP4 suite <sup>55</sup>. The  
752 crystals present a highly anisotropic diffraction (between 2.85Å and 4.25Å resolution  
753 according to the axes). The anisotropy of Kucc-hDNA-pAPLF crystals was treated with the

754 STARANISO program (<http://staraniso.globalphasing.org/>). The software performs an  
755 anisotropic cut-off of merged intensity data, a Bayesian estimation of the structure  
756 amplitudes, and applies an anisotropic correction to the data. The structure of the Ku70/80-  
757 hDNA-pAPLF was determined by molecular replacement with the program MOLREP using  
758 the structure of Ku70/80 (pdb 1JEY) without the DNA coordinates<sup>3</sup>. Four molecules of Ku  
759 were consecutively positioned. Electron density for the hDNA was clearly visible in the  
760 position previously reported by Walker. Refinement was performed using BUSTER<sup>56</sup> and  
761 PHENIX<sup>57</sup>. The models were built with Coot<sup>58</sup>. After DNA building, an electron density was  
762 visible on the Ku80 vWA near the Ku80 amino acids identified by Grundy et al by  
763 mutagenesis. The final statistics are presented below. In the final model, the following regions  
764 of Ku70/Ku80 are not visible: Ku70 1-33, 535-544 and Ku80 1-5, 170-181, 190-191, 543-  
765 551.. The quality of the model was assessed using Molprobit<sup>59</sup>.

766 The crystal structures of the Ku70/80-hDNA complexed with the long and short pXLF were  
767 solved by molecular replacement. Firstly, the coordinates of Ku70/80 and the hairpin DNA  
768 present in pdb 1JEY were used as model. We positioned two molecules in the asymmetric  
769 unit with clear electron density except on the vWA region of Ku80 (region aa 6 to 242). We  
770 then performed the molecular replacement with the same coordinates deleted of the Ku80  
771 vWA region. The electron density for the secondary structure elements of the Ku80 vWA was  
772 clearly visible though at a different position than in 1JEY. A second molecular replacement  
773 step was performed to position the vWA domain of the two Ku80 molecules in the  
774 asymmetric unit. The structure of the Ku70/80-DNA-short pXLF was solved in a similar  
775 manner. An electron density was visible in an internal position of Ku80 vWA that could be  
776 attributed to pXLF. The final statistics are presented in Table1. In the final model with the  
777 long pXLF, the following regions of Ku70/Ku80 are not visible (Ku70 1-34, 535-554 and  
778 Ku80 543-544). In the final model with the short pXLF, the following regions of Ku70/Ku80  
779 are not visible (Ku70 1-34; 535-554 and Ku80 171-194, 300-301, 542-544).

780 An additional electron density was observed in the long and short pXLF complex structures  
781 located close to the extreme N-terminus of Ku80 and to the hairpin DNA. This electron  
782 density was successfully modeled with the first missing residues of Ku80, the TEV site  
783 sequence preceding (ENLYFQG) and seven histidines from the 10-His tag. To evaluate the  
784 influence of the tag present on the N-terminus of Ku80 on the Ku-XLF interaction, we  
785 prepared a digested form of Ku with the TEV protease (Ku<sup>TEV</sup>) and measured by ITC its  
786 interaction with the X-KBM. The ITC shows that thermodynamic parameters of the Ku<sup>TEV</sup> are

787 similar to the Ku and that the tag does not significantly influence the binding of the XLF  
788 motif.

789

790

791

792 **Small-angle Xray scattering:** Several data of Ku<sub>cc</sub>-hDNA complex with a protein-DNA ratio  
793 (1:1.2) were collected alone or in presence of pXLF or pAPLF peptides. Data were collected  
794 on the SWING beamline (SOLEIL synchrotron) at a 1.8m sample-detector distance. The  
795 complexes were prepared at 1.0, 3.0 and 5mg/mL and spun for 10 minutes at 13000 rpm prior  
796 to SAXS analysis to eliminate aggregates. Volumes of 40μL of each sample and buffer (20  
797 mM Tris pH 8.0, 150mM NaCl, 5mM β-mercaptoethanol) were injected into the SAXS  
798 capillary cell and collected continuously, with a frame duration of 0.5 s and a dead time  
799 between frames of 0.5 s. Data reduction to absolute units, frame averaging and subtraction  
800 were done using FOXTROT<sup>60</sup>, a dedicated home-made application. All subsequent data  
801 processing and analysis steps were carried out with PRIMUS and other programs of the  
802 ATSAS suite<sup>61</sup>. The program GNOM<sup>62</sup> was used to compute the pair-distance distribution  
803 functions,  $P(r)$ .

804

#### 805 **Isothermal titration Calorimetry (ITC)**

806 Interactions between Ku70/80 wild-type and Ku70/80cc and the different peptides containing  
807 the KBM and XLM motifs were determined by isothermal titration calorimetry (ITC) using a  
808 VP-ITC calorimeter (Malvern). Prior to measurements, all solutions were degassed under  
809 vacuum. The reaction cell of the ITC (volume 1.8 mL) was loaded with Ku heterodimers  
810 alone or complexed with DNA or peptides for competition experiments. Proteins were  
811 extensively dialyzed against buffer I (20 mM Tris, pH 8.0, 150 mM NaCl, and 5 mM β-  
812 mercaptoethanol). Peptides and DNA were prepared at high concentrations. The syringe (290  
813 μL) was filled with the different peptides at concentration between 20 μM to 200 μM. The Ku  
814 heterodimer present in the cell was titrated by automatic injections of 6-10μL of the different  
815 peptides. Enthalpy  $\Delta H$  (in kcal.mol<sup>-1</sup>), stoichiometry of the reaction  $N$ , and association  
816 constant  $K_a$  (in M<sup>-1</sup>) were obtained by nonlinear least-squares fitting of the experimental data  
817 using the single set of independent binding sites model of the Origin software provided with  
818 the instrument. The free energy of binding ( $\Delta G$ ) and the entropy ( $\Delta S$ ) were determined using  
819 the classical thermodynamic formula,  $\Delta G = - RT \ln(K_a)$  and  $\Delta G = \Delta H - T\Delta S$ . All binding

820 experiments were performed in duplicate or triplicate at 25°C. Control experiments were  
821 performed with peptides injected into the buffer to evaluate the heat of the dilution.

822

### 823 **EMSA:**

824 Binding reactions (10 µL) were performed by incubating the annealed oligonucleotides  
825 (oligonucleotides used in this study are listed in Table Sxx and indicated in the figure  
826 legends) at a final concentration of 25 nM, with the indicated final concentrations of proteins  
827 in 75 mM KCl, 10 mM Tris (pH 7.5), 0.5 mM EDTA, 0.5 mM DTT, 0.5 mg/mL acetylated-  
828 BSA, and 5% glycerol. Reactions were incubated at room temperature for 1 hr and  
829 fractionated by 6% PAGE (29%/1% [w/v] Acrylamide:Bis-acrylamide) in 0.53 standard Tris-  
830 borate-EDTA (TBE) buffer at 80 V for 45 min to 1 hr. After electrophoresis, DNA was  
831 visualized using a ChemiDoc MP imaging system (Bio-Rad), either by direct detection of the  
832 fluorescently labeled DNA (FAM) or after staining with 0.2 mg/mL EtBr. Data were  
833 processed and quantified with the Image Lab software version 5.2.1 (Bio-Rad).

834

### 835 **switchSENSE measurements :**

836 All switchSENSE measurements were carried out on a DRX 2400 instrument, using a multi-  
837 purpose 48bp chip (both Dynamic Biosensors GmbH; Planegg, Germany). The sample and  
838 running buffer was Tris 10mM pH 7.4, 140mM NaCl, 0.05% Tween20, 50µM EDTA, 50µM  
839 EGTA. In all kinetics experiments, complementary DNA to the strand immobilized on the  
840 chip was first hybridized on the measurement electrodes. On top of the 48bp complementary  
841 to the chip, this DNA carried a 32bp-overhang hybridized with its 32bp complementary  
842 strand. Second, the Ku70/80 protein was associated at 100nM for 3min (reaching saturation)  
843 to the immobilized dsDNA. Finally, the association and dissociation of the XLF protein was  
844 measured using triplicates of each concentration, at a flow rate of 2mL/min. For each mutant,  
845 the surface was not regenerated in between concentrations – Ku does not dissociate in the  
846 timescale of the XLF measurement, and XLF fully dissociates for each concentration. The  
847 electrodes were only regenerated between the measurements of different mutants of XLF.  
848 Kinetics values were determined using Origin software.

849

### 850 **Cell lines and cell culture**

851 U2OS human osteosarcoma cells and immortalized BuS cells (derived from an XLF-  
852 deficient patient, gift from Jean-Pierre de Villartay, Institut Imagine, Paris, France) were  
853 grown in DMEM and RPMI, respectively. Media were supplemented with 10% fetal calf

854 serum (Eurobio), 125 U/ml penicillin and 125 µg/ml streptomycin. Cells were maintained at  
855 37°C in a 5% CO<sub>2</sub> humidified incubator. Cell lines were tested negative for mycoplasma by  
856 PCR. All culture media and antibiotics were from Invitrogen. When necessary (conditional  
857 expression of shRNA against Ku80), doxycyclin (Sigma-Aldrich) was added to the medium at  
858 a 4 µg/ml final concentration.

859

#### 860 **Expression vectors**

861 See Supplementary Data Set 1

862

#### 863 **Cell transfection and transduction**

864 Production of lentiviral particles in HEK-293T cells and transduction of U2OS and BuS  
865 cells were performed as previously described<sup>31</sup>. Transduced cells were used as populations  
866 without clonal selection, except when indicated otherwise.

867

#### 868 **Plasmid recircularization assay**

869 U2OS cell populations expressing wild-type or mutated Ku80 protein were seeded to near  
870 confluence in 6-well plates and incubated overnight at 37°C. Cells were then transfected with  
871 1 µg BamHI-linearized pEGFP-N1 plasmid (Clontech) and 1 µg pmCherry-C1-3NLS circular  
872 plasmid as a transfection control (gift from Dyche Mullins (Addgene #58476),<sup>63</sup>). Cells were  
873 split 24 h later, incubated at 37°C for two additional days and analyzed by flow cytometry on  
874 a Fortessa X-20 (BD Biosciences). For each cell population, the integrated GFP fluorescence  
875 signal was normalized to mCherry signal. End-joining activity was set to 100% for Ku80-WT  
876 expressing cells. Results were plotted as the mean values of four independent experiments  
877 ±s.d.

878

#### 879 **Distal End-Joining assay**

880 U2OS-EJ5 cells (a kind gift from Jeremy Stark, City of Hope, Duarte, USA) containing  
881 one integrated copy of a GFP reporter cassette which allows to measure rejoining of two  
882 tandem I-SceI cut sites separated by a ~1.8-kb insert, were modified as follows. The cells  
883 were first transduced with lentiviruses produced from pLV-tTR-KRAB and pLV3-Tet-RFP-  
884 ISceI-GRLBD to allow inducible expression and nuclear translocation of the I-SceI yeast  
885 meganuclease. The resulting cells were then transduced with lentiviruses prepared from  
886 pLVTHM2-shKu80. A positive clone enabling doxycyclin-dependent conditional knockdown  
887 of Ku80 expression and subsequent cell death was isolated. These cells were further

888 transduced with lentiviruses produced from pLV3-HA-Ku80-shR-(WT or mutants) plasmids  
889 to replace, in the presence of doxycyclin, endogenous Ku80 expression by expression of the  
890 various Ku80 constructs described in the study. To perform Distal End-Joining assay, the  
891 different U2OS-EJ5 modified cell populations were seeded onto 6-well plates and incubated  
892 at 37°C for 24 h. Dexamethasone (Sigma-Aldrich) was then added to a final concentration of  
893 250 nM. Cells were washed 24 h later, further incubated at 37°C for two to three days and  
894 analyzed by flow cytometry on a Fortessa X-20 analyzer (BD Biosciences). The fraction of  
895 GFP-positive cells was measured and normalized to 100% for Ku80-WT expressing cells.  
896 Results were plotted as the mean values of seven independent experiments  $\pm$ s.d.

897

### 898 **Ionizing irradiation and cell survival analysis**

899 Three to six thousand cells per well were seeded in 6-well plates. Plated cells were  
900 exposed 24 h later to various doses of X-ray using a Faxitron RX-650 device (130 kV, 5 mA,  
901 dose rate 0.5 Gy.min<sup>-1</sup>). Six to seven days later, cells were washed with PBS, stained 10 min  
902 with crystal violet (0.1% aqueous solution). Stained cells were extensively washed with water  
903 and plates were air dried. Staining was dissolved with 10% acetic acid solution and absorption  
904 was measured at 570 nm (Ultrospec-3000 spectrophotometer, Pharmacia Biotech). Results  
905 were plotted as mean values of 5-10 independent experiments  $\pm$ s.d. using Microsoft Excel  
906 software.

907

### 908 **Live-cell microscopy and micro-irradiation**

909 U2OS or BuS cells were seeded in 35-mm glass-bottom culture dishes (MatTek) two  
910 days prior laser irradiation. Experiments were carried out with a Zeiss LSM-710 confocal  
911 laser scanning microscope equipped with a coherent chameleon Vision-II tunable laser (690-  
912 1080 nm), a 40X/1.3 oil immersion objective and a heated environmental chamber set at 37°C  
913 in 5% CO<sub>2</sub> atmosphere. ECFP was excited using biphotonic laser at 800 nm (1.5% of  
914 maximum power). Confocal image series were recorded with a frame size of 512×512 pixels.  
915 Nuclei micro-irradiation was carried out at 800 nm at 20% of maximum power (mean max  
916 power was 3070 mW) in rectangle of 15  $\mu$ m<sup>2</sup> area during 50  $\mu$ s pixel dwell time. Before and  
917 after micro-irradiation, confocal image series of one mid z-section were recorded at 1.94 s  
918 time interval (typically 9 pre-irradiation and 40-45 post-irradiation frames). For evaluation of  
919 the recruitment kinetics, fluorescence intensities of the irradiated region were corrected for  
920 total nuclear loss of fluorescence over the time course and normalized to the pre-irradiation  
921 value. Data from micro-irradiation of individual cells obtained in several independent

922 experiments performed on different days were averaged, analyzed and displayed using  
923 PRISM software. *p* values at last time point were calculated using unpaired Student's t-test.

924

### 925 **Protein extraction and immunoblotting**

926 Sub-confluent cells from 60-mm culture dishes were harvested and washed with PBS.  
927 Pellets were resuspended in 100  $\mu$ l of lysis buffer (50 mM HEPES.KOH pH 7.5, 450 mM  
928 NaCl, 1% Tritin-X100, 1 mM EDTA, 1 mM DTT, protease-phosphatase Halt Inhibitor  
929 cocktail (Pierce, Thermo Scientific)) and processed by four freeze/thaw cycles. After  
930 centrifugation at 14000 g for 10 min, protein concentration was measured in the supernatant  
931 with the Bradford assay (Bio-Rad). Proteins from 50  $\mu$ g of cell extracts were separated in 4-  
932 15% Mini-Protean TGX precast polyacrylamide gels (Bio-Rad) and transferred to PVDF  
933 membrane (Millipore). Immunoblotting analysis was performed with the following  
934 antibodies: anti-Ku70 monoclonal antibody (clone N3H10 from NeoMarkers), anti-Ku80  
935 monoclonal antibody (clone 111 from Thermo Fisher Scientific), anti-DNA-PKcs monoclonal  
936 antibody (clone 18.2 from Abcam), anti-XLF polyclonal antibody (Bethyl Laboratories), anti-  
937 APLF polyclonal antibody (SK3595, kind gift from K.W. Caldecott, University of Sussex,  
938 Brighton, UK, see <sup>25</sup>), anti- $\beta$ -Actin monoclonal antibody (clone AC-15 from Ambion).

939

### 940 **Cell culture and preparation for super-resolution imaging**

941 BuS cells were grown in RPMI medium with 10% FBS and 100 U/mL Penicillin-  
942 Streptomycin. For super-resolution imaging, cells were cultured on glass coverslips for 24  
943 hours, followed by serum starvation for 48 hours. The cells were then released into full  
944 medium for 4 hours so that most of the cells were in G1 phase.

945 The synchronized cells were then washed twice with PBS, and permeabilized with 0.5%  
946 Triton X-100 in CSK buffer (10 mM Hepes, 200 mM Sucrose, 100 mM NaCl, and 3 mM  
947 MgCl<sub>2</sub>, pH=7.4) for 10 minutes <sup>64</sup>. Cells were then fixed with paraformaldehyde (4%) for 20  
948 mins, and blocked in block solution (2% glycine, 2% BSA, 0.2% gelatin, and 50 mM NH<sub>4</sub>Cl  
949 in PBS) overnight at 4°C.

950 Fixed cells were then immunostained with validated monoclonal antibodies: XLF was stained  
951 with XLF-antibody (3D6, NBP2-03275, NOVUS) for 1 hour at room temperature, followed  
952 by Alexa Fluor 647 conjugated goat-anti-mouse 2nd antibody staining (ab 150115, abcam) for  
953 30 minutes at room temperature. Cells were then stained with Alexa Fluor conjugated anti-  
954 Ku80 antibody (EPR3467, ab202659, abcam) for 1 hour at room temperature.

955 Cells were then mounted onto microscope glass slide, and imaged with freshly mixed imaging  
956 buffer (1 mg/mL glucose oxidase, 0.02 mg/mL catalase, 10% glucose, and 100 mM  
957 cycteanube (MEA)).

958

### 959 **Microscope and Single-Molecule Localization imaging**

960 Super-resolution imaging was performed on a custom-built optical imaging platform based on  
961 a Leica DMI 300 inverse microscope, equipped with a 488 nm (OBIS, Coherent) and a 639  
962 nm laser line (MRL-FN-639-800, CNI). Laser lines were reflected into an HCX PL APO 63X  
963 NA=1.47 OIL CORR TIRF Objective (Zeiss) by a penta-edged dichroic beam splitter  
964 (FF408/504/581/667/762-Di01-22x29), and the emitted fluorescence was further extended by  
965 a 2X lens tube (Diagnostic Instruments), filtered by single-band filters (Semrock FF01-531/40  
966 and FF01-676/37 for Alexa Fluor 488 and Alexa Fluor 647, respectively), and collected onto  
967 a sCMOS camera (Prim95B, Photometrics). A 405 nm Laser line (MDL-III-405-150, CNI)  
968 was also equipped to reactivate Alexa Fluor 647 fluorophores.

969 For super-resolution imaging, the 488 and 639 laser lines were adjusted to  $\sim 1.0$  and  $1.5$   
970 kW/cm<sup>2</sup>, and a Highly Inclined and Laminated Optical sheet (HILO) illumination mode for  
971 sample excitation. Alexa Fluor 488 and Alexa Fluor 647 were sequentially excited and their  
972 emitted fluorescence was also sequentially collected by switching the single-band filters in a  
973 filter wheel. The emitted photons were collected onto a sCMOS camera (Photometrics Prime  
974 95B), and a minimum of 2000 frames at 33 Hz were recorded for each image stack.

975

### 976 **Reconstruction from Single-Molecule Localization image to super-resolution image**

977 Each image of the image stack was first of all box-filtered with the box size of 4 times of the  
978 FWHM of a 2D gaussian point spread function (PSF). Considering the patterned noise for  
979 sCMOS camera construction, each pixel was weighted by the inverse of its noise variance  
980 during filtering. The low-pass filtered image was then extracted as the background from the  
981 raw image. The local maximums of the image were then recognized and segmented into  
982 single PSFs for 2D gaussian fit.

983 2D gaussian fit of each individual PSF was performed by GPU using the Maximum  
984 Likelihood Estimation (MLE) algorithm. Briefly, the likelihood function of each pixel was  
985 constructed by convolving the Poisson distribution of the shot noise governed by the photons  
986 emitted from fluorophores nearby, and the gaussian distribution of the readout noise of the  
987 camera itself<sup>65</sup>. The offsets, variance, and analog-to-digital conversion factor of each pixel of  
988 the camera was calibrated beforehand. The position, amplitude, sigma, and background of



989 each individual 2D gaussian PSF were addressed by maximizing the likelihood function of  
990 each pixel given its readout digital value. The fitting accuracy was estimated by Cramér-Rao  
991 lower bound (CRLB) and the accuracy of Alexa Fluor 488 and Alexa Fluor 647 in this work  
992 are ~ 17 and 13 nm, respectively.

993

#### 994 **Alignment of images of different colors**

995 Mapping of the two colors was performed using a polynomial mapping algorithm. Briefly, a  
996 spatially separated and broad-spectrum fluorescent bead (TetraSpec, ThermoFisher) slide was  
997 imaged on both color channels. Mass centers of each same bead but in different channels were  
998 recorded pairwise. The coefficient of each term in a 2nd order polynomial function was  
999 optimized by training the polynomial regression using all the recorded bead's centers. The  
1000 optimized polynomial function was then used for two color channels alignment. We note that  
1001 choosing the proper order of polynomial function for optimization depends on the number of  
1002 training beads, and higher order but not enough trainees would lead to overfit.

1003

#### 1004 **Pair-Correlation analysis**

1005 Coordinates localized within ~2.5 times of the averaged localization uncertainty, and from  
1006 consecutive frames were considered as artificial blinking and grouped as one coordinate. This  
1007 coordinate was calculated by taking the 1/var weighted average of all the coordinates within  
1008 the group, where var is the localization uncertainty of each coordinate. The grouped  
1009 coordinates were then rendered onto a pixelized image of 5 nm/pixel, and the pair-correlation  
1010 was performed on this image by series 2D Fourier and inverse Fourier transfers<sup>66</sup>.

1011 The correlation was then fitted into two correlation terms: the correlation among coordinates  
1012 within localization uncertainty, and that among the coordinates that form a cluster/molecular  
1013 assemble/filament. Considering the size of XLF 'clusters' is not randomly distributed, we  
1014 fitted the second term into a normal distribution and interpret the fitted sigma as the apparent  
1015 radius of the averaged XLF 'cluster' radius (Figure XC) across the image.

1016

#### 1017 **Data availability**

1018 Crystal structures are deposited at the pdb with the following codes Ku-DNA-pAPLF (6ERF),  
1019 Ku-DNA-pXLF (6ERH) and Ku-DNA-pXLFshort (6ERG)







1020 Source data for figures 3, 4 and 5 are available with the paper online







1021 Other data that support the findings of this study are available from the corresponding author  
1022 upon reasonable request.

1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060

## METHODS-ONLY-REFERENCES


53. Bieniossek, C., Imasaki, T., Takagi, Y. & Berger, I. MultiBac: expanding the research toolbox for multiprotein complexes. *Trends Biochem Sci* **37**, 49-57 (2012).
54. Kabsch, W. XDS. *Acta Crystallogr D Biol Crystallogr* **66**, 125-32 (2010).
55. Winn, M.D. et al. Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol Crystallogr* **67**, 235-42 (2011).
56. Smart, O.S. et al. Exploiting structure similarity in refinement: automated NCS and target-structure restraints in BUSTER. *Acta Crystallogr D Biol Crystallogr* **68**, 368-80 (2012).
57. Afonine, P.V. et al. Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**, 352-67 (2012).
58. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).
59. Chen, V.B. et al. MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).
60. Perez, J. & Nishino, Y. Advances in X-ray scattering: from solution SAXS to achievements with coherent beams. *Curr Opin Struct Biol* **22**, 670-8 (2012).
61. Petoukhov, M.V. et al. New developments in the ATSAS program package for small-angle scattering data analysis. *J Appl Crystallogr* **45**, 342-350 (2012).
62. Svergun, D.I. Small-angle X-ray and neutron scattering as a tool for structural systems biology. *Biol Chem* **391**, 737-43 (2010).
63. Belin, B.J., Lee, T. & Mullins, R.D. DNA damage induces nuclear actin filament assembly by Formin -2 and Spire-(1/2) that promotes efficient DNA repair. *Elife* **4**, e07735 (2015).
64. Britton, S., Coates, J. & Jackson, S.P. A new method for high-resolution imaging of Ku foci to decipher mechanisms of DNA double-strand break repair. *J Cell Biol* **202**, 579-95 (2013).
65. Huang, F. et al. Video-rate nanoscopy using sCMOS camera-specific single-molecule localization algorithms. *Nat Methods* **10**, 653-8 (2013).
66. Veatch, S.L. et al. Correlation functions quantify super-resolution images and estimate apparent clustering due to over-counting. *PLoS ONE* **7**, e31457 (2012).

















**a**


A-KBM WT:



A-KBM W189G:


**b****c****d****e****f****g**









# Sup Figure 1

**a**





**b**

## APLF-like KBM motifs




**c**

## XLF-like KBM motifs



**d**








## PAXX KBM motifs











## Supplementary Figure 1

**(a)** Scheme of the interactions between Ku70-Ku80 and the NHEJ factors containing an A-KBM (APLF, CYREN), an X-KBM (XLF) and both KBMs (WRN). The interaction of PAXX with Ku70 through its C-terminus is also represented. **(b-d)** Logo motif of the A-KBM, X-KBM and PAXX motifs obtained from multiple sequences alignment of these proteins as indicated (Crooks, G.E. *et al.*, WebLogo: a sequence logo generator. *Genome Res* 14, 1188-90 (2004)).

**a****b****c****d****e****f****g**




## Supplementary Figure 2

**(a)** SDS gel showing purified Ku and XLF proteins as indicated. FL: full-length; cc: C-terminal truncation. **(b)** Electron density of peptide pAPLF. **(c)** Electron density of peptide pXLF. **(d)** DNA interactions with Ku in presence of X-KBM of XLF. Ku70-Ku80-hDNA-X-KBM (colored) compared to Ku70/Ku80/hDNA (PDB 1JEY, grey). Front view of Ku70-Ku80-hDNA-X-KBM showing the major deviation of hDNA molecules because of the conformational change of Ku80. **(d-e)** ITC analyses: representative thermograms and isotherms of titration corresponding to selected measurements from Table 2, as indicated.










### Supplementary Figure 3

**(a-e) EMSA analyses:** (a, b) Gel shift assays with XLF and Ku at 20nM (a) or 200nM concentrations (b) in presence of a 50bp DNA with a FAM in 5'. (c) pXLF interaction with DNA as a control of the competition experiment in Figure 2e. (d-e) The pAPLF and PAXX Cter do not compete with the Ku-XLF complex. **(f-h) switchSENSE analyses:** (f) Scheme of the switchSENSE measurement flow: 1) Ku is bound to an 80bp nanolever with a fluorescent probe at position 48; 2) A washing step removes non-specifically bound Ku molecules; 3) XLF is then injected for real-time associations and dissociations at different concentrations followed by normalized changes in the fluorescence. (g) Binding kinetics of the Ku protein on the 80mer double-stranded DNA prior to the interaction with XLF, shown as changes in the dynamic response upwards (between 0 and 4 $\mu$ s). The dynamic response reflects the speed of the switching DNA, which decreases upon binding of the Ku analyte. The dissociation is represented only for one minute, to show that no dissociation of Ku from the DNA occurs while the XLF kinetics is measured. (h) Kinetic analyses of (LW) and (LE)XLF interactions. Solid grey lines represent raw data (from 1 to 8  $\mu$ M; light grey to dark grey; averages of triplicates). Global fitting was performed, following a single-exponential function (solid orange lines) yielding kinetic rate constants;  $k_{ON} = 4.9 \pm 0.5 \cdot 10^4 \text{ M}^{-1}\text{s}^{-1}$  and  $k_{OFF} = 4.8 \pm 0.5 \cdot 10^{-2} \text{ s}^{-1}$  for XLF(LE) and  $k_{ON} = 1.9 \pm 1.1 \cdot 10^5 \text{ M}^{-1}\text{s}^{-1}$  and  $k_{OFF} = 8.4 \pm 0.6 \cdot 10^{-2} \text{ s}^{-1}$  for XLF(LW).



#### Supplementary Figure 4


**(a)** Variations among the sequence motifs observed for the C-terminal tail of XLF and for the seven positions of the Ku80 XBM pocket in various clades of the eukaryotic phylogenetic tree. 10 clades are represented summarizing the properties of 60 Mammalia, 36 Sauria, 31 Neopterygii, 62 Ecdysozoa, 9 Lophotrochozoa, 5 Cnidaria, 150 Fungi, 5 Ciliophora and 4 Dictyostellida sequences of XLF and Ku80. For each clade, web logos of the last 25 C-terminal amino-acids of XLF sequences are represented on top and the web logo of the X-KBM site positions is squared in green. A red star indicates the clades in which the position of Ku80 E133 was conserved as an acidic residue whereas a grey star points out that the acidic character of the residue was not maintained and was generally switched to a hydrophobic residue as observed in Ku70. **(b)** Superimposition of human Ku70 and Ku80 structures (PDB: 1JEQ) colored in orange and green, respectively, and focused on the region surrounding Ku80 E133 position in the X-KBM site. The red star points out the location of Ku80E133. Residues labelled and shown as sticks are the spatial neighbours of Ku80 E133. The side chain of Ku80 E133 is buried in the hydrophobic core of Ku80 and is not involved in any hydrogen bond or salt-bridge interaction resulting in a predicted pKa above 9.1 in the absence of XLF. **(c)** Molecular modelling of the interaction between Ku80 and the C-terminus of WRN containing an A-KBM in tandem with a X-KBM. The position of WRN motifs were deduced from the crystal structures presented here with APLF and XLF KBMs. The orientation of the KBMs and the size of the linker between WRN KBMs are compatible with a simultaneous binding of both WRN motifs to Ku80.





## Supplementary Figure 5

**(a-d)** Western blot of U2OS cell extracts. **(a)** Whole cell extracts of U2OS shKu80 (U2OS-Ku80<sup>KD</sup>) cells treated with doxycyclin for the indicated time were denatured and separated on 10% SDS-PAGE gel followed by electrotransfer on membrane. The membranes were blotted with the antibodies as indicated. **(b)** Whole cell extracts of U2OS shAPLF cells expressing WT or mutant CFP-X-KBM as indicated were processed as in (a). **(c)** Whole cell extracts of U2OS-Ku80<sup>KD</sup> cells treated with doxycyclin for 7 days and expressing WT or mutant HA-Ku80 as indicated were processed as in (a). **(d)** Whole cell extracts of U2OS-Ku80<sup>KD</sup> cells treated with doxycyclin for 7 days and expressing WT or mutant HA-Ku80 as indicated were processed as in (a). Uncropped blot images are shown in Supplementary Data Set 1. **(e)** Wild-type (WT) or mutant CFP-(A-KBM) and mCherry-Ku70 simultaneous behaviour after 800 nm pulsed-laser nuclear micro-irradiation assessed in U2OS cells by live cell-imaging at 0 s and 50 s post-irradiation. The white rectangle and arrows mark irradiated areas. **(f-h)** Dynamics of wild-type (WT) and mutant CFP-tagged full-length XLF at laser-damaged sites in BuS cells. Images were obtained at 1.94 s intervals and fluorescence intensities at the damage sites and in undamaged area were quantified. Mean values of the relative fluorescence with SEM were calculated from 20 independent measurements for each of WT and G296W XLF in (f), from 45, 40 and 20 independent measurements for each of WT, S299A and S299E XLF in (g) and from 45, 36, 20 and 20 independent measurements for each of WT, F298G, F298G/S299A, and F298G/S299E XLF in (h), respectively. *p* values at last time point: (f) WT vs G296W *p*<0.0001; (g) WT vs S>A *p*=0.2785; WT vs S>E *p*<0.0001. (h) WT vs F>G *p*<0.0001; WT vs FS>GA *p*<0.0001; WT vs FS>GE *p*<0.0001. **(i)** Dynamics of L115D and L115D/L233E CFP-tagged full-length mutant XLF at laser-damaged sites in BuS cells as in (f). Mean values of the relative fluorescence with SEM were calculated from 11 independent measurements for each of L115D±shAPLF and L115D/L233E XLF conditions. *p* values at last time point: L115D vs L115D±shAPLF *p*=0.6113; L115D vs L115D/L233E *p*=0.0002.



### Supplementary Figure 6

**(a-b)** Dynamics of CFP-(A-KBM) (a) and (X-KBM) (b) at laser damaged sites in U2OS cells expressing wild-type (WT), E133M or Q162E mutant Ku80 as in Figure 3 b). Mean values of the relative fluorescence with SEM were calculated from 20, 23 and 22 independent measurements for A-KBM with WT, E133M or Q162E mutant Ku80 in (a) and from 48, 29 and 29 independent measurements for X-KBM with WT, E133M or Q162E mutant Ku80 in (b), respectively.  $p$  values at last time point: (a) WT vs E133M  $p=0.831$ ; WT vs Q162E  $p=0.59519$ ; (b) WT vs E133M  $p=0.0003$ ; WT vs Q162E  $p=0.0111$ . **(c-d)** Dynamics of wild-type (WT) and

mutant CFP-Ku80 at laser damaged sites in U2OS cells. Mean values of the relative fluorescence with SEM were calculated from 25, 24, 20 and 15 independent measurements for WT, I112R, E133M and Q162E mutant Ku80 in (c) and from 25 and 26 independent measurements for WT or I112R/E133M mutant Ku80 in (d), respectively.  $p$  values at last time point: (c) WT vs Q162E  $p=0.9252$ ; WT vs I112R  $p=0.2734$ ; WT vs E133M  $p=0.1101$ . (d) WT vs I112R-E133M  $p=0.5362$ . **(e-f)** Analysis of XLF foci in U2OS cells by super-resolution. (e) Statistics of XLF foci size: each plot represents the average XLF foci size (indicated as radius translated from the correlation radius) in one nucleus. Box's height displays the standard deviation with the mean value labelled in the middle. 87, 110, 64, and 79 nuclei were taken in account for WT, E133M, I112R, and E133M-I112R double-mutant, respectively. The  $p$ -values were obtained by the t-test; (f) Statistics of the Cross-Pair-Correlation between Ku and XLF: Ku and XLF were stained with antibodies labelled by different fluorophores (Alexa488 conjugated rabbit anti-Ku80, abcam198586, Alexa647 conjugated goat anti-mouse secondary + Mouse anti-XLF, NBP2-03275), and dual-colour super-resolution imaging was performed to examine the cross-correlation between Ku and XLF foci within each nucleus. Each plot represents the cross-correlation amplitude calculated across one nucleus. Box's height displays the standard deviation with the mean value labelled in the middle. 83, 107, 57, and 72 nuclei were taken in account for WT, E133M, I112R, and E133M-I112R mutants respectively. The  $p$ -value were obtained by the t-test.

## Expression vectors

All lentiviral vectors derived from pLVTHM (Addgene plasmid #12247) and pLV-tTR-KRAB-Red (Addgene plasmid #12250) plasmids. Both were gifts from Didier Trono <sup>1</sup>.

The pLVTHM2 vector was obtained by digesting pLVTHM with PmeI/SpeI and inserting the PBXS linker (i.e. preannealed PBXS-F/PBXS-R pair of oligonucleotides) in order to remove the GFP coding sequence. To generate lentiviral vectors for conditional expression of shRNA, pLVTHM2 was digested with MluI/ClaI and the following pairs of preannealed oligonucleotides were inserted by ligation: shKu80-F/shKu80-R (target sequence designed by Denis Biard, CEA-DSV, France, personal communication) or shAPLF-F/shAPLF-R (target sequence from <sup>2</sup>) to knockdown the expression of Ku80 or APLF, respectively.

The pLV-tTR-KRAB vector was obtained from pLV-tTR-KRAB-Red by replacing the DsRed coding sequence by the XBES linker at XmaI/SpeI restriction sites.

The pLV-Red vector derived from pLV-tTR-KRAB-Red by replacing the tTR-KRAB coding sequence by the PEKBBMX linker at PmeI/XmaI restriction sites.

The pLV3 vector derived from pLV-tTR-KRAB-Red through the following modifications: first, the NsiI/Kpn2I fragment was replaced by the nPNk linker. The tTR-KRAB coding sequence was then removed by PmeI/XmaI digestion and replaced by the PEKBBMX linker. Finally, the IRES-DsRed fragment was removed by PmeI/SpeI digestion and replaced by the PKXMBBES linker.

To express ECFP-tagged A-KBM (APLF P177 to E193) or X-KBM (XLF S287 to S299) motifs, the A-KBM linker or X-KBM linker, respectively, was inserted into the Acc65I/BamHI restriction sites of the pECFP-C1 plasmid (Clontech) and the AgeI/BamHI fragment (ECFP-A-KBM or ECFP-X-KBM) was subcloned into the Kpn2I/BamHI restriction sites of pLV3. The ECFP-tagged A-KBM-W189G, X-KBM-L297W and X-KBM-L297E expressing vectors were obtained by amplifying by PCR the corresponding cDNAs using the pECFP-A-KBM or pECFP-X-KBM plasmid as a template, as well as CMV-F as forward primer and Bam-A-KBM-W189G-R, XLF-L297W-Bam-R or XLF-L297E-Bam-R as reverse primer, respectively. The PCR fragments were digested with AgeI/BamHI and inserted into Kpn2I/BamHI restriction sites of pLV3.

To express full-length ECFP-tagged XLF protein, human XLF cDNA (a gift from Jean-Pierre de Villartay, Institut Imagine, Paris, France) was amplified by PCR using the XLF-Hind-F and XLF-Bam-R primers. The resulting PCR fragment was digested with HindIII/BamHI and inserted into pECFP-C1. The ECFP-XLF coding fragment was then excised with AgeI/BamHI and inserted into pLV3 at Kpn2I/BamHI restriction sites. ECFP-XLF-L297W, -L297E, -F298G, -S299A, -S299E, -F298G-S299A, and -F298G-S299E mutants were expressed by PCR amplification of the corresponding XLF coding sequences using the pECFP-XLF vector as a template, Kpn2-MCS-F as forward primer, and XLF-L297W-Bam-R, XLF-L297E-Bam-R, XLF-F298G-Bam-R, XLF-S299A-Bam-R, XLF-S299E-Bam-R, XLF-FS298GA-Bam-R, or XLF-FS298GE-Bam-R, as reverse primer, respectively. The resulting fragments were digested with Kpn2I/BamHI and inserted into pLV3-ECFP-XLF to replace the XLF wild-type coding sequence. The XLF-L115D coding sequence was obtained by overlapping PCR mutagenesis on pLV3-ECFP-XLF template using ECFP-Cter-F and XLF-Bam-R oligos as outer primers, and XLF-L115D-F and XLF-L115D-R as mutated inner primers. The PCR product (XLF-L115D) was then digested with Kpn2I/BamHI and inserted into pLV3-ECFP-XLF to replace the XLF-WT coding sequence. The XLF-L115D-L297E double mutant coding sequence was obtained as above, except that the XLF-Bam-R outer primer was replaced by XLF-L297E-Bam-R.

Lentiviral vectors expressing untagged full-length XLF proteins (WT, L297W or L297E) were obtained by subcloning Kpn2I/BamHI fragments from the respective pLV3-ECFP-XLF into the pLV-Red vector.

The ECFP-XRCC4 expressing vector was obtained by excision of a Kpn2I/BamHI fragment containing the XRCC4 coding sequence from the pEGFP-C1-FLAG-XRCC4 plasmid (gift from Steve Jackson (Addgene #46959), <sup>3</sup>). The resulting fragment was inserted into pLV3-ECFP-XLF to replace the XLF coding sequence.

To obtain shRNA-resistant human Ku80 expression vector, HA-Ku80 coding sequence was amplified by overlapping PCR from pICE-Puro-HA-Ku80 (a kind gift from Sébastien Britton, IPBS, Toulouse, France) with Kpn2-HA-F and pICE-Xba-R as outer primers, and Ku80-shRes-F and Ku80-shRes-R as inner primers that introduce silent mutations in the shRNA target sequence. The resulting fragment was then inserted into pLV3 after digestion with Kpn2I/MluI. The Ku80-L112R expressing pLV3 vector was constructed by overlapping PCR from pLV3-HA-Ku80-shR using pLV-F and pLV-R oligonucleotides as outer primers and Ku80-L112R-F and Ku80-L112R-R as mutated inner primers. The PCR product was then digested with Kpn2I/MluI and inserted into pLV3. Other Ku80 single mutant constructs

(E133M and Q162E) were obtained similarly by using the corresponding pairs of inner primers (Ku80-E133M-F/Ku80-E133M-R and Ku80-Q162E-F/Ku80-Q162E-R, respectively). The pLV3-HA-Ku80-shR-L112R-E133M double mutant expressing vector was obtained as above for the E133M single mutant construct, except that pLV3-HA-Ku80-shR-I112R was used as a template for PCR reactions.

Expression vectors for ECFP-tagged WT or mutants Ku80 were obtained by amplifying the ECFP coding sequence by PCR from the pECFP-C1 plasmid with the Pme-Koz-ECFP-F and pme-CFP-80-R primers. The PCR fragment was then inserted at the PmeI restriction site in the various pLV3-HA-Ku80-shR plasmids by use of the Hot-Fusion strategy <sup>4</sup>.

To generate pLV3-mCherry-FLAG-Ku70 vector allowing expression of an mCherry-tagged Ku70 protein, the FLAG-Ku70 coding sequence was amplified by PCR using Kpn2-FLAG-F and Mlu-Ku70-R primers, digested with Kpn2I/MluI and cloned into pLV3. The mCherry coding sequence was then inserted at the PmeI restriction site by Hot-Fusion cloning <sup>5</sup> following PCR amplification using Pme-Koz-ECFP-F and Kpn2-pme-mCh-R primers and the pmCherry-C1-3NLS plasmid as a template (gift from Dyche Mullins (Addgene #58476), <sup>6</sup>).

The pLV3-Tet-RFP-ISceI-GRLBD lentiviral vector for conditional expression of I-SceI was prepared as follows : pLV3 was first modified by inserting at the PacI restriction site the Tet-Pac-F/Tet-Pac-R pre-annealed linker which contains two tetracyclin operator DNA elements. The resulting pLV3-Tet plasmid was then digested by Kpn2I/BamHI to receive the AgeI/BamHI fragment from the pISceI-GR-RFP plasmid (gift from Tom Misteli (Addgene #17654), <sup>7</sup>) which contains the coding sequence of DsRed-ISceI-GRLBD.

All oligonucleotides were purchased from Eurofins Genomics (Ebersberg, Germany). Restriction and modifying enzymes (Phusion and T4 DNA Ligase) were from ThermoFisher Scientific (Illkirch, France). All constructs were checked by sequencing (Eurofins Genomics).

**Oligonucleotides used as linkers** (alphabetical order; sequences 5' to 3')

<b>A-KBM-F</b>	GTACC CCA ATC CTT GCC GAG AGG AAA AGA ATC CTT CCA ACT TGG ATG TTA GCA GAA TAG
<b>A-KBM-R</b>	GATCCTA TTC TGC TAA CAT CCA AGT TGG AAG GAT TCT TTT CCT CTC GGC AAG GAT TGG G
<b>nPNk-F</b>	CTCCATCGATCGCCATGGTGA
<b>nPNk-R</b>	CCGGTCACCATGGCGATCGATGGAGTGCA

<b>PBXS-F</b>	AAAC CGTACG GATATC T CCCGGG TC A
<b>PBXS-R</b>	CTAGT GA CCCGGG A GATATC CGTACG GTTT
<b>PEKBBMX-F</b>	AAACTACGGGATC GAATTC CTCGCT TCCGGA CTTCGT GGATCC ACTCTC CGTACG ACTGCT ACGCGT ACTTCAC
<b>PEKBBMX-R</b>	CCGGTGAAGT ACGCGT AGCAGT CGTACG GAGAGT GGATCC ACGAAG TCCGGA AGCGAG GAATTC GATCCCGTAGTTT
<b>PKXMBBES-F</b>	AAACTACGG GATC TCCGGA CACCTT CCCGGG TCACTC ACGCGT CTCATT GGATCC CGTACG GAATTC A
<b>PKXMBBES-R</b>	CTAGT GAATTC CGTACG GGATCC AATGAG ACGCGT GAGTGA CCCGGG AAGGTG TCCGGA GATC CCGTAGTTT
<b>shAPLF-F</b>	CGCGTCCCC GAA GAA ATC TGC AAA GAT A TTCAAGAGA T ATC TTT GCA GAT TTC TTC TTTTTGGAAAT
<b>shAPLF-R</b>	CGATTTCCAAAAA GAA GAA ATC TGC AAA GAT A TCTCTTGAA T ATC TTT GCA GAT TTC TTC GGGGA
<b>shKu80-F</b>	CGCGTCCCC G AAC AAG GAT GAG ATT GCT TTCAAGAGA AGC AAT CTC ATC CTT GTT C TTTTTGGAAAT
<b>shKu80-R</b>	CGATTTCCAAAAA G AAC AAG GAT GAG ATT GCT TCTCTTGAA AGC AAT CTC ATC CTT GTT C GGGGA
<b>TetO-Pac-F</b>	TCCCTATCAGTGATAGAGATCTCCCTATCAGTGATAGAGAAT
<b>TetO-Pac-R</b>	TCTCTATCACTGATAGGGAGATCTCTATCACTGATAGGGAAT
<b>XBES-F</b>	CCGGG GGATCC CTCGAG GAATTC A
<b>XBES-R</b>	CTAGT GAATTC CTCGAG GGATCC C
<b>X-KBM-F</b>	GTACC TCA AAG GTC AAG AGG AAG AAG CCA AGG GGT CTC TTC AGT TAG
<b>X-KBM-R</b>	GATCCTA ACT GAA GAG ACC CCT TGG CTT CTT CCT CTT GAC CTT TGA G

### Oligonucleotides used as PCR primers (alphabetical order; sequences 5' to 3')

<b>Bam-A-KBM-W189G-R</b>	CGTACGGGATC CTA TTC TGC TAA CAT CCC AGT TGG AAG GAT TCT TTT CCT C
<b>CMV-F</b>	GTAGGCGTGTACGGTGGGAGG
<b>ECFP-Cter-F</b>	C ATG GTC CTG CTG GAG TTC GTG
<b>Kpn2-HA-F</b>	CTCTGC TCCGGA GCCACC ATG TAC CCC TAC GAT GTG C
<b>Kpn2-FLAG-F</b>	CTCTCGTCCGGAGCCGCACC ATG GAC TAC AAG GAT G
<b>Kpn2-MCS-AKF-F</b>	CTCGCTTCCGGACTCAGATCTCGAGCTC
<b>Kpn2-pme-mCh-R</b>	GGTGCGGCTCCGGAGATCCCGTAGTTTGGACTTGTACAGCTCGTCCATGCCG
<b>Ku80-E133M-F</b>	GAG GCA TAT TAT GAT ATT CAC TGA CCT CAG CAG CCG ATT C
<b>Ku80-E133M-R</b>	GGT CAG TGA ATA TCA TAA TAT GCC TCT TCT CAA ACT TCT TTC CTA TTG
<b>Ku80-I112R-F</b>	C TTC CTG GAT GCA CTA AGA GTG AGC ATG GAT GTG ATT CAA C
<b>Ku80-I112R-R</b>	G AAT CAC ATC CAT GCT CAC TCT TAG TGC ATC CAG GAA GTC
<b>Ku80-Q162E-F</b>	CAT CTC CCT GGA ATT CTT CTT GCC TTT CTC ACT TGG C

<b>Ku80-Q162E-R</b>	GGC AAG AAG AAT TCC AGG GAG ATG TCA CAT TTC TTC AAG C
<b>Ku80-shRes-F</b>	GCT GAA AAT AAA GAC GAA ATC GCC TTA GTC CTG TTT GGT ACA GAT GGC
<b>Ku80-shRes-R</b>	GAC TAA GGC GAT TTC GTC TTT ATT TTC AGC AAA CAC CTG TCG CTG TAC
<b>Mlu-Ku70-R</b>	CTCTGCACGCG TCA GTC CTG GAA GTG CTT GGT GAG GGC
<b>pICE-Xba-R</b>	CAGCGGGTTTA TCTAGA CTGCAG ACGCGT GC
<b>pLV-F</b>	CCGATCACGAGACTAGCCTCGAGG
<b>pLV-R</b>	CCAGTCAATCTTTCACAAATTTTGTAAATCCAGAGG
<b>pme-CFP-80-R</b>	CATGGTGGCTCCGGAGATCCCGTAGTTTGACTTGTACAGCTCGTCCATGCCG
<b>Pme-Koz-CFP-F</b>	CGATCACGAGACTAGCCTCGAGGTTTAAACGCCGCCACCATGGTGAGCAAGG GC
<b>XLF-Bam-R</b>	CTCTC GGATC CTA ACT GAA GAG ACC CCT TGG CTT CTT CCT CTT GAC C
<b>XLF-F298G-Bam-R</b>	CTCTC GGATC CTA ACT GCC GAG ACC CCT TGG CTT CTT CCT CTT GAC C
<b>XLF-FS298GA-Bam-R</b>	CTCTCGGATC CTA AGC GCC GAG ACC CCT TGG CTT CTT CCT CTT GAC C
<b>XLF-FS298GE-Bam-R</b>	CTCTCGGATC CTA CTC GCC GAG ACC CCT TGG CTT CTT CCT CTT GAC C
<b>XLF-Hind-F</b>	CTCTCAAGCTTCCGCCACC ATG GAA GAA CTG GAG CAA GGC CTG
<b>XLF-L115D-F</b>	G CGA AGT GAG CTC TCT GGC GAC CCC TTC TAT TGG AAT TTC C
<b>XLF-L115D-R</b>	G GAA ATT CCA ATA GAA GGG GTC GCC AGA GAG CTC ACT TCG C
<b>XLF-L297E-Bam-R</b>	CTCTC GGATC CTA ACT GAA CTC ACC CCT TGG CTT CTT CCT CTT GAC C
<b>XLF-L297W-Bam-R</b>	CTCTC GGATC CTA ACT GAA CCA ACC CCT TGG CTT CTT CCT CTT GAC C
<b>XLF-S299A-Bam-R</b>	CTCTCGGATC CTA AGC GAA GAG ACC CCT TGG CTT CTT CCT CTT GAC C
<b>XLF-S299E-Bam-R</b>	CTCTCGGATC CTA CTC GAA GAG ACC CCT TGG CTT CTT CCT CTT GAC C


## References

1. Wiznerowicz, M. & Trono, D. Conditional suppression of cellular genes: lentivirus vector-mediated drug-inducible RNA interference. *J Virol* **77**, 8957-61 (2003).
2. Iles, N., Rulten, S., El-Khamisy, S.F. & Caldecott, K.W. APLF (C2orf13) is a novel human protein involved in the cellular response to chromosomal DNA strand breaks. *Mol Cell Biol* **27**, 3793-803 (2007).
3. Britton, S., Coates, J. & Jackson, S.P. A new method for high-resolution imaging of Ku foci to decipher mechanisms of DNA double-strand break repair. *J Cell Biol* **202**, 579-95 (2013).
4. Fu, C., Donovan, W.P., Shikapwashya-Hasser, O., Ye, X. & Cole, R.H. Hot Fusion: an efficient method to clone multiple DNA fragments as well as inverted repeats without ligase. *PLoS ONE* **9**, e115318 (2014).



5. Frit, P., Barboule, N., Yuan, Y., Gomez, D. & Calsou, P. Alternative end-joining pathway(s): Bricolage at DNA breaks. *DNA Repair (Amst)* **17**, 81-97 (2014).
6. Belin, B.J., Lee, T. & Mullins, R.D. DNA damage induces nuclear actin filament assembly by Formin -2 and Spire-(1/2) that promotes efficient DNA repair. [corrected]. *Elife* **4**, e07735 (2015).
7. Soutoglou, E. et al. Positional stability of single double-strand breaks in mammalian cells. *Nat Cell Biol* **9**, 675-82. (2007).

Uncropped image related to Figure 2e



Uncropped scans related to Suppl. Fig. 5a-d

