

LETTER • OPEN ACCESS

## Urban versus rural health impacts attributable to PM<sub>2.5</sub> and O<sub>3</sub> in northern India

To cite this article: Alexandra Karambelas *et al* 2018 *Environ. Res. Lett.* **13** 064010

View the [article online](#) for updates and enhancements.

# Environmental Research Letters



## LETTER

# Urban versus rural health impacts attributable to PM<sub>2.5</sub> and O<sub>3</sub> in northern India

### OPEN ACCESS

#### RECEIVED

8 March 2018

#### REVISED

30 April 2018

#### ACCEPTED FOR PUBLICATION

3 May 2018

#### PUBLISHED

25 May 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Alexandra Karambelas<sup>1,7</sup>, Tracey Holloway<sup>2</sup>, Patrick L Kinney<sup>3</sup>, Arlene M Fiore<sup>4</sup>, Ruth DeFries<sup>5</sup>, Gregor Kieseewetter<sup>6</sup> and Chris Heyes<sup>6</sup>

<sup>1</sup> The Earth Institute, Columbia University, New York, NY, United States of America

<sup>2</sup> Nelson Institute Center for Sustainability and the Global Environment, University of Wisconsin—Madison, Madison, WI, United States of America

<sup>3</sup> School of Public Health, Boston University, Boston, MA, United States of America

<sup>4</sup> Department of Earth and Environmental Sciences and Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY, United States of America

<sup>5</sup> Department of Ecology, Evolution, and Environmental Biology, Columbia University, New York, NY, United States of America

<sup>6</sup> Air Quality and Greenhouse Gases, International Institute for Applied Systems Analysis, Laxenburg, Austria

<sup>7</sup> Author to whom any correspondence should be addressed.

E-mail: [ak4040@columbia.edu](mailto:ak4040@columbia.edu)

**Keywords:** emissions, air quality, India, health, rural, urban

Supplementary material for this article is available [online](#)

## Abstract

Ambient air pollution in India contributes to negative health impacts and early death. Ground-based monitors often used to quantify health impacts are located in urban regions, yet approximately 70% of India's population lives in rural communities. We simulate high-resolution concentrations of fine particulate matter (PM) and ozone from the regional Community Multi-scale Air Quality model over northern India, including updated estimates of anthropogenic emissions for transportation, residential combustion and location-based industrial and electrical generating emissions in a new anthropogenic emissions inventory. These simulations inform seasonal air quality and health impacts due to anthropogenic emissions, contrasting urban versus rural regions. For our northern India domain, we estimate 463 200 (95% confidence interval: 444 600–482 600) adults die prematurely each year from PM<sub>2.5</sub> and that 37 800 (28 500–48 100) adults die prematurely each year from O<sub>3</sub>. This translates to 5.8 deaths per 10 000 attributable to air pollution out of an annual rate of 72 deaths per 10 000 (8.1% of deaths) using 2010 estimates. We estimate that the majority of premature deaths resulting from PM<sub>2.5</sub> and O<sub>3</sub> are in rural (383 600) as opposed to urban (117 200) regions, where we define urban as cities and towns with populations of at least 100 000 people. These findings indicate the need for rural monitoring and appropriate health studies to understand and mitigate the effects of ambient air pollution on this population in addition to supporting model evaluation.

## 1. Introduction

Worldwide, more than 7 million adult premature deaths each year are attributable to ambient air pollution (Lim *et al* 2012, Forouzanfar *et al* 2015, 2016). An additional estimated 600 000 children are annually subject to premature mortality resulting from long-term ambient air pollution exposure (UNICEF 2016). Human health impacts from air pollution disproportionately affect populations in countries undergoing rapid industrialization, population growth, urbanization, and motorization. In India, annual premature

mortality estimates due to ambient air pollution are estimated between 580 000 (Ghude *et al* 2016) to more than one million people (Health Effects Institute 2017). Premature deaths due to air pollution can bring significant monetary losses to the Indian economy (Ghude *et al* 2016). However, many health analyses fail to reflect differences across urban versus rural populations, important considering India's population is largely rural.

The Global Burden of Disease (GBD) studies seek to quantify health effects (mortality and morbidity) due to risk factors, including environmental risks such

as ambient outdoor air pollution (Brauer *et al* 2012, Lim *et al* 2012, Lelieveld *et al* 2015, Forouzanfar *et al* 2016). Estimates to ambient air pollution risk-attributable mortality and morbidity are calculated using baseline mortalities for specific diseases and dose-response relationships derived from cohort studies. However, evaluating ambient air pollution impacts requires accurate quantitative estimates of concentrations at the surface, where people are subject to harmful exposure effects.

Pollutant exposures are estimated using a combination of atmospheric chemistry models, satellite observations, and *in situ* measurements. Traditionally, surface measurements reflect the best recording of ambient concentrations to determine exposure-response relationships. Yet, in India, surface monitoring is largely limited to urban regions as opposed to rural villages, and available observations supported by the Central Pollution Control Board (CPCB) in India do not have an extensive history (Central Pollution Control Board 2008, ENVIS Centre on Control of Pollution 2016). We can fill in some of these gaps using long-term satellite tropospheric column averages, which show spatial variations in increased pollution over time e.g. Duncan *et al* (2015) that may have slowed in recent years (Hilboll *et al* 2017). Previous health impact assessments in India have used satellite observations to infer surface concentrations and health impacts of PM<sub>2.5</sub> at the district level (Chowdhury and Dey 2016). Advanced chemical transport models are increasingly used to focus on pollution across India regionally, from evaluating and developing NO<sub>x</sub> emissions inventories (Ghude *et al* 2013, Jena *et al* 2015) to testing modeled O<sub>3</sub> sensitivities in the region (Surendran *et al* 2015, Sharma *et al* 2016), in addition to estimating premature mortality due to air pollution (Ghude *et al* 2016). However, some models may be too coarse to resolve large topographic gradients that affect modeled airflow and thus pollutant distributions, something particularly important in northern India due to the Himalayas. Yet the current capacity of atmospheric chemistry models to produce surface concentrations remains largely limited by our understanding of emissions inventories.

Emissions remain challenging to quantify as a result of ongoing industrialization, motorization, and urbanization in India. Many different emissions inventories for use in atmospheric chemistry models have been used for understanding air quality impacts in India (Zhang *et al* 2009, Amann *et al* 2011, EC-JRC/PBL 2011, Kurokawa *et al* 2013, Guttikunda and Jawahar 2014, Pandey *et al* 2014, Sadavarte and Venkataraman 2014), however emissions estimates remain highly uncertain in magnitude—NO<sub>x</sub> and SO<sub>2</sub> range between 5–6.75 Tg and 7–9 Tg per year respectively—and location (Saikawa *et al* 2017). In addition estimating rural emissions by extrapolation from urban data can be problematic. Karambelas *et al* (2018) found large low biases across modeled concentrations evaluated with

satellite and surface observations, demonstrating the importance of having representative inventories and high-resolution chemical transport models. Emissions inventories are invaluable to air quality modeling and for estimating human health impacts due to pollution exposure, playing a vital role in advancing air quality and health understanding in India.

In this study, we present a high-resolution anthropogenic emissions inventory to identify pollution impacts on urban and rural populations. We update anthropogenic emissions from the Greenhouse Gas-Air Pollution Interactions and Synergies (GAINS) model (0.5° by 0.5°) using urban and rural population densities and emissions activity information for predominant anthropogenic sectors—domestic combustion and transportation. Applying this new updated emissions inventory at a high horizontal resolution (12 km by 12 km), the US Environmental Protection Agency (EPA) Community Multi-scale Air Quality (CMAQ) model, yields premature mortality estimates that are higher in rural regions compared to urban areas with populations greater than 100 000 people. Results demonstrate the sensitivity of health impact calculations to emissions inventories, and the value of expanding surface monitoring in rural India to improve air quality and accuracy of health impact assessments of the region.

## 2. Methods

### 2.1. Modeling

Air quality modeling was performed using the regional CMAQ model version 5.1. Previously, CMAQ has been used in India to assess O<sub>3</sub> regimes (Sharma *et al* 2016, Sharma and Khare 2017) and evaluate urban and rural emissions uncertainties (Karambelas *et al* 2018). We completed simulations for four seasonally representative months using 2010 meteorology and emissions: January, April, July, and October for winter, pre-monsoon spring, monsoon, and post-monsoon fall respectively. Model processes include surface- and upper-level emissions, photolysis, gas- and particle-phase chemistry, deposition, and dispersion Byun and Schere (2006). Simulations include the Carbon Bond 05 (CB05) chemical mechanism (Yarwood *et al* 2005) and AERO 6 aerosol mechanism, the inclusion of windblown dust (Dong *et al* 2015) with enhancements following Karambelas *et al* (2018), and in-line lightning NO<sub>x</sub> production (Allen *et al* 2012). Boundary and initial conditions are from a larger, 36 km by 36 km domain (Karambelas *et al* 2018). Meteorology is simulated using the Weather Research and Forecasting (WRF) model v3.2 and Preprocessing System with European Center for Medium-Range Weather Forecasting ERA-Interim globally gridded reanalysis data (Dee *et al* 2011). WRF is used to interpolate weather data from an 80 km resolution over 60 vertical layers at 6 hour increments

to our model domain over 36 vertical sigma layers from the surface to approximately 150 hPa using the Grell cumulus parameterization Grell and Devenyi (2002). For use in CMAQ, WRF output was ultimately preprocessed with the Meteorology-Chemistry Interface Processor (MCIP). Planetary boundary height, temperature, and precipitation from MCIP and evaluated with TRMM show MCIP captures seasonality in meteorology for the region, though underestimates precipitation (supplemental figure 1 available at [stacks.iop.org/ERL/13/064010/mmedia](http://stacks.iop.org/ERL/13/064010/mmedia)).

## 2.2. Emissions

We use biogenic, biomass burning, and anthropogenic emissions for year 2010 gridded to 12 by 12 km and vertically distributed (supplemental table 1). Monthly biogenic emissions are from the Community Land Model with the Model of Emissions and Gases from Nature (Guenther *et al* 2006) (downloaded from <http://lar.wsu.edu/megan/guides.html>). Monthly biomass burning emissions are from the Global Fire Emissions Database including small fires (GFED v4.1s) (van Der Werf *et al* 2010) and were vertically distributed following methods used in Karambelas *et al* (2018). Annual total anthropogenic emissions for 2010 are from the GAINS Model gridded globally following the ECLIPSE (Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants) project version 5a (Stohl *et al* 2015, Klimont *et al* 2017). In addition to regridding anthropogenic emissions from the global 0.5 degree by 0.5 degree to 12 km by 12 km over northern India, we distribute anthropogenic emissions vertically according to Simpson *et al* (2012). Updates according to urban and rural activities and population distributions where cities and towns greater than 100 000 people are designated as urban were included for the domestic combustion and transportation sectors to provide enhanced detail at the higher domain resolution. Emissions from electricity generation were also updated based on power plant location information. More details on population data and energy sector emissions updates can be found in the supplemental information. Our urban designation is different than used for population distribution in India, that states an urban population must have (1) at least a population of 5000; (2) 75% of the male working population is employed in non-agricultural work; and (3) the density is at least 400 people per square kilometer. Our urban-rural designations result in a distribution of 79% rural and 21% urban, and is approximately 2% more rural (less urban) than the 2001 population distribution off of which our data is based.

Solid fuel based cooking is one of the main sources of primary PM emissions in India, particularly among rural communities where biomass is freely available and access to clean fuels like liquid petroleum gas (LPG) or cookstoves that use them may be limited in availability and cost. Recent studies found residential combustion emissions are attributable for

20%–25% of premature deaths associated with ambient PM<sub>2.5</sub> air pollution (Conibear *et al* 2018, GBD MAPS Working Group 2018). We reallocate emissions from the domestic cooking sector using state totals and gridded population data following greater LPG consumption in urban regions and more traditional biomass burned in rural regions. Information on distinctions between urban and rural domestic cooking fuel use were from Pachauri *et al* (2013), and updated emissions were gridded according to urban and rural population activity fractions across each state. We did not include other socio-demographic factors such as proximity to forest edge (Winijkul *et al* 2016a, 2016b), yet we recognize this may be a relevant factor in emissions allocation.


Transportation emissions were adjusted according to vehicle type: mopeds, motor-cycles, light duty cars, light duty trucks, buses, and heavy duty trucks. Based on best available information from Indian transportation research on vehicle population composition in the Mumbai metropolitan region (Shirgaonkar 2012) and personal communications (with Dr. Jens Borken, AIR, IIASA in July 2015), emissions for urban and rural regions per vehicle type are distributed according to ratios in supplemental table 2. Equal distribution is given to urban and rural mopeds due to their accessibility for both low-income and middle-class households. Urban populations operate more motor-cycles, light duty cars and trucks, while buses and heavy-duty trucks are operated in greater frequency in rural areas. Assumptions regarding vehicle fleet age are not included in the redistribution. Urban and rural emissions were gridded based on respective population distributions.

The final emissions inventory included the updated sectors merged with non-updated sectors including agriculture, non-road, industry, and other sectors.

## 2.3. Estimating premature mortality

We use parameters from integrated exposure response functions from the GBD 2013 study (Cohen *et al* 2016, Forouzanfar *et al* 2015, World Health Organization 2016) to calculate the relative risk associated with diseases attributable to an excess concentration of ambient PM<sub>2.5</sub>. The attributable fraction of cause-specific mortality related to excess ambient air pollution is multiplied by our population dataset to estimate the number of premature deaths per grid cell.

We estimate the effects of long-term exposure to PM<sub>2.5</sub> on deaths due to chronic obstructive pulmonary disease (COPD), ischemic heart disease (IHD), lung cancer, and stroke. Baseline mortality rate estimates for these outcomes in India are from the World Health Organization (WHO) (World Health Organization 2017) and have been previously used for ambient air pollution health assessments in India (Chowdhury and Dey 2016, Ghude *et al* 2016). Our work follows methods similar to Ghude *et al* (2016), Burnett *et al* (2014) and Apte *et al* (2015). We estimate annual premature



mortality ( $\Delta M_j$ ) attributable to estimated annual average PM<sub>2.5</sub> concentrations as a result of disease  $j$ , where instances of IHD and stroke are age-dependent, following:

$$\Delta M_j = Y_j \times \frac{RR_{j-1}}{RR_j} \times P \quad (1)$$

where  $Y_j$  is the baseline mortality rate for a particular disease and  $P$  is total population. We apply the same baseline mortality rates across India, but we recognize that there are limitations to this method. We use relative risk tables parameters from GBD (2013) in calculating relative risk for disease  $j$  using an integrated exposure response (IER) function following Burnett *et al* (2014):

$$RR_j = 1 + \alpha_j \times [1 - \exp(-\gamma_j(\Delta PM_{2.5})^{\delta_j})] \quad (2)$$

where  $\alpha$ ,  $\gamma$ , and  $\delta$  are disease-dependent parameters (Burnett *et al* 2014, Apte *et al* 2015).  $\Delta PM_{2.5}$  is the excess PM<sub>2.5</sub> beyond an annual average counterfactual minimum risk concentration commonly placed between 5.8–8.0  $\mu\text{g m}^{-3}$  (Chowdhury and Dey 2016, Ghude *et al* 2016). Other recent estimates have used nonlinear power law functions derived from cohort studies to estimate relative risk attributable to PM<sub>2.5</sub> in India which result in lower premature mortality estimates compared to using IER methodology (Chowdhury and Dey 2016) and references therein, and we include analysis using this method in the supplemental information. We calculate the 95% confidence interval of disease-specific relative risk at each grid cell to define a range of mortality estimates attributable to PM<sub>2.5</sub> exposure.

We estimate premature deaths from COPD due to long-term exposure of estimated annual average concentrations of maximum daily 8 hour average (MDA8) O<sub>3</sub> following Ghude *et al* (2016) and Ostro (2004).

We use the relative risk calculation:


$$RR = \left[ \frac{X + 1}{X_0 + 1} \right]^{\delta} \quad (3)$$

where the minimum-risk concentration,  $X_0$ , is 37.6 ppb (range: 33.3–41.9 ppb) and the factor  $\delta$  is 0.1521, both derived from the GBD assessment (Lim *et al* 2012). This minimum-risk concentration range was used to estimate minimum and maximum mortality impacts. To estimate  $\Delta M$  attributable to annual average MDA8 O<sub>3</sub>, we substitute RR from equation (3) into equation (1). Premature mortality for both PM<sub>2.5</sub> and O<sub>3</sub> is estimated for adults 25 years and older, meaning our estimates are likely conservative considering they do not take into account infant, child, and young adult mortality. Finally, we apply grid-cell specific population distributions to assess differences across urban and rural regions, with urban defined as towns and cities with greater than 100 000 people in total respectively.

### 3. Results

#### 3.1. Updated annual emissions for northern India

Domain-wide 2010 annual emissions of anthropogenic NO<sub>x</sub>, SO<sub>2</sub>, and primary PM<sub>2.5</sub> (elemental carbon [EC], organic carbon [OC]) indicate spatial variability across the Indo-Gangetic Plain. Emissions vary across the region, with many instances of localized high values indicative of regions of relatively higher population density. Emissions of NO<sub>x</sub> (figure 1(a)) and SO<sub>2</sub> (figure 1(b)) exhibit similar distributions, with annual totals in the north India region (model domain) of 2.3 Tg N (2.6 Tg N) and 6.3 Tg (6.9 Tg) respectively, where there are many concentrated regions of emissions and fewer emissions from area sources, however this is much more pronounced among the SO<sub>2</sub> emissions due to power plants being a main source. Emissions of NO<sub>x</sub> are slightly greater than reported in a recent



total India 2005 inventory of 1.9 TgN derived using satellite tropospheric columns (Ghude *et al* 2013), but emissions of both NO<sub>x</sub> and SO<sub>2</sub> are less than the range (5–6.5 TgN and 7–9.5 Tg SO<sub>2</sub> respectively for 2010) of a recent comparative analysis (Saikawa *et al* 2017), as expected considering our simulations cover the northernmost half of India.


Particulate emissions are largely dominated by the domestic combustion sector, emissions from which are strongly dependent on population density and availability of combustible materials. Total PM<sub>2.5</sub> emissions for the north India domain (2.7 Tg) are about one-third of those used in recent studies covering all of India (9–9.1 Tg) (Venkataraman *et al* 2017, GBD MAPS Working Group 2018). Emissions indicate a high number of localized peaks in PM<sub>2.5</sub> emissions, and, unlike NO<sub>x</sub> and SO<sub>2</sub> emissions, broad regions of lower

levels of emissions (figure 1(c)). Variations in emissions magnitude indicate relatively densely populated villages among low-populated rural areas and the effect these different population distributions have on total emissions. Adjusting emissions according to urban-rural population and activity distribution information resolves detailed urban and rural differences for more precise air quality modeling.

### 3.2. Estimated annual concentrations of surface PM<sub>2.5</sub> and O<sub>3</sub>

Estimated annual average concentrations of PM<sub>2.5</sub> and MDA8 O<sub>3</sub> using four seasonally representative months are shown in figure 2; monthly average concentrations are shown in supplemental figure 2. Modeled average concentrations for PM<sub>2.5</sub> and MDA8 O<sub>3</sub> are 25.5  $\mu\text{g m}^{-3}$  and 41.7 ppb in northern India respectively (figures 2(a) and (b) respectively). Concentrations of PM<sub>2.5</sub> in the New Delhi National Capital Region are much higher (167.4  $\mu\text{g m}^{-3}$ ). Concentrations of both pollutants are relatively higher along the Indo-Gangetic Plain (IGP), corresponding with population density. South of the IGP, annual mean concentrations of PM<sub>2.5</sub> are comparatively low, between 30 and 40  $\mu\text{g m}^{-3}$  with some localized instances of concentrations greater than 50  $\mu\text{g m}^{-3}$  including near Ahmedabad to the west and Kolkata to the east. Localized instances of higher concentrations are predominantly from primary PM<sub>2.5</sub> species such as elemental and organic carbon (supplemental figure 3). Ozone concentrations exhibit local minima in highly populated urban areas, for instance in Ahmedabad and most noticeably Delhi, where concentrations average 34.3 ppb as a result of NO<sub>x</sub> titration due to modeled high NO<sub>2</sub> concentrations in excess of 20 ppb.

We evaluate model performance for 2010 with available observations from India's CPCB. Modeled O<sub>3</sub> exhibits a minimal high bias (NMB = +3.5%) across nine monitor locations. However at Delhi monitor locations model biases are actually low on average in comparison due to an overestimation of modeled O<sub>3</sub> depletion from NO<sub>x</sub> titration. Average high O<sub>3</sub> biases exhibited across northern India is exhibited in other CMAQ (Sharma *et al* 2016 and WRF-Chem Ghude *et al* 2016) analyses. Compared to two sites in Delhi, modeled daily PM<sub>2.5</sub> concentrations exhibit a slight high bias of +6.0%. Despite low model biases, normalized mean errors are quite high (O<sub>3</sub>: 83.4%; PM<sub>2.5</sub> 45.7%) and spatial correlations are poor (O<sub>3</sub>: -0.29; PM<sub>2.5</sub>: +0.41). Evaluation with CPCB measurements is limited due to challenges such as unavailable data and the influence of very local sources on measured concentrations. Finally, to determine if the model can represent annual average values of PM<sub>2.5</sub> and O<sub>3</sub> from the four month simulations, we compare our modeled values with long-term average concentrations at available monitors from 1 January 2010–31 December 2014 (detailed in the supplemental information). The four-month




average concentrations compare well with observations at nine monitor locations in Delhi for  $\text{PM}_{2.5}$ — $163 \mu\text{g m}^{-3}$  modeled and  $160 \mu\text{g m}^{-3}$  observed—and at 23 monitor locations in the IGP for annual average  $\text{O}_3$ —22 ppb modeled and 21 ppb observed. Evaluation metrics for  $\text{NO}_2$ ,  $\text{SO}_2$ , and  $\text{PM}_{10}$  for 2010 are shown in supplemental table 3.

### 3.3. Adult premature mortality in northern India

Using representative annual average concentrations of  $\text{PM}_{2.5}$  and  $\text{O}_3$ , we calculate premature mortality attributable to these pollutants within northern India.

Estimated population-weighted annual average concentrations of  $\text{PM}_{2.5}$  are  $72.1 \mu\text{g m}^{-3}$  in northern India. We estimate 463 200 (95% CI: 444 600–482 600) annual premature adults deaths resulting from excess  $\text{PM}_{2.5}$  pollution in northern India. Concentrations and adult premature deaths from COPD, IHD, stroke, and LC are greatest along IGP (figure 3(a)), where

most grid cells reflect annual premature deaths between 40 and 75 adults and several localized hotspots estimate more than 300 adults deaths attributable to  $\text{PM}_{2.5}$  (i.e. in New Delhi). In far northwestern India where the population and emission density is low, there are some cells where zero deaths are attributable to  $\text{PM}_{2.5}$ . Estimated annual average modeled concentrations in Delhi and Kolkata are  $152.7 \mu\text{g m}^{-3}$  and  $144.4 \mu\text{g m}^{-3}$ , respectively, well above the Indian National Ambient Air Quality Standard (NAAQS) of  $40 \mu\text{g m}^{-3}$  (annual average) and the WHO guideline of  $10 \mu\text{g m}^{-3}$ . The populations of Delhi and Kolkata are 16.8 and 4.5 million people according to the 2011 Census, respectively. Using the average concentration and total populations of Delhi (National Capital Territory) and Kolkata, we find that approximately 12 200 (11 700–12 800) and 3300 (3600–3900) premature deaths occur each year in Delhi and Kolkata, respectively, as a result of excess  $\text{PM}_{2.5}$ . Our estimate of premature mortality



in Delhi is about 50% greater than the 8300 deaths from Amann *et al* (2017), likely due to our average modeled concentration being greater by nearly  $30 \mu\text{g m}^{-3}$  and differences in model resolution, emissions, and population datasets. Values for other cities atop the WHO most polluted cities list are included in the Supplemental Information.

Larger numbers of premature deaths attributable to MDA8 O<sub>3</sub> are evident across the IGP than in central India (figure 3(b)). The average population weighted MDA8 O<sub>3</sub> concentration in northern India is 56.5 ppb. Exposure to O<sub>3</sub> contributes to approximately 37 600 (28 500–48 100) adult lives lost annually. In Delhi, the MDA8 O<sub>3</sub> concentration of 34.3 ppb is lower than both the O<sub>3</sub> ambient air quality standards of both India and the WHO (50 ppb) due to NO<sub>x</sub> titration, while the average MDA8 O<sub>3</sub> concentration in Kolkata (59.5 ppb) is in excess of these standards. Ozone-attributable COPD deaths in these cities are an estimated 100 and 315, respectively. In contrast to PM<sub>2.5</sub>, there are many grid cells across India where fewer than five premature adult deaths are attributable to O<sub>3</sub>, suggesting that PM<sub>2.5</sub> rather than O<sub>3</sub> plays a larger health risk to the northern India population.

### 3.3.1. Urban and rural adult premature deaths

Using urban and rural population distributions, we find that excess PM<sub>2.5</sub> and O<sub>3</sub> pollution affects rural populations three to five times larger, with annual premature adult deaths in rural regions of 383 600 adults (figure 3(c)) compared to 117 200 deaths in urban regions

(figure 3(d)). An estimated 352 400 premature deaths in rural regions are attributable to excess PM<sub>2.5</sub>, compared to an estimated 110 800 adults in urban regions. Rural deaths correspond with a greater quantity of primary particulate emissions (2.3 Tg of elemental and organic carbon) versus urban (0.48 Tg). An urban-rural effect is found for O<sub>3</sub> as well, where the number of premature deaths in rural (urban) regions due to O<sub>3</sub> pollution is 31 200 (6400) adults. However, per total urban and rural population and in comparison to the total death rate of 72 people per 10 000, urban (rural) deaths attributable to ambient air pollution are 6.4 (5.6) per 10 000, such that ambient pollution in 2010 contributed to 8.1% of deaths.

Ambient PM<sub>2.5</sub>-attributable premature deaths are greatest in rural regions for all diseases (figure 4). By far the greatest cause of death as a result of PM<sub>2.5</sub> is from ischemic heart disease (IHD), causing an estimated 160 900 premature deaths of adults annually. Total deaths from PM<sub>2.5</sub>-attributable IHD is lower in urban areas at 48 000 premature deaths. Stroke contributes to the second greatest cause of PM<sub>2.5</sub> related premature mortality, with an estimated 133 500 (41 000) people in rural (urban) areas. COPD due to both PM<sub>2.5</sub> and O<sub>3</sub> affects a smaller number of people in both rural and urban regions (80 200 rural and 25 300 urban annually), and lung cancer has the lowest impact according to our estimates (9000 rural and 2900 urban annually). All of these estimates were based only on population exposure with no weight given to economic measures such as access to healthcare, however this may play



a considerable role in survival rate against a disease (Chowdhury and Dey 2016).

### 3.4. Limitations

This study has several limitations. One limitation is deficiencies in other emissions sectors, for instance, data is particularly lacking for the industrial sector. Further, emissions inventories in India remain highly uncertain. Recently, an emissions inventory comparison noted large uncertainties, on the order of 150%–325%, across several inventories for different pollutants from the domestic combustion sector (Saikawa *et al* 2017). Updates to support reducing uncertainties in the domestic combustion sector have been included in this study, yet still requires considerable research.

A second limitation is using four model months to represent annual average concentrations. Each month was chosen to represent a unique air quality condition: January for the polluted winter due to inversions, April to represent spring windblown dust, July to simulate air quality conditions during the monsoon, and October to represent the onset of the agricultural biomass burning season in northwestern India. However, we note above that the average concentrations estimated using these four months ( $\text{PM}_{2.5}$ :  $163 \mu\text{g m}^{-3}$ ,  $\text{O}_3$ : 22 ppb) are comparable to annual average long-term observations from the CPCB ( $\text{PM}_{2.5}$ :  $160 \mu\text{g m}^{-3}$ ,  $\text{O}_3$ : 21 ppb).

Finally, there are limitations in using the IER methodology to estimate premature mortality in India. The concentration-response functions for ambient air pollution used here were developed in the context of the GBD 2013 study (Forouzanfar *et al* 2015) based on epidemiological studies in comparatively low-pollution regions including the US and Europe. Applying these functions to estimate health effects in a high pollution region such as India relies upon extrapolation to higher levels based on studies from the passive and active smoking literatures (Burnett *et al* 2014) and references therein. The parameters used in the IERs underlie considerable uncertainty, and recent updates in the context of GBD studies since 2010 led to distinctly different estimates of premature deaths. However, in the absence of long-term air pollution cohort studies at high concentrations, using published and documented exposure-response relationships remains the best approach for estimating effects in India, and yields findings that can promote awareness of health implications and the need for observational studies in India to obtain more accurate representation of underlying health status, relevant time spent outdoors, access to healthcare, and genetic predisposition.

## 4. Conclusion

An estimated 463 200 (95% CI: 444 600–482 600) and 37 800 (28 500–48 100) premature deaths occurred

annually in 2010 as a result of excess  $\text{PM}_{2.5}$  and  $\text{O}_3$  concentrations, respectively, in northern India. Compared to prior studies that estimate annual adult premature mortality due to  $\text{PM}_{2.5}$  (between 486 100 and 1 100 000 (Chowdhury and Dey 2016, Ghude *et al* 2016, GBD MAPS Working Group 2018)) and/or  $\text{O}_3$  (12 000 (Ghude *et al* 2016)) across all of India, our estimates from this study over northern India are slightly lower for  $\text{PM}_{2.5}$  (463 200 premature deaths) as expected due to the smaller domain we use in our study, yet are much higher for  $\text{O}_3$  (37 800 premature deaths). The discrepancy between these totals may reflect our use of MDA8  $\text{O}_3$  concentrations to consider effects from peak  $\text{O}_3$  exposure used for setting air quality standards, whereas Ghude *et al* (2016) used bias-corrected annual average concentrations to estimate average exposure levels. Finally, we note that our estimates are for adult premature mortality, though it is estimated that up to 6% of childhood premature deaths between the ages of five and fifteen are the result of lower respiratory infections attributable to ambient air pollution exposure (Institute for Health Metrics and Evaluation 2016).

We conduct model simulations to generate estimated annual average ambient concentrations of  $\text{PM}_{2.5}$  and  $\text{O}_3$  from the CMAQ model using a new high-resolution anthropogenic emissions inventory updated according to urban and rural population and activity distribution measures for transportation and domestic combustion. Separating total premature deaths according to respective urban and rural areas, we find that 6.4 (5.6) deaths per 10 000 people occur in urban (rural) areas attributable to ambient  $\text{PM}_{2.5}$  and  $\text{O}_3$  concentrations. However, despite the greater rate of deaths per 10 000 people that occur in urban areas, the total amount of premature deaths attributable to air pollution in rural areas (383 600 deaths) is much larger in magnitude than those that occur in urban populations annually (117 200). Our results (76% rural) are in line with those in a modeling study from the Health Effect Institute, that found 75% of premature deaths attributable to air pollution affect rural populations (GBD MAPS Working Group 2018), with a large fraction of premature mortality in rural populations due to residential combustion. We note this similarity in urban-to-rural ratios despite the present study using (1) a more common global anthropogenic emissions inventory with updates as opposed to an inventory developed from prior studies over India, and (2) a higher resolution regional chemical transport model. Rural populations are burdened by ambient air pollution three to five times greater than urban populations, indicating a need for surface air pollution monitoring in rural regions.

This work supports the growing body of research using modeled ambient concentrations to estimate premature mortality associated with exposure to air pollution in India by using high-resolution detailed updates for urban and rural emissions for

transportation and domestic combustion in addition to location-specific power plant emissions. Ultimately, it is challenging to gain a full understanding of the impacts on adult mortality in northern India without significant measurements available to validate air quality model results. Satellite tropospheric vertical column depths (VCDs) can potentially fill in the spatial gaps (Chowdhury and Dey 2016), yet model data are often necessary to derive surface estimates from the VCDs to use in health impacts assessments (Ghude *et al* 2016). Future efforts would benefit from an increased network of surface observations to assess more rigorously air quality model performance and to build confidence in applying these models to quantify human health impacts in the region. With surface observations available in rural regions, we can also better characterize ambient air pollution and the skill of high-resolution emissions inventories to provide the scientific basis for selecting effective emission control strategies to improve air quality in India.

### Acknowledgments

We thank three anonymous reviewers for providing valuable feedback. A K acknowledges J Borken for guidance on transportation emissions, W Schoepp for providing gridded urban and rural population data, and The Earth Institute at Columbia University for supporting this research.

### ORCID iDs

Alexandra Karambelas  <https://orcid.org/0000-0003-1886-7000>

Chris Heyes  <https://orcid.org/0000-0001-5254-493X>

### References

- Allen D J, Pickering K E, Pinder R W, Henderson B H, Appel K W and Prados A 2012 Impact of lightning-NO on eastern United States photochemistry during the summer of 2006 as determined using the CMAQ model *Atmos. Chem. Phys.* **12** 1737–58
- Amann M *et al* 2011 Cost-effective control of air quality and greenhouse gases in Europe: modeling and policy applications *Environ. Model. Softw.* **26** 1489–501
- Amann M *et al* 2017 Managing future air quality in megacities: a case study for Delhi *Atmos. Environ.* **161** 99–111
- Apte J S, Marshall J D, Cohen A J and Brauer M 2015 Addressing global mortality from ambient PM<sub>2.5</sub> *Environ. Sci. Technol.* **49** 8057–66
- Brauer M *et al* 2012 Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution *Env. Sci Technol.* **46** 652–60
- Burnett R T *et al* 2014 An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure *Environ. Health Perspect.* **122** 397–403
- Byun D and Schere K L 2006 review of the governing equations, computational algorithms, and other components of the models-3 community multiscale air quality (CMAQ) modeling system *Appl. Mech. Rev.* **59** 51
- Central Pollution Control Board 2008 National Air Quality Monitoring Programme ([www.cpcb.nic.in/air.php](http://www.cpcb.nic.in/air.php)) (Accessed: 1 January 2016)
- Chowdhury S and Dey S 2016 Cause-specific premature death from ambient PM<sub>2.5</sub> exposure in India: estimate adjusted for baseline mortality *Environ. Int.* **91** 283–90
- Cohen A *et al* 2016 private communication
- Conibear L, Butt E W, Knotte C, Arnold S R and Spracklen D V 2018 Residential energy use emissions dominate health impacts from exposure to ambient particulate matter in India *Nat. Commun.* **9** 1–9
- Dee D P *et al* 2011 The ERA-Interim reanalysis: Configuration and performance of the data assimilation system *Q. J. R. Meteorol. Soc.* **137** 553–97
- Dong X, Fu J S, Huang K, Tong D and Zhuang G 2015 Model development of dust emission and heterogeneous chemistry within the community multiscale air quality modeling system and its application over East Asia *Atmos. Chem. Phys. Discuss.* **15** 35591–643
- Duncan B N *et al* 2015 A space-based, high-resolution view of notable changes in urban NO<sub>x</sub> pollution around the world (2005–2014) *J. Geophys. Res. Atmos.* **121** 976–96
- ENVIS Centre on Control of Pollution 2016 Air Quality: National Air Quality Monitoring Programme (NAMP) ([http://cpcbenviis.nic.in/air\\_pollution\\_main.html](http://cpcbenviis.nic.in/air_pollution_main.html)) (Accessed: 1 January 2016)
- European Commission and Joint Research Centre (JRC)/Netherlands Environmental Assessment (PBL) 2011 Emission Database for Global Atmospheric Research (EDGAR), release version 4.2
- Forouzanfar M H *et al* 2015 Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013 *Lancet* **386** 2287–323
- Forouzanfar M H *et al* 2016 Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the global burden of disease Study 2015 *Lancet* **388** 1659–724
- GBD MAPS Working Group 2018 *Burden of Disease Attributable to Major Air Pollution Sources in India* (Boston, MA: Health Effects Institute)
- Ghude S D, Pfister G G, Jena C, Van Der R J A, Emmons L K and Kumar R 2013 Satellite constraints of nitrogen oxide (NO<sub>x</sub>) emissions from India based on OMI observations and WRF-Chem simulations *Geophys. Res. Lett.* **40** 423–8
- Ghude S D, Chate D M M, Jena C, Beig G, Kumar R, Barth M C C, Pfister G G G, Fadnavis S and Pithani P 2016 Premature mortality in India due to PM<sub>2.5</sub> and ozone exposure *Geophys. Res. Lett.* **43** 4650–8
- Grell G A and Devenyi D 2002 A generalized approach to parameterizing convection combining ensemble and data assimilation techniques *Geophys. Res. Lett.* **29** 10–13
- Guenther a, Karl T, Harley P, Wiedinmyer C, Palmer P I and Geron C 2006 Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature) *Atmos. Chem. Phys.* **6** 107–73
- Guttikunda S K and Jawahar P 2014 Atmospheric emissions and pollution from the coal-fired thermal power plants in India *Atmos. Environ.* **92** 449–60
- Health Effects Institute 2017 State of Global Air 2017
- Hilboll A, Richter A and Burrows J P 2017 NO<sub>2</sub> pollution over India observed from space and the impact of rapid economic growth, and a recent decline *Atmos. Chem. Phys. Discuss.* **20** 1–18

- Institute for Health Metrics and Evaluation 2016 GBD Compare Viz Hub (<https://vizhub.healthdata.org/gbd-compare/>) (Accessed: 25 April 2018)
- Jena C, Ghude S D, Beig G, Chate D M, Kumar R, Pfister G G, Lal D M, Surendran D E, Fadnavis S and van der R J A 2015 Inter-comparison of different NO<sub>x</sub> emission inventories and associated variation in simulated surface ozone in Indian region *Atmos. Environ.* **117** 61–73
- Karambelas A, Holloway T, Kiesewetter G and Heyes C 2018 Constraining the uncertainty in emissions over India with a regional air quality model evaluation *Atmos. Environ.* **174** 194–203
- Klimont Z, Kupiainen K, Heyes C, Purohit P, Cofala J and Rafaj P 2017 Global anthropogenic emissions of particulate matter including black carbon *Atmos. Chem. Phys.* **17** 8681–723
- Kurokawa J, Ohara T, Morikawa T, Hanayama S, Janssens-Maenhout G, Fukui T, Kawashima K and Akimoto H 2013 Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: regional emission inventory in ASia (REAS) version 2 *Atmos. Chem. Phys.* **13** 11019–58
- Lelieveld J, Evans J S, Fnais M, Giannadaki D and Pozzer A 2015 The contribution of outdoor air pollution sources to premature mortality on a global scale *Nature* **525** 367–71
- Lim S S *et al* 2012 A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the global burden of disease study 2010 *Lancet* **380** 2224–60
- Ostro B 2004 *Outdoor Air Pollution: Assessing the Environmental Burden of Disease at National and Local Levels* (Geneva: World Health Organization)
- Pachauri S, van Ruijven B J, Nagai Y, Riahi K, van Vuuren D P, Brew-Hammond A and Nakicenovic N 2013 Pathways to achieve universal household access to modern energy by 2030 *Environ. Res. Lett.* **8** 8
- Pandey A, Sadavarte P, Rao A V and Venkataraman C 2014 Trends in a technology-linked multi-pollutant inventory of Indian energy-use emissions: II. Residential, agricultural and informal industry sectors *Atmos. Environ.* **99** 341–52
- Sadavarte P and Venkataraman C 2014 Trends in multi-pollutant emissions from a technology-linked inventory for India: I. Industry and transport sectors *Atmos. Environ.* **99** 353–64
- Saikawa E *et al* 2017 Uncertainties in emissions estimates of greenhouse gases and air pollutants in India and their impacts on regional air quality *Environ. Res. Lett.* **12** 065002
- Sharma S and Khare M 2017 Simulating ozone concentrations using precursor emission inventories in Delhi - National Capital Region of India *Atmos. Environ.* **151** 117–32
- Sharma S, Chatani S, Mahtta R, Goel A and Kumar A 2016 Sensitivity analysis of ground level ozone in India using WRF-CMAQ models *Atmos. Environ.* **131** 29–40
- Shirgaonkar M 2012 The rapid rise of middle-class vehicle ownership in Mumbai *Thesis* (Berkeley, CA: University of California)
- Simpson D *et al* 2012 The EMEP MSC-W chemical transport model—Technical description *Atmos. Chem. Phys.* **12** 7825–65
- Stohl A *et al* 2015 Evaluating the climate and air quality impacts of short-lived pollutants *Atmos. Chem. Phys. Discuss.* **15** 15155–241
- Surendran D E, Ghude S D, Beig G, Emmons L K, Jena C, Kumar R, Pfister G G and Chate D M 2015 Air quality simulation over South Asia using hemispheric transport of air pollution version-2 (HTAP-v2) emission inventory and model for ozone and related chemical tracers (MOZART-4) *Atmos. Environ.* **122** 357–72
- UNICEF 2016 *Clear the Air for Children - The Impact of Air Pollution on Children* (New York: UNICEF)
- Venkataraman C *et al* 2017 Source influence on emission pathways and ambient PM<sub>2.5</sub> pollution over India 2015–50 *Atmos. Chem. Phys. Discuss.* [in review](#)
- van Der Werf G R, Randerson J T, Giglio L, Collatz G J, Mu M, Kasibhatla P S, Morton D C, Defries R S, Jin Y and Van Leeuwen T T 2010 Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires 1997–2009 *Atmos. Chem. Phys.* **10** 11707–35
- Winijkul E, Fierce L and Bond T C 2016a Emissions from residential combustion considering end-uses and spatial constraints: Part I, methods and spatial distribution *Atmos. Environ.* **125** 126–39
- Winijkul E, Fierce L and Bond T C 2016b Emissions from residential combustion considering end-uses and spatial constraints: Part I, methods and spatial distribution *Atmos. Environ.* **125** 126–39
- World Health Organization 2016 *Ambient Air Pollution: A Global Assessment of Exposure and Burden of Disease* (Geneva: World Health Organization)
- World Health Organization 2017 Mortality Raw Data ([www.who.int/healthinfo/statistics/mortality\\_rawdata/en/%0A](http://www.who.int/healthinfo/statistics/mortality_rawdata/en/%0A)) (Accessed: 6 December 2017)
- Yarwood G, Rao S, Yocke M and Whitten G Z 2005 *Updates to the Carbon Bond chemical mechanism: CB05 Final Report to the US EPA, RT-0400675*
- Zhang Q *et al* 2009 Asian emissions in 2006 for the NASA INTEX-B mission *Atmos. Chem. Phys.* **9** 5131–53