

# Loughborough University Institutional Repository

---

## *Cooling athletes with a spinal cord injury*

This item was submitted to Loughborough University's Institutional Repository by the/an author.

**Citation:** GRIGGS, K.E., PRICE, M.J. and GOOSEY-TOLFREY, V.L., 2015. Cooling athletes with a spinal cord injury. *Sports Medicine*, 45 (1), pp.9-21.

**Additional Information:**

- The final publication is available at Springer via <http://dx.doi.org/10.1007/s40279-014-0241-3>

**Metadata Record:** <https://dspace.lboro.ac.uk/2134/17075>

**Version:** Accepted for publication

**Publisher:** © Springer

**Rights:** This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the published version.

1 **Cooling Athletes with a Spinal Cord Injury**

2 *Katy E. Griggs<sup>1</sup>, Michael J. Price<sup>2</sup> and Victoria L. Goosey-Tolfrey<sup>1</sup>*

3 <sup>1</sup>Peter Harrison Centre for Disability Sport, School of Sport, Exercise and Health Sciences,  
4 Loughborough University, Loughborough, UK

5 <sup>2</sup>Department of Biomolecular and Sports Science, Coventry University, Coventry, UK

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1 **Acknowledgements**

2 The authors thank the Peter Harrison Centre for Disability Sport for their ongoing support.  
3 The authors have no conflicts of interest that are directly relevant to the content of this review.

4 **Corresponding Author**

5 Prof. Victoria Goosey-Tolfrey

6 Peter Harrison Centre for Disability Sport,

7 School of Sport, Exercise and Health Sciences

8 Loughborough University,

9 Loughborough,

10 UK

11 Email: [V.L.Tolfrey@lboro.ac.uk](mailto:V.L.Tolfrey@lboro.ac.uk)

12 Telephone: 01509 226386

13 **Short title/ Running head**

14 Cooling Athletes with a Spinal Cord Injury

15 **Key Points**

- 16 • According to the studies reviewed, wearing an ice vest during intermittent sprint  
17 exercise both reduces thermal strain and enhances performance.
- 18 • A combination of pre-cooling and cooling during exercise or half-time cooling may  
19 be an effective strategy.
- 20 • Cooling strategies for athletes with a spinal cord injury should be individualized to  
21 account for the level and completeness of the lesion while also considering the  
22 specific regulations and logistics of the sport involved.

23

24

25

## 1 **Abstract**

2

3 Cooling strategies that help prevent a reduction in exercise capacity whilst exercising in the  
4 heat have received considerable research interest over the past three decades, especially in the  
5 lead up to a relatively hot Olympic and Paralympic Games. Progressing into the next  
6 Olympic/Paralympic cycle the host, Rio de Janeiro, could again present an environmental  
7 challenge for those athletes competing. Despite the interest and vast array of research into  
8 cooling strategies for the able-bodied athlete, less is known regarding the application of these  
9 cooling strategies in the thermoregulatory impaired spinal cord injured (SCI) athletic  
10 population. Individuals with a spinal cord injury (SCI) have a reduced afferent input to the  
11 thermoregulatory centre and a loss of both sweating capacity and vasomotor control below  
12 the level of the spinal cord lesion. The magnitude of this thermoregulatory impairment is  
13 proportional to the level of the lesion. For instance, individuals with high level lesions  
14 (tetraplegia) are at a greater risk of heat illness than individuals with lower level lesions  
15 (paraplegia) at a given exercise intensity. Therefore, cooling strategies may be highly  
16 beneficial in this population group, even in moderate ambient conditions (~21°C). This  
17 review was undertaken to examine the scientific literature that addresses the application of  
18 cooling strategies in individuals with a SCI. Each method is discussed in regards to the  
19 practical issues associated with the method and the potential underlying mechanism. For  
20 instance, site specific cooling would be more suitable for an athlete with a SCI than whole  
21 body water immersion, due to the practical difficulties of administering this method in this  
22 population group. From the studies reviewed, wearing an ice vest during intermittent sprint  
23 exercise has been shown to decrease thermal strain and improve performance. These  
24 garments have also been shown to be effective during exercise in the able-bodied. Drawing  
25 on additional findings from the able-bodied literature the combination of methods used prior  
26 to and during exercise and/or during rest periods/half-time may increase the effectiveness of a  
27 strategy. However, due to the paucity of research involving athletes with a SCI, it is difficult  
28 to establish an optimal cooling strategy. Future studies are needed to ensure that research  
29 outcomes can be translated into meaningful performance enhancements by investigating  
30 cooling strategies under the constraints of actual competition. Cooling strategies which meet  
31 the demands of intermittent wheelchair sports need to be identified, with particular attention  
32 to the logistics of the sport.

# 1 Introduction

2

3 Prolonged exercise in high ambient temperatures and/or high humidity has a detrimental  
4 effect on physical performance in able-bodied individuals.<sup>[1]</sup> Cooling strategies that reduce  
5 the attenuation of athletic performance have been investigated over the past three decades.  
6 There has been increasing interest in cooling strategies, particularly in the lead up to a  
7 relatively hot Olympic Games, such as Athens 2004 (30-38°C, 40-48% relative humidity)<sup>[2]</sup>  
8 or Beijing 2008 (20-30°C, 63-90% relative humidity).<sup>[3]</sup> Progressing into the next  
9 Olympic/Paralympic cycle, the Rio de Janeiro 2016 Games will again present a unique  
10 challenge for the athlete, with the need to prepare and adapt to the potential environmental  
11 conditions of 19-29°C and ~70% relative humidity.<sup>[4]</sup>

12 Despite the interest in the application of cooling strategies for the able-bodied athlete,  
13 comparatively little is known regarding this for a wheelchair bound athletic population,  
14 specifically athletes with a spinal cord injury (SCI). Individuals with a SCI have a reduced  
15 afferent input to the thermoregulatory centre<sup>[5-7]</sup> and a loss of both sweating capacity and  
16 vasomotor control below the level of the spinal cord lesion.<sup>[5, 8, 9]</sup> As blood redistribution and  
17 sweating are two major thermoregulatory effectors, this suggests that individuals with a SCI  
18 have compromised thermoregulation and a greater risk of heat illness than able-bodied  
19 individuals.<sup>[10]</sup>

20 The magnitude of the thermoregulatory impairment is proportional to the level and  
21 completeness of the lesion. For instance, exercising for 60-90 minutes at 60% peak oxygen  
22 uptake ( $\dot{V}O_{2\text{peak}}$ ) in 15-25°C, athletes with a thoracic, lumbar or sacral SCI (paraplegia) may  
23 experience an increase in core temperature ( $T_{\text{core}}$ ) of ~1°C, similar to their able-bodied  
24 counterparts.<sup>[11, 12]</sup> For the same exercise intensity in hot conditions (35-40°C), a greater  
25 increase in  $T_{\text{core}}$  over time may be apparent, when compared to the able-bodied, signifying an  
26 imbalance between heat loss and heat gain, though exercise can still be maintained.<sup>[13]</sup> Above  
27 exercise intensities of 60%  $\dot{V}O_{2\text{peak}}$  in hot conditions, sweating and heat production may  
28 become unbalanced causing an increased risk of heat injury for athletes with paraplegia.<sup>[10]</sup>  
29 Compared to individuals with paraplegia or an incomplete spinal injury, individuals with a  
30 cervical SCI (tetraplegia) and/or a complete injury possess a smaller area of sensate skin, a  
31 lesser amount of afferent input regarding their thermal state and a reduced efferent response.<sup>[9,</sup>  
32 <sup>14]</sup> Limited data is available for athletes with tetraplegia regarding heat strain in relation to

1 exercise intensity, yet their high level lesions predispose them to a greater degree of thermal  
2 strain than athletes with paraplegia.<sup>[11, 15]</sup>

3 In an athletic setting, it is imperative that the appropriate training practices, including cooling  
4 strategies, are applied to support athletic performance for athletes, but more importantly for  
5 those with a SCI, that the risk of heat injury is also considered. The purpose of this review is  
6 to examine the scientific literature addressing application of cooling strategies to individuals  
7 with a SCI with specific reference to the practical and logistical issues associated with each  
8 method.<sup>[8]</sup> The potential underlying mechanism for improved heat tolerance for each method  
9 will also be addressed.

### 10 **1.1 Methods**

11

12 A comprehensive literature search was conducted using the search engines Web of Science,  
13 SPORTDiscus, MEDLINE and Biological Sciences. Key terms employed included “cooling,”  
14 with “sport,” “Paralympic,” “wheelchair sport,” “exercise,” “performance,” with “spinal cord  
15 lesion,” “spinal cord injury,” “tetraplegia,” “quadriplegia” and “paraplegia.” The last day of  
16 the literature search was 09 May 2014. The inclusion criteria comprised studies that reported  
17 a population group of individuals with a SCI, use of cooling strategies with exercise and  
18 articles that were published in English language peer reviewed journals. Figure 1 provides a  
19 flow diagram of the literature search. By assessing the eligibility of the articles, in regards to  
20 the inclusion criteria, nine articles were selected for review (Table 1).

21

22

23


24

25

26

27

28



10 Figure 1. Overview of study selection process. n indicates number of studies. SCI spinal cord  
11 injury.

*Insert Table 1 here*



## 1 2 Cooling Methods

2

3 The majority of exercise-related cooling methods are applied externally, either in isolation or  
4 as a combination of methods. Recent studies have considered internal application by  
5 ingestion of a cooling medium (i.e. ice slurry).<sup>[16, 17]</sup> Widespread use of various cooling  
6 methods and techniques in research in the able-bodied population include; cold air  
7 exposure,<sup>[18, 19]</sup> wearing of ice vests,<sup>[20-22]</sup> whole body water immersion,<sup>[23-25]</sup> water  
8 immersion of the extremities<sup>[26, 27]</sup> and ice slurry or cold water ingestion.<sup>[16, 17]</sup> These methods  
9 influence the body in different ways by either reducing  $T_{\text{core}}$  and mean skin temperature ( $T_{\text{sk}}$ )  
10 (water immersion); reducing  $T_{\text{core}}$  with no effect on  $T_{\text{sk}}$  (ice slurry or cold water ingestion) or  
11 primarily reducing  $T_{\text{sk}}$  (ice vests, cold air exposure, water sprays). Methods differ greatly in  
12 terms of their physiological effects, cooling power and, importantly, the timing of application.  
13 Moreover, the duration of the cooling strategy employed is also crucial to ensure maximum  
14 benefit to the athlete's performance without compromising athlete comfort or pre-event/half-  
15 time routine. Depending on the method used, the time required to achieve a significant  
16 reduction in  $T_{\text{core}}$  has been shown to range from 15-60 minutes of cooling.<sup>[28, 29]</sup> In assessing  
17 the practicality of a cooling method for an athletic population the effectiveness, sporting  
18 regulations, staffing/assistance, logistics, cost and athlete comfort need to be considered.  
19 Although, there is still debate regarding which strategy provides the greatest performance  
20 benefit in the able-bodied population, recent meta-analysis data has shown that pre-cooling,  
21 especially mixed method cooling,<sup>[30]</sup> can improve subsequent intermittent and prolonged  
22 exercise performance in the heat.<sup>[31]</sup> During exercise, meta-analysis data has revealed the use  
23 of ice vests to have the largest effect on performance.<sup>[30]</sup>

24

25

*Insert Table 2 here*

## 2.1 Hand/Foot cooling

An alternative to whole body water immersion is to submerge, in water, inactive or active body parts. An advantage of water immersion, even if only partial immersion, is that the thermal conductivity of water is 27 times greater than the thermal conductivity of air at the same temperature, causing greater heat loss in water compared to air.<sup>[32]</sup> For instance, cooling of the hands or feet for 20 minutes in 10°C water has been shown to remove heat of 100-150 W.<sup>[33, 34]</sup> Consequently, a reduction in  $T_{sk}$  and possibly  $T_{core}$ , can be achieved without directly cooling active muscle tissue enabling greater potential for heat storage.<sup>[35]</sup> Compared to the practically difficult method of whole body water immersion, this site-specific cooling, in particular cooling of the extremities, is more accessible for athletes with a SCI. Furthermore, cooling of the extremities in water is a self-limiting method of cooling, whereby the peripheral blood flow controls the degree of cooling which is in turn driven by the level of  $T_{core}$ .<sup>[27]</sup> This method of cooling thus relies on an elevated  $T_{core}$  in order for the cooler venous blood in the hands and feet to flow directly to the core through the superficial veins.<sup>[27, 33, 34]</sup> Without this drive for blood flow, vasoconstriction occurs and heat loss is prevented<sup>[27]</sup> and further elevations in heat storage will occur.

To assess if cooling hyperthermic athletes with a SCI following 60 minutes of steady state exercise in ~31°C improved subsequent 1 km time trial performance, Goosey-Tolfrey et al.<sup>[36]</sup> demonstrated that immersing the hands in 10°C water for 10 minutes elicited a 0.4°C reduction in insulated aural temperature ( $T_{ac}$ ). Time to complete 1 km was improved by 20.5 seconds, yet this was non-significant. A statistically significant performance improvement may have been masked by the diverse range of disabilities examined within this study (Table 1), which included both athletes with tetraplegia and paraplegia. During exercise, substantial thermoregulatory differences have been observed between high (tetraplegia) and low (paraplegia) lesion levels, indicating that the thermal responses, and as a consequence, performance, of athletes with paraplegia and tetraplegia in this study may have been different.<sup>[11]</sup> As the chosen performance trial is not a standard race distance for wheelchair athletes whether this small reduction in performance has practical significance in a competition setting has yet to be determined.

In comparison to the able-bodied control group, the athletes with a SCI in the study by Goosey-Tolfrey et al.<sup>[36]</sup> experienced a smaller increase in  $T_{ac}$  from the steady state exercise (2.0°C vs 1.3°C for the able-bodied and wheelchair athletes, respectively) and a smaller reduction in  $T_{ac}$

1 from cooling the hands, due to the lower thermal drive for cutaneous blood flow and thus  
2 cooling. The authors also noted that although thermal sensation was slightly reduced following  
3 hand cooling, the wheelchair athletes experienced a lack of grip and feelings of numbness of the  
4 hands, indicating that immersing the hands for 10 minutes in 10°C water may not be appropriate  
5 for wheelchair court sports where hand dexterity is of paramount importance. Many athletes in  
6 wheelchair sports wear gloves making hand cooling before or during play practically difficult.  
7 Consequently, cooling inactive body regions, such as the feet, may be more suitable, to ensure  
8 upper body movement and hand dexterity needed for wheelchair propulsion are not inhibited.

9 Cooling devices which target the heat-exchanging arteriovenous anastomoses in the feet have  
10 been developed.<sup>[37]</sup> Such devices increase the transfer of heat by alternating between negative  
11 and neutral pressure to facilitate blood return from the lower extremities and apply a heat sink to  
12 the skin surface, increasing the extraction of heat from the circulating blood.<sup>[37]</sup> One of these  
13 foot cooling devices was used by Hagobian et al.<sup>[37]</sup> to cool individuals with a SCI during 45  
14 minutes of arm cranking at 66%  $\dot{V}O_{2peak}$  in 32°C conditions. The increase in tympanic  
15 temperature ( $T_{ty}$ ) was attenuated by 0.6°C in the foot cooling condition (1.0°C vs. 1.6°C,  
16 respectively). It should be noted that  $T_{ty}$  was not attenuated when negative pressure was not  
17 applied, potentially due to a reduction in cutaneous blood flow in the feet. However, the thermal  
18 responses of the participants in Hagobian et al.<sup>[37]</sup> may have been accentuated as participants  
19 were untrained (arm crank  $\dot{V}O_{2peak}$   $14.4 \pm 2.9$  ml·kg<sup>-1</sup>·min<sup>-1</sup>). Therefore, they may have  
20 experienced a lower dissipation of heat and greater increases in  $T_{core}$  compared to trained  
21 individuals,<sup>[10]</sup> who could potentially benefit from partial acclimation.<sup>[38]</sup> Similarly to Goosey-  
22 Tolfrey et al.<sup>[36]</sup> the authors did not specify the number of participants with tetraplegia or  
23 paraplegia. Due to the lack of a performance measure used in this study, the effect on  
24 performance and practicality of using the device during competition is currently unknown.  
25 Future application of this method could investigate the use of this device post exercise or in a  
26 clinical setting.

## 27           2.2 Water Sprays

28  
29 Players in wheelchair court sports with a SCI tend to use water sprays during rest periods to cool  
30 themselves due to their ease of use. The concept of using water sprays is to partially mimic the  
31 sweat response that an able-bodied individual would experience.<sup>[39]</sup> Despite this, only one study  
32 to date has investigated the use of water sprays in individuals with a SCI, in particular athletes

1 with paraplegia. Pritchett et al.<sup>[39]</sup> conducted an incremental arm cranking protocol involving  
2 seven minute stages interspersed with 1 minute breaks when participants sprayed themselves *ad*  
3 *libitum*. The protocol was terminated voluntarily or when oesophageal temperature ( $T_{es}$ ) rose by  
4 more than  $0.2^{\circ}\text{C}/\text{min}^{-1}$ . There were no differences between water spraying and no cooling in  
5 heart rate (HR),  $T_{es}$ , rectal temperature ( $T_{re}$ ),  $T_{sk}$ , mean total work performed, rating of perceived  
6 exertion (RPE) or thermal sensation. Although the lack of positive findings may be due to the  
7 limited skin surface area covered by the water sprays (back of head, neck, forearm and face), due  
8 to the obstruction of clothing, this reflects the typical water spraying habits of wheelchair  
9 athletes. Interestingly, it has been suggested that a critical body surface cooling area of ~40% is  
10 needed to attenuate a rise in  $T_{core}$ ,<sup>[40]</sup> though this is rarely achieved by cooling methods used by  
11 athletes. It is unknown which factor plays the greatest role in a cooling method's ability to  
12 dissipate heat; the amount of body surface area cooled or the specific body location.<sup>[41]</sup>

13 The absence of positive findings in the work of Pritchett et al.<sup>[39]</sup> may also be due to a number of  
14 limitations of the study. Although the authors stated that the protocol and ambient conditions  
15 ( $22^{\circ}\text{C}$ , 45-50% relative humidity) were used to maintain high ecological validity, the exercise  
16 mode of arm cranking is not reflective of wheelchair court sports of an intermittent nature.  
17 Wheelchair propulsion has been shown to elicit lower physiological and thermal strain than arm  
18 cranking, potentially because the convective cooling currents generated by the arm action used  
19 to push the wheelchair increases the potential for peripheral heat loss.<sup>[42]</sup> Therefore, employing  
20 an exercise mode which does not reflect actual game play would consequently alter the  
21 ecological validity of the protocol. Secondly, players only experienced a slight increase in  $T_{core}$   
22 by the end of the protocol ( $0.5^{\circ}\text{C}$ ). This could be because exercising at an air temperature ( $22^{\circ}\text{C}$ )  
23 often classified as moderate,<sup>[43]</sup> trained individuals with paraplegia may be able to regulate heat  
24 loss similarly to an able-bodied counterpart in the absence of external heat gains.<sup>[10]</sup> As a high  
25  $T_{core}$  is crucial for a cooling strategy to be effective, a more demanding exercise protocol, akin to  
26 a wheelchair basketball match, may have elicited a larger cooling improvement from water  
27 spraying. On this note, whether a similar outcome would be observed for players with tetraplegia  
28 is unknown.

29 Remarkably, Pritchett et al.<sup>[39]</sup> found no improvement in perceptual measures (thermal sensation  
30 and RPE) in the cooling condition. Evidence for the responsiveness of the face to cooling has  
31 shown a two to fivefold suppression of thermal discomfort in the face compared to cooling other  
32 body regions,<sup>[44]</sup> indicating that water spraying the face may be beneficial for thermal comfort.  
33 Although considered to be a practical method frequently used by many wheelchair athletes,

1 caution is needed if thermal comfort is improved but there is no equivalent reduction in  $T_{\text{core}}$ , as  
2 athletes could potentially override signs of high levels of heat strain, putting them at risk of  
3 hyperthermia and heat illness.

### 4           **2.3 Cooling Garments**

5  
6 Lightweight cooling garments have received considerable interest in the able-bodied athletic  
7 population.<sup>[20-22, 29, 45-47]</sup> An advantage of these garments is their potential use prior to or during  
8 exercise. However, a disadvantage is the trade-off between a lightweight garment with a  
9 sufficient cooling capacity, in relation to the amount and type of coolant contained in the  
10 garment. Recent developments in design features include; close-fitting designs, silica based gels  
11 to promote greater evaporative cooling, phase change materials that operate close to normal skin  
12 temperature<sup>[48]</sup> and wet cooling to increase heat transfer between torso and vest.<sup>[35]</sup> Cooling  
13 garments typically weigh ~1-4 kg,<sup>[28]</sup> bulkier cooling systems would be counter-active to the  
14 technology of the modern sports wheelchair, produced to reduce mass and improve  
15 aerodynamics.<sup>[49]</sup>

16 In the able-bodied athletic population, research into the use of cooling garments has  
17 predominantly focused on ice vests as a pre-cooling method prior to endurance exercise.<sup>[35]</sup> Pre-  
18 cooling using an ice vest, usually results in relative reductions in  $T_{\text{sk}}$ <sup>[29]</sup> and body temperature,<sup>[50]</sup>  
19 in addition to skin blood flow,<sup>[50]</sup> with no change in  $T_{\text{core}}$ .<sup>[50, 51]</sup> This suggests that the transfer of  
20 heat mainly occurs at the skin and less so at the core. The use of cooling garments prior to  
21 intermittent exercise has produced mixed results, being dependent upon the ambient condition,  
22 with positive effects found in warm environments (30-34°C).<sup>[52, 53]</sup> The use of ice vests as a pre-  
23 cooling method has been shown to improve performance by 4.8%<sup>[53]</sup> but also provide limited  
24 efficacy compared to ice slurry ingestion.<sup>[54]</sup> Yet, meta-analysis data has found wearing ice vests  
25 during exercise has a significantly larger effect (ES = 4.64) than other cooling techniques, such  
26 as cold water ingestion (ES = 1.75) and cooling packs (ES = 0.39).<sup>[30]</sup> Research into wearing  
27 these garments during sport is, however, still limited, due to sporting regulations and difficulty  
28 in incorporating such garments into athletic clothing worn during competition.

29 Although not as heavily researched, cooling devices for the head and neck, such as cooling hats,  
30 neckbands and headpieces, are also commercially available.<sup>[55-58]</sup> The rationale for garments that  
31 cool the head is based on the concept that this body region contains a high concentration of  
32 thermoreceptors,<sup>[59]</sup> as demonstrated by the head feeling warmer than the rest of the body in

1 warm environments.<sup>[60]</sup> It should be noted, however, that the influence of head cooling and the  
2 extent to which brain temperature increases during exercise in the heat is unknown and needs to  
3 be further investigated.<sup>[61, 62]</sup> The neck area is a region of high thermosensitivity for both  
4 sudomotor and discomfort responses<sup>[44]</sup> yet few studies in the able-bodied have investigated  
5 cooling this region during exercise. The limited number of studies that have investigated the use  
6 of cooling neck collars have found a performance improvement whilst exercising in the heat  
7 (~30°C) and wearing the collars for 90 minutes during exercise (75 minutes at 60%  $\dot{V}O_{2peak}$   
8 followed by a 15 minute time trial).<sup>[57, 58]</sup> However, a shorter cooling exposure (during a 15  
9 minute time trial) had no effect on time trial performance.<sup>[57]</sup> This suggests that the small surface  
10 area of cooling provided by the neck collars may only have an ergogenic effect under high levels  
11 of thermal strain and when worn for a long duration, below which a performance benefit cannot  
12 be achieved.<sup>[57]</sup>

13 In the athletic SCI population, the use of a refrigerated headpiece and an ice vest used to cool  
14 during 30 minutes of wheelchair exercise at 10 km race pace (~70% maximal aerobic power) has  
15 shown to provide no thermal, perceptual or performance benefit compared to no cooling.<sup>[49]</sup> The  
16 lack of a performance benefit observed may be due to the cooling potential of the two garments  
17 (75 W and 117 W for the headpiece and ice vest, respectively) being considerably lower than the  
18 metabolic heat generated during the exercise (704-766 W). From thermal manikin work, this  
19 commercially available vest has been found to extract only ~70 W of heat from the torso over 45  
20 minutes at 26°C.<sup>[45]</sup> A greater cooling potential may have been achieved with additional ice  
21 packets in the vest covering a greater skin surface area.<sup>[49]</sup> However, this would increase the  
22 mass of the 3 kg garment, potentially hindering athletic performance. Practically, the length of  
23 time the ice vest stays cold also needs to be considered as once the ice strips of the vest have  
24 melted, cooling effectiveness is reduced. Therefore, applying this cooling method prior to  
25 exercise may be more practical and increase cooling effectiveness when compared to wearing  
26 the vest for a longer period of time during exercise.

27 Webborn et al.<sup>[63, 64]</sup> demonstrated promising results using ice vests prior to and during  
28 intermittent arm cranking exercise for athletes with tetraplegia. Twenty minutes of pre-cooling  
29 (PRE) and cooling during exercise (DUR) (warm-up and intermittent sprint protocol (ISP)) using  
30 an ice vest was shown to elicit lower  $T_{core}$  and thermal sensation throughout exercise, compared  
31 to no cooling. Despite  $T_{core}$  being greater in DUR throughout exercise, compared to PRE, a  
32 lower rate of increase resulted in similar  $T_{core}$  at the end of exercise. Pre-cooling demonstrated

1 central cooling through an offset in the absolute increase in  $T_{\text{core}}$ . Upon removal of the vest  $T_{\text{sk}}$   
2 increased at a greater rate than the other conditions, due to the greater thermal gradient between  
3 the environment and the skin, but still remained lower than the no cooling condition. In contrast,  
4 in DUR, initial peripheral cooling may have been apparent as direct heat transfer from the body  
5 core to the periphery was facilitated, demonstrated by the lower rate of increase in  $T_{\text{core}}$  resulting  
6 in a cooling of the skin. Therefore, whether cooling before or during exercise is the most  
7 appropriate form of cooling may depend on whether peripheral or core cooling is of importance.  
8 In terms of sprint performance, whilst there were no differences in overall work done during the  
9 14 sprint protocol,<sup>[63]</sup> the extended protocol of 30 sprints exhibited substantial differences  
10 between the conditions.<sup>[64]</sup> Athletes were able to complete more sprints in PRE and DUR than no  
11 cooling, with the greatest number of sprints completed in the latter. Peak power output  
12 significantly decreased in PRE (~15%) and the no cooling condition (~13%) over time, whilst  
13 this was not observed for DUR (~3%). The peak power output of the initial sprints in DUR were  
14 lower than the other conditions, which could be linked to the ability of the athletes to exercise  
15 for longer in DUR. Consequently, during the initial sprints the metabolic processes of the muscle  
16 will not have been stressed as greatly in DUR compared to the other two conditions, potentially  
17 allowing better recovery between sprints. This implies that cooling during exercise with an ice  
18 vest may not be ideal when the initial power output of sprints is essential to the overall  
19 performance. Participants also commented on the initial coldness of the vest, which may have  
20 affected their initial sprint performance from a psychological perspective.

21 To date, only one study has investigated the use of a cooling garment in a field based  
22 environment. Trbovich et al.<sup>[65]</sup> studied the use of a cooling vest containing phase change  
23 materials (reported to maintain at 15°C for 2-3 hours) during 60 minutes of wheelchair  
24 basketball or wheelchair rugby (21.1-23.9°C environmental temperature) in athletes with  
25 paraplegia and tetraplegia, respectively. A greater rise in  $T_{\text{core}}$  was experienced by the athletes  
26 with tetraplegia at the end of the 60 minutes compared to athletes with paraplegia, yet the  
27 cooling vest had no effect on  $T_{\text{core}}$  for either group. Even though this study may appear novel by  
28 exploring a cooling strategy in a field based environment for athletes with a SCI, a number of  
29 limitations are apparent which affects the application of the findings. Firstly, there was no  
30 control of or recording of exercise intensity. Secondly, although the sports are both of an  
31 intermittent nature using court sports wheelchairs, the kinematics of the sports differ,<sup>[66, 67]</sup> and a  
32 player's function and classification affects their speed and distance covered,<sup>[67]</sup> limiting the  
33 comparative nature of the study. Finally, irrespective of any thermal benefit, it is unclear if any



1 performance benefit was provided by the cooling vest, or whether  $T_{sk}$  or perceptual responses  
2 were affected.

3 To ensure an athlete gets the most benefit from a cooling strategy, methods need to be practiced  
4 prior to major competitions to enable the athlete to elicit a coupling between the potential for  
5 heat storage and thermal sensation.<sup>[16]</sup> Diaper et al.<sup>[55]</sup> investigated pre-cooling using an ice vest  
6 and the combination of pre-cooling using an ice vest and cooling hats and neckbands during  
7 intermittent exercise ( $\sim 30^{\circ}\text{C}$  environmental temperature) for a wheelchair tennis player prior to a  
8 Paralympic Games. The player exhibited a lower  $T_{core}$ , thermal sensation and faster average peak  
9 speed using the combination of methods compared to pre-cooling only.<sup>[55]</sup> However, the study  
10 did not measure any other thermal or physiological parameters or provide a control condition to  
11 determine whether pre-cooling only had any thermal benefit compared to no cooling. The  
12 thermoregulatory and physiological effects of pre-cooling have been shown to wane after 30-40  
13 minutes,<sup>[68, 69]</sup> therefore at the end of the 60 minute exercise protocol, the pre-cooling condition  
14 may have had limited benefit compared to no cooling. During a similar protocol and ambient  
15 conditions, Goosey-Tolfrey et al.<sup>[56]</sup> demonstrated the combined use of cooling hats and  
16 neckbands versus no cooling elicited lower RPE and thermal sensation during intermittent sprint  
17 exercise.<sup>[56]</sup> Although 42% less water was consumed during the cooling condition and sweat  
18 rates did not differ between conditions, athletes did maintain euhydration. Limited findings can  
19 be drawn in respect to the thermal benefit of this cooling strategy, as no thermal measures were  
20 taken, yet practically the cooling hats could be dual purpose (additional sun protection) during  
21 outdoor sports.

## 22 **3 Timing of cooling application**

### 23 **3.1 Pre-cooling**

24

25 Pre-cooling is based on the concept of starting exercise with a lowered body temperature to  
26 widen the temperature margin before reaching a critical limiting  $T_{core}$  contributing to fatigue.<sup>[70]</sup>  
27 It is generally accepted, in the able-bodied, that cooling prior to endurance and intermittent  
28 exercise elicits a performance improvement<sup>[31, 35, 53]</sup> ( $ES = 0.52$  and  $ES = 0.43$ , respectively),<sup>[53]</sup>  
29 yet performance benefits are limited when cooling occurs prior to sprint exercise ( $ES = 0.03$ ).<sup>[53]</sup>  
30 The extent of the performance improvement may also be dependent on the exercise duration,  
31 with the effect of pre-cooling decreasing for exercise durations over 60 minutes, potentially due

1 to the limited duration of the cooling effect.<sup>[53]</sup> To enhance the cooling effect, a combination of  
2 methods could be used. Applying these methods simultaneously could also limit the time before  
3 an event that needs to be dedicated to pre-cooling.<sup>[35]</sup>

4 To maintain a physiological advantage and enhance performance pre-cooling needs to be  
5 provided as close to the start of exercise as possible and potentially during the warm-up.<sup>[20, 63]</sup> In  
6 light of this, previous cooling research in the able-bodied has sometimes failed to account for the  
7 effect of athletes' warm-up prior to competition thus limiting its application in a field  
8 environment. Nevertheless, pre-cooling during a warm-up will depend on the body area cooled  
9 and the method of cooling. A heavy cooling garment worn during the warm-up may also affect  
10 the metabolic cost of exercise, with the intensity of the warm-up needing to be adjusted  
11 accordingly.<sup>[20]</sup> Lastly, the cooling intensity of the strategy is particularly important as a sharp  
12 decrease in  $T_{\text{core}}$ ,  $T_{\text{sk}}$  or muscle temperature prior to exercise may have a detrimental effect on  
13 performance.<sup>[53, 71, 72]</sup>

14 Two of the three studies reviewed utilising cooling with ice vests prior to intermittent sprint  
15 exercise observed a performance and thermal benefit compared to no cooling.<sup>[63, 64]</sup> A  
16 combination of pre-cooling using ice vests and cooling hats and neckbands during exercise, to  
17 attenuate the rise in  $T_{\text{core}}$  and thermal sensation, has also been shown to be effective as a case  
18 study.<sup>[55]</sup> Therefore, there may be some justification for pre-cooling using ice vests in athletes  
19 with a SCI, though further work is greatly needed.

### 20 **3.2 Cooling during exercise/ rest periods**

21  
22 To offset the gains in heat throughout exercise, cooling during exercise has been utilised,  
23 predominantly through the use of ice vests. As previously mentioned, the difficulty with this  
24 cooling technique is to avoid an adverse effect on performance due to the size, coverage and  
25 weight of the cooling garment. Hence, due to this complexity, cooling during exercise has  
26 received less research attention than pre-cooling. Although research in the able-bodied shows  
27 that it could be just as effective as pre-cooling in improving performance.<sup>[30]</sup>

28 In addition to cooling during exercise, cooling provided during rest periods between multiple  
29 bouts of exercise could be viable for many team sports i.e. wheelchair courts sports. Strategies  
30 employed during these rest periods need to ensure minimal influence on athlete preparation. Due  
31 to the limited time athletes are allowed during rest periods/half-time (2-15 minutes depending on

1 the wheelchair court sport) this may not allow for enough cooling exposure, but may provide  
2 some benefit compared to no cooling at all. Therefore, to enhance the cooling effect, a  
3 combination of both pre-cooling and cooling during rest periods/half-time may be more effective  
4 at reducing thermal strain by offsetting heat storage during the subsequent bout of exercise.<sup>[73]</sup>

5 Cooling during exercise was utilised in seven reviewed studies<sup>[37, 49, 55, 56, 63-65]</sup>  
6 with mixed results. Either performance was improved or  $T_{\text{core}}$  reduced when using ice vests,  
7 cooling hats and neckbands during intermittent exercise for trained wheelchair athletes.<sup>[55, 63, 64]</sup>  
8 Though, in a field based environment a cooling vest had no effect on  $T_{\text{core}}$  during a game of  
9 wheelchair basketball or wheelchair rugby.<sup>[65]</sup> During continuous exercise, foot cooling was  
10 thermally beneficial to untrained individuals, yet a refrigerated headpiece and ice vest provided  
11 no benefit to trained wheelchair athletes, due to the low cooling potential of the garments.<sup>[37, 49]</sup>  
12 In the two studies that applied cooling during rest periods, the use of water sprays had no  
13 thermal or performance effect whilst hand cooling provided more positive findings, though as  
14 previously mentioned, this method may elicit practical issues for athletes with a SCI.<sup>[36, 39]</sup>

#### 15 **4 Summary of cooling methods and timing of cooling application**

16

17 In comparison to the advice that can be gathered from the literature for able-bodied athletes it is  
18 difficult to establish the optimal cooling method for athletes with a SCI. This is predominantly  
19 due to the lack of studies that have investigated the use of cooling strategies in this population  
20 leading to a lack of evidence for each method (Table 2). Nevertheless, from the studies reviewed  
21 it would appear that wearing an ice vest during intermittent sprint exercise has beneficial effects  
22 both in terms of decreasing thermal strain and enhancing performance,<sup>[64]</sup> though contrasting  
23 findings have been observed during steady state exercise.<sup>[49]</sup> Drawing on findings from the able-  
24 bodied, the use of ice vests has been shown to be a feasible method of pre-cooling, with a 4.8%  
25 improvement in performance<sup>[53]</sup> in addition to improved performance from wearing the garments  
26 during exercise in the heat.<sup>[30]</sup> A combination of pre-cooling and cooling during exercise or half-  
27 time cooling might also be an effective strategy.<sup>[30]</sup>

28 Although not currently studied in athletes with a SCI, the consumption of cold beverages, in  
29 particular ice slurries, have been shown to be an effective pre-cooling method in the able-bodied  
30 by lowering  $T_{\text{core}}$  and improving endurance performance in the heat.<sup>[74]</sup> Further investigation into  
31 this internal method is warranted but could be studied as a practical cooling alternative for

1 athletes with a SCI. The specific sporting regulations and logistics of administering any cooling  
2 method for athletes with a SCI in wheelchair court sports needs to be carefully considered.

### 3 **5 Limitations and future research**

4  
5 The majority of studies reviewed had a SCI group consisting of individuals with paraplegia and  
6 tetraplegia, potentially affecting the groups mean thermal response, due to an individual's  
7 thermoregulatory response being proportional to lesion level.<sup>[10]</sup> Thermoregulatory responses  
8 may also depend on the completeness of the injury<sup>[75]</sup> with incomplete lesions resulting in a  
9 greater amount of sensory information about their thermal state and a greater capacity to  
10 sweat.<sup>[64]</sup> Therefore, studies should separate individuals with paraplegia and tetraplegia in  
11 population groups and note the number of individuals with incomplete lesions, providing  
12 justification for their inclusion in the sample.

13 Overall the studies reviewed neglected to employ a standard measure of performance specific to  
14 the sport, making it difficult to determine if there is a clear performance benefit associated with a  
15 cooling strategy. For application in wheelchair court sports future research should consider  
16 employing an ISP that reflects the nature and exercise intensity of the sport. The mode of  
17 exercise also needs to be appropriate as differences in thermoregulation are elicited depending  
18 on the modality, i.e. arm crank vs. wheelchair ergometry.<sup>[42]</sup>

19 Only one of the eight studies reviewed mentioned the use of a warm-up and the application of  
20 cooling during this time. Often an athlete's warm-up is performed at intensities (~40-60%  
21  $\dot{V}O_{2max}$ , able-bodied research)<sup>[76]</sup> that result in an increase in  $T_{core}$ . Therefore, excluding a warm-  
22 up from the experimental design could affect the practical application of the findings of the  
23 reviewed studies. Prospective studies need to ensure the inclusion of a warm-up and, if pre-  
24 cooling is being administered, the wearing of cooling garments during this time, if appropriate.

25 Future studies need to be as closely matched as possible to the intended environmental playing  
26 conditions. In relation to wheelchair court sports, athletes with a SCI predominantly compete in:  
27 wheelchair tennis, wheelchair rugby or wheelchair basketball. The latter two sports are played  
28 indoors, suggesting that it should be possible to closely control the ambient playing conditions.  
29 How strictly these indoor conditions are regulated is unclear, i.e. air conditioned, with anecdotal  
30 evidence suggesting playing conditions vary greatly between training centres and competition

1 venues. In addition to this, the increased potential for thermal strain owing to the intermittent  
2 sprint nature of the sports and the impaired thermoregulatory response of individuals with a SCI  
3 suggest that cooling strategies could be highly beneficial for athletes with a SCI in these sports.  
4 In wheelchair tennis, guidelines are put in place by the International Tennis Federation (ITF) to  
5 ensure wheelchair players are not put at an increased risk of heat injury. Guidelines include a  
6 wet bulb globe temperature index (WBGT) limit of 28°C, provision of shade and ice buckets on  
7 court and scheduling of matches morning or late afternoon/evening.<sup>[77]</sup> It should be noted that to  
8 implement a structured pre-cooling strategy in this sport may be potentially difficult due to the  
9 unknown start time caused by previous extended play. In all wheelchair sports, due to the  
10 practical problems of travelling to venues, athletes with a SCI may arrive at a match with an  
11 elevated  $T_{\text{core}}$ , especially in high ambient temperatures and/or humidity. Without sufficient  
12 cooling or recovery this may limit their temperature margin before reaching a critical  $T_{\text{core}}$  during  
13 exercise.

14 Future studies are required to establish the transferability of practical cooling strategies in a  
15 laboratory to the field environment.<sup>[35]</sup> To achieve this, future work should attempt to replicate  
16 actual competition and practical limitations experienced in “real-life” situations. This includes  
17 participants of a highly trained and motivated nature to ensure research findings can be  
18 translated into meaningful performance enhancements<sup>[35]</sup> in the athletic SCI population.

## 19 **6 Practical recommendations**

20

21 While research is lacking and inconclusive, several practical recommendations can be suggested.

- 22 • Sport scientists must adopt an individualised approach and refine the cooling strategy  
23 based on the sport and the athlete’s level and completeness of lesion.
- 24 • The sporting regulations, effectiveness, practicality, staffing/assistance, logistics, cost  
25 and athlete comfort of a cooling strategy need to be considered. For example, hand  
26 cooling may be impractical for some athletes with a SCI due to their limited hand grip  
27 function and the use of gloves during play.
- 28 • Athletes and coaches must trial cooling strategies prior to their use in competition to  
29 determine the suitability of a strategy. For instance, the appropriate fit of an ice vest

1 would need to be determined to ensure upper body mobility is not affected and  
2 abdominal binding/strapping can still be used.

- 3 • The cooling power of a cooling method must be considered and where possible a  
4 combination of practical and portable cooling methods should be used to increase the  
5 effectiveness of a cooling strategy. One example would be to use an ice vest prior to a  
6 match and water sprays during rest periods/half-time. Which combination of cooling is  
7 the most effective would heavily depend on the practicality of the methods for the  
8 particular sport.
- 9 • Coaches and support staff need to be educated on the signs and symptoms of heat illness.  
10 This will enable them to recognise when a player needs to stop playing, be removed from  
11 the environment and cooled with an appropriate cooling strategy. Notably, Webborn et  
12 al.<sup>[78]</sup> stopped two participants during the ISP as their  $T_{core}$  had reached the safety limit  
13 (39.3°C or a 2°C increase from rest). Therefore, it is crucial that coaches and support  
14 staff can identify players with heat illness symptoms, such as extreme fatigue and  
15 dizziness, which will greatly affect their decision making ability during play.
- 16 • It is imperative that sports venues are air conditioned and ice is provided at the  
17 competition venues. Water sprays and fans should also be provided for use during time-  
18 out/ half-time. Both techniques should lower perceptual responses of thermal sensation  
19 and comfort, whilst fanning should increase heat loss by convection.<sup>[79]</sup> Although, if  
20 athletes are substituted for a long period of time they should be careful not to overcool,  
21 especially as athletes with a SCI possess a smaller surface area for temperature  
22 perception.
- 23 • Athletes must ensure that they are hydrated and ingest adequate fluid during play.  
24 However, to avoid autonomic dysreflexia athletes should ensure they void their bladder  
25 prior to play.
- 26 • An acclimation strategy must be considered for athletes who compete outdoors or who  
27 are likely to spend considerable time transferring to sports venues (e.g., competitor at the  
28 Paralympic Games). Partial heat acclimation may be possible for athletes with a SCI. For  
29 instance, Castle et al.<sup>[80]</sup> demonstrated a lower resting  $T_{core}$  and increased plasma volume,  
30 but no improvement in sweat response, following 7 days of heat acclimation in athletes  
31 with a SCI.

- Medications and sleep deprivation may affect an athlete's heat tolerance<sup>[43]</sup> and should be addressed with the athlete to minimise the risk when the athlete is travelling or competing in a warm environment.

## 7 Conclusion

Due to the thermoregulatory impairment of athletes with a SCI, employing cooling strategies in this population group could be beneficial to offset any decrement in performance. From the studies reviewed wearing an ice vest during intermittent sprint exercise has a beneficial effect on both decreasing thermal strain and enhancing performance. These garments have also been shown to be effectively used prior to and during exercise in the able-bodied population. Drawing on additional findings from the able-bodied, a combination of pre-cooling and cooling during exercise or half-time cooling might increase the effectiveness of a strategy.<sup>[30, 31]</sup> However, to date, there has been a lack of studies addressing fundamental issues regarding which cooling strategy is the most appropriate in terms of method, timing, duration, and body region cooled in a competition environment for athletes with a SCI. Future studies should employ cooling strategies under constraints of actual competition with highly trained athletes with a SCI and appropriate exercise protocols to ensure research findings can be translated into meaningful performance improvements. The specific sporting regulations and logistics of administering a cooling method in wheelchair court sports need to be carefully considered. Sport scientists and coaches should take an individualised approach when refining a strategy and conduct their own tests of a cooling strategy prior to its use in competition.

## 1 **References**

- 2 1. Galloway SD, Maughan RJ. Effects of ambient temperature on the capacity to perform  
3 prolonged cycle exercise in man. *Med Sci Sports Exerc.* 1997;29(9):1240-9.
- 4 2. Nielsen B. Olympics in Atlanta: A fight against physics. *Med Sci Sports Exerc.*  
5 1996;28(6):665-8.
- 6 3. Borresen J. Environmental considerations for athletic performance at the 2008 Beijing  
7 Olympic Games. *Int Sport Med J.* 2008;9(2):44-55.
- 8 4. INMET. Rio 2016. Climate Information [online]. Available from URL:  
9 <http://www.rio2016.com/pregamestraining/en/climate-information/rio-de-janeiro> [Accessed 10  
10 Jun 2013]
- 11 5. Freund PR, Brengelmann GL, Rowell LB, et al. Attenuated skin blood flow response to  
12 hyperthermia in paraplegic men. *J Appl Physiol.* 1984;56(4):1104-9.
- 13 6. Rawson RO, Hardy JD. Sweat inhibition by cutaneous cooling in normal sympathectomized  
14 and paraplegic man. *J Appl Physiol.* 1967;22(2):287-91.
- 15 7. Tam HS, Darling RC, Cheh HY, et al. Sweating response: A means of evaluating the set-point  
16 theory during exercise. *J Appl Physiol.* 1978;45(3):451-8.
- 17 8. Hopman MT. Circulatory responses during arm exercise in individuals with paraplegia. *Int J*  
18 *Sports Med.* 1994;15(3):126-31.
- 19 9. Normell LA. Distribution of impaired cutaneous vasomotor and sudomotor function in  
20 paraplegic man. *Scand J Clin Lab Invest Suppl.* 1974;138:25-41.
- 21 10. Price MJ. Thermoregulation during exercise in individuals with spinal cord injuries. *Sports*  
22 *Med.* 2006;36(10):863-79.
- 23 11. Price MJ, Campbell IG. Thermoregulatory responses of spinal cord injured and able-bodied  
24 athletes to prolonged upper body exercise and recovery. *Spinal Cord.* 1999;37(11):772-9.



- 1 12. Price MJ, Campbell IG. Thermoregulatory responses of paraplegic and able-bodied athletes  
2 at rest and during prolonged upper body exercise and passive recovery. *Eur J Appl Physiol*  
3 *Occup Physiol.* 1997;76(6):552-60.
- 4 13. Hopman MTE, Binkhorst RA. Heat balance during exercise in a hot and cold environment in  
5 spinal cord injured individuals. In: van der Woude, L.H.V, Hopman MTE, van Kemenade CH,  
6 editors. *Biomedical aspects of wheelchair propulsion: the state of the art II.* Amsterdam: IOS  
7 Press; 1998. pg. 333-5.
- 8 14. Guttman L, Silver J, Wyndham CH. Thermoregulation in spinal man. *J Physiol.*  
9 1958;142(3):406-19.
- 10 15. Price MJ, Campbell IG. Effects of spinal cord lesion level upon thermoregulation during  
11 exercise in the heat. *Med Sci Sports Exerc.* 2003;35(7):1100-7.
- 12 16. Ross ML, Garvican LA, Jeacocke NA, et al. Novel precooling strategy enhances time trial  
13 cycling in the heat. *Med Sci Sports Exerc.* 2011;43(1):123-33.
- 14 17. Siegel R, Mate J, Brearley MB, et al. Ice slurry ingestion increases core temperature capacity  
15 and running time in the heat. *Med Sci Sports Exerc.* 2010;42(4):717-25.
- 16 18. Lee DT, Haymes EM. Exercise duration and thermoregulatory responses after whole body  
17 precooling. *J Appl Physiol.* 1995;79(6):1971-6.
- 18 19. Olschewski H, Bruck K. Thermoregulatory, cardiovascular, and muscular factors related to  
19 exercise after precooling. *J Appl Physiol.* 1988;64(2):803-11.
- 20 20. Arngrimsson SA, Petitt DS, Stueck MG, et al. Cooling vest worn during active warm-up  
21 improves 5-km run performance in the heat. *J Appl Physiol.* 2004;96(5):1867-74.
- 22 21. Duffield R, Dawson B, Bishop D, et al. Effect of wearing an ice cooling jacket on repeat  
23 sprint performance in warm/humid conditions. *Br J Sports Med.* 2003;37(2):164-9.
- 24 22. Duffield R, Marino FE. Effects of pre-cooling procedures on intermittent-sprint exercise  
25 performance in warm conditions. *Eur J Appl Physiol.* 2007;100(6):727-35.

- 1 23. Booth J, Marino F, Ward JJ. Improved running performance in hot humid conditions  
2 following whole body precooling. *Med Sci Sports Exerc.* 1997;29(7):943-9.
- 3 24. Booth J, Wilsmore BR, Macdonald AD, et al. Whole-body pre-cooling does not alter human  
4 muscle metabolism during sub-maximal exercise in the heat. *Eur J Appl Physiol.*  
5 2001;84(6):587-90.
- 6 25. Kay D, Taaffe DR, Marino FE. Whole-body pre-cooling and heat storage during self-paced  
7 cycling performance in warm humid conditions. *J Sports Sci.* 1999;17(12):937-44.
- 8 26. Grahn DAD, Murray JVJ, Heller HCH. Cooling via one hand improves physical  
9 performance in heat-sensitive individuals with multiple sclerosis: A preliminary study. *BMC*  
10 *Neurology.* 2008;8:14.
- 11 27. House JR, Holmes C, Allsopp AJ. Prevention of heat strain by immersing the hands and  
12 forearms in water. *J R Nav Med Serv.* 1997;83(1):26-30.
- 13 28. Barwood MJ, Davey S, House JR, et al. Post-exercise cooling techniques in hot, humid  
14 conditions. *Eur J Appl Physiol.* 2009;107(4):385-96.
- 15 29. Quod MJ, Martin DT, Laursen PB, et al. Practical precooling: Effect on cycling time trial  
16 performance in warm conditions. *J Sports Sci.* 2008;26(14):1477-87.
- 17 30. Bongers CCWG, Thijssen DHJ, Veltmeijer MTW, et al. Precooling and percooling (cooling  
18 during exercise) both improve performance in the heat: A meta-analytical review. *Br J Sports*  
19 *Med.* Epub 2014 Apr 19.
- 20 31. Tyler CJ, Sunderland C, Cheung SS. The effect of cooling prior to and during exercise on  
21 exercise performance and capacity in the heat: A meta-analysis. *Br J Sports Med.* Epub 2013  
22 Aug 14.
- 23 32. Cheung SS. *Advanced environmental exercise physiology.* Champaign, IL: Human Kinetics;  
24 2010.
- 25 33. Livingstone SD, Nolan RW, Cattroll SW. Heat loss caused by immersing the hands in water.  
26 *Aviat Space Environ Med.* 1989;60(12):1166-71.

- 1 34. Livingstone SD, Nolan RW, Keefe AA. Heat loss caused by cooling the feet. *Aviat Space*  
2 *Environ Med.* 1995;66(3):232-7.
- 3 35. Ross M, Abbiss C, Laursen P, et al. Precooling methods and their effects on athletic  
4 performance : A systematic review and practical applications. *Sports Med.* 2013;43(3):207-25.
- 5 36. Goosey-Tolfrey V, Swainson M, Boyd C, et al. The effectiveness of hand cooling at  
6 reducing exercise-induced hyperthermia and improving distance-race performance in wheelchair  
7 and able-bodied athletes. *J Appl Physiol.* 2008;105(1):37-43.
- 8 37. Hagobian TA, Jacobs KA, Kiratli BJ, et al. Foot cooling reduces exercise-induced  
9 hyperthermia in men with spinal cord injury. *Med Sci Sports Exerc.* 2004;36(3):411-7.
- 10 38. Mora-Rodriguez R. Influence of aerobic fitness on thermoregulation during exercise in the  
11 heat. *Exerc Sport Sci Rev.* 2012;40(2):79-87.
- 12 39. Pritchett RC, Bishop PA, Yang Z, et al. Evaluation of artificial sweat in athletes with spinal  
13 cord injuries. *Eur J Appl Physiol.* 2010;109(1):125-31.
- 14 40. Kume M, Yoshida T, Tsuneoka H, et al. Relationship between body surface cooling area,  
15 cooling capacity, and thermoregulatory responses wearing water perfused suits during exercise  
16 in humans. *Jpn J Phys Fit Sports Med.* 2009;58(1):109-21.
- 17 41. Young AJ, Sawka MN, Epstein Y, et al. Cooling different body surfaces during upper and  
18 lower body exercise. *J Appl Physiol.* 1987;63(3):1218-23.
- 19 42. Price M, Campbell I. Thermoregulatory and physiological responses of wheelchair athletes  
20 to prolonged arm crank and wheelchair exercise. *Int J Sports Med.* 1999;20(7):457-63.
- 21 43. American College of Sports Medicine, Armstrong LE, Casa DJ, et al. American college of  
22 sports medicine position stand. exertional heat illness during training and competition. *Med Sci*  
23 *Sports Exerc.* 2007;39(3):556-72.
- 24 44. Cotter JD, Taylor NA. The distribution of cutaneous sudomotor and alliesthesial  
25 thermosensitivity in mildly heat-stressed humans: An open-loop approach. *J Physiol.*  
26 2005;565(Pt 1):335-45.

- 1 45. Bogerd N, Psikuta A, Daanen HAM, et al. How to measure thermal effects of personal  
2 cooling systems: Human, thermal manikin and human simulator study. *Physiol Meas.*  
3 2010;31(9):1161-8.
- 4 46. Hornery DJ, Papalia S, Mujika I, et al. Physiological and performance benefits of halftime  
5 cooling. *J Sci Med Sport.* 2005;8(1):15-25.
- 6 47. Cotter JD, Sleivert GG, Roberts WS, et al. Effect of pre-cooling, with and without thigh  
7 cooling, on strain and endurance exercise performance in the heat. *Comp Biochem Physiol A*  
8 *Mol Integr Physiol.* 2001;128(4):667-77.
- 9 48. Tate M, Forster D, Mainwaring DE. Influence of garment design on elite athlete cooling.  
10 *Sports Technology.* 2010;1(2-3):117-24.
- 11 49. Armstrong LEL, Maresh CMC, Riebe DD, et al. Local cooling in wheelchair athletes during  
12 exercise-heat stress. *Med Sci Sports Exerc.* 1995;27(2):211-6.
- 13 50. Bogerd N, Perret C, Bogerd CP, et al. The effect of pre-cooling intensity on cooling  
14 efficiency and exercise performance. *J Sports Sci.* 2010;28(7):771-9.
- 15 51. Minett GM, Duffield R, Marino FE, et al. Volume-dependent response of precooling for  
16 intermittent-sprint exercise in the heat. *Med Sci Sports Exerc.* 2011;43(9):1760-9.
- 17 52. Cheung S, Robinson A. The influence of upper-body pre-cooling on repeated sprint  
18 performance in moderate ambient temperatures. *J Sports Sci.* 2004;22(7):605-12.
- 19 53. Wegmann M, Faude O, Poppendieck W, et al. Pre-cooling and sports performance: A meta-  
20 analytical review. *Sports Med.* 2012;42(7):545-64.
- 21 54. Jones PR, Barton C, Morrissey D, et al. Pre-cooling for endurance exercise performance in  
22 the heat: A systematic review. *BMC Med.* 2012;10:166,7015-10-166.
- 23 55. Diaper NJ, Goosey-Tolfrey V. A physiological case study of a paralympic wheelchair tennis  
24 player: Reflective practise. *J Sports Sci Med.* 2009;8(2):300-7.
- 25 56. Goosey-Tolfrey VL, Diaper NJ, Crosland J, et al. Fluid intake during wheelchair exercise in  
26 the heat: Effects of localized cooling garments. *Int J Sports Physiol Perform.* 2008;3(2):145-56.

- 1 57. Tyler CJ, Wild P, Sunderland C. Practical neck cooling and time-trial running performance  
2 in a hot environment. *Eur J Appl Physiol.* 2010;110(5):1063-74.
- 3 58. Tyler CJ, Sunderland C. Neck cooling and running performance in the heat: Single versus  
4 repeated application. *Med Sci Sports Exerc.* 2011;43(12):2388-95.
- 5 59. Nakamura M, Yoda T, Crawshaw LI, et al. Regional differences in temperature sensation  
6 and thermal comfort in humans. *J Appl Physiol.* 2008;105(6):1897-906.
- 7 60. Arens E, Zhang H, Huizenga C. Partial- and whole-body thermal sensation and comfort—  
8 part I: Uniform environmental conditions. *J Therm Biol.* 2006;31(1–2):53-9.
- 9 61. Brengelmann GL. Specialized brain cooling in humans? *FASEB J.* 1993;7(12):1148-53.
- 10 62. Cabanac M, Caputa M. Natural selective cooling of the human brain: Evidence of its  
11 occurrence and magnitude. *J Physiol.* 1979;286:255-64.
- 12 63. Webborn N, Price MJ, Castle PC, et al. Effects of two cooling strategies on thermoregulatory  
13 responses of tetraplegic athletes during repeated intermittent exercise in the heat. *J Appl Physiol.*  
14 2005;98(6):2101-7.
- 15 64. Webborn N, Price MJ, Castle P, et al. Cooling strategies improve intermittent sprint  
16 performance in the heat of athletes with tetraplegia. *Br J Sports Med.* 2010;44(6):455-60.
- 17 65. Trbovich M, Ortega C, Schroeder J, et al. Effect of a cooling vest on core temperature in  
18 athletes with and without spinal cord injury. *Top Spinal Cord Inj Rehabil.* 2014;20(1):70-80.
- 19 66. Coutts KD. Dynamics of wheelchair basketball. *Med Sci Sports Exerc.* 1992;24(2):231-4.
- 20 67. Sarro KJ, Misuta MS, Burkett B, et al. Tracking of wheelchair rugby players in the 2008  
21 demolition derby final. *J Sports Sci.* 2010;28(2):193-200.
- 22 68. Hessemer V, Langusch D, Bruck LK, et al. Effect of slightly lowered body temperatures on  
23 endurance performance in humans. *J Appl Physiol.* 1984;57(6):1731-7.
- 24 69. Wilson TE, Johnson SC, Petajan JH, et al. Thermal regulatory responses to submaximal  
25 cycling following lower-body cooling in humans. *Eur J Appl Physiol.* 2002;88(1-2):67-75.

- 1 70. Quod MJ, Martin DT, Laursen PB. Cooling athletes before competition in the heat:  
2 Comparison of techniques and practical considerations. *Sports Med.* 2006;36(8):671-82.
- 3 71. Faulkner SH, Ferguson RA, Gerrett N, et al. Reducing muscle temperature drop post warm-  
4 up improves sprint cycling performance. *Med Sci Sports Exerc.* 2013;45(2):359-63.
- 5 72. Sleivert GG, Cotter JD, Roberts WS, et al. The influence of whole-body vs. torso pre-cooling  
6 on physiological strain and performance of high-intensity exercise in the heat. *Comp Biochem*  
7 *Physiol A Mol Integr Physiol.* 2001;128(4):657-66.
- 8 73. Price MJ, Boyd C, Goosey-Tolfrey VL. The physiological effects of pre-event and mid-event  
9 cooling during intermittent running in the heat in elite female soccer players. *Appl Physiol Nutr*  
10 *Metab.* 2009;34(5):942-9.
- 11 74. Siegel R, Laursen PB. Keeping your cool: Possible mechanisms for enhanced exercise  
12 performance in the heat with internal cooling methods. *Sports Med.* 2012;42(2):89-98.
- 13 75. Theisen D. Cardiovascular determinants of exercise capacity in the paralympic athlete with  
14 spinal cord injury. *Exp Physiol.* 2012;97(3):319-24.
- 15 76. Woods K, Bishop P, Jones E. Warm-up and stretching in the prevention of muscular injury.  
16 *Sports Med.* 2007;37(12):1089-99.
- 17 77. International Tennis Federation. International tennis federation, regulations for wheelchair  
18 tennis 2013. [online]. Available from URL: <http://www.itftennis.com/media/137505/137505.pdf>  
19 [Accessed 2013 Jun 13]
- 20 78. Webborn N, Price MJ, Castle P, et al. Cooling strategies improve intermittent sprint  
21 performance in the heat of athletes with tetraplegia. *Br J Sports Med.* 2010;44(6):455-60.
- 22 79. Kato M, Sugeno J, Matsumoto T, et al. The effects of facial fanning on thermal comfort  
23 sensation during hyperthermia. *Pflugers Arch.* 2001;443(2):175-9.
- 24 80. Castle PC, Kularatne BP, Brewer J, et al. Partial heat acclimation of athletes with spinal cord  
25 lesion. *Eur J Appl Physiol.* 2013;113(1):109-15.

