

**1 Epidemiological Evaluation of Meniscal Ramp Lesions in 3214 ACL-Injured Knees**  
**2 from the (X) Database: A Risk Factor Analysis and Study of Secondary Meniscectomy**  
**3 Rates Following 769 Ramp Repairs.**

4

5

6

7 **Abstract**

8 **Background:** Ramp lesions are characterized by disruption of the peripheral  
9 meniscocapsular attachments of the posterior horn of the medial meniscus. Ramp repair  
10 performed at the time of ACL reconstruction has been shown to improve knee biomechanics.

11 **Hypothesis/Purpose:** Primary objectives of this study were to evaluate the **incidence** and  
12 risk factors for ramp lesions in a large series of patients undergoing ACL reconstruction,  
13 Secondary objectives were to determine the re-operation rate for failure of ramp repair,  
14 defined by subsequent re-operations for partial medial meniscectomy

15 **Study Design:** Case series

16 **Methods:** All patients underwent trans-notch posteromedial compartment evaluation of the  
17 knee during ACL reconstruction. Ramp repair was performed if a lesion was detected.  
18 Potentially important risk factors were analyzed for their association with ramp lesions. A  
19 secondary analysis of all patients who underwent ramp repair and had a minimum follow-up  
20 of two years was undertaken in order to determine the secondary partial meniscectomy rate  
21 for failed ramp repair.

22 **Results:** The overall **incidence** of ramp lesions in the study population was 23.9% (769 ramp  
23 lesions in 3214 patients). Multivariate analysis demonstrated that the presence of ramp  
24 lesions was significantly associated with the following risk factors: male gender, patients  
25 aged under 30 years, revision ACLR, chronic injuries, pre-operative side-to-side laxity >6  
26 mm and the presence of concomitant lateral meniscus tears. The secondary meniscectomy  
27 rate was 10.8% at a mean follow up of 45.6 months (24.2-66.2). Patients who underwent  
28 ACLR + ALLR had a greater than 2-fold reduction in the risk of reoperation for failure of  
29 ramp repair as compared with patients who underwent isolated ACLR (hazard ratio, 0.457;  
30 95%CI, 0.226-0.864;  $P = .021$ )

31 **Conclusion:** There is a high **incidence** of ramp lesions in patients undergoing ACLR. The  
32 identification of important risk factors for ramp lesions in this study in an individual patient  
33 should help raise an appropriate index of suspicion and prompt posteromedial compartment  
34 evaluation. The overall secondary partial meniscectomy rate after ramp repair is 10.8%.  
35 Anterolateral ligament reconstruction appears to confer a protective effect on the ramp repair  
36 performed at the time of ACLR and results in a significant reduction in secondary  
37 meniscectomy rates.

38

39 **Key Terms:** Ramp lesions. ACL, ACLR, ALL, ALLR, Meniscus. Meniscus repair

40

41 **What is known about the subject:**

42 Ramp lesions are important because they have adverse effects on the stability and  
43 biomechanics of the ACL injured knee. It is known that isolated ACLR fails to restore this  
44 fully in the presence of a ramp lesion, but that when ramp repair is performed concurrently,  
45 normal stability can be restored.

46

47 Although previous studies have reported the **incidence** of ramp lesions in the ACL injured  
48 knee, the majority have had very small sample sizes and therefore it is difficult to hold great  
49 confidence that they reliably estimate the true **incidence**. The same comment can be made  
50 regarding previously reported risk factors for ramp lesions.

51

52 There is very little published in the literature regarding failure rates of ramp repair.

53

54

55

56 **What this study adds to existing knowledge:**

57 To the knowledge of the authors this study is considerably larger than any other  
58 epidemiological and risk factor evaluation of ramp lesions in ACL injured knees. It is our  
59 opinion that the size of this series helps to give confidence that the **incidence** of ramp lesions  
60 seen in this study is a reliable estimate of the true **incidence**. Furthermore, this study has been  
61 able to confirm that many previously reported potential risk factors are significantly  
62 associated with ramp lesions but refute others which have been proposed on the basis of  
63 studies that were likely hindered by small sample sizes.

64

65 This study also adds to existing knowledge by reporting secondary meniscectomy rates after  
66 ramp repair and also demonstrating that anterolateral ligament reconstruction confers a  
67 protective effect on ramp repairs, as evidence by a significant reduction in secondary  
68 meniscectomy rates.

69

70

71 **Introduction**

72 Meniscal ramp lesions are typically associated with anterior cruciate ligament (ACL)  
73 deficiency. They are characterized by a disruption or tear of the peripheral meniscocapsular  
74 attachments of the posterior horn of the medial meniscus.<sup>7</sup> The term “ramp lesion” was first  
75 attributed to this injury pattern by Strobel in the 1980’s,<sup>44</sup> and is useful for differentiating this  
76 particular tear morphology from other types of longitudinal posterior horn tear. Despite the  
77 long history of recognition of ramp lesions, it is evident that the risk factors for developing  
78 this type of injury, the **incidence**, and the outcomes of treatment remain incompletely defined.  
79 This is partly due to the small populations evaluated in previous reports. As a result, the study  
80 of ramp lesions continues to be a subject of great interest.<sup>33,35</sup>

81

82 The biomechanical importance of these lesions has been demonstrated by cadaveric studies  
83 that have performed posterior meniscocapsular sectioning in the ACL-deficient knee. These  
84 studies have demonstrated that ramp lesions are associated with increases in both anterior  
85 tibial translation and external rotation.<sup>1,29,43</sup> More importantly, from the perspective of  
86 clinical applicability, these studies have also demonstrated restoration of knee biomechanics  
87 after meniscocapsular lesion repair.<sup>1,43</sup> It is therefore considered important to identify these  
88 lesions in order to repair them when necessary. However, it should be noted that historically  
89 these injuries were probably under-appreciated because pre-operative examination<sup>50</sup> and  
90 imaging modalities<sup>3,6,15,20,37</sup> have a low sensitivity for ramp lesions. Furthermore, a  
91 substantial number of these lesions may also be missed at the time of arthroscopic evaluation,  
92 particularly if this is performed using standard anterior portal viewing only.<sup>40</sup> In order to  
93 minimize the risk of missed diagnoses of ramp lesions, it is imperative to undertake a  
94 systematic arthroscopic examination, including that of the posteromedial compartment.

95

96 The primary objectives of this study were to evaluate the **incidence** of ramp lesions in a large  
97 series of patients undergoing posteromedial compartment evaluation at the time of ACL  
98 reconstruction, and also to determine the risk factors associated with ramp lesions. The  
99 secondary objectives of this study were to determine the re-operation rate for failure of ramp  
100 repair, defined by subsequent re-operations for partial medial meniscectomy of the repaired  
101 posterior horn, at a minimum follow-up of 2 years.

102

103 **Methods**

104 Institutional review board approval (IRB COS-RGDS-2018-03-003) was granted for this  
105 study and all patients gave valid consent to participate. A retrospective analysis of  
106 prospectively collected data from the XXXX (anonymized for review) study group database  
107 was conducted. All patients who underwent arthroscopic primary or revision anterior cruciate  
108 ligament reconstruction (ACLR) between September 2012 and March 2018 were considered  
109 for study eligibility. Patients were only excluded if they underwent major concomitant  
110 surgery (for example multiligament reconstruction and/or high tibial osteotomy) or had other  
111 types of medial meniscal lesions (including root tears, horizontal tears, radial tears or vertical  
112 tears more centrally located than the red-white zone).

113

114 Preoperatively, all patients had sustained an ACL tear, diagnosed on the basis of clinical  
115 examination and magnetic resonance imaging (MRI). The patients had been unable to resume  
116 their previous levels of activity because of instability symptoms and therefore underwent  
117 ACL reconstruction. The decision to use a particular graft type for ACLR was based on  
118 patient factors/choice and the evolving indications for performing a concomitant anterolateral  
119 ligament reconstruction (ALLR) during the study period. This decision was taken  
120 preoperatively and was independent of the status of the MM. ~~Indications for ALLR included~~  
121 ~~one or more of the following criteria: grade 3 pivot shift, high level of sporting activity,~~  
122 ~~participation in pivoting sports, deep lateral femoral notch sign on radiographs, associated~~  
123 ~~Second fracture, chronic ACL rupture (>3months after injury), and patients < 25 years old.~~

124

125 Surgical Technique


126 All surgical procedures were performed by a single surgeon (Y) with the patient positioned in  
127 the standard arthroscopy position, a lateral support at the level of a padded tourniquet, and a

128 foot post to allow the knee to be maintained at 90 degrees of flexion when required. Meniscal  
129 and chondral lesions were addressed prior to ACLR.

130

131 *Posteromedial compartment evaluation*

132 All patients underwent a systematic arthroscopic exploration of the knee as previously  
133 described.<sup>40</sup> In order to assess the posteromedial compartment, trans-notch visualization was  
134 performed with the arthroscope placed in the anterolateral portal. Visualization of the  
135 posterior horn medial meniscocapsular attachment was optimized by the application of tibial  
136 internal rotation (Fig 1).<sup>47</sup>



137

138 *Figure 1. Intra-operative images from a Right knee. All images taken with 30 degree*  
139 *arthroscope placed through the anterolateral portal: A) Standard view of the medial*  
140 *compartment, the ramp lesion is not visualised B) The probe is placed in order to*  
141 *demonstrate the location in the notch between the medial femoral condyle (MFC) and the*  
142 *posterior cruciate ligament (PCL) through which the arthroscope will subsequently be*  
143 *advanced into the posteromedial compartment, C) Placing the knee in approximately 30*  
144 *degrees flexion and valgus allows opening of this space and facilitates passage of the*  
145 *arthroscope into the posteromedial compartment, D) View of posteromedial compartment*



146 *shows the ramp lesion; Visualization was optimized by the application of tibial internal*  
147 *rotation*

148

149


150 Using the same methodology as Liu et al, the menisco-capsular attachments and meniscus  
151 were evaluated by probing using either a needle or an arthroscopy hook inserted through a  
152 posteromedial portal.<sup>47</sup> For the purposes of differentiating from other types of meniscal  
153 lesion, a ramp lesion was defined as a medial meniscocapsular tear of the posterior horn of  
154 the medial meniscus. The rationale for including only ramp repairs performed through a  
155 posteromedial portal was based on reports from several authors that different tear types are  
156 associated with different failure rates.<sup>18,25,32,34</sup>

157

### 158 *Ramp Repair*

159 If a ramp lesion was identified, a shaver was inserted through the posteromedial portal and  
160 both surfaces of the tear were prepared (Fig 2).

161



162


163 *Figure 2. Posteromedial compartment evaluation in a Right knee. Trans-notch view obtained*  
164 *with arthroscope placed through anterolateral portal: A) Needle localisation of portal is*  
165 *performed, B) 11-blade scalpel is used to create the portal under direct vision, C) A shaver is*  
166 *inserted and both surfaces of the tear are debrided to encourage healing, D) Appearance of*  
167 *the tear after preparation is completed*

168

169

170 A 25° suture hook (SutureLasso; Arthrex) loaded with a No. 0 absorbable monofilament  
171 suture (PDS; Ethicon) was then inserted, and between one and three separate sutures were  
172 used to perform a repair. After passage, the sutures were tied using a sliding knot and half  
173 hitches. A satisfactory repair was confirmed by evaluation with an arthroscopic probe placed  
174 through the anteromedial portal (Fig 3).

175



176

177 *Figure 3. Ramp repair performed in a Right knee. Trans-notch view of posteromedial*  
178 *compartment obtained with arthroscope placed through anterolateral portal: A) 20 degree*  
179 *left suture hook (Arthrex, Naples, USA) is inserted via the posteromedial portal, B) Suture*

180 *hook passed through meniscocapsular junction into the tear. This allows the hook to be*  
181 *repositioned and then passed into the meniscus body, C) The suture hook is passed into the*  
182 *meniscus body. The 0-PDS suture is then advanced and retrieved through the posteromedial*  
183 *portal after which it is tied, D) The ramp lesion has been repaired, two 0-PDS sutures have*  
184 *been placed using the steps demonstrated. They have been tied with a sliding knot and half*  
185 *hitches via the posteromedial portal under direct vision*

186

187

188 *ACLR with or without concomitant ALLR.*

189 ACLR was performed either as an isolated procedure or in conjunction with ALLR. The ACL  
190 grafts used included quadrupled semitendinosus tendons;<sup>41</sup> bone-patellar tendon-bone<sup>13</sup>  
191 quadrupled hamstring tendons (4HT) or in the case of combined ACL+ALL grafts (HT) a  
192 tripled semitendinosus with a single strand of gracilis.<sup>22</sup> In those cases where an ALLR was  
193 performed independently of the ACL graft, the ALL reconstruction was performed with  
194 gracilis autograft. Our current indications for ALLR include a grade III pivot shift, associated  
195 Second fracture, chronic ACL rupture, high levels of sporting activity, participation in  
196 pivoting sports (eg, soccer, rugby, handball, basketball), patients  $\leq 25$  years old, preoperative  
197 side-to-side laxity  $>6$  mm, lateral femoral notch sign on plain radiographs, and patients  
198 undergoing revision ACL reconstruction.

199

200

201 *Rehabilitation*


202 All patients underwent the same post-operative rehabilitation. This comprised immediate  
203 brace-free mobilization, weight bearing as tolerated, and a restricted range of motion from 0-  
204 90° for the first 4 weeks postoperatively.<sup>30</sup> Full extension and quadriceps activation were key

205 elements of the early physiotherapy. Return to sports was allowed gradually with non-  
206 pivoting sports at 4 months, pivoting non-contact sports at 6 months and pivoting contact  
207 sports at 8-9 months.

208

209 *Follow-up*

210 Postoperative evaluation was conducted by a sports physician, independent of the primary  
211 surgeons at 3 and 6 weeks, and 3, 6 and 12 and 24 months. Only those patients who had a  
212 minimum follow up of two-years and underwent ramp repair were included in the analyses of  
213 secondary meniscectomy rates. In this subgroup, all patients were contacted at final follow-  
214 up by an investigator (Z), independent of the primary surgeon to determine if they had  
215 undergone ipsilateral re-operation for secondary meniscectomy. If further surgery had been  
216 undertaken, the operative records were obtained and reviewed. For the purposes of this study  
217 the term “secondary meniscectomy” was used to describe failure of ramp repair defined by a  
218 re-operation for partial medial meniscectomy involving the previously repaired posterior  
219 horn. A flowchart of included patient is presented in Fig 4.



220

221 *Figure 4. Flowchart of included patients*

222

223

224

225

226 *Epidemiological and Risk Factor Analysis of Ramp Lesions*

227 The epidemiology of ramp lesions was characterized by their **incidence** stratified by key  
228 demographic parameters. Potentially important risk factors were evaluated for association  
229 with ramp lesions and this included gender, body mass index, primary or revision ACLR,  
230 age, time between injury and surgery, type of sport (contact vs non-contact); associated  
231 lateral meniscus tears and; pre-operative side-to-side laxity difference (<6 mm vs >6mm).

232

233 *Statistical Analysis*

234 All calculations were made with SAS for Windows (v 9.4; SAS Institute Inc), with the level  
235 of statistical significance set at  $p < 0.05$ . Descriptive data analysis was conducted depending  
236 on the nature of the considered criteria. **For quantitative data this included number of**  
237 **observed (and missing, if any) values, mean, standard-deviation, median, first and third**  
238 **quartiles, and minimum and maximum. For qualitative data this included the number of**  
239 **observed (and missing, if any) values, and the number and percentage of patients per class. A**  
240 **multivariate logistic regression was performed in order to identify predictive factors of ramp**  
241 **lesions. The factors considered in the multivariate analysis were selected by the way of an**  
242 **univariate approach, including statistically significant effects at the 20% threshold. Moreover,**  
243 **the incidence of such lesions, stratified by time interval from injury to surgery, was described**  
244 **and graphically displayed. The characteristics of patients with ramp lesions were compared**  
245 **between the two groups, defined according to the type of surgery (isolated ACL or ACL +**  
246 **extra articular reconstruction), using the Chi-Square or Fisher exact tests and the Student's t-**

247 test for the qualitative and quantitative data, respectively. The time to secondary  
248 meniscectomy was analyzed considering Kaplan-Meier approach and adjusted Cox model.  
249

250 RESULTS

251 3214 patients undergoing ACL reconstruction were included in the study. A ramp lesion was  
 252 identified and repaired in 769 patients (23.9%). Patient characteristics associated with both  
 253 the presence and absence of associated ramp lesions are presented in Table 1.

254

255 *Table 1 Individual characteristics of patients with or without an associated ramp lesion*

|                           |                 | RAMP lesion        | No RAMP lesion     |
|---------------------------|-----------------|--------------------|--------------------|
| Gender                    | n               | 769                | 2445               |
|                           | Male            | 610 (26.2%)        | 1721 (73.8%)       |
|                           | Female          | 159 (18%)          | 724 (82%)          |
| Age at injury (years)     | n (d.m.)        | 758 (11)           | 2412 (33)          |
|                           | <= 20           | 255 (27.2%)        | 683 (72.8%)        |
|                           | 20 - 30         | 321 (26.2%)        | 900 (73.8%)        |
|                           | 30 - 40         | 128 (21.3%)        | 472 (78.7%)        |
|                           | > 40            | 54 (13.1%)         | 357 (86.9%)        |
| BMI (kg/m <sup>2</sup> )  | n               | 769                | 2445               |
|                           | Mean (SD)       | 23.96 (3.00)       | 23.89 (3.34)       |
|                           | Median (Q1; Q3) | 23.6 (21.8 ; 25.7) | 23.5 (21.6 ; 25.6) |
|                           | Min ; Max       | 17.3 ; 38.6        | 14.6 ; 41.3        |
| Time from injury (months) | n (d.m.)        | 758 (11)           | 2412 (33)          |
|                           | <= 3            | 326 (21.6%)        | 1183 (78.4%)       |
|                           | 3 - 6           | 175 (24.6%)        | 535 (75.4%)        |
|                           | 6 - 12          | 100 (24.6%)        | 306 (75.4%)        |
|                           | 12 - 24         | 49 (25.1%)         | 146 (74.9%)        |
|                           | > 24            | 108 (30.8%)        | 242 (69.2%)        |
| ACLR revision             | n               | 769                | 2445               |

|                         |                   | RAMP lesion | No RAMP lesion |
|-------------------------|-------------------|-------------|----------------|
| Cause of rupture        | Yes               | 120 (37.4%) | 201 (62.6%)    |
|                         | No                | 649 (22.4%) | 2244 (77.6%)   |
|                         | n                 | 769         | 2445           |
|                         | Contact sport     | 528 (25.7%) | 1526 (74.3%)   |
|                         | Non-contact sport | 241 (20.8%) | 919 (79.2%)    |
| Laxity (mm)             | n                 | 769         | 2445           |
|                         | <= 6              | 346 (21%)   | 1300 (79%)     |
|                         | > 6               | 423 (26.9%) | 1145 (73.1%)   |
| Lateral meniscus lesion | n                 | 769         | 2445           |
|                         | Yes               | 297 (33.8%) | 582 (66.2%)    |
|                         | No                | 472 (20.2%) | 1863 (79.8%)   |

256

257

258


### 259 *Risk Factors for Ramp Lesions*

260 Multivariate analyses were performed in order to investigate the association of potential risk  
 261 factors with the occurrence of ramp lesions (Table 2). These analyses demonstrate that male  
 262 gender, age < 30 years, revision ACLR, side-to-side laxity difference greater than 6mm, and  
 263 the presence of a lateral meniscal tear are all significant risk factors for ramp lesions.  
 264 Although the **incidence** of ramp lesions in contact sports (**25.7%**) was higher than non-  
 265 contact sports (**20.8%**) this was not significant in a multivariate analysis ( $P = .247$ ).

266



267 A significantly higher **incidence** of ramp lesions was observed in patients with chronic ACL  
268 ruptures compared to acute ACL ruptures (26% vs 21.6%;  $P = .0037$ ). Specifically, there was  
269 a significant increase in the **incidence** of ramp lesions in the groups with greater chronicity  
270 for all time intervals studied, up to 60 months (Table 3). Regression analysis demonstrates  
271 the correlation between time since injury and the increasing **incidence** of ramp lesions (Fig 5)



272

273 *Figure 5.* Scatter plot of the incidence of ramp lesions identified in patients undergoing  
274 surgery at the following time intervals since injury:  $\leq 3$  months,  $\leq 6$  months,  $\leq 12$  months,  $\leq$   
275 24 months,  $\leq 36$  months,  $\leq 48$  months and  $\leq 60$  months. The linear regression line and  
276 corresponding 95% confidence limits are shown.

277

278 Table 2 Multivariate *logistic regression* analysis of the association of potentially important  
 279 risk factors with ramp lesions<sup>a</sup>

| Risk factor               | Comparison                               | OR<br>(N= 3170) | OR IC95%       | P value         |
|---------------------------|--|-----------------|----------------|-----------------|
| Gender                    | Male vs Female                           | 1.498           | [1.228; 1.836] | <b>&lt;.001</b> |
| Age at injury (years)     | <= 30 years vs > 30 years                | 1.609           | [1.33; 1.952]  | <b>&lt;.001</b> |
| Time from injury (months) |  |                 |                | <b>0.002</b>    |
|                           | <b>]12, 24] months vs ]6, 12] months</b> | 0.965           | [0.64; 1.442]  |                 |
|                           | <b>]3, 6] months vs ]6, 12] months</b>   | 0.979           | [0.733; 1.312] |                 |
|                           | <b>]12, 24] months vs ]3, 6] months</b>  | 0.985           | [0.671; 1.43]  |                 |
|                           | <b>]12, 24] months vs &lt;= 3 months</b> | 1.248           | [0.865; 1.774] |                 |
|                           | <b>]3, 6] months vs &lt;= 3 months</b>   | 1.266           | [1.019; 1.569] |                 |
|                           | <b>]6, 12] months vs &lt;= 3 months</b>  | 1.293           | [0.99; 1.681]  |                 |
|                           | <b>&gt; 24 months vs ]6, 12] months</b>  | 1.313           | [0.944; 1.829] |                 |
|                           | <b>&gt; 24 months vs ]3, 6] months</b>   | 1.342           | [0.998; 1.799] |                 |
|                           | <b>&gt; 24 months vs ]12, 24] months</b> | 1.361           | [0.909; 2.058] |                 |
|                           | <b>&gt; 24 months vs &lt;= 3 months</b>  | 1.698           | [1.296; 2.218] |                 |
| ACLR revision?            | Yes vs No                                | 1.821           | [1.41; 2.344]  | <b>&lt;.001</b> |
| Laxity (mm)               | > 6 mm vs <= 6 mm                        | 1.190           | [1.002; 1.413] | <b>0.047</b>    |
| Lateral meniscus lesion?  | Yes vs No                                | 1.905           | [1.594; 2.276] | <b>&lt;.001</b> |
| Cause of rupture          | Contact vs Non-contact sport             |                 |                | 0.257           |

280 <sup>a</sup>*Bolded P values indicate statistical significance; ACLR : Anterior cruciate ligament reconstruction*

281

282

283

284

285

286

287

288

289 *Table 3 The incidence of ramp lesions in the study population, stratified by class of time*  
 290 *interval between injury and ACLR*

| Time From Injury   | Total number of Patients | Patients with ramp lesions<br>n (%) | P Value* |
|--------------------|--------------------------|-------------------------------------|----------|
| ≤3 mo <sup>a</sup> | 1509                     | 326 (21.6%)                         | 0.0037   |
| >3 mo              | 1661                     | 432 (26%)                           |          |
| ≤6 mo              | 2219                     | 501(22.6%)                          | 0.0072   |
| >6 mo              | 951                      | 257 (27%)                           |          |
| ≤12 mo             | 2625                     | 601 (22.9%)                         | 0.0032   |
| >12 mo             | 545                      | 157 (28.8%)                         |          |
| ≤24 mo             | 2820                     | 650 (23%)                           | 0.0012   |
| >24 mo             | 350                      | 108 (30.9%)                         |          |
| ≤36 mo             | 2927                     | 682 (23.3%)                         | 0.0051   |
| >36 mo             | 243                      | 76 (31.3%)                          |          |
| ≤48 mo             | 2970                     | 692 (23.3%)                         | 0.0019   |
| >48 mo             | 200                      | 66 (33%)                            |          |
| ≤60 mo             | 3006                     | 711 (23.7%)                         | 0.1433   |
| >60 mo             | 164                      | 47 (28.7%)                          |          |

291 <sup>a</sup>3 months after injury was defined as a time between acute anterior cruciate ligament rupture and chronic  
 292 injury; \* *Chi-square test*

293

294

295

296

297 *Secondary meniscectomy rate with a minimum of 2-years of follow-up*

298 Of those patients who underwent ramp repair, 465 had a minimum post-operative  
 299 follow-up of two years and were considered eligible for the secondary meniscectomy  
 300 analysis. However, 49 (10.5%) were lost to follow-up despite attempts to contact them by  
 301 telephone, mail and via their primary care physician. The final subgroup population therefore  
 302 comprised 416 patients with a mean follow up of 45.6 months (range 24.2-66.2 months). At  
 303 final follow up, 45 patients (10.8%) had undergone reoperation for partial medial  
 304 meniscectomy at a mean delay of 21.5 months (3.9-66.2).

305

306 This subgroup of 416 patients was further divided into 2 groups: isolated ACLR (n=225) and  
 307 ACLR + ALLR (n=191) (Table 4). Figure 6 shows the cumulative survivorship of MM  
 308 repairs derived from Kaplan-Meier analysis, with reoperation for medial meniscectomy as an  
 309 endpoint. At both 24 and 48 months follow-up, rates of failure of ramp repair were  
 310 significantly lower for patients who underwent combined ACLR + ALLR compared to those  
 311 who underwent isolated ACLR ( $P = .0178$ ). Patients who underwent ACLR + ALLR had a  
 312 greater than 2-fold reduction in the risk of reoperation for failure of ramp repair as compared  
 313 with patients who underwent isolated ACLR (hazard ratio, 0.457; 95%CI, 0.226-0.864;  $P =$   
 314  $.021$ ).


315

316 *Table 4 Kaplan-Meier Rates of Medial Meniscus Repair Failure by Follow-up Period<sup>a</sup>*

| Time point | Overall                     |                  | Isolated ACLR               |                   | ACLR + ALLR                 |                  | Log-Rank test:<br><i>P</i> Value |
|------------|-----------------------------|------------------|-----------------------------|-------------------|-----------------------------|------------------|----------------------------------|
|            | Rate of second meniscectomy | 95%CI            | Rate of second meniscectomy | 95%CI             | Rate of second meniscectomy | 95%CI            |                                  |
| 1 years    | 4.08%                       | [2.59% ; 6.39%]  | 5.90%                       | [3.54% ; 9.76%]   | 1.96%                       | [0.74% ; 5.14%]  | <b>0.0178</b>                    |
| 2 years    | 6.67%                       | [4.68% ; 9.46%]  | 9.40%                       | [6.29% ; 13.93%]  | 3.50%                       | [1.68% ; 7.20%]  |                                  |
| 3 years    | 8.97%                       | [6.56% ; 12.20%] | 11.88%                      | [8.29% ; 16.87%]  | 5.57%                       | [3.00% ; 10.21%] |                                  |
| 4 years    | 11.26%                      | [8.33% ; 15.13%] | 14.82%                      | [10.55% ; 20.62%] | 6.66%                       | [3.64% ; 12.01%] |                                  |
| 5 years    | 11.26%                      | [8.33% ; 15.13%] | 14.82%                      | [10.55% ; 20.62%] | 6.66%                       | [3.64% ; 12.01%] |                                  |

317 <sup>a</sup>Values are expressed as mean percentage (95%). Bold indicates statistical significance,  $P < 0.05$ . ACLR,  
 318 anterior cruciate ligament reconstruction; ALLR, anterolateral ligament reconstruction.

319



320

321

322 *Figure 6. Kaplan-Meier Survivorship with reoperation for secondary partial medial*  
 323 *meniscectomy (as previously defined) as an endpoint. Numbers at risk with 95% CI. ACLR,*  
 324 *anterior cruciate ligament reconstruction, ALLR, anterolateral ligament reconstruction*

325

326 DISCUSSION

327 A key finding of this study was that the **incidence** of ramp lesions was 23.9% in ACL  
328 deficient knees. Previous authors have reported rates of diagnosis between 9% to  
329 30%,<sup>6,10,11,15,20,36</sup> but it has been unclear how reliably this data can be used to estimate the true  
330 **incidence** of ramp lesions due to the majority of studies including only a small number of  
331 patients. Bollen et al. reported a rate of 9.3%, following arthroscopic examination, in a  
332 prospective series of 183 ACL reconstructions.<sup>6</sup> Di Vico et al. reported a rate of 9.6% in a  
333 series of 115 patients who underwent ACL reconstruction.<sup>11</sup> Liu et al. reported a **incidence** of  
334 16.6% in a series of 868 patients with ACL injury<sup>20</sup> and more recently, Seil et al. reported a  
335 rate of 24% in 224 patients.<sup>36</sup> These variations in **incidence** may also be related to the  
336 diagnostic techniques used. Specifically, pre-operative examination of knee laxity under  
337 anesthesia has been shown to be ineffective at predicting the presence of ramp lesions.<sup>50</sup>  
338 Imaging is also unreliable and a number of studies have reported difficulty identifying these  
339 lesions with MRI, which has a high specificity, but a moderate sensitivity, leading to an  
340 underestimation the true **incidence**.<sup>3,6,10,15,20,37</sup> For example, Bollen et al reported that pre-  
341 operative MRI failed to detect a single ramp lesion in a group of eleven knees with  
342 arthroscopically confirmed lesions.<sup>6</sup>

343

344 In the current study, a systematic evaluation of the posteromedial compartment was  
345 undertaken in all 3214 knees. This is an important point when considering **incidence** data  
346 because, as reported in a previous series, many (approximately 17%) ramp lesions were only  
347 identified after probing the tear through a posteromedial portal in conjunction with a minimal  
348 debridement of the superficial soft tissue layer.<sup>40</sup> These hidden lesions are highly likely to be  
349 missed if arthroscopic examination is only conducted through standard anterior portals.

350

351 The other major findings of this study relate to the evaluation of risk factors associated with  
352 ramp lesions. This study has confirmed previous findings from other authors that male  
353 gender, younger age (<30 years), a concomitant lateral meniscus lesion and chronicity, are  
354 significantly associated with ramp lesions.<sup>20,36</sup> However, the findings of the current study,  
355 based on multivariate analysis, disputed previous work by Seil et al,<sup>36</sup> which suggested that  
356 contact sports injuries were an important risk factor for ramp lesions. It could be the case that  
357 the discrepancy between studies is a result of the difference in sample sizes.

358

359 In any case, there are a number of risk factors which should be emphasized because they have  
360 now been demonstrated to be of significance by several authors. This increases the  
361 confidence in the strength of evidence and highlights the need for posteromedial  
362 compartment evaluation in patients with these characteristics. Delay between injury and  
363 ACLR is significantly associated with increasing **incidence** of ramp lesions over time. In  
364 1984, Woods and Chapman reported on arthroscopic assessment of a series of 234 knees with  
365 ACL rupture. Although not defined as ramp lesions, they found posterior meniscocapsular  
366 disruptions of the medial meniscus occurred in 20 of 112 (17.8%) acute cases (<3 months)  
367 versus 31 of 122 (25.4%) at an average time of 37 months.<sup>50</sup> Liu et al. demonstrated that with  
368 increased time delay between ACL injury and surgery, the **incidence** of ramp lesions  
369 increased up until 24 months.<sup>20</sup> Church et al. equally found an increased number of all types  
370 of meniscal lesions after 12 months, recommending early ACL reconstruction to avoid these  
371 injuries.<sup>9</sup> Other series have also found an association between medial meniscal tears and  
372 increased time to surgery.<sup>9,17,28,46</sup>

373 Gender and age are also important risk factors identified by numerous authors. In the current  
374 study, the male gender was associated with a significantly higher **incidence** of ramp lesions  
375 (27%) compared to females (19%). Liu et al. similarly observed a significantly increased rate

376 in males (18.56% versus female patients 11.97%).<sup>20</sup> Seil et al. reported an increased rate of  
377 27% for males versus 17% for females, although this difference did not reach significance  
378 due to a small sample size.<sup>36</sup> The current study also demonstrated that there was also a  
379 significantly higher **incidence** of ramp lesions in patients under the age of 30. Similarly,  
380 results are found in previously published data. Malatray et al. found that the prevalence of  
381 ACL-associated ramp lesions in children and adolescents is similar to adult populations.<sup>23</sup> Liu  
382 et al. also found that those younger than 30 years of age had a significantly higher **incidence**  
383 of ramp lesions.<sup>20</sup>

384

385 The current study also identified several new significant risk factors, including revision  
386 ACLR. This finding may be explained by either a failure to repair a ramp lesion at the first  
387 surgery or by chronic residual laxity following ACLR leading to a new lesion. Similarly, a  
388 pre-operative anteroposterior side-to-side laxity difference greater than 6mm, was also found  
389 to be an important newly recognized association. However, it is unclear whether this  
390 excessive laxity may predispose to ramp lesions or whether it is simply a reflection of the  
391 role of the medial meniscus as a secondary restraint to anterior laxity of the knee,<sup>1</sup> with the  
392 abnormality being a consequence of a ramp lesion rather than the cause. Another explanation  
393 may be that a high-energy mechanism or injury is often involved in ramp lesions.<sup>5</sup> Other risks  
394 factors previously reported in the literature, but not evaluated in the present study, were a  
395 complete rupture versus partial<sup>36</sup> and a higher medial tibial slope.<sup>39</sup>

396

397 The importance of clearly defining risk factors is in aiding surgeons to hold an appropriate  
398 index of suspicion for ramp lesions, prompt them to perform a posteromedial compartment  
399 evaluation, and identify and repair injuries in order to restore knee stability. When ramp  
400 lesions are overlooked in an ACL reconstruction, anterior and rotatory instability



401 persists<sup>1,24,43</sup> but meniscocapsular repair has been demonstrated to restore normal knee  
402 biomechanics.<sup>1,43</sup>

403

404 If ramp repair is to be advocated in a large proportion of patients undergoing ACLR it is  
405 important to understand the secondary meniscectomy rate. In this study, it was found to be  
406 10.8% at a mean follow up of 45.6 months. These results are in keeping with previous  
407 reports.<sup>16,48</sup> However, a new finding is that the secondary meniscectomy rate after ramp  
408 repair was significantly lower after combined ACLR + ALLR reconstruction compared to  
409 isolated ACLR ( $P = .0178$ ). The combined procedure was associated with a greater than two-  
410 fold reduction in the failure rate of RR ( $P = .021$ ). This supports the results of a previous  
411 study, which demonstrated the protective effect of ALLR on medial meniscal repairs.<sup>42</sup>

412

413 Meniscal healing after repair remains a topical issue.<sup>27</sup> In 1983, Hamberg et al. reported high  
414 healing rates (84%) with suture repair of a series of 43 peripheral medial meniscal tears using  
415 an open posteromedial approach.<sup>14</sup> More recent studies of arthroscopic repair using all-inside  
416 techniques with suture hook<sup>2</sup> or fast-fix anchors<sup>19</sup> have reported good functional results,  
417 with complete healing of 84.3% of tears. A comparison of all-inside repair with outside-in  
418 repair showed similar meniscal healing rates (71.4% vs.70.6%) at a mean follow-up of 36  
419 months.<sup>8</sup> Some authors have suggested that not all ramp meniscal lesions need repair.<sup>12,38,50</sup>

420 Liu et al. reported that stable ramp lesions can be treated with abrasion and trephination alone  
421 with equivalent results to repair.<sup>21</sup> Unfortunately, these studies are limited by relatively small  
422 samples size, and do not present conclusions about the optimal treatment. Pujol et al, in  
423 systematic review, evaluated ten studies in which meniscal tears were left in-situ during  
424 ACLR.<sup>31</sup> Tears were generally left if they were deemed stable on arthroscopic probing or  
425 were less than 10mm in size. Using the endpoint of significant pain or meniscectomy at

426 follow-up, medial meniscal tears left in-situ failed in 10-66% of the cases (mean 14.8%).  
427 They concluded that repair of stable peripheral tears should always be performed to decrease  
428 the risk of postoperative pain or subsequent meniscectomy. In our practice, we therefore aim  
429 to repair all ramp lesions. If the surgeon is already creating a posteromedial portal to perform  
430 abrasion and trephination, a meniscal repair through the same portal is relatively easily  
431 performed with minimal additional risk.

432

### 433 *Limitations*

434 The limitations of a retrospective study design are well recognized. Despite that, this  
435 methodology has advantages, particularly allowing a large sample size, which has been a  
436 limitation of previous studies. This study did not include an assessment of functional  
437 outcomes or a comparison with a control group, for example patients undergoing non-  
438 operative treatment of ramp lesions, or tear debridement without repair. In addition, the study  
439 methodology did not include routine second-look arthroscopy, MRI or clinical functional  
440 evaluation of all patients at final follow-up. This may have resulted in missed diagnoses of  
441 both ramp lesions and of failed ramp repair. However, routine second look arthroscopy is  
442 now rarely reported in the literature due to the unnecessary risk to the patient and evidence  
443 that arthroscopic findings often do not correlate with patient symptoms.<sup>4,45</sup> Furthermore,  
444 performing routine follow-up MRI for the entire series of patients in order to evaluate the  
445 healing of the meniscus was not economically or technically feasible in such a large  
446 population. However, all patients were contacted by telephone at final follow up and those  
447 who had symptoms were recalled for these investigations and assessment. Failure of a ramp  
448 lesion repair was instead based on the hard end-point of patients who underwent subsequent  
449 re-operation of the posterior horn of the medial meniscus. Previous studies have defined  
450 failure of meniscal repair by the presence of osteoarthritis, abnormal MRI, clinical symptoms

451 or subsequent meniscal surgery.<sup>26,27,49</sup> Another limitation is that we have not reported upon  
452 the possible etiology or size of ramp lesions that underwent repair and then secondary partial  
453 meniscectomy. Although it would have been interesting to study this the relevant data was  
454 not recorded or available due to the retrospective study design. A further study limitation is  
455 that the results of this study cannot be extrapolated to patients with ACL injury who undergo  
456 non-operative treatment as they were not evaluated arthroscopically in this study.

457

#### 458 CONCLUSION

459 The high incidence of ramp lesions identified in this study, along with description of  
460 important risk factors, allows an appropriate index of suspicion to be held for these injuries at  
461 the time of ACLR and prompt posteromedial compartment evaluation in order to reduce the  
462 rate of missed diagnoses.

463 The overall secondary meniscectomy rate after ramp repair was 10.8% in this series but this  
464 was significantly lower in those patients who underwent ACLR and anterolateral ligament  
465 reconstruction, the latter appearing to confer a protective effect.

466

467

468

469

470 References

471

- 472 1. Ahn JH, Bae TS, Kang K-S, Kang SY, Lee SH. Longitudinal tear of the medial  
473 meniscus posterior horn in the anterior cruciate ligament-deficient knee significantly  
474 influences anterior stability. *Am J Sports Med.* 2011;39(10):2187-2193.  
475 doi:10.1177/0363546511416597.
- 476 2. Ahn JH, Lee YS, Yoo JC, Chang MJ, Koh KH, Kim MH. Clinical and second-look  
477 arthroscopic evaluation of repaired medial meniscus in anterior cruciate ligament-  
478 reconstructed knees. *Am J Sports Med.* 2010;38(3):472-477.  
479 doi:10.1177/0363546509348102.
- 480 3. Arner JW, Herbst E, Burnham JM, et al. MRI can accurately detect meniscal ramp  
481 lesions of the knee. *Knee Surg Sports Traumatol Arthrosc.* 2017;25(12):3955-3960.  
482 doi:10.1007/s00167-017-4523-9.
- 483 4. Asahina S, Muneta T, Yamamoto H. Arthroscopic meniscal repair in conjunction with  
484 anterior cruciate ligament reconstruction: factors affecting the healing rate. *Arthroscopy.*  
485 1996;12(5):541-545.
- 486 5. Bisson LJ, Kluczynski MA, Hagstrom LS, Marzo JM. A prospective study of the  
487 association between bone contusion and intra-articular injuries associated with acute  
488 anterior cruciate ligament tear. *Am J Sports Med.* 2013;41(8):1801-1807.  
489 doi:10.1177/0363546513490649.
- 490 6. Bollen SR. Posteromedial meniscocapsular injury associated with rupture of the anterior  
491 cruciate ligament: a previously unrecognised association. *J Bone Joint Surg Br.*  
492 2010;92(2):222-223. doi:10.1302/0301-620X.92B2.22974.

- 493 7. Chahla J, Dean CS, Moatshe G, et al. Meniscal Ramp Lesions: Anatomy, Incidence,  
494 Diagnosis, and Treatment. *Orthop J Sports Med.* 2016;4(7):2325967116657815.  
495 doi:10.1177/2325967116657815.
- 496 8. Choi N-H, Kim T-H, Victoroff BN. Comparison of arthroscopic medial meniscal suture  
497 repair techniques: inside-out versus all-inside repair. *Am J Sports Med.*  
498 2009;37(11):2144-2150. doi:10.1177/0363546509339010.
- 499 9. Church S, Keating JF. Reconstruction of the anterior cruciate ligament: timing of  
500 surgery and the incidence of meniscal tears and degenerative change. *J Bone Joint Surg*  
501 *Br.* 2005;87(12):1639-1642. doi:10.1302/0301-620X.87B12.16916.
- 502 10. DePhillipo NN, Cinque ME, Chahla J, Geeslin AG, Engebretsen L, LaPrade RF.  
503 Incidence and Detection of Meniscal Ramp Lesions on Magnetic Resonance Imaging in  
504 Patients With Anterior Cruciate Ligament Reconstruction. *Am J Sports Med.*  
505 2017;45(10):2233-2237. doi:10.1177/0363546517704426.
- 506 11. Di Vico G, Di Donato SL, Balato G, et al. Correlation between time from injury to  
507 surgery and the prevalence of ramp and hidden lesions during anterior cruciate ligament  
508 reconstruction. A new diagnostic algorithm. *Muscles Ligaments Tendons J.*  
509 2018;7(3):491-497. doi:10.11138/mltj/2017.7.3.491.
- 510 12. Duchman KR, Westermann RW, Spindler KP, et al. The Fate of Meniscus Tears Left In  
511 Situ at the Time of Anterior Cruciate Ligament Reconstruction: A 6-Year Follow-up  
512 Study From the MOON Cohort. *Am J Sports Med.* 2015;43(11):2688-2695.  
513 doi:10.1177/0363546515604622.

- 514 13. Garofalo R, Mouhsine E, Chambat P, Siegrist O. Anatomic anterior cruciate ligament  
515 reconstruction: the two-incision technique. *Knee Surg Sports Traumatol Arthrosc.*  
516 2006;14(6):510-516. doi:10.1007/s00167-005-0029-y.
- 517 14. Hamberg P, Gillquist J, Lysholm J. Suture of new and old peripheral meniscus tears. *J*  
518 *Bone Joint Surg Am.* 1983;65(2):193-197.
- 519 15. Hatayama K, Terauchi M, Saito K, Aoki J, Nonaka S, Higuchi H. Magnetic Resonance  
520 Imaging Diagnosis of Medial Meniscal Ramp Lesions in Patients With Anterior  
521 Cruciate Ligament Injuries. *Arthroscopy.* February 2018.  
522 doi:10.1016/j.arthro.2017.12.022.
- 523 16. Jan N, Sonnery-Cottet B, Fayard J-M, Kajetanek C, Thaunat M. Complications in  
524 posteromedial arthroscopic suture of the medial meniscus. *Orthop Traumatol Surg Res.*  
525 2016;102(8, Supplement):S287-S293. doi:10.1016/j.otsr.2016.08.008.
- 526 17. Kennedy J, Jackson MP, O'Kelly P, Moran R. Timing of reconstruction of the anterior  
527 cruciate ligament in athletes and the incidence of secondary pathology within the knee.  
528 *J Bone Joint Surg Br.* 2010;92(3):362-366. doi:10.1302/0301-620X.92B3.22424.
- 529 18. Laurendon L, Neri T, Farizon F, Philippet R. Prognostic factors for all-inside meniscal  
530 repair. A 87-case series. *Orthop Traumatol Surg Res.* 2017;103(7):1017-1020.  
531 doi:10.1016/j.otsr.2017.05.025.
- 532 19. Li W-P, Chen Z, Song B, Yang R, Tan W. The FasT-Fix Repair Technique for Ramp  
533 Lesion of the Medial Meniscus. *Knee Surg Relat Res.* 2015;27(1):56-60.  
534 doi:10.5792/ksrr.2015.27.1.56.

- 535 20. Liu X, Feng H, Zhang H, Hong L, Wang XS, Zhang J. Arthroscopic prevalence of ramp  
536 lesion in 868 patients with anterior cruciate ligament injury. *Am J Sports Med.*  
537 2011;39(4):832-837. doi:10.1177/0363546510388933.
- 538 21. Liu X, Zhang H, Feng H, Hong L, Wang X-S, Song G-Y. Is It Necessary to Repair  
539 Stable Ramp Lesions of the Medial Meniscus During Anterior Cruciate Ligament  
540 Reconstruction? A Prospective Randomized Controlled Trial. *Am J Sports Med.*  
541 2017;45(5):1004-1011. doi:10.1177/0363546516682493.
- 542 22. Löcherbach C, Zayni R, Chambat P, Sonnery-Cottet B. Biologically enhanced ACL  
543 reconstruction. *Orthop Traumatol Surg Res.* 2010;96(7):810-815.  
544 doi:10.1016/j.otsr.2010.06.007.
- 545 23. Malatray M, Raux S, Peltier A, Pfirrmann C, Seil R, Chotel F. Ramp lesions in ACL  
546 deficient knees in children and adolescent population: a high prevalence confirmed in  
547 intercondylar and posteromedial exploration. *Knee Surg Sports Traumatol Arthrosc.*  
548 March 2017. doi:10.1007/s00167-017-4471-4.
- 549 24. Mariani PP. Posterior horn instability of the medial meniscus a sign of posterior  
550 meniscotibial ligament insufficiency. *Knee Surg Sports Traumatol Arthrosc.*  
551 2011;19(7):1148-1153. doi:10.1007/s00167-011-1424-1.
- 552 25. Morgan CD, Wojtys EM, Casscells CD, Casscells SW. Arthroscopic meniscal repair  
553 evaluated by second-look arthroscopy. *Am J Sports Med.* 1991;19(6):632-637;  
554 discussion 637-638. doi:10.1177/036354659101900614.
- 555 26. Nepple JJ, Dunn WR, Wright RW. Meniscal repair outcomes at greater than five years:  
556 a systematic literature review and meta-analysis. *J Bone Joint Surg Am.*  
557 2012;94(24):2222-2227. doi:10.2106/JBJS.K.01584.

- 558 27. Noyes FR, Chen RC, Barber-Westin SD, Potter HG. Greater than 10-year results of red-  
559 white longitudinal meniscal repairs in patients 20 years of age or younger. *Am J Sports*  
560 *Med.* 2011;39(5):1008-1017. doi:10.1177/0363546510392014.
- 561 28. Papastergiou SG, Koukoulis NE, Mikalef P, Ziogas E, Voulgaropoulos H. Meniscal  
562 tears in the ACL-deficient knee: correlation between meniscal tears and the timing of  
563 ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(12):1438-1444.  
564 doi:10.1007/s00167-007-0414-9.
- 565 29. Peltier A, Lording T, Maubisson L, Ballis R, Neyret P, Lustig S. The role of the  
566 meniscotibial ligament in posteromedial rotational knee stability. *Knee Surg Sports*  
567 *Traumatol Arthrosc.* 2015;23(10):2967-2973. doi:10.1007/s00167-015-3751-0.
- 568 30. Perkins B, Gronbeck KR, Yue RA, Tompkins MA. Similar failure rate in immediate  
569 post-operative weight bearing versus protected weight bearing following meniscal repair  
570 on peripheral, vertical meniscal tears. *Knee Surg Sports Traumatol Arthrosc.* August  
571 2017. doi:10.1007/s00167-017-4665-9.
- 572 31. Pujol N, Beaufils P. Healing results of meniscal tears left in situ during anterior cruciate  
573 ligament reconstruction: a review of clinical studies. *Knee Surg Sports Traumatol*  
574 *Arthrosc.* 2009;17(4):396-401. doi:10.1007/s00167-008-0711-y.
- 575 32. Pujol N, Panarella L, Selmi TAS, Neyret P, Fithian D, Beaufils P. Meniscal healing  
576 after meniscal repair: a CT arthrography assessment. *Am J Sports Med.*  
577 2008;36(8):1489-1495. doi:10.1177/0363546508316771.
- 578 33. Reider B. Ramped Up. *Am J Sports Med.* 2017;45(5):1001-1003.  
579 doi:10.1177/0363546517700092.



- 580 34. Rubman MH, Noyes FR, Barber-Westin SD. Arthroscopic repair of meniscal tears that  
581 extend into the avascular zone. A review of 198 single and complex tears. *Am J Sports*  
582 *Med.* 1998;26(1):87-95. doi:10.1177/03635465980260013301.
- 583 35. Seil R. Editorial Commentary: Medial Meniscal Ramp Lesions: Lessons Learned From  
584 the Past in the Pursuit of Evidence. *Arthroscopy.* 2018;34(5):1638-1640.  
585 doi:10.1016/j.arthro.2018.02.032.
- 586 36. Seil R, Mouton C, Coquay J, et al. Ramp lesions associated with ACL injuries are more  
587 likely to be present in contact injuries and complete ACL tears. *Knee Surg Sports*  
588 *Traumatol Arthrosc.* June 2017. doi:10.1007/s00167-017-4598-3.
- 589 37. Seil R, VanGiffen N, Pape D. Thirty years of arthroscopic meniscal suture: What's left  
590 to be done? *Orthop Traumatol Surg Res.* 2009;95(8 Suppl 1):S85-96.  
591 doi:10.1016/j.otsr.2009.09.004.
- 592 38. Shelbourne KD, Rask BP. The sequelae of salvaged nondegenerative peripheral vertical  
593 medial meniscus tears with anterior cruciate ligament reconstruction. *Arthroscopy.*  
594 2001;17(3):270-274. doi:10.1053/jars.2001.19978.
- 595 39. Song G-Y, Liu X, Zhang H, et al. Increased Medial Meniscal Slope Is Associated With  
596 Greater Risk of Ramp Lesion in Noncontact Anterior Cruciate Ligament Injury. *Am J*  
597 *Sports Med.* 2016;44(8):2039-2046. doi:10.1177/0363546516640516.
- 598 40. Sonnery-Cottet B, Conteduca J, Thaunat M, Gunepin FX, Seil R. Hidden lesions of the  
599 posterior horn of the medial meniscus: a systematic arthroscopic exploration of the  
600 concealed portion of the knee. *Am J Sports Med.* 2014;42(4):921-926.  
601 doi:10.1177/0363546514522394.

- 602 41. Sonnery-Cottet B, Freychet B, Murphy CG, Pupim BHB, Thauinat M. Anterior Cruciate  
603 Ligament Reconstruction and Preservation: The Single-Anteromedial Bundle Biological  
604 Augmentation (SAMBBA) Technique. *Arthrosc Tech.* 2014;3(6):e689-693.  
605 doi:10.1016/j.eats.2014.08.007.
- 606 42. Sonnery-Cottet B, Saithna A, Blakeney WG, et al. Anterolateral Ligament  
607 Reconstruction Protects the Repaired Medial Meniscus : A Comparative Study of 383  
608 Anterior Cruciate Ligament Reconstruction From the SANTI Study Group With a  
609 Minimum Follow-up of 2 Years. *Am J Sports Med.* 2018.  
610 doi:10.1177/0363546518767659.
- 611 43. Stephen JM, Halewood C, Kittl C, Bollen SR, Williams A, Amis AA. Posteromedial  
612 Meniscocapsular Lesions Increase Tibiofemoral Joint Laxity With Anterior Cruciate  
613 Ligament Deficiency, and Their Repair Reduces Laxity. *Am J Sports Med.*  
614 2016;44(2):400-408. doi:10.1177/0363546515617454.
- 615 44. Strobel M. Menisci. In: *Fett HM, Flechtner P, eds Manual of Arthroscopic Surgery New*  
616 *York: Springer.* 1988:171-178.
- 617 45. Tachibana Y, Sakaguchi K, Goto T, Oda H, Yamazaki K, Iida S. Repair integrity  
618 evaluated by second-look arthroscopy after arthroscopic meniscal repair with the FasT-  
619 Fix during anterior cruciate ligament reconstruction. *Am J Sports Med.* 2010;38(5):965-  
620 971. doi:10.1177/0363546509356977.
- 621 46. Tandogan RN, Taşer O, Kayaalp A, et al. Analysis of meniscal and chondral lesions  
622 accompanying anterior cruciate ligament tears: relationship with age, time from injury,  
623 and level of sport. *Knee Surg Sports Traumatol Arthrosc.* 2004;12(4):262-270.  
624 doi:10.1007/s00167-003-0398-z.

625 47. Thaunat M, Fayard JM, Guimaraes TM, Jan N, Murphy CG, Sonnery-Cottet B.  
626 Classification and Surgical Repair of Ramp Lesions of the Medial Meniscus. *Arthrosc*  
627 *Tech.* 2016;5(4):e871-e875. doi:10.1016/j.eats.2016.04.009.

628 48. Thaunat M, Jan N, Fayard JM, et al. Repair of Meniscal Ramp Lesions Through  
629 a Posteromedial Portal During Anterior Cruciate Ligament Reconstruction: Outcome  
630 Study With a Minimum 2-Year Follow-up. *Arthroscopy.* 2016;32(11):2269-2277.  
631 doi:10.1016/j.arthro.2016.02.026.

632 49. Westermann RW, Duchman KR, Amendola A, Glass N, Wolf BR. All-Inside Versus  
633 Inside-Out Meniscal Repair With Concurrent Anterior Cruciate Ligament  
634 Reconstruction: A Meta-regression Analysis. *Am J Sports Med.* 2017;45(3):719-724.  
635 doi:10.1177/0363546516642220.

636 50. Woods GW, Chapman DR. Repairable posterior menisco-capsular disruption in anterior  
637 cruciate ligament injuries. *Am J Sports Med.* 1984;12(5):381-385.  
638 doi:10.1177/036354658401200509.

639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655

656 Figure Legends:

657

658 Figure 1. Intra-operative images from a Right knee. All images taken with 30 degree  
659 arthroscope placed through the anterolateral portal: A) Standard view of the medial  
660 compartment, the ramp lesion is not visualised B) The probe is placed in order to demonstrate  
661 the location in the notch between the medial femoral condyle (MFC) and the posterior  
662 cruciate ligament (PCL) through which the arthroscope will subsequently be advanced into  
663 the posteromedial compartment, C) Placing the knee in approximately 30 degrees flexion and  
664 valgus allows opening of this space and facilitates passage of the arthroscope into the  
665 posteromedial compartment, D) View of posteromedial compartment shows the ramp lesion;  
666 Visualization was optimized by the application of tibial internal rotation

667

668 Figure 2. Posteromedial compartment evaluation in a Right knee. Trans-notch view obtained  
669 with arthroscope placed through anterolateral portal: A) Needle localisation of portal is  
670 performed, B) 11-blade scalpel is used to create the portal under direct vision, C) A shaver is  
671 inserted and both surfaces of the tear are debrided to encourage healing, D) Appearance of  
672 the tear after preparation is completed

673

674 Figure 3. Ramp repair performed in a Right knee. Trans-notch view of posteromedial  
675 compartment obtained with arthroscope placed through anterolateral portal: A) 20 degree left  
676 suture hook (Arthrex, Naples, USA) is inserted via the posteromedial portal, B) Suture hook  
677 passed through meniscocapsular junction into the tear. This allows the hook to be  
678 repositioned and then passed into the meniscus body, C) The suture hook is passed into  
679 meniscus body. The 0-PDS suture is then advanced and retrieved through the posteromedial  
680 portal after which it is tied, D) The ramp lesion has been repaired, two 0-PDS sutures have

681 been placed using the steps demonstrated. They have been tied with a sliding knot and half  
682 hitches via the posteromedial portal under direct vision

683

684 Figure 4. Flowchart of included patients

685

686 Figure 5. Increasing incidence of ramp lesions with increasing time from initial ACL injury  
687 to surgery. Two models of curve estimation of the regression analysis between the incidence  
688 of ramp lesion and time interval from anterior cruciate ligament (ACL) injury to surgery.

689

690

691 Figure 6. Kaplan-Meier Survivorship with reoperation for secondary partial medial  
692 meniscectomy (as previously defined) as an endpoint. Numbers at risk with 95% CI. ACLR,  
693 anterior cruciate ligament reconstruction, ALLR, anterolateral ligament reconstruction.

694

695